



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

W. H. Greulich
1922

Lane Medical Library
Stanford University Medical Center

Gift

of

W.H. Greulich, M.D.

LANE MEDICAL LIBRARY
STANFORD UNIV MED CTR

SEP 17 1991

STANFORD, CA 94305

Am. N. Guelish
- 1922 -

The
MEDICAL DEPARTMENT
OF THE UNITED STATES ARMY
IN THE WORLD WAR

VOLUME XV
STATISTICS

PART ONE
ARMY ANTHROPOLOGY

BASED ON OBSERVATIONS MADE ON
DRAFT RECRUITS, 1917-1918, AND ON
VETERANS AT DEMOBILIZATION, 1919

PREPARED UNDER THE DIRECTION OF

M. W. IRELAND
Surgeon General of the Army

By
CHARLES B. DAVENPORT
Carnegie Institution of Washington
(Formerly Major S. C., U. S. A.)

AND
ALBERT G. LOVE
Major, M. C., U. S. A.



WASHINGTON :: GOVERNMENT PRINTING OFFICE :: 1921

LANE MEDICAL DEPARTMENT

LETTER OF TRANSMISSION.

I have the honor to submit herewith a portion of the history of the MEDICAL DEPARTMENT OF THE UNITED STATES ARMY IN THE WORLD WAR. The portion submitted is entitled, "*Army Anthropology*," and is Part One of Volume XV, on the subject of STATISTICS.

The various parts of this history, irrespective of sequence in volume numbers, will be published from time to time in such order as material becomes available.

MERRITTE W. IRELAND,
Surgeon General, United States Army.

The SECRETARY OF WAR.

EDITORIAL BOARD.*

Colonel CHARLES LYNCH, M. C., *Editor-in-Chief.*

Dr. LOY MCAFEE, C. S., *Assistant Editor-in-Chief.*

MEMBERS.

Colonel BAILEY K. ASHFORD, M. C.,
Colonel FRANK BILLINGS, M. C.,
Colonel THOMAS R. BOGGS, M. C.,
Colonel GEORGE E. BREWER, M. C.,
Colonel ROGER BROOKE, M. C.,
Colonel W. P. CHAMBERLAIN, M. C.,
Colonel C. F. CRAIG, M. C.,
Colonel HAVEN EMERSON, M. C.,
Brigadier General JOHN M. T. FINNEY, M. D.,
Colonel J. H. FORD, M. C.,
Lieutenant-Colonel FIELDING H. GARRISON, M. C.,
Colonel H. L. GILCHRIST, M. C.,
Brigadier General JEFFERSON KEAN, M. D.,
Lieutenant-Colonel A. G. LOVE, M. C.,
Colonel JAMES F. MCKERNON, M. C.,
Colonel C. F. MORSE, M. C.,
Colonel R. T. OLIVER, D. C.,
Colonel CHARLES R. REYNOLDS, M. C.,
Colonel THOMAS W. SALMON, M. C.,
Lieutenant-Colonel G. E. DE SCHWEINITZ, M. C.,
Colonel J. F. SILER, M. C.,
Brigadier General W. S. THAYER, M. D.,
Colonel A. D. TUTTLE, M. C.,
Colonel WILLIAM H. WELCH, M. C.,
Colonel E. P. WOLFE, M. C.,
Lieutenant-Colonel CASEY A. WOOD, M. C.,
Colonel HANS ZINSSER, M. C.

* The highest rank held during the World War has been used in the case of each officer.

PREFACE.

The anthropological data contained in this study were collected at the time of the selective service draft of 1917 and 1918 and at the demobilization during the late summer and fall of 1919. The principal data concerning stature, weight, and chest circumference were taken from physical examination schedules (Form 1010 P. M. G. O.)^a for the first million selective service recruits, and, with special reference to men found with selected diseases or defects, also for the second million.

The preliminary study of the result of the physical examinations of approximately the first million drafted men sent to mobilization camps was published in Bulletin No. 11, Surgeon General's Office, March, 1919. The complete study of approximately 2,000,000 drafted men who were sent to the mobilization camps, and of the 549,099 who were rejected by the local boards as totally and permanently unfit, mentally or physically, for the military service, was published in *Defects Found in Drafted Men*, War Department, Surgeon General's Office, 1920.

The anthropological data contained in this work relative to the draft recruits were taken from the same source (Form 1010 P. M. G. O.). A preliminary study of the physical dimensions of the men with the selected diseases for the first million draft recruits was made in 1919. Such diseases and defects were selected as, it was anticipated, might show some deviation from the normal of the physical dimensions. Subsequently, similar data were collected for the second million. Accordingly, the results of the men with the special diseases or defects among the first and second million draft recruits were tabulated and the constants were calculated separately as well as combined. Such a procedure has certain advantages, especially in enabling one to make a comparison between the first and the second million,^b and to secure a criterion as to the constancy and significance of the findings. In the second million recruits there were found more cases of pulmonary tuberculosis, goiter of both types, errors of refraction, tachycardia, varicose veins, hernia, underweight, and congenital defects, and less, or about the same, of various cardiac disorders, varicocele, hemorrhoids, flat-foot, and "defective physical development."

Unfortunately, no provision was made on the physical examination forms for recording color, nativity, age, or occupation.

Acknowledgment is made of the very kind and hearty cooperation of the entire office of the Provost Marshal General, and thanks are especially due to Colonel James Easby-Smith, Colonel Frank H. Wigmore, and Colonel Frank R. Keefer, of that office. An excellent study containing material for the Civil War

^a The earlier form used was Form 14 P. M. G. O.

^b In the tables and illustrations throughout this publication the "first million draft recruits" are designated by the symbol P₁ and the "second million draft recruits" by the symbol P₂.

draft recruits, draft substitutes, and late volunteers, similar to that published in *Defects Found in Drafted Men*, 1920, and in this work, was prepared after the close of the Civil War by Colonel J. H. Baxter, Medical Corps, Chief Medical Officer, Provost Marshal General's Bureau, in the office of the Provost Marshal General, and published in a two-volume work in 1875, under the title of "Statistics, Medical and Anthropological."

The part of this work that is based on the measurements of approximately 100,000 troops at demobilization has also an interesting history. Having in mind the study made by Dr. B. A. Gould, of the United States Sanitary Commission, on the physique of the Civil War volunteer recruits and troops at demobilization in 1865, and recognizing the importance of special anthropometry to the Army, to science, and to the Nation at large, an effort was made by the National Academy of Science from the summer of 1917 to secure authorization for special measurements, but in the stress of the preparation for warfare and during the war itself, authorization was not deemed advisable by the military authorities. However, an order to measure returning soldiers, to secure data for the fashioning of uniforms, was obtained from the Secretary of War during the latter half of 1919.

Thanks are due to Dr. Charles D. Walcott, Secretary of the Smithsonian Institution, to Colonel William H. Welch, M. C., of Johns Hopkins University Medical School, to Brigadier General Edward L. Munson, Morale Branch of the General Staff, for their continued efforts to secure the necessary authorization for the measurements, and to Colonel A. J. Dougherty, of the Equipment Branch of the General Staff, who finally secured the authorization for the work.

Thanks are also due to The Adjutant General for the permission granted to remove records of physical examinations to the Medical Record Section of the Surgeon General's Office for use in collecting statistical data; to the chief clerk of that office, Mr. Thomas A. O'Brien, for his advice and assistance in arranging the details for the use of the records, and to Mr. John N. Manning, principal clerk, Medical Record Section, Adjutant General's Office, for his very kind assistance in expediting the transfer of the records to and from the Surgeon General's Office.

Acknowledgement is made of the services of Mr. Louis R. Sullivan, anthropologist (formerly second lieutenant, Sanitary Corps), for his careful and painstaking work in the preparation of Tables 17, 18, 19, and 20 (sections of the United States, with the "groups" of them).

Especial mention must be made of the services of the civilian anthropologists and anatomists who supervised the work of taking the measurements of soldiers at the camps during the heat of the summer and early autumn of 1919, frequently at considerable self-sacrifice in other ways.^c The good quality of the results are evidence of the effectiveness of the service they rendered.

Acknowledgement is also made of the assistance rendered by the clerical personnel of the Medical Record Section, Surgeon General's Office; to Mr. John W. Beath for his care in the supervision of the preparation of most of the large statistical tables; to Miss Anna T. Buckley and Mrs. Lillian K. Taylor for their

^c See pp. 56 and 57 for the list of the names of the supervising anthropologists and anatomists, and of the camps where the measurements were taken.

exceptionally excellent and accurate work in calculating the constants of the large tables in the text and appendix; to Dr. Thomas J. Griffith, Miss Martha E. Burton and Miss Viola M. Rose for their careful and painstaking work in supervising the coding of the data on the statistical cards; to Second Lieutenant Glendon H. Armstrong, S. C., for his conscientious and painstaking work in supervising the tabulation of the material for the draft recruits; to Miss Helen R. Markley for her equally excellent work in supervising the tabulation of the data for special measurements of the 100,000 demobilized men; to Mrs. Blanche E. Moore for the preparation of the majority of the graphs; and, indeed, to the entire clerical force of the Medical Record Section, Surgeon General's Office, who cooperated efficiently and intelligently, both during the last year of the war and afterward, in making this report as accurate and valuable as possible.

Acknowledgement is also due to Miss Miriam Kortright, of the Carnegie Institution's Station for Experimental Evolution, Cold Spring Harbor, Long Island, N. Y., who assisted in the calculation of many of the smaller text tables.

TABLE OF CONTENTS.

	Page.
Preface	3
Introduction	34
1. Importance of anthropology in the Army	34
2. Stature	34
3. Weight	36
4. Chest circumference	37
5. Build	37
6. Other dimensions	38
7. The general comparative picture of white and Negro troops	39
8. Correlations	42
9. Distribution of eye color	42
10. Distribution of hair color	43
11. Physical dimensions in relation to disease	43
SECTION I.—PHYSICAL MEASUREMENTS.	
A. The importance of anthropometry in the Army	45
B. History of the anthropological work in connection with the Army, 1917-1919	49
I. Anthropometric work in connection with the draft recruits	49
II. Anthropometric work in connection with demobilization	52
1. Orders issued relative to special uniform measurements	53
2. Supervising personnel and camps where measurements were taken	56
3. Apparatus used	57
4. Directions for taking and recording measurements	57
5. Directions for use of record on "descriptive" face of form	59
6. Specifications for arrangements required at camp and for taking measurements there	59
7. Statistical treatment of data	60
C. Results of the standard Army physical measurements	64
I. Age of recruits	64
II. Stature	66
1. General discussion	66
2. Mean stature	67
3. Comparison of mean stature with Civil War records	67
4. Comparison of mean stature in various countries	68
5. Frequency distribution	71
6. Standard deviation	73
7. Mean stature from different States	74
a. Recruits	74
b. Demobilized men	75
8. Comparison of stature of recruits and veterans, by States	77
9. Comparison of stature of recruits from various States, 1863-64 and 1918-19	78
10. Average stature of recruits from different sections	100
11. High and low standard deviations in the different sections	105
12. Mean stature, by groups, of similar sections	105
13. The frequency distribution of statures in the groups of sections	111
14. Comparison of stature in eight European races, of men at demobilization	113
15. Comparison of color races	117

C. Results of the standard Army physical measurements—Continued.	Page.
III. Weight.....	117
1. General discussion.....	117
2. Method.....	118
3. Mean weight.....	119
4. The frequency distribution of weight.....	120
a. Recruits, 1917-18.....	120
b. At demobilization.....	120
5. The standard deviation of the weight series.....	121
6. Mean weight for the different States.....	121
7. Increase in weight at demobilization over mobilization.....	122
8. Mean weight of recruits from the different sections.....	124
9. Mean weight for the different groups.....	127
10. Comparison of weight in eight European races of men at demobilization.....	135
11. Comparison of weight of the color races.....	136
IV. Chest circumference.....	136
1. General discussion.....	136
2. Methods of measurement.....	137
3. Mean chest circumference at expiration.....	138
4. Comparison with Civil War data.....	138
5. Comparison with other countries.....	139
6. Distribution of frequencies of various classes of chest circumference.....	140
7. The frequency distribution of chest circumference by States.....	140
8. Mean chest circumference by sections.....	144
9. Standard deviations of chest circumference by sections.....	147
10. Mean chest circumference by groups of sections.....	149
11. Mean chest circumference of the eight European races of men at demobilization.....	152
12. Chest circumference of men of the color races.....	156
V. Build.....	162
1. Importance of the index of build.....	162
2. Method of determining.....	163
3. Index of build for mean stature and weight.....	164
4. The index of build of Civil and World War veterans for each inch of stature.....	164
5. Distribution of index of build by States.....	165
6. Comparison of index of build in recruits, 1917-1918, and in veterans of 1919 and 1864-1865.....	168
7. Index of build by sections.....	170
8. Index of build by groups of sections.....	173
9. The mean index of build of eight European races of men at demobilization.....	173
10. The mean index of build of color races.....	174
11. Explanation of Plates XIV-XIX.....	174
VI. Pignet's index of robustness.....	186
D. Special anthropological measurements.....	190
1. Sitting height.....	190
(a) General discussion.....	190
(b) Mean sitting height.....	190
(c) Standard deviation.....	191
(d) Comparison of eight European races.....	191
(e) Comparison of color races.....	193
2. Span.....	193
(a) General discussion.....	193
(b) Mean span.....	193
(c) The comparison of eight European races.....	193
(d) Comparison of the color races.....	196

D. Special anthropological measurements—Continued.

	Page.
3. Height of sternal notch.....	196
(a) General discussion.....	196
(b) Mean height of sternal notch.....	197
(c) Comparison of eight European races.....	197
(d) Comparison of color races.....	199
4. Height of pubic arch.....	199
(a) General discussion.....	199
(b) Mean pubic height.....	200
(c) Standard deviation of height of pubic arch.....	200
(d) Comparison of eight European races.....	200
(e) Comparison of color races.....	202
5. Neck circumference.....	202
(a) General discussion.....	202
(b) Mean neck circumference.....	202
(c) Standard deviation of neck circumference.....	203
(d) Comparison of eight European races.....	203
(e) Comparison of color races.....	203
6. Breadth of shoulder.....	203
(a) General discussion.....	203
(b) Mean shoulder breadth.....	204
(c) Standard deviation of shoulder breadth.....	204
(d) Comparison of eight European races.....	205
(e) Comparison of color races.....	207
7. Transverse diameter of the chest.....	207
(a) General discussion.....	207
(b) Mean transverse chest diameter.....	208
(c) Standard deviation of transverse chest diameter.....	208
(d) Comparison of eight European races.....	208
8. Antero-posterior diameter of the chest, and thoracic index.....	208
(a) General discussion.....	208
(b) Mean antero-posterior chest diameter.....	209
(c) Standard deviation of antero-posterior chest diameter.....	209
(d) Comparison of eight European races.....	209
9. Waist circumference.....	213
(a) General discussion.....	213
(b) Mean waist circumference.....	213
(c) Standard deviation of waist circumference.....	213
(d) Comparison of eight European races.....	213
(e) Comparison of white and colored races.....	214
10. Transverse diameter of the pelvis.....	215
(a) General discussion.....	215
(b) Mean transverse diameter of the pelvis.....	216
(c) Standard deviation of transverse diameter of the pelvis.....	216
(d) Comparison of eight European races.....	216
(e) Comparison of color races.....	219
11. Arm length.....	219
(a) General discussion.....	219
(b) Mean arm length.....	219
(c) Standard deviation of arm length.....	220
12. Forearm length.....	220
13. Leg length.....	220
(a) General discussion.....	220
(b) Mean leg length.....	221
(c) Standard deviation of leg length.....	221
(d) Comparison of eight European races.....	221
(e) Comparison of color races.....	224

D. Special anthropological measurements—Continued.

	Page.
14. Knee height.....	224
(a) General discussion.....	224
(b) Mean knee height.....	224
(c) Standard deviation of knee height.....	225
15. Thigh circumference.....	227
(a) General discussion.....	227
(b) Mean thigh circumference.....	227
(c) Standard deviation of thigh circumference.....	227
(d) Comparison of eight European races.....	227
16. Calf circumference.....	230
(a) General discussion.....	230
(b) Mean calf circumference.....	230
(c) Standard deviation of calf circumference.....	230
(d) Comparison of eight European races.....	230
17. Suprapatella circumference.....	232
(a) General discussion.....	232
(b) Mean suprapatella circumference.....	232
(c) Standard deviation of suprapatella circumference.....	232
18. Knee patella circumference.....	232
(a) General discussion.....	232
(b) Mean knee patella circumference.....	232
(c) Standard deviation of knee patella circumference.....	232
19. Comparison of dimensions of white and Negro troops.....	232
(a) Comparison of means of whites and negroes.....	232
(b) Span.....	232
(c) Leg length.....	233
(d) Arm length.....	233
(e) Pubic height.....	233
(f) Knee height.....	233
(g) Forearm.....	233
(h) Sternal notch.....	233
(i) Sitting height.....	233
20. General comparison of other color races.....	236
21. Comparison of the somatic proportions in the eight European races.....	242
(a) General discussion.....	242
(b) Stature.....	243
(c) Weight.....	243
(d) Index of build.....	244
(e) Summary.....	244
E. Correlations between measurements.....	253
1. Correlations between measurements for white troops and Negro troops.....	253
(a) General description of tables.....	253
(b) Correlation between stature and weight.....	254
(c) Correlation between stature and chest circumference.....	256
(d) Correlation between weight and chest circumference.....	258
(e) Correlation between stature and waist circumference.....	258
2. Correlation between measurements for white troops.....	258
(a) Correlation between chest circumference and transverse diameter of pelvis between cristæ.....	258
(b) Correlation between stature and sitting height.....	259
(c) Correlation between stature and height of sternal notch.....	259
(d) Correlation between stature and height of pubic arch.....	259
(e) Correlation between stature and span.....	260
(f) Correlation between chest circumference and weight.....	260
(g) Correlation between chest circumference and neck circumference.....	260

E. Correlations between measurements—Continued.

2. Correlation between measurements for white troops—Continued.	Page.
(h) Correlation between transverse and antero-posterior diameters of the chest.....	261
(i) Correlation between chest circumference and transverse pelvic diameter.....	261
(j) Correlation between waist circumference and transverse pelvic diameter.....	262
(k) Correlation between arm length and forearm.....	262
(l) Correlation between leg length and knee height.....	263
(m) Correlation between leg length and waist circumference.....	264
3. Correlation between measurements for Negro troops.....	264
(a) Correlation between stature and sitting height.....	264
(b) Correlation between stature and height of sternal notch.....	265
(c) Correlation between stature and height of pubic arch.....	265
(d) Correlation between stature and knee height.....	265
(e) Correlation between stature and span.....	265
(f) Correlation between chest circumference and weight.....	265
(g) Correlation between chest circumference and sitting height.....	266
(h) Correlation between chest circumference and neck circumference.....	266
(i) Correlation between transverse and antero-posterior diameters of the chest.....	266
(j) Correlation between chest circumference and transverse diameter of pelvis.....	267
(k) Correlation between waist circumference and transverse diameter of pelvis.....	267
(l) Correlation between arm length and forearm.....	267
(m) Correlation between leg length and knee height.....	268
4. Comparison of correlation between whites and negroes.....	268

F. Patterns for uniforms.....

1. Measurements for blouses.....	271
(a) General discussion.....	271
(b) Chest circumference.....	271
(c) Weight.....	272
2. Measurements for breeches.....	274
3. Dimensions of manikins.....	276
4. Sizes and proportions of men in the distribution zones, Quartermaster Corps.....	276

G. Distribution of eye color.....	280
(a) Clear blue eyes.....	280
(b) Blue eyes with brown spots.....	282
(c) Brown eyes.....	282
(d) Eye color in eight European races.....	284
(e) Comparison with Civil War data.....	285

H. Distribution of hair color.....	288
(a) General discussion.....	288
(b) Flaxen hair.....	288
(c) Dark brown hair.....	289
(d) Red hair.....	289
(e) Comparison with Civil War data.....	291
(f) By Quartermaster Corps distribution zones.....	293
(g) Hair color in eight European races.....	293

SECTION II.

Height, weight, and chest circumference of recruits in relation to various diseases and defects.....	296
I. Introductory.....	296
II. Standard of measurements of drafted men.....	296
(a) Stature.....	296
(b) Weight.....	296

Height, weight, and chest circumference of recruits in relation to various diseases and defects—Continued.

	Page.
III. Physical examination standards.....	297
(a) Stature and weight.....	297
(b) Chest circumference.....	298
IV. The dimensions of men with specific defects and diseases.....	299
1. Pulmonary tuberculosis.....	299
(a) Stature.....	299
(b) Weight.....	300
(c) Chest circumference.....	301
(d) Robustness.....	301
2. Simple goiter.....	304
(a) Stature.....	304
(b) Weight.....	305
(c) Chest circumference.....	305
(d) Robustness.....	305
3. Exophthalmic goiter.....	308
(a) Stature.....	308
(b) Weight.....	308
(c) Chest circumference.....	309
(d) Robustness.....	309
4. Myopia.....	314
(a) Stature.....	314
(b) Weight.....	314
(c) Chest circumference.....	315
(d) Robustness.....	315
5. Hyperopia.....	315
(a) Stature.....	315
(b) Weight.....	318
(c) Chest circumference.....	318
(d) Robustness.....	319
6. Astigmatism.....	319
(a) Stature.....	319
(b) Weight.....	320
(c) Chest circumference.....	322
(d) Robustness.....	322
7. Hypertrophic tonsillitis.....	323
(a) Stature.....	323
(b) Weight.....	323
(c) Chest circumference.....	326
(d) Robustness.....	327
8. Tachycardia, simple.....	327
(a) Stature.....	327
(b) Weight.....	330
(c) Chest circumference.....	330
(d) Robustness.....	331
9. Cardiac hypertrophy.....	331
(a) Stature.....	331
(b) Weight.....	331
(c) Chest circumference.....	334
(d) Robustness.....	334
10. Mitral insufficiency.....	335
(a) Stature.....	335
(b) Weight.....	335
(c) Chest circumference.....	338
(d) Robustness.....	338

I. Height, weight, and chest circumference of recruits in relation to various diseases and defects—Continued.

Page.

IV. The dimensions of men with specific defects and diseases—Continued.

11. Mitral stenosis.....	338
(a) Stature.....	338
(b) Weight.....	339
(c) Chest circumference.....	339
(d) Robustness.....	339
12. Valvular diseases of the heart (unclassified).....	342
(a) Stature.....	342
(b) Weight.....	342
(c) Chest circumference.....	342
(d) Robustness.....	345
13. Varicose veins and varicocele.....	345
(a) Stature.....	345
(b) Weight.....	346
(c) Chest circumference.....	347
(d) Robustness.....	348
14. Hemorrhoids.....	353
(a) Stature.....	353
(b) Weight.....	353
(c) Chest circumference.....	356
(d) Robustness.....	356
15. Asthma.....	356
(a) Stature.....	356
(b) Weight.....	356
(c) Chest circumference.....	357
(d) Robustness.....	360
16. Defective and deficient teeth.....	360
(a) Stature.....	360
(b) Weight.....	360
(c) Chest circumference.....	363
(d) Robustness.....	363
17. Hernia.....	363
(a) Stature.....	363
(b) Weight.....	364
(c) Chest circumference.....	364
(d) Robustness.....	369
18. Enlarged inguinal rings.....	369
(a) Stature.....	369
(b) Weight.....	369
(c) Chest circumference.....	369
(d) Robustness.....	369
19. Flat-foot.....	370
(a) Stature.....	370
(b) Weight.....	370
(c) Chest circumference.....	370
(d) Robustness.....	370
20. Defective physical development.....	372
(a) Stature.....	372
(b) Weight.....	372
(c) Chest circumference.....	375
(d) Robustness.....	375
21. Underweight.....	375
(a) Stature.....	375
(b) Weight.....	376
(c) Chest circumference.....	376
(d) Robustness.....	376

Height, weight, and chest circumference of recruits in relation to various diseases and defects—Continued.

IV. The dimensions of men with specific defects and diseases—Continued.	Page.
22. Overweight and obesity.....	379
(a) Stature.....	379
(b) Weight.....	379
(c) Chest circumference.....	379
(d) Robustness.....	379
23. Cryptorchidism, hypospadias, anorchism, and monorchism.....	379
(a) Stature.....	379
(b) Weight.....	382
(c) Chest circumference.....	382
(d) Robustness.....	382
V. Summary: Bodily dimensions in relation to diseases.....	382
References.....	413

LIST OF TABLES IN TEXT.

Table.

1. Approximate average stature of principal races represented in the United States, arranged in order of size.....	47
2. Ages of soldiers (officers and men) serving in the Civil and World War.....	64
3. Average stature of European males of various countries.....	68
4. Stature, its mean standard deviation and coefficient of variation for men (and in part for women also) for certain especially studied groups.....	69
5. Average stature of adult males of various nationalities in the United States in the Civil War period.....	69
6. Frequency distribution of stature, by classes, at mobilization and demobilization (white and Negro troops), 1917-1919.....	70
7. Stature of Army conscripts and recruits, in inches, as determined by Laplace-Charlier frequency curves.....	70
8. Calculated frequency distribution of statures of men of the United States (Civil War period), France, Belgium, and Italy.....	71
9. Comparison of frequency distribution of statures, United States recruits, Civil and World Wars.....	72
10. Mean stature at each age, 18 to 25 years, United States Army recruits, 1906-1915.....	73
11. Distribution of stature and weight, draft recruits, 1917-1918.....	74
12. Distribution of stature and weight in 6,359 American-born Civil War draft recruits.....	74
13. Mean height by States, first million draft recruits; States arranged in order of standing, with proportional weight and chest circumference (expiration) for each inch of height.....	75
14. Mean stature by States, of soldiers at demobilization, 1919.....	76
15. Increase in stature of soldiers at demobilization, 1919, over stature of recruits, 1917-1918.....	76
16. Comparison of stature, in inches, of native and foreign-born white and colored draft recruits, United States, 1917-1918, and white recruits of the Civil War.....	79
17. Characteristics in composition of the population of the various sections of the United States.....	80
18. List of counties comprised in each "section".....	88
19. Consolidation of similar sections; the series and their constituent groups.....	98
20. Consolidation of similar sections; the groups and their composition out of sections.....	98
21. Mean height by sections; sections arranged in order of standing, with proportional weight and chest circumference (expiration) for each inch of height; also the standard deviation for each height; first million draft recruits.....	101
22. Mean height by groups of sections; groups arranged in order of standing, with proportional weight and chest circumference (expiration) for each inch of height; also the standard deviation for each height; first million draft recruits.....	108

Table.	Page.
23. Height distribution shown by groups of sections, first million draft recruits.....	109
Section A: Absolute numbers.....	109
Section B: Ratios per 1,000.....	109
24. The mean stature and standard deviation in each of eight European races.....	113
25. Comparative frequency distribution of height in each of eight European races, demobilization, 1919.....	116
Section A: Absolute numbers.....	116
Section B: Proportional ratios per 1,000.....	116
26. The mean stature in five color races, demobilization, 1919.....	117
27. Weights associated with statures, with the standard deviations and the coefficient of variation for each, in various classes of American males.....	118
28. Average weight of adult males of various nationalities.....	120
29. Frequency distribution of various classes of weight (per mille) at mobilization, 1917-1918, and demobilization, 1919.....	121
30. Average weight, by States, at mobilization, 1917-1918, and demobilization, 1919 (in pounds); States arranged in order of standing, with proportional weight for each inch of height, and chest circumference (expiration) for each pound of weight for the first million draft recruits.....	122
31. States arranged in order of difference of weight at mobilization, 1917-1918, and demobilization, 1919.....	123
32. Comparative view of mean height and mean weight of men from different States: (a) first million draft recruits (white and colored), 1917-1918; (b) 100,000 demobilized troops (white and colored), 1919; and (c) mean height for Civil War volunteer recruits.....	124
33. Mean weight by section; sections arranged in order of standing, with proportional weight for each inch of height and chest circumference (expiration) for each pound of weight; also the standard deviation for each weight; first million draft recruits...	125
34. Mean weight by groups of sections; groups arranged in order of standing, with proportional weight for each inch of height and chest circumference (expiration) for each pound of weight; also the standard deviation for each weight; first million draft recruits.....	132
35. Weight distribution shown by groups of sections, first million draft recruits.....	133
Section A: Absolute numbers.....	133
Section B: Ratios per 1,000.....	134
36. Mean weight and standard deviation in each of eight European races.....	135
37. Comparative frequency distribution of weight in each of eight European races, demobilization, 1919.....	135
Section A: Absolute numbers.....	135
Section B: Proportional ratios per 1,000.....	135
38. Mean weight in five color races, with the standard deviation for the white and Negro troops, demobilization, 1919.....	136
39. Frequency and proportional distribution of chest circumference (expiration) at mobilization, 1917-1918, and of chest circumference (at rest) at demobilization, 1919.....	139
40. Mean chest circumference (expiration), by States; States arranged in order of standing, with proportional chest circumference (at expiration) in inches for each inch of height and each pound of weight; also the proportional weight in pounds for each inch of chest circumference; first million draft recruits.....	142
41. Chest circumference (expiration) native American white draft recruits of the Civil War.....	142
42. Relative chest circumference (mean chest circumference divided by mean stature), by States, arranged in order of standing; first million draft recruits.....	144
43. Mean chest circumference (expiration) by sections; sections arranged in order of standing, with proportional chest circumference (expiration) in inches for each inch of height and each pound of weight; also the standard deviation for each chest circumference; first million draft recruits.....	145

Table.	Page.
44. The standard deviation of chest circumference (expiration) by sections, arranged in order of standing; first million draft recruits.....	148
45. Mean chest circumference (expiration) by groups of sections; groups arranged in order of standing, with proportional chest circumference (expiration) in inches for each inch of height and each pound of weight; also the standard deviation for each chest circumference; first million draft recruits.....	150
46. Relative chest circumference, by groups of sections (chest circumference divided by stature), first million draft recruits.....	150
47. Distribution of chest circumference (expiration) shown by groups of sections, first million draft recruits.....	151
Section A: Absolute numbers.....	151
Section B: Ratios per 1,000.....	152
48. Absolute and relative chest circumference (at rest) of eight European races, with standard deviation and the coefficient of variation for each, demobilization, 1919, (cms.).....	154
49. Comparative frequency distribution of chest circumference (rest) in each of eight European races, demobilization, 1919.....	155
Section A: Absolute numbers.....	155
Section B: Proportional ratios per 1,000.....	155
50. Mean and relative chest circumference (rest), white and Negro troops, demobilization, 1919 (cms.).....	156
51. Various heights, weights, and chest circumference (expiration), shown for the United States, with rates per 1,000 of each first million draft recruits.....	157
52. Height and weight classes; mean weight and standard deviation for each height; also the mean height and standard deviation for each weight; derived from summation of sections (Table I), first million draft recruits; height in inches and weight in pounds.....	158
53. Height and chest circumference (expiration) classes; mean chest circumference (expiration) and the standard deviation for each height; also the mean height and the standard deviation for each chest circumference; derived from summation of sections (Table II), first million draft recruits; chest circumference and height in inches..	158
54. Weight and chest circumference (expiration) classes; mean chest circumference (expiration) and the standard deviation, for each weight; also the mean weight and the standard deviation for each chest circumference; derived from summation of sections (Table III), first million draft recruits; weight in pounds and chest circumference in inches.....	162
55. Index of build calculated by three methods (based on Table I, first million draft recruits); modal weight used.....	164
56. Index of build calculated by three methods (based on Table I, first million draft recruits); average weight used.....	164
57. Index of build, Civil War veterans (white troops) and World War veterans (white and Negro troops).....	165
58. Index of build at mobilization by States, 1917-1918.....	166
59. Index of build at demobilization, by States, 1919.....	167
60. Increase in index of build at demobilization, 1919, over mobilization, 1917-1918....	168
61. Comparison of index of build of men at demobilization in 1865 and 1919 (weight divided by the first power of height).....	170
62. Comparison of index of build of men at demobilization, 1865 and 1919 (weight multiplied by 1,000 divided by the square of height).....	170
63. Index of build of recruits by sections, 1917-1918.....	172
64. Index of build by groups of sections, 1917-1918.....	173
65. Index of build of eight races (weight multiplied by 1,000 divided by the square of the height).....	173

Table.	Page.
66. Index of build of color races (weight multiplied by 1,000 divided by the square of the height).....	174
67. Comparison of Pignet's index for men of various heights, with average chest and weight for certain percentages of the men of each height.....	188
68. Absolute and relative <i>sitting heights</i> , with standard deviations and the coefficient of variations in eight races, demobilization, 1919.....	191
69. Comparative frequency distribution of <i>sitting height</i> in each of eight European races..	192
Section A: Absolute numbers.....	192
Section B: Ratios per 1,000.....	192
70. Absolute and relative <i>sitting height</i> in five color races, demobilization, 1919.....	193
71. Absolute and relative <i>span</i> , with the standard deviation in eight European races, demobilization, 1919.....	194
72. Comparative frequency distribution of <i>span</i> in each of eight European races, demobilization, 1919.....	195
Section A: Absolute numbers.....	195
Section B: Ratio per 1,000.....	195
73. Absolute and relative <i>span</i> in five color races, demobilization, 1919.....	196
74. Absolute and relative height of <i>sternal notch</i> in eight European races, demobilization, 1919.....	197
75. Comparative frequency distribution of height of <i>sternal notch</i> in each of eight European races, demobilization, 1919.....	198
Section A: Absolute numbers.....	198
Section B: Ratios per 1,000.....	198
76. Absolute and relative height of <i>sternal notch</i> in five color races, demobilization, 1919..	199
77. Absolute and relative height of <i>pubic arch</i> in eight European races, demobilization, 1919.....	200
78. Comparative frequency distribution of height of <i>pubic arch</i> in each of eight races, demobilization, 1919.....	201
Section A: Absolute numbers.....	201
Section B: Proportional ratios per 1,000.....	201
79. Absolute and relative height of <i>pubic arch</i> in five color races, demobilization, 1919...	202
80. Absolute and relative neck circumference of white and Negro troops, demobilization, 1919.....	203
81. Absolute and relative <i>shoulder breadth</i> with the standard deviation and the coefficient of variation in eight European races, demobilization, 1919.....	206
82. Comparative frequency distribution of <i>shoulder breadth</i> in each of eight European races, demobilization, 1919.....	206
Section A: Absolute numbers.....	206
Section B: Proportional ratios per 1,000.....	206
83. Absolute and relative <i>shoulder breadth</i> in five color races, demobilization, 1919.....	207
84. Thoracic index of various races (from Martin, 1914, p. 277).....	209
85. Absolute and relative <i>transverse diameter of the chest</i> with the standard deviation; also <i>antero-posterior diameter of the chest</i> ; with the thoracic index multiplied by 1,000 (transverse diameter divided by the antero-posterior diameter) in eight European races; demobilization, 1919.....	210
86. Comparative frequency distribution of the <i>transverse diameter of the chest</i> in each of eight European races.....	211
Section A: Absolute numbers.....	211
Section B: Proportional ratios per 1,000.....	211
87. Comparative frequency distribution of <i>antero-posterior diameter of the chest</i> in each of eight European races, demobilization, 1919.....	212
Section A. Absolute numbers.....	212
Section B. Proportional ratios per 1,000.....	212

Table.	Page.
88. Absolute and relative <i>waist circumference</i> , with the standard deviation and the coefficient of variation in eight European races, demobilization, 1919.....	214
89. Comparative frequency distribution of <i>waist circumference</i> in each of eight European races, demobilization, 1919.....	214
Section A: Absolute numbers.....	214
Section B: Proportional ratios per 1,000.....	214
90. Absolute <i>transverse diameter of the pelvis</i> by sections, demobilization, 1865.....	216
91. Absolute and relative <i>transverse pelvic diameter</i> , with the standard deviation and the coefficient of variation in eight European races, demobilization, 1919.....	217
92. Comparative frequency distribution of <i>transverse pelvic diameter</i> in each of eight European races, demobilization, 1919.....	218
Section A: Absolute numbers.....	218
Section B: Proportional ratios per 1,000.....	218
93. Absolute and relative <i>transverse diameter of the pelvis</i> in five color races, demobilization, 1919.....	219
94. Absolute and relative <i>leg length</i> with the standard deviation and the coefficient of variation in eight European races, demobilization, 1919.....	222
95. Comparative frequency distribution of <i>leg length</i> in each of eight European races, demobilization, 1919.....	223
Section A: Absolute numbers.....	223
Section B: Proportional ratios per 1,000.....	223
96. Absolute and relative <i>leg length</i> in five color races, demobilization, 1919.....	224
97. Absolute and relative <i>knee height</i> with the standard deviation and the coefficient of variation in eight European races, demobilization, 1919.....	225
98. Comparative frequency distribution of <i>knee height</i> in each of eight European races, demobilization, 1919.....	226
Section A: Absolute numbers.....	226
Section B: Proportional ratios per 1,000.....	226
99. Absolute and relative <i>thigh circumference</i> with the standard deviation and the coefficient of variation in eight European races, demobilization, 1919.....	228
100. Comparative frequency distribution of <i>thigh circumference</i> in each of eight European races, demobilization, 1919.....	229
Section A: Absolute numbers.....	229
Section B: Proportional ratios per 1,000.....	229
101. Absolute and relative <i>calf circumference</i> with the standard deviation, and the coefficient of variation in eight European races, demobilization, 1919.....	231
102. Comparative frequency distribution of <i>calf circumference</i> in each of eight European races, demobilization, 1919.....	231
Section A: Absolute numbers.....	231
Section B: Proportional ratios per 1,000.....	231
103. Summary of <i>dimensions</i> of approximately 100,000 white troops, demobilization, 1919..	234
104. Summary of <i>dimensions</i> of approximately 6,000 colored troops, demobilization, 1919..	234
105. Average <i>dimension</i> in color races, demobilization, 1919.....	235
106. Relative <i>dimensions</i> in color races, demobilization, 1919. Percentage rates.....	235
107. Comparative frequency distribution of measurements in color races at demobilization, 1919.....	236
Section A: Stature.....	236
Section B: Weight.....	236
Section C: Span.....	237
Section D: Height of sternal notch.....	237
Section E: Sitting height.....	238
Section F: Height of pubic arch.....	238
Section G: Leg length.....	239
Section H: Knee height.....	239
Section I: Shoulder breadth.....	240
Section J: Transverse pelvic diameter.....	240

Table.	Page.
108. Approximate number of men measured in eight races, demobilization, 1919.....	242
109. Comparative measurements at demobilization, Civil and World War, white and colored troops.....	243
110. Absolute <i>dimensions</i> in eight races, demobilization, 1919.....	243
111. Index of build in eight European races, obtained by dividing the weight by the stature, and also by the square of the stature, demobilization, 1919.....	244
112. Relative dimensions in eight races, demobilization, 1919.....	245
113. Comparative weight (in pounds) of men of different statures, among white soldiers of 1865, at demobilization, and white and colored soldiers at demobilization, 1919....	254
114. Correlation between <i>stature and chest circumference</i> , Civil War recruits (Baxter II, p. 166).	256
115. Correlation between <i>stature and chest circumference</i> , recruits, 1917-1918.....	256
116. Correlations, summary for white and colored troops, demobilization, 1919.....	269
117. Dimensions associated with the " <i>blouse</i> " groups—white troops, demobilization, 1919..	273
118. Dimensions associated with the " <i>blouse</i> " groups—colored troops, demobilization, 1919.	273
119. Dimensions associated with the " <i>breeches</i> " groups—white troops, demobilization, 1919..	274
120. Dimensions associated with the " <i>breeches</i> " groups—colored troops, demobilization, 1919.	275
121. " <i>Blouse</i> " and " <i>breeches</i> " groups, white and Negro troops; designation of each group; basic measurements adopted; and proportional number of each group of the total number of men measured at demobilization.....	275
122. Dimensions of 21 manikins, in centimeters—white troops.....	276
123. Absolute and relative number of veterans with <i>clear blue eyes</i> , by States of nativity, in order of incidence, demobilization, 1919.....	281
124. Absolute and relative number of veterans with <i>blue eyes with brown spots</i> , by States of nativity, in order of incidence, demobilization, 1919.....	282
125. Absolute and relative number of veterans with <i>light brown eyes</i> , by States of nativity, in order of incidence, demobilization, 1919.....	283
126. Absolute and relative number of veterans with <i>dark brown eyes</i> , by States of nativity, in order of incidence, demobilization, 1919.....	283
127. Comparative frequency distribution of <i>eye color</i> in each of eight European races, demobilization, 1919.....	284
Section A: Absolute numbers.....	284
Section B: Race distribution per 1,000 of each eye color.....	284
Section C: Eye color distribution per 1,000 of each race.....	284
128. Color of eyes—proportional number for different States in 1865.....	285
129. Color of eyes—proportional number for different nativities in the United States, 1865..	285
130. Comparative frequency distribution of <i>eye color</i> by Quartermaster Corps distribution zones, based on nativity of demobilized troops, 1919.....	287
Section A: Absolute numbers.....	287
Section B: Eye color distribution per 1,000 of each zone.....	287
131. Absolute and relative number of veterans with <i>flaxen hair</i> , by States of nativity, in order of incidence, demobilization, 1919.....	288
132. Absolute and relative number of veterans with <i>red hair</i> , by States of nativity, in order of incidence, demobilization, 1919.....	289
133. Absolute and relative number of veterans with <i>light brown hair</i> , by States of nativity, in order of incidence, demobilization, 1919.....	290
134. Absolute and relative number of veterans with <i>medium brown hair</i> , by States of nativity, in order of incidence, demobilization, 1919.....	290
135. Absolute and relative number of veterans with <i>dark brown hair</i> , by States of nativity, in order of incidence, demobilization, 1919.....	291
136. Comparative frequency distribution of <i>hair color</i> in each of eight races, demobilization, 1919.....	292
Section A: Absolute numbers.....	292
Section B: Hair color, distribution per 1,000 of each race.....	292

Table.	Page.
137. Comparative frequency distribution of <i>hair color</i> by Quartermaster Corps distribution zones, based on nativity of demobilized troops.....	294
Section A: Absolute numbers.....	294
Section B: Hair color, distribution per 1,000 of each zone.....	294
138. Standards, height, weight, chest circumference and mobility of chest, adopted for draft recruits, United States Army, 1917.....	297
139. Correlation between height and weight in recruits with <i>tuberculosis, pulmonary</i> , first and second million draft recruits.....	302
140. Correlation between height and chest circumference (expiration) in recruits with <i>tuberculosis, pulmonary</i> , first and second million draft recruits.....	303
141. Correlation between height and weight in recruits with <i>goiter, simple</i> , first and second million draft recruits.....	306
142. Correlation between height and chest circumference (expiration) in recruits with <i>goiter, simple</i> , first and second million draft recruits.....	307
143. Correlation between height and weight in recruits with <i>exophthalmic goiter</i> , first and second million draft recruits.....	310
144. Correlation between height and chest circumference (expiration) in recruits with <i>exophthalmic goiter</i> , first and second million draft recruits.....	311
145. Correlation between height and weight in recruits with <i>myopia</i> , first and second million draft recruits.....	312
146. Correlation between height and chest circumference (expiration) in recruits with <i>myopia</i> , first and second million draft recruits.....	313
147. Correlation between height and weight in recruits with <i>hyperopia</i> , first and second million draft recruits.....	316
148. Correlation between height and chest circumference (expiration) in recruits with <i>hyperopia</i> , first and second million draft recruits.....	317
149. Correlation between height and weight in recruits with <i>astigmatism</i> , first and second million draft recruits.....	321
150. Correlation between height and chest circumference (expiration) in recruits with <i>astigmatism</i> , first and second million draft recruits.....	322
151. Correlation between height and weight in recruits with <i>tonsillitis, hypertrophic</i> , first and second million draft recruits.....	324
152. Correlation between height and chest circumference (expiration) in recruits with <i>tonsillitis, hypertrophic</i> , first and second million draft recruits.....	325
153. Correlation between height and weight in recruits with <i>tachycardia</i> , first and second million draft recruits.....	328
154. Correlation between height and chest circumference (expiration) in recruits with <i>tachycardia</i> , first and second million draft recruits.....	329
155. Correlation between height and weight in recruits with <i>cardiac hypertrophy</i> , first and second million draft recruits.....	332
156. Correlation between height and chest circumference (expiration) in recruits with <i>cardiac hypertrophy</i> , first and second million draft recruits.....	333
157. Correlation between height and weight in recruits with <i>mitral insufficiency</i> , first and second million draft recruits.....	336
158. Correlation between height and chest circumference (expiration) in recruits with <i>mitral insufficiency</i> , first and second million draft recruits.....	337
159. Correlation between height and weight in recruits with <i>mitral stenosis</i> , first and second million draft recruits.....	340
160. Correlation between height and chest circumference (expiration) in recruits with <i>mitral stenosis</i> , first and second million draft recruits.....	341
161. Correlation between height and weight in recruits with <i>valvular diseases of the heart (unclassified)</i> , first and second million draft recruits.....	343
162. Correlation between height and chest circumference (expiration) in recruits with <i>valvular diseases of the heart (unclassified)</i> , first and second million draft recruits.....	344

Table.	Page.
163. Correlation between height and weight in recruits with <i>varicose veins</i> , first and second million draft recruits.....	349
164. Correlation between height and chest circumference (expiration) in recruits with <i>varicose veins</i> , first and second million draft recruits.....	350
165. Correlation between height and weight in recruits with <i>varicocele</i> , first and second million draft recruits.....	351
166. Correlation between height and chest circumference (expiration) in recruits with <i>varicocele</i> , first and second million draft recruits.....	352
167. Correlation between height and weight in recruits with <i>hemorrhoids</i> , first and second million draft recruits.....	354
168. Correlation between height and chest circumference (expiration) in recruits with <i>hemorrhoids</i> , first and second million draft recruits.....	355
169. Correlation between height and weight in recruits with <i>asthma</i> , first and second million draft recruits.....	358
170. Correlation between height and chest circumference (expiration) in recruits with <i>asthma</i> , first and second million draft recruits.....	359
171. Correlation between height and weight in recruits with <i>defective and deficient teeth</i> , first and second million draft recruits.....	361
172. Correlation between height and chest circumference (expiration) in recruits with <i>defective and deficient teeth</i> , first and second million draft recruits.....	362
173. Correlation between height and weight in recruits with <i>hernia</i> , first and second million draft recruits.....	365
174. Correlation between height and chest circumference (expiration) in recruits with <i>hernia</i> , first and second million draft recruits.....	366
175. Correlation between height and weight in recruits with <i>enlarged inguinal rings</i> , first and second million draft recruits.....	367
176. Correlation between height and chest circumference (expiration) in recruits with <i>enlarged inguinal rings</i> , first and second million draft recruits.....	368
177. Correlation between height and weight in recruits with <i>flat-foot</i> , first and second million draft recruits.....	371
178. Correlation between height and weight in recruits with <i>defective physical development</i> , first and second million draft recruits.....	373
179. Correlation between height and chest circumference (expiration) in recruits with <i>defective physical development</i> , first and second million draft recruits.....	374
180. Correlation between height and weight in recruits with <i>underweight</i> , first and second million draft recruits.....	377
181. Correlation between height and chest circumference (expiration) in recruits with <i>underweight</i> , first and second million draft recruits.....	378
182. Correlation between height and weight in recruits with <i>cryptorchidism, hypospadia, anorchism, and monorchism</i> , first and second million draft recruits.....	380
183. Correlation between height and chest circumference (expiration) in recruits with <i>cryptorchidism, hypospadia, anorchism, and monorchism</i> , first and second million draft recruits.....	381
184. Height distribution by <i>special diseases</i> or defects for first and second million draft recruits.....	398
Section A: Absolute numbers.....	398
Section B: Ratios per 1,000.....	399
185. Weight distribution by <i>special diseases</i> or defects for first and second million draft recruits.....	400
Section A: Absolute numbers.....	400
Section B: Ratios per 1,000.....	401
186. Chest circumference (expiration) distribution by <i>special diseases</i> or defects for first and second million draft recruits.....	402
Section A: Absolute numbers.....	402
Section B: Ratios per 1,000.....	403

Table.	Page.
187. Summary of mean dimensions, height, weight, and chest circumference (expiration), for men found with <i>certain diseases or defects</i> shown for the first and second million draft recruits separately, as well as for the combination of the two; standard deviation with the probable error for each; the coefficient of variation and the probable error for each; also the correlation between height and weight, and height and chest with the probable error for each.	404
188. The mean stature and weight of recruits found with the <i>specific diseases</i> and defects among the first and second million draft recruits, arranged in the descending order of the means.	408
189. The index of build (weight multiplied by 1,000, divided by the stature squared) and Pignet's index of recruits found with the <i>specific diseases and defects</i> , arranged in order of standing, first and second million draft recruits.	408
190. Variability (standard deviation and coefficient of variability) of stature associated with various defects and diseases, first and second million draft recruits.	409
191. Variability (standard deviation and coefficient of variability) of weight associated with various diseases and defects among the first and second million draft recruits, arranged in descending order of size.	409
192. Relative weight (weight divided by the height) and relative chest circumference (chest circumference at expiration divided by the height and also by the weight) for men found with <i>special defects and diseases</i> in the first and second million draft recruits, 1917-1918.	410
193. Table for converting centimeters into inches.	410
194. Table for converting inches into centimeters.	411

LIST OF TABLES IN APPENDIX.

Table.	
I. Correlation between <i>height and weight</i> for first million draft recruits.	417
Section A: Absolute numbers derived from summation of sections (weight in pounds and height in inches).	417
Section B: Ratios per 1,000 of the separate weights shown for each height (weight in pounds and height in inches).	418
Section C: Ratios per 1,000 of the separate heights shown for each weight (height in inches and weight in pounds).	419
II. Correlation between <i>height and chest circumference</i> (expiration), first million draft recruits.	420
Section A: Absolute numbers derived from summation of sections (chest in inches and height in inches).	420
Section B: Ratios per 1,000 of the separate chest measurements to each height (chest in inches and height in inches).	421
Section C: Ratios per 1,000 of the separate heights to each chest measurement (height in inches and chest in inches).	421
III. Correlation between <i>weight and chest circumference</i> (expiration), first million draft recruits.	422
Section A: Absolute numbers derived from summation of sections (weight in pounds and chest in inches).	422
Section B: Ratios per 1,000 of the separate weights to each chest measurement (weight in pounds and chest in inches).	422
Section C: Ratios per 1,000 of the separate chest measurements to each weight (chest in inches and weight in pounds).	423
IV. Mean height by groups and component sections, arranged in order of standing, with proportional weight and chest circumference (expiration) for each inch of height; also standard deviation for each height, first million draft recruits (height and chest in inches and weight in pounds).	424

Table.	Page.
V. Mean weight by groups and component sections, arranged in order of standing, with proportional weight for each inch of height, and chest circumference (expiration) for each pound of weight; also standard deviation for each weight, first million draft recruits (height and chest in inches and weight in pounds).....	430
VI. Mean chest by groups and component sections, arranged in order of standing, with proportional chest circumference (expiration) for each inch of height and each pound of weight; also standard deviation for each chest measurement, first million draft recruits (height and chest in inches and weight in pounds).....	436
VII. Correlation between height and weight, <i>group 1, agricultural—North, native white 73% +</i> first million draft recruits (weight in pounds and height in inches).....	442
VIII. Correlation between height and chest circumference (expiration), <i>group 1, agricultural—North, native white 73% +</i> first million draft recruits (chest in inches and height in inches).....	443
IX. Correlation between weight and chest circumference (expiration), <i>group 1, agricultural—North, native white 73% +</i> , first million draft recruits (weight in pounds).....	444
X. Correlation between height and weight, <i>group 2, agricultural—mixed foreign and native white, North</i> , first million draft recruits (weight in pounds and height in inches).....	445
XI. Correlation between height and chest circumference (expiration), <i>group 2, agricultural, mixed foreign and native white, North</i> , first million draft recruits (chest in inches and height in inches).....	446
XII. Correlation between weight and chest circumference (expiration), <i>group 2, agricultural, mixed foreign and native white, North</i> , first million draft recruits (weight in pounds and chest circumference in inches).....	447
XIII. Correlation between height and weight, <i>group 3, agricultural, native white, South</i> , first million draft recruits (weight in pounds and height in inches).....	448
XIV. Correlation between height and chest circumference (expiration), <i>group 3, agricultural, native white, South</i> , first million draft recruits (chest in inches).....	449
XV. Correlation between weight and chest circumference (expiration), <i>group 3, agricultural, native white, South</i> , first million draft recruits (weight in pounds and chest circumference in inches).....	450
XVI. Correlation between height and weight, <i>group 4, agricultural, Negro, 45% +</i> first million draft recruits (weight in pounds and height in inches).....	451
XVII. Correlation between height and chest circumference (expiration), <i>group 4, agricultural, Negro, 45% +</i> , first million draft recruits (chest in inches).....	453
XVIII. Correlation between weight and chest circumference (expiration), <i>group 4, agricultural, Negro, 45% +</i> , first million draft recruits (weight in pounds)....	453
XIX. Correlation between height and weight, <i>group 5, eastern manufacturing</i> , first million draft recruits (weight in pounds and height in inches).....	454
XX. Correlation between height and chest circumference (expiration), <i>group 5, eastern manufacturing</i> , first million draft recruits (chest in inches and height in inches).....	455
XXI. Correlation between weight and chest circumference (expiration), <i>group 5, eastern manufacturing</i> , first million draft recruits (weight in pounds and chest circumference in inches).....	456
XXII. Correlation between height and weight, <i>group 6, commuter</i> , first million draft recruits (weight in pounds and height in inches).....	457
XXIII. Correlation between height and chest circumference (expiration), <i>group 6, commuter</i> , first million draft recruits (chest in inches and height in inches).....	458
XXIV. Correlation between weight and chest circumference (expiration), <i>group 6, commuter</i> , first million draft recruits (weight in pounds and chest circumference in inches).....	459

Table.	Page.
XXV. Correlation between height and weight, <i>group 7, mining</i> , first million draft recruits (weight in pounds and height in inches).....	460
XXVI. Correlation between height and chest circumference (expiration), <i>group 7, mining</i> , first million draft recruits (chest in inches and height in inches)...	461
XXVII. Correlation between weight and chest circumference (expiration), <i>group 7, mining</i> , first million draft recruits (weight in pounds and chest circumference in inches).....	462
XXVIII. Correlation between height and weight, <i>group 8, sparsely settled, not more than 3 per square mile</i> , first million draft recruits (weight in pounds and height in inches)	463
XXIX. Correlation between height and chest circumference (expiration), <i>group 8, sparsely settled, not more than 3 per square mile</i> , first million draft recruits (chest in inches and height in inches).....	464
XXX. Correlation between weight and chest circumference (expiration), <i>group 8, sparsely settled, not more than 3 per square mile</i> , first million draft recruits (weight in pounds and chest circumference in inches).....	465
XXXI. Correlation between height and weight, <i>group 9, desert</i> , first million draft recruits (weight in pounds and height in inches).....	466
XXXII. Correlation between height and chest circumference (expiration), <i>group 9, desert</i> , first million draft recruits (chest in inches and height in inches).....	467
XXXIII. Correlation between weight and chest circumference (expiration), <i>group 9, desert</i> , first million draft recruits (weight in pounds and chest circumference in inches).....	468
XXXIV. Correlation between height and weight, <i>group 10, maritime</i> , first million draft recruits (weight in pounds and height in inches).....	469
XXXV. Correlation between height and chest circumference (expiration), <i>group 10, maritime</i> , first million draft recruits (chest in inches and height in inches)...	470
XXXVI. Correlation between weight and chest circumference (expiration), <i>group 10, maritime</i> , first million draft recruits (weight in pounds and chest circumference in inches).....	471
XXXVII. Correlation between height and weight, <i>group 11, mountain</i> , first million draft recruits (weight in pounds and height in inches).....	472
XXXVIII. Correlation between height and chest circumference (expiration), <i>group 11, mountain</i> , first million draft recruits (chest in inches and height in inches)...	473
XXXIX. Correlation between weight and chest circumference (expiration), <i>group 11, mountain</i> , first million draft recruits (weight in pounds and chest circumference in inches).....	474
XL. Correlation between height and weight, <i>group 12, mountain whites</i> , first million draft recruits (weight in pounds and height in inches).....	475
XLI. Correlation between height and chest circumference (expiration), <i>group 12, mountain whites</i> , first million draft recruits (chest in inches and height in inches).....	476
XLII. Correlation between weight and chest circumference (expiration), <i>group 12, mountain whites</i> , first million draft recruits (weight in pounds and chest in inches).....	477
XLIII. Correlation between height and weight, <i>group 13, Indian, sparsely settled</i> , first million draft recruits (weight in pounds and height in inches).....	478
XLIV. Correlation between height and chest circumference (expiration), <i>group 13, Indian, sparsely settled</i> , first million draft recruits (chest in inches and height in inches).....	479
XLV. Correlation between weight and chest circumference (expiration), <i>group 13, Indian, sparsely settled</i> , first million draft recruits (weight in pounds and chest circumference in inches).....	480
XLVI. Correlation between height and weight, <i>group 14, Mexican, sparsely settled</i> , first million draft recruits (weight in pounds and height in inches).....	481

Table.	Page.
XLVII. Correlation between height and chest circumference (expiration), <i>group 14, Mexican, sparsely settled, first million draft recruits (chest in inches and height in inches)</i>	482
XLVIII. Correlation between weight and chest circumference (expiration) <i>group 14, Mexican, sparsely settled, first million draft recruits (weight in pounds and chest circumference in inches)</i>	483
XLIX. Correlation between height and weight, <i>group 15, native whites of Scotch origin, first million draft recruits (weight in pounds and height in inches)</i>	484
I. Correlation between height and chest circumference (expiration), <i>group 15, native whites of Scotch origin, first million draft recruits (chest in inches and height in inches)</i>	485
II. Correlation between weight and chest circumference (expiration), <i>group 15, native whites of Scotch origin, first million draft recruits (weight in pounds and chest circumference in inches)</i>	486
III. Correlation between height and weight, <i>group 16, Russian 10% +, first million draft recruits (weight in pounds and height in inches)</i>	487
IV. Correlation between height and chest circumference (expiration), <i>group 16, Russian 10% +, first million draft recruits (chest in inches and height in inches)</i>	488
V. Correlation between weight and chest circumference (expiration), <i>group 16, Russian 10% +, first million draft recruits (weight in pounds and chest circumference in inches)</i>	489
VI. Correlation between height and weight, <i>group 17, Scandinavian, 10%, first million draft recruits (weight in pounds and height in inches)</i>	490
VII. Correlation between height and chest circumference (expiration), <i>group 17, Scandinavian, 10%, first million draft recruits (chest in inches and height in inches)</i>	491
VIII. Correlation between weight and chest circumference (expiration), <i>group 17, Scandinavian, 10%, first million draft recruits (weight in pounds and chest circumference in inches)</i>	492
IX. Correlation between height and weight, <i>group 18, Finn, 10%, first million draft recruits (weight in pounds and height in inches)</i>	493
X. Correlation between height and chest circumference (expiration), <i>group 18, Finn, 10%, first million draft recruits (chest in inches and height in inches)</i> ..	494
XI. Correlation between weight and chest circumference (expiration), <i>group 18, Finn, 10%, first million draft recruits (weight in pounds and chest circumference in inches)</i>	495
XII. Correlation between height and weight, <i>group 19, French Canadian, 10%, first million draft recruits (weight in pounds and height in inches)</i>	496
XIII. Correlation between height and chest circumference (expiration), <i>group 19, French Canadian, 10%, first million draft recruits (chest in inches and height in inches)</i>	497
XIV. Correlation between weight and chest circumference (expiration), <i>group 19, French Canadian, 10%, first million draft recruits (weight in pounds and chest circumference in inches)</i>	498
XV. Correlation between height and weight, <i>group 20, German and Scandinavian, 10% +, first million draft recruits (weight in pounds and height in inches)</i> ..	499
XVI. Correlation between height and chest circumference (expiration), <i>group 20, German and Scandinavian, 10% +, first million draft recruits (chest in inches and height in inches)</i>	500
XVII. Correlation between weight and chest circumference (expiration), <i>group 20, German and Scandinavian, 10% +, first million draft recruits (weight in pounds and chest circumference in inches)</i>	501
XVIII. Correlation between height and weight, <i>group 21, German and Austrian, 20% +, first million draft recruits (weight in pounds and height in inches)</i> ..	502

Table.	Page.
LXVIII. Correlation between height and chest circumference (expiration), <i>group 21, German and Austrian, 20%+</i> , first million draft recruits (chest in inches and height in inches).....	503
LXIX. Correlation between weight and chest circumference (expiration), <i>group 21, German and Austrian, 20%+</i> , first million draft recruits (weight in pounds and chest circumference in inches).....	504
LXX. Correlation between height and weight, <i>group 22, German and Austrian, 15%+</i> , first million draft recruits (weight in pounds and height in inches)...	505
LXXI. Correlation between height and chest circumference (expiration), <i>group 22, German and Austrian, 15%+</i> , first million draft recruits (chest in inches and height in inches).....	506
LXXII. Correlation between weight and chest circumference (expiration), <i>group 22, German and Austrian, 15%+</i> , first million draft recruits (weight in pounds and chest circumference in inches).....	507
LXXIII. Comparative frequency distribution of <i>color</i> races by Quartermaster Corps distribution zones, demobilization.....	507
LXXIV. Correlation between <i>height and weight</i> in white and colored troops, demobilization (weight in pounds and height in centimeters).....	508
LXXV. Correlation between <i>height and waist circumference</i> , white and colored troops, demobilization (height in centimeters and waist in centimeters).....	509
LXXVI. Correlation between <i>leg length and knee height</i> , white troops, demobilization (leg length in centimeters and knee height in centimeters).....	510
LXXVII. Correlation between <i>chest circumference (rest) and weight</i> , white troops, demobilization (chest circumference in centimeters and weight in pounds).....	511
LXXVIII. Correlation between <i>chest circumference (rest) and neck circumference</i> , white troops, demobilization (chest circumference (rest) in centimeters and neck circumference in centimeters).....	512
LXXIX. Correlation between <i>chest circumference (rest) and transverse pelvic diameter</i> , white troops, demobilization (chest circumference (rest) in centimeters and transverse pelvic diameter in centimeters).....	513
LXXX. Correlation between <i>chest transverse and chest antero-posterior</i> , white troops, demobilization (chest transverse in centimeters and chest antero-posterior in centimeters).....	514
LXXXI. Correlation between <i>waist circumference and transverse pelvis</i> , white troops, demobilization (waist circumference in centimeters and transverse pelvis in centimeters).....	515
LXXXII. Correlation between <i>length of arm and forearm</i> , white troops, demobilization (arm length in centimeters and forearm in centimeters).....	516
LXXXIII. Correlation between <i>height and sitting height</i> , white troops, demobilization (height in centimeters and sitting height in centimeters).....	517
LXXXIV. Correlation between <i>height and span</i> , white troops, demobilization (height in centimeters and span in centimeters).....	518
LXXXV. Correlation between <i>height and sternal notch</i> , white troops, demobilization (height in centimeters and sternal notch in centimeters).....	518
LXXXVI. Correlation between <i>height and height of pubic arch</i> , white troops, demobilization (height in centimeters and pubic arch in centimeters).....	520
LXXXVII. Correlation between <i>height and sitting height</i> , colored troops, demobilization (height in centimeters and sitting height in centimeters).....	521
LXXXVIII. Correlation between <i>height and span</i> , colored troops, demobilization (height in centimeters and span in centimeters).....	522
LXXXIX. Correlation between <i>height and sternal notch</i> , colored troops, demobilization (height in centimeters and sternal notch in centimeters).....	523
XC. Correlation between <i>height and height of pubic arch</i> , colored troops, demobilization (height in centimeters and pubic height in centimeters).....	524

Table.	Page.
XCI. Correlation between <i>height and knee height</i> , colored troops, demobilization (height in centimeters and knee height in centimeters).....	525
XCII. Correlation between <i>leg length and knee height</i> , colored troops, demobilization (leg length in centimeters and knee height in centimeters).....	526
XCIII. Correlation between <i>chest circumference (rest) and weight</i> , colored troops, demobilization (chest circumference (rest) in centimeters and weight in pounds).....	526
XCIV. Correlation between <i>chest circumference (rest) and neck circumference</i> , colored troops, demobilization (chest circumference (rest) in centimeters and neck circumference in centimeters).....	527
XCV. Correlation between <i>chest circumference (rest) and transverse pelvis</i> , colored troops, demobilization (chest circumference in centimeters and transverse pelvis in centimeters).....	528
XCVI. Correlation between <i>chest transverse and chest antero-posterior</i> , colored troops, demobilization (chest transverse in centimeters and chest antero-posterior in centimeters).....	529
XCVII. Correlation between <i>waist circumference and transverse diameter of pelvis</i> , colored troops, demobilization (waist circumference in centimeters and transverse pelvis in centimeters).....	530
XCVIII. Correlation between <i>arm length and forearm</i> , colored troops, demobilization (arm length in centimeters and forearm in centimeters).....	531
XCIX. Correlation between <i>chest circumference and sitting height</i> , white troops, demobilization (bases of construction of blouse groups shown by double lines) (sitting height in centimeters and chest circumference in centimeters).....	532
C. Association between blouse groups and <i>weight</i> , white troops, demobilization (chest circumference (rest) in centimeters and weight in pounds).....	533
CI. Association between blouse groups and <i>shoulder width</i> , white troops, demobilization (chest circumference (rest) in centimeters and shoulder width in centimeters).....	534
CII. Association between blouse groups and <i>chest transverse diameter</i> , white troops, demobilization (chest circumference (rest) in centimeters and chest transverse in centimeters).....	535
CIII. Association between blouse groups and <i>chest diameter, antero-posterior</i> , white troops, demobilization (chest circumference (rest) in centimeters and chest antero-posterior in centimeters).....	536
CIV. Association between blouse groups and <i>transverse pelvic diameter</i> , white troops, demobilization (chest circumference (rest) in centimeters and transverse pelvic diameter in centimeters).....	537
CV. Association between blouse groups and <i>neck circumference</i> , white troops, demobilization (chest circumference (rest) in centimeters and neck circumference in centimeters).....	538
CVI. Association between blouse groups and <i>total arm length</i> , white troops, demobilization (chest circumference (rest) in centimeters and arm length in centimeters).....	539
CVII. Correlation between <i>chest circumference (rest) and sitting height</i> , colored troops, demobilization (sitting height in centimeters and chest circumference in centimeters).....	540
CVIII. Association between blouse groups and <i>weight</i> , colored troops, demobilization (chest circumference (rest) in centimeters and weight in pounds).....	541
CIX. Association between blouse groups and <i>shoulder width</i> , colored troops, demobilization (chest circumference (rest) in centimeters and shoulder width in centimeters).....	542
CX. Association between blouse groups and <i>transverse diameter of chest</i> , colored troops, demobilization (chest circumference (rest) in centimeters and transverse chest in centimeters).....	543

Table.	Page.
CXI. Association between blouse groups and <i>antero-posterior diameter of chest</i> , colored troops, demobilization (chest circumference (rest) in centimeters and antero-posterior in centimeters).....	544
CXII. Association between blouse groups and <i>transverse pelvic diameter</i> , colored troops, demobilization (chest circumference (rest) in centimeters and transverse pelvis in centimeters).....	545
CXIII. Association between blouse groups and <i>neck circumference</i> , colored troops, demobilization (chest circumference (rest) in centimeters and neck circumference in centimeters).....	546
CXIV. Association between blouse groups and <i>total arm length</i> , colored troops, demobilization (chest circumference (rest) in centimeters and arm length in centimeters).....	547
CXV. Correlation between <i>waist circumference and leg length</i> , white troops, demobilization (waist circumference (rest) in centimeters and leg length in centimeters).....	548
CXVI. Association between breeches groups and <i>transverse pelvic diameter</i> , white troops, demobilization (waist circumference in centimeters and transverse pelvis in centimeters).....	549
CXVII. Association between breeches groups and <i>knee height</i> , white troops, demobilization (waist circumference in centimeters and knee height in centimeters).....	550
CXVIII. Association between breeches groups and <i>thigh circumference</i> , white troops, demobilization (waist circumference in centimeters and thigh circumference in centimeters).....	551
CXIX. Association between breeches groups and <i>suprapatella circumference</i> , white troops, demobilization (waist circumference in centimeters and suprapatella circumference in centimeters).....	552
CXX. Association between breeches groups and <i>patella circumference</i> , white troops, demobilization (waist circumference in centimeters and patella circumference in centimeters).....	553
CXXI. Association between breeches groups and <i>calf circumference</i> , white troops, demobilization (waist circumference in centimeters and calf circumference in centimeters).....	554
CXXII. Correlation between <i>waist circumference and leg length</i> , colored troops, demobilization (waist circumference in centimeters and leg length in centimeters).....	555
CXXIII. Association between breeches groups and <i>transverse diameter of pelvis</i> , colored troops, demobilization (waist circumference in centimeters and transverse pelvis in centimeters).....	556
CXXIV. Association between breeches groups and <i>knee height</i> , colored troops, demobilization (waist circumference in centimeters and knee height in centimeters).....	557
CXXV. Association between breeches groups and <i>circumference of thigh</i> , colored troops, demobilization (waist circumference in centimeters and thigh circumference in centimeters).....	558
CXXVI. Association between blouse groups and <i>suprapatella circumference</i> , colored troops, demobilization (waist circumference in centimeters and suprapatella circumference in centimeters).....	559
CXXVII. Association between breeches groups and <i>circumference at patella</i> , colored troops, demobilization (waist circumference in centimeters and patella circumference in centimeters).....	560
CXXVIII. Association between breeches groups and <i>circumference of calf</i> , colored troops, demobilization, (waist circumference in centimeters and calf circumference in centimeters).....	561
CXXIX. Comparative frequency distribution of <i>blouse groups by States of nativity</i> , white troops, absolute numbers (chest circumference (rest) in centimeters, sitting height in centimeters).....	562

TABLE OF CONTENTS.

29

Table.	Page.
CXXX. Comparative frequency distribution of <i>blouse groups by States of nativity, colored troops, absolute numbers (chest circumference (rest) in centimeters, sitting height in centimeters)</i>	563
CXXXI. Comparative frequency distribution of <i>breeches groups by States of nativity, white troops, absolute numbers (waist circumference in centimeters, leg length in centimeters)</i>	564
CXXXII. Comparative frequency distribution of <i>breeches groups by States of nativity, colored troops, absolute numbers (waist circumference in centimeters, leg length in centimeters)</i>	565
CXXXIII. Comparative frequency distribution of <i>height by States, white and colored troops, at demobilization (height in centimeters)</i>	566
CXXXIV. Comparative frequency distribution of <i>height by Q. M. C. distribution zones, demobilization</i>	568
Section A: Absolute numbers.....	568
Section B: Proportional number of the various heights in each 1,000 for a zone.....	568
Section C: Proportional number of each 1,000 heights in the various distribution zones.....	569
CXXXV. Comparative frequency distribution of <i>weight by States, white and colored troops, at demobilization (weight in pounds)</i>	570
CXXXVI. Comparative frequency distribution of <i>chest circumference (rest) by Q. M. C. distribution zones, white and colored troops, at demobilization</i>	571
Section A: Absolute numbers.....	571
Section B: Proportional number of the various chest circumferences (rest) to each 1,000 for a zone.....	571
Section C: Proportional number of each 1,000 chest circumferences (rest) in the various distribution zones.....	571
CXXXVII. Comparative frequency distribution of <i>waist circumference by Q. M. C. distribution zones, white and colored troops, at demobilization</i>	572
Section A: Absolute numbers.....	572
Section B: Comparative number of the various circumferences to each 1,000 for a zone.....	572
Section C: Proportional number of each 1,000 waist circumferences in the various distribution zones.....	572
CXXXVIII. Comparative frequency of <i>eye color</i> in the various States, nativity of demobilized men.....	573
CXXXIX. Comparative frequency of <i>hair color</i> in various States, nativity of demobilized men.....	574

LIST OF PLATES.

Plate.	Page.
I. Comparative stature and measurements, white (93,185) and colored (6,264) soldiers, demobilization (measurements in inches).....	41
II. Measurement card for clothing patterns, demobilization, 1919.....	61
III. Statistical perforated cards, medical and anthropological data.....	63
1. First million draft recruits, 1917-1918.	
2. Second million draft recruits, 1918.	
3. Demobilized troops, 1919.	
IV. Age distribution, Civil War volunteers, and World War troops, officers and enlisted men, ratios per 1,000; also average age, Civil and World War.....	65
V. Map showing the distribution of the various States into sections.....	97

Plate.		Page.
VI.	Height distribution by groups of sections (P_1), ratios per 1,000.....	114
	1. United States average.	
	2. Agricultural, native white, North, 73% +.	
	3. Agricultural, native white, South.	
	4. Agricultural, Negroes, 45% +.	
	5. Indian.	
	6. Mexican.	
VII.	Height distribution by groups of sections (P_1), ratios per 1,000.....	115
	1. Native white, Scotch origin.	
	2. Russian, 10% +.	
	3. Scandinavian, 10%.	
	4. Finn, 10%.	
	5. French-Canadian, 10%.	
	6. German and Austrian, 15% +.	
VIII.	Weight distribution by groups of sections (P_1), ratios per 1,000.....	130
	1. United States average.	
	2. Agricultural, native white, 73% +.	
	3. Agricultural, native white, South.	
	4. Agricultural, Negroes, 45% +.	
	5. Indian.	
	6. Mexican.	
IX.	Weight distribution by groups of sections (P_1), ratios per 1,000.....	131
	1. Native white, Scotch origin.	
	2. Russian, 10% +.	
	3. Scandinavian, 10%.	
	4. Finn, 10%.	
	5. French-Canadian, 10%.	
	6. German and Austrian, 15% +.	
X.	Chest (expiration) distribution by groups of sections (P_1), ratios per 1,000.....	153
	1. United States average.	
	2. Agricultural, native white, 73% +.	
	3. Agricultural, native white, South.	
	4. Agricultural, Negroes, 45% +.	
	5. Indian.	
	6. Mexican.	
	7. Native, white, Scotch origin.	
	8. Russian, 10% +.	
	9. Scandinavian, 10%.	
	10. Finn, 10%.	
	11. French-Canadian, 10%.	
	12. German and Austrian, 15% +.	
XI.	Weight distribution by height (P_1), ratios per 1,000.....	159
	1. Height, 62 inches.	
	2. Height, 63 inches.	
	3. Height, 64 inches.	
	4. Height, 65 inches.	
	5. Height, 66 inches.	
	6. Height, 67 inches.	
XII.	Weight distribution by height (P_1), ratios per 1,000.....	160
	1. Height, 68 inches.	
	2. Height, 69 inches.	
	3. Height, 70 inches.	
	4. Height, 71 inches.	
	5. Height, 72 inches.	
	6. Height, 73 inches.	

Plate.	Page.
XIII. Chest (expiration) distribution by height (P_1), ratios per 1,000.	161
1. Height, 62 inches.	
2. Height, 63 inches.	
3. Height, 64 inches.	
4. Height, 65 inches.	
5. Height, 66 inches.	
6. Height, 67 inches.	
7. Height, 68 inches.	
8. Height, 69 inches.	
9. Height, 70 inches.	
10. Height, 71 inches.	
11. Height, 72 inches.	
12. Height, 73 inches.	
XIV. Height, weight, and chest (expiration) measurements showing the proportionate measurement of two of them to the total of the third (P_1).....	177
1. Height with relative weight and chest (expiration).	
2. Weight with relative height and chest (expiration).	
3. Chest (expiration) with relative height and weight.	
XV. Total and proportionate measurements by States (P_1); height with relative weight and chest (expiration).....	178
XVI. Total and proportionate measurements by States (P_1); weight with relative height and chest (expiration).....	179
XVII. Total and proportionate measurement by States (P_1); chest (expiration), with relative height and weight.....	180
XVIII. Total and proportionate measurement, groups of sections (P_1).....	181
1. Height with relative weight and chest (expiration).	
2. Weight with relative height and chest (expiration).	
3. Chest (expiration) with relative height and weight.	
XIX. Total and proportionate measurement by each section (P_1); height with relative weight and chest (expiration).....	182
XX. Distribution, height, weight, and chest (expiration) measurement; States of nativity, draft recruits, 1917-1918, and demobilized troops, 1919.....	184
1. Mean height.	
2. Mean weight divided by mean height.	
3. Mean weight.	
4. Mean weight divided by chest (expiration), in pounds.	
5. Mean chest (expiration).	
6. Mean chest (expiration) divided by mean weight (inches).	
7. Lines of equal index of chest circumference.	
8. Lines of equal height (inches).	
XXI. Pignet's index, men of various heights (P_1), percentage distribution of each height.....	189
XXII. Average dimensions, eight races, demobilization, 1919. Measurements in centimeters; weight in pounds.....	246
1. Span.	
2. Stature.	
3. Weight.	
4. Sternal notch.	
XXIII. Average dimensions, eight races, demobilization, 1919 (measurements in centimeters).....	247
1. Sitting height.	
2. Chest circumference.	
3. Pubic arch.	
4. Waist circumference.	

Plate.	Page.
XXIV. Average dimensions, eight races, demobilization, 1919 (measurements in centimeters).....	248
1. Leg length.	
2. Thigh circumference.	
3. Knee height.	
4. Calf circumference.	
XXV. Average dimensions, eight races, demobilization, 1919 (measurements in centimeters).....	249
1. Shoulder width.	
2. Pelvic diameter, transverse.	
3. Chest diameter, transverse.	
4. Chest circumference, antero-posterior.	
XXVI. Relative dimensions, eight races, demobilization, 1919 (percentage rates based on measurements in centimeters).....	250
1. Span.	
2. Sternal notch.	
3. Sitting height.	
4. Pubic height.	
XXVII. Relative dimensions, eight races, demobilization, 1919 (percentage rates based on measurements in centimeters).....	251
1. Waist circumference.	
2. Leg length.	
3. Thigh circumference.	
4. Knee height.	
XXVIII. Relative dimensions, eight races, demobilization, 1919 (percentage rates based on measurements in centimeters).....	252
1. Shoulder width.	
2. Calf circumference.	
3. Pelvic diameter, transverse.	
4. Chest antero-posterior transverse.	
XXIX. Correlations, white and negro troops, demobilization, 1919.....	270
XXX. Distribution, hair and eye color, demobilization, 1919, States of nativity.....	295
1. Flaxen and red hair.	
2. Clear blue eyes and blue eyes with brown spots.	
3. Light and medium brown hair.	
4. Light brown eyes.	
5. Dark brown hair.	
6. Dark brown eyes.	
7. Lines of equal proportion of flaxen hair.	
8. Lines of equal proportion of clear blue eyes.	
XXXI. Height distribution by special diseases or defects (P_1 and P_2), ratios per 1,000....	386
1. Tuberculosis, pulmonary.	
2. Goiter, simple.	
3. Exophthalmic goiter.	
4. Asthma.	
5. Astigmatism.	
6. Hyperopia.	
XXXII. Height distribution by special diseases or defects (P_1 and P_2), ratios per 1,000....	387
1. Valvular diseases of the heart (unclassified).	
2. Hemorrhoids.	
3. Varicose veins.	
4. Varicocele.	
5. Defective and deficient teeth.	
6. Hernia.	

TABLE OF CONTENTS.

33

Plate.	Page.
XXXIII. Height distribution by special diseases or defects (P_1 and P_2), ratios per 1,000...	388
1. Myopia.	
2. Tonsillitis, hypertrophic.	
3. Cardiac hypertrophy.	
4. Mitral insufficiency.	
5. Mitral stenosis.	
6. Tachycardia, simple.	
XXXIV. Weight distribution by special diseases or defects (P_1 and P_2), ratios per 1,000..	389
1. Tuberculosis, pulmonary.	
2. Goiter, simple.	
3. Exophthalmic goiter.	
4. Asthma.	
5. Astigmatism.	
6. Hyperopia.	
XXXV. Weight distribution by special diseases or defects (P_1 and P_2), ratios per 1,000..	390
1. Valvular diseases of the heart (unclassified).	
2. Hemorrhoids.	
3. Varicose veins.	
4. Varicocele.	
5. Defective or deficient teeth.	
6. Hernia.	
XXXVI. Weight distribution by special diseases or defects (P_1 and P_2), ratios per 1,000..	391
1. Myopia.	
2. Tonsillitis, hypertrophic.	
3. Cardiac hypertrophy.	
4. Mitral insufficiency.	
5. Mitral stenosis.	
6. Tachycardia, simple.	
XXXVII. Chest (expiration) distribution by special diseases or defects (P_1 and P_2) ratios per 1,000.....	392
1. Tuberculosis, pulmonary.	
2. Goiter, simple.	
3. Exophthalmic goiter.	
4. Asthma.	
5. Astigmatism.	
6. Hyperopia.	
7. Myopia.	
8. Tonsillitis, hypertrophic.	
9. Cardiac hypertrophy.	
XXXVIII. Chest (expiration) distribution by special diseases or defects (P_1 and P_2) ratios per 1,000.....	393
1. Mitral insufficiency.	
2. Mitral stenosis.	
3. Tachycardia, simple.	
4. Valvular diseases of the heart (unclassified).	
5. Hemorrhoids.	
6. Varicose veins.	
7. Varicocele.	
8. Defective and deficient teeth.	
9. Hernia.	
XXXIX. Height measurement, special diseases (P_1 and P_2) in relation to the average, inducted men (P_1).....	394
XL. Weight measurement, special diseases (P_1 and P_2) in relation to the average, inducted men (P_1).....	395
XLI. Chest measurement (expiration), special diseases (P_1 and P_2), in relation to the average, inducted men (P_1).....	396
XLII. Total and proportionate measurements, special diseases (P_1 and P_2).....	397

INTRODUCTION.

This study involves the analysis of the three standard physical measurements of the Army, taken on 1,000,000 recruits; with special reference to physical defects, taken on 2,000,000 recruits; and of a set of 17 other measurements made of 100,000 troops at demobilization, for the purpose of securing dimensions for uniforms. The whole study gives an insight into the sizes and proportions of the American male population, ages 21 to 30 years, and is a study of dimensions with reference to health and development, to geographical distribution and environment, and to race and color.

The data were gathered partly at local and camp boards on the occasion of the selective draft, and partly on special order from the War Department to secure detailed measurements of 100,000 troops at the time of demobilization. The statistical work was done by the Medical Record Section of the Surgeon General's Office.

1. IMPORTANCE OF ANTHROPOLOGY IN THE ARMY.

For over a century armies have prescribed limits of size for recruits on various grounds. It is urged that small men (under 60 inches) can not carry the prescribed equipment. Men over 78 inches are more apt to suffer from circulatory and other diseases. The size of men has a relation to the standard food ration. This differs in the English and Italian Armies because of the difference in body size of the soldiers. Troops in an Army camp containing a large proportion of South Italians and Polish Jews from New York city should use a different average amount of food per man than those composed mostly of Scandinavians. The length of leg is important for the classification of troops which are required to make long marches. A knowledge of proportions of facial features is essential to gas-mask manufacturers. A knowledge of the size and proportions of the body is essential to the proper cutting of uniforms. It will also aid in detecting pulmonary tuberculosis and cardiac disorders, as well as thyroid and other diseases. A knowledge of racial characteristics is often necessary to decide on classification when military organizations are being formed on racial lines, such as Negro regiments, Slavic legions, etc. And finally, the whole system of identification, whether by finger prints or by Bertillon's proportions, belongs to the field of anthropology.

2. STATURE.

The mean stature of the first million recruits, ages 21 to 30 years, inclusive, and including white and colored, is 67.49 inches (1,714 mm.). The 100,000 troops measured at demobilization measured 67.72 inches tall (1,720 mm.). The gain of 6 millimeters on the average was partly because they were older, partly because they were straighter, partly because some of the shorter divisions were not included in the hundred thousand, and partly because some short men were rejected when examined for mobilization.

Comparing the average stature of recruits with those of the Civil War, after combining the figures of Baxter,¹ 1875, and Gould,² 1869, due allowance being

NOTE.—Superior figures refer to literature and other documents cited, p. 417.

made for the number of men recorded in each case, we find that it is practically the same, being 67.502 in the Civil War, and 67.49 in 1917-1918. We might conclude, then, that the mean stature of men of military age has changed little in the United States in the last 50 years. But this conclusion might be hasty, for the men of 1917-1918 were taken from all parts of the United States, while those of 1864-1865 largely excluded the Southern States, and since the men from these States are exceptionally tall, their inclusion tends to raise the mean stature.

Taking the figures from Gould,² 1869, the ages of volunteers showed a greater proportion of men below the ages of 24—that is, of those who had not obtained their maximum growth—than in 1917-1918. This again tends to raise the average stature of 1917-1918 over that of 1861-1865. Baxter,¹ gives a higher average age for draft recruits, namely, 27.307.

It is reasonable to suppose that since this country has received a very large number of immigrants of prevailingly low stature from the southern part of Europe, during the last 50 years the average stature of the population of the country should show a decrease. Such, however, is very difficult to demonstrate mathematically, since the methods used in the recruiting of the two armies, at the two periods, differed so materially. Indeed, the question whether the physique of our young men has changed in the last 50 years thus unqualified has little meaning. Had the racial constitution of the population remained constant—that is, had there been no heavy immigration—then the question would have more meaning; but in view of the tremendous immigration, amounting in some years to nearly a million persons, the physical changes of the racial constitution of our stock have been so great as to mask entirely any slight alteration that may have occurred in the physique of the stock of 50 years ago, through either improvement or deterioration of environmental or economic conditions.

From the different States men differ much in stature. The Texans are tallest, having an average stature nearly 1 inch above the national average. The mean of the Southern States is taller than the average, while the men of Connecticut, Pennsylvania, New York, Massachusetts, and New Jersey have an average short stature. They are the States with many immigrants from southeastern Europe. Among the North Central States, Kansas, Idaho, Oregon, Nebraska, South Dakota, Iowa, and Minnesota have high average statures. At demobilization the greatest increase in average stature was found in the Southern States, which had apparently greater room for improvement; at least in absolute millimeters. The average stature of veterans from Massachusetts, District of Columbia, and Indiana had not increased.

The average stature of the men from different sections revealed points of even greater interest. At the head of the list stands the mountain section of North Carolina, with a mean stature of 68.67 inches, nearly 1.2 inches above the national average. The inhabitants of this section are largely descendants of the early Scotch settlers (a tall race) in Cape Fear River basin. The next tallest mean man is found in the Ozark mountain region, 68.64 and 68.63

inches, and then come the Texas sections, averaging about 68.47. At the top of the northern sections is northern Minnesota with its "big Swedes." Other sections with tall average stature are Mississippi, the mountain sections of Tennessee and Kentucky, other parts of North Carolina, Western Kansas, Oklahoma, Arkansas, California, and Nebraska. At the other extreme are Rhode Island, New York City, the mining area of eastern Pennsylvania, Philadelphia, the manufacturing towns of northeast Massachusetts, eastern New Jersey, and all parts of Connecticut. The inhabitants of these mining and manufacturing sections are not small because of the injurious somatic effect of the miner's and manufacturer's occupations; for the miners of Idaho average far above the mean of the country, and the inhabitants of the flour mill "twin cities" of Minnesota average three-tenths of an inch above the mean of the country. On the other hand, an agricultural section of eastern Pennsylvania has a population that is seven-tenths of an inch below the mean of the country.

Combining sections, the mountain whites have the greatest mean stature and a low variability; they "run tall" fairly uniformly. Next comes the prevailing white agricultural group of the South, then two groups with a large Mexican and Indian population, then the German-Scandinavian groups and those low-land sections with many native whites of Scotch origin. The shortest group is that containing many French Canadians. The next taller is the eastern manufacturing group with its great numbers of representatives of the short races. Of the eight European races that were most numerous in the examination at demobilization the Scotch were the tallest (67.93 inches), next the English (67.75), then the Germans (67.73), the Irish (67.46), the Polish (66.70), French (66.37), Hebrew (65.71), and Italians (65.03). At demobilization the stature of the whites had increased over mobilization from 67.49 inches to 67.71; the Negroes were 67.70 and the Indians were 67.52 tall at demobilization; the Chinese 67.37; and the Japanese 67.30. At demobilization the Negroes were found to be more variable in stature than the whites as 6.91 is to 6.66 centimeters.

3. WEIGHT.

The mean weight of the first million recruits was 141.54 pounds, which is slightly higher than the mean weight (136.05 pounds) of a few thousand "white American" recruits measured at the time of the Civil War (Baxter, Vol. II, p. 15).^{*} At demobilization troops weighed, on the average, 3 pounds more than did recruits and showed about the same increase that veterans showed over recruits in Civil War times. At demobilization in 1919 there was reduced variability in the weight. The soldiers had increased 2 per cent in weight and diminished 2 per cent in variability; the fine physical conditions of army life tended to raise the weight to a uniform high level.

The greatest weight is found in men from the extreme north. The following States stand at the head of the list: Alaska, South and North Dakota, Minnesota, Oregon, Montana, Washington, Nevada, and Idaho. The men from these States are not the tallest, but as we shall see later they are the stockiest. This stocky condition is not entirely racial; it is probably the reaction of the body

^{*} The figures cited are for draft substitutes and late volunteers, as well as for draft recruits *per se*.

to climatic conditions. Just as the Eskimos are robust, so whites in Alaska and the Dakotas tend to become so. Also, under army conditions men from Alaska gained on the average 11.5 pounds. However, the number of men from Alaska examined was small and the averages probably untrustworthy. The average increase for the whole country was only 3.4 pounds. In general, the men from the Southern States showed an increase of weight above the average of the entire country.

The "French-Canadian" sections comprised recruits of the least weight; the eastern manufacturing groups came next, largely because they contained so many small men. That conditions of life were not the principal cause of the low weight is indicated by the fact that the recruits from commuter (suburban) groups also showed a low average weight. Of the Europeans at demobilization, Germans show the greatest average weight, South Italians and Hebrews the least. The Scotch are the most variable in weight, the Poles the least. The Negro troops are slightly heavier and show a greater variability in weight than the white troops. The Japanese weigh the least of all color races. In the southern sections those containing a large proportion of colored men show relatively less obesity than those containing a small proportion of them.

4. CHEST CIRCUMFERENCE.

The mean circumference of the deflated chest of the first million recruits was 33.22 inches. At demobilization uninflated chests of the veterans measured, on the average, 34.94 inches. These results are not strictly comparable, however, as the chest was not measured in the same phase of expansion in the two sets. Despite this there is evidence that the mean chest girth of the veterans had increased about 1 inch. The same thing happened in the Civil War. The recruits from the Northwestern States showed the greatest chest circumference, those from the Southern States, Rhode Island, and the District of Columbia the least. In relative chest circumference Connecticut stands first, partly because of the racial composition of her population. Indeed, all States which have many representatives of the stocky Mediterranean race stand high in this regard. The tall Southerners stand very low in the series of relative chest girth. For the groups, the largest mean chest circumference is found in such as are occupied by the Finns, agricultural Russians, French-Canadians, German-Austrians, and Scandinavians. Scotch sections and the southern white show the smallest relative chest girth. The chest girth of the Negro troops was relatively somewhat less than that of whites.

5. BUILD.

The best index of build is debatable. The square of stature as a base is probably the most satisfactory. On this basis recruits of 1917-1918 showed a much slenderer build than veterans of the World War. Recruits, and also veterans from Alaska and the extreme northwest, revealed the stoutest build; recruits from the southeast the slenderest. Recruits from Colorado, New Mexico, and Arizona had a slender build on account of the presence of so many tuberculous persons in those States, many of whom had gone there on account of the disease. The absolute increase in the index of build of veterans over

recruits is, for the whole United States, about 0.5. For Colorado it is 1.4, an increase of 4.3 per cent. This may mean a weeding out of the tuberculous or it may mean an extraordinary reaction to the outdoor life of the Army, or both. Some of the Southern States show more than the average increase of build, some less. The Western States show more increase than the Eastern. New Hampshire gave a reduction of index amounting to 0.60, and Florida and Connecticut also a clear decrease.

As compared with Civil War veterans, recruits from our Eastern States show a stouter build; from States west of the Alleghenies, a slenderer build. Of all the sections, men from Alaska, the Finns, the Scandinavians, and those of the North Central States show the heaviest build. The sections with many orientals and Indians also show stout builds. The slightest build is found in the Ozark region and among the mountaineers of the southern Appalachian Mountains. Certain sections of New Mexico and Colorado come low in the list. Of the eight European races, the Poles have the heaviest build and the Scotch and "Irish" the slightest. The condition found in the "Irish" is probably influenced by the Scotch who live in north Ireland. Of the color races, the whites have the slenderest build; the Indian and Chinese the stoutest.

6. OTHER DIMENSIONS.

(a) *Sitting height*.—This is relatively shorter in the Nordic races than in the Mediterraneans. For the color races, it is least in the Negro troops.

(b) *Span*.—Span is slightly greater than stature on the average, but individuals differ greatly in this respect; in some the span is 15 per cent greater, in others 15 per cent less than the mean. For the color races the Negroes have the greatest span in relation to stature, 105 per cent, the white troops least (102) per cent. In relation to sitting height the span of Negro troops is 207 per cent, that of white troops 194 per cent.

(c) *Sternal notch*.—Among the European races the sternal notch is relatively the highest in the Irish (83 per cent), who consequently have the shortest head and neck. It is relatively lowest in the French (81.8 per cent), who have the longest head and neck. It is high among Negro troops (82.8 per cent) and low among whites in general (82.1 per cent).

(d) *Height of pubic arch*.—This dimension measured nearly the physiological length of the leg. In white troops it is about 50.5 per cent of total stature. Among the French the proportion rises to 50.9; among the Italians it falls to 50.1. The Negro troops have relatively long legs (52 per cent) and the Chinese short legs (50.3 per cent).

(e) *Neck circumference*.—This measurement for white troops gives a mean of 35.98 centimeters, or 14.16 inches. The mean man wears about a 14½ or 15 inch collar. The Negro troops have an average neck circumference about 1 per cent larger than that of the white troops.

(f) *The breadth of shoulder*.—The breadth of shoulder is measured between the deltoid muscles. In whites it is 41.8 centimeters, or 24.3 per cent of stature. This is nearly 0.3 centimeters greater than the shoulder breadth of Civil War veterans. In Negro troops the shoulder breadth is about 1 centimeter

more, and the coefficient of variation is much less. Of the eight European races, the Poles have the broadest shoulders, the French the narrowest, the Italians the greatest ratio of shoulder breadth to height, and the French the least. Orientals and Indians have a relatively greater shoulder breadth than whites, but only the Chinese have it as great as the Negro.

(g) *Chest diameter*.—The shape of the chest is given by the thoracic index (transverse diameter $\times 100 \div$ antero-posterior diameter). The Hebrews have the relatively deepest chests (index 131.9), the English the broadest index (134.6). In general, the Nordic races have broad and shallow chests; the Hebrews, Mediterranean, and Celtic races have narrow but deep chests.

(h) *Waist circumference*.—The mean waist circumference of the whites is 77.87 centimeters, or 45.3 per cent of stature. This relative waist girth is greatest among Italians, next among Poles, Hebrews, French, and German, and least among Irish, English, and Scotch. Absolutely the Germans have the largest waists, but not so large a chest girth as the Poles.

(i) *Transverse diameter of the pelvis*.—The human pelvis, like that of the anthropoids, is relatively broad as compared with other mammals. The most striking fact about it is the small breadth in the Negro (16.5 per cent of height) and the great breadth in the Chinese (17.5 per cent of height). Whites are intermediate.

(j) *Leg length*.—The mean leg length is 2.7 centimeters longer for the Negro troops than white. Similarly, arm span is 5.2 centimeters greater. If the Negro race is more like the simians in arm length than whites are, it is less like the simians in leg length, for the simians have long arms but short legs. Similarly, the relative leg length is greatest (43.3 per cent) in the Negro, except for the Japanese (43.4 per cent), and least in the Chinese (41.4 per cent). Of the eight European races the Scotch and Germans have the greatest relative leg length (41.54 per cent) and French and Italians the least (41.06 per cent and 41.07 per cent, respectively).

(k) *Thigh circumference*.—This averages 52.71 for white troops and 54.08 for Negro. It is relatively greatest among Italians and least among Scotch.

(l) *Calf circumference*.—This averages for whites 34.09 centimeters, for Negro troops 34.71 centimeters; but in relation to thigh circumference, calf circumference is somewhat less in Negro than in white troops. Many African tribes are characterized by relatively slender calf.

7. THE GENERAL COMPARATIVE PICTURE OF WHITE AND NEGRO TROOPS.

Tables 103 and 104 give the differences in means and standard deviations of 20 dimensions of white and Negro troops. The results of these tables are shown graphically in Plate I. From the tables and the figure it appears that whereas the average height of white and Negro soldiers is practically the same, the Negro men exceeded, on the average, the white men in the following dimensions:

(a) *Span*.—The total span of the Negroes is about 3 per cent greater than that of white men.

(b) *Leg length*.—Since the lengths of arm and leg are correlated in animals generally, it is in accordance with expectation to find that the leg is longer in the Negro than in the white troops, showing an excess of about 3 per cent.

(c) *Arm length*.—As this constitutes an important part of the span, we may expect, as we find, that arm length will be greater in the Negro than in the white troops.

(d) *Pubic height*.—This measures the physiological length of leg and shows about the same excess in Negroes as leg length.

(e) *Knee height*.—As a component of leg length, knee height shows a slight excess in Negro over white troops.

(f) *Forearm*.—This, as in the total arm length, shows an excess in the Negro troops.

(g) *Sternal notch*.—This is slightly greater in Negro than in white troops. Consequently the height of neck and head together must be less in Negro than in white troops.

(h) *Sitting height*.—Since the total height is the same and the leg length greater in Negro than in white troops, it is clear that sitting height must be less in Negro than in white troops, and such proves to be the case. This smaller sitting height is due in part to the smaller length of head-and-neck in Negro troops as compared with white troops, but also the length of the trunk from the gluteal fold to sternal notch is relatively less in Negro than in white troops.


In contrast with the vertical dimensions the circumferences and diameters show for the most part relatively slight differences between white and Negro troops, largely because they are smaller dimensions. However, certain differences are clearly shown. The circumference of the trunk, whether taken at chest or at waist, is slightly less in Negro than in white troops. The transverse diameter of the pelvis is strikingly less in Negro troops. The breadth of the shoulder, however, is somewhat greater in Negro than in white troops, and the same is true of the circumference of the neck, thigh, and calf.

Despite approximately the same height, Negro troops weighed nearly 5 pounds more than white troops. The index of build of the Negro troops was about 32.7 as compared with 31.6 for white troops.

The general comparative picture we get of the white troops (including a great variety of races) and the Negro troops is this: The Negro troops have relatively longer legs and arms, shorter trunk, narrower pelvis, more nearly circular ellipse of cross-section of the chest; larger, shorter neck; more nearly parallel outlines of the trunk, larger leg girth, and a greater weight than the whites. The waist is less marked because of the relatively small transverse diameter of the pelvis and chest and the greater circumference of the waist. The Negro seems more powerfully developed from the pelvis down and the white more powerfully developed in the chest.

In summary, then, the main differences of shape between Negro and white troops are that the former have relatively longer appendages, shorter trunk, head, and neck, broader shoulders, narrower pelvis, and greater girth of neck, length of thigh and calf, than the latter.

DEMOGRAPHIC INFORMATION



WHITE (b)(6) (b)(7)(C) COLORED (b)(6) (b)(7)(C)

PLATE I.

8. CORRELATIONS.

Correlation indicates similarity of variation; thus the right and left sides of the body are correlated in their variation. The variations of the right arm and leg lengths are correlated less closely than the same organ on the two sides of the body. The larger the correlation the closer is the physiological or developmental interdependence. Considering white troops only (which were the more numerous), the correlations calculated in order of size are given below. It is to be noted that the maximum correlation (approached by the correlation between the two sides of the body) is 1. The minimum is 0. The departure from 0 marks the relative strength of correlations. Probable errors are omitted; none exceeds 0.0021.

Stature and sternal notch.....	0.857
Stature and span.....	.794
Stature and pubic arch.....	.696
Stature and sitting height.....	.663
Weight and chest circumference.....	.660
Arm and forearm.....	.584
Neck girth and chest girth.....	.506
Stature and knee height.....	.436
Leg length and knee height.....	.418
Pelvic diameter and waist girth.....	.351
Chest girth and pelvic diameter.....	.307
Transverse and antero-posterior diameter of chest.....	.271
Chest circumference and sitting height.....	.242
Waist circumference and leg length.....	.159

It will be noticed that the high correlations are often between the measurement of a whole organ and a part of it, like stature and height of sternal notch or of pubic arch. But stature and span are not of this kind, nor weight and chest circumference. However, arm length (span) and leg length vary together and leg length is an element of stature; consequently span varies with stature. When the two dimensions are not closely related, as in waist circumference and leg length, the index is low.

9. DISTRIBUTION OF EYE COLOR.

Eye color is a rough index of race. The fair-skinned, blond-haired people of Europe belong to the "Nordic" race, and have clear blue eyes. The Mediterranean peoples have dark skin, hair, and eyes. The States with the largest proportion of blue eyes have the largest Nordic element. Alaska, with only seven measured, and Wisconsin, with 1,441, lead with 54 per cent; Maine and Vermont also have a large proportion and stand high (probably because of their French-Canadian blood). Then come Minnesota and Oregon. At the bottom of the list stands Florida with only 9 per cent of clear blue eyes. The Negroes, Cubans, and West Indians in its population have dark eyes. Next above Florida comes Georgia and then Nevada, Alabama, Tennessee, South Carolina, and other Southern States, with many Negroes in the population. Roughly, the proportion of clear blue eyes diminishes with latitude. Of the eight European races, Irish and Scotch have the highest percentage of blue eyes (clear blue and blue with brown spots combined), 73 and 71 per cent,

respectively. Polish and English have about 66 per cent, German 65 per cent, French 49, Hebrew 37, and Italian 20 per cent. For the United States as a whole the percentage of blue eyes seems to have dropped from 45 per cent in Civil War times to 38 per cent 55 years later. Blue eyes are passing.

10. DISTRIBUTION OF HAIR COLOR.

Since no measure was applied to hair color, the results are not closely comparable *inter se*. In general, the States with the largest proportion of blue eyes have the largest proportion of blond or flaxen hair. Oregon leads with 28 per cent flaxen hair, Montana comes next with 23 per cent, Utah next with 14 per cent, and then Minnesota and South Dakota with 10 per cent each. The Gulf States show less than 1 per cent.

11. PHYSICAL DIMENSIONS IN RELATION TO DISEASE.

A special study has been made of stature, weight, and chest circumference, with the interrelation of the three measurements of recruits, found with certain diseases and defects. A close relation is found between the physique and defects. Tall men are especially prone to varicose veins, varicocele, pulmonary tuberculosis, cardiac disorders (both functional and organic), and goiter (both simple and exophthalmic). A very high percentage of men with low stature were found with defective teeth and refractive errors of the eye. Heavy weight was found in men with varicose veins and flat-foot; the weight was slightly above the average for those with simple goiter and hypertrophic tonsillitis, while for both organic and functional diseases of the heart and tuberculosis, as well as errors of refraction, the weight was below the average. Chest circumference above the average was found in men with varicose veins and asthma; for the first condition, the large chest was associated with great stature and weight; for the latter with low stature and weight, and hence it seems that large chest was a result of the disease itself. Small chest circumferences were found specially in men with tuberculosis, organic and functional diseases of the heart, and errors of refraction.

Considering the three measurements in the relation of the one to the other, the following points are noted: Men with varicose veins are tall, heavy, and large-chested; with varicocele and hemorrhoids, tall, small-chested, and underweight; with pulmonary tuberculosis and all cardiac disorders, both organic and functional, tall, small chest, and of low weight; with both goiters, the stature is above normal and the chest is small, but for the exophthalmic form the weight is low, while for the simple it is normal. Men with hypertrophied tonsils have normal build; those with relaxed inguinal rings and hernia were slightly below the average in stature and slightly below weight, with relatively small chest; those with flat-foot have low statures, but are very heavy; those with errors of refraction have low stature and low weight, but relatively normal chest. Asthmatic cases show low stature and abnormally low weight, but markedly hypertrophied chest. Men with defective and deficient teeth and congenital genital defects are short, underweight, and small-chested.

The population with different sizes of stature, weight, chest circumference, and build show diverse variability. High variability results when two or more dissimilar classes are combined in one group. Thus myopics who are average-sized combined with a short racial group make a very variable size group. Men in early stages of asthma make of asthmatics a group very variable in chest circumference. Where size and defect are intimately bound together as cause and effect, variability is low. Weight and pulmonary tuberculosis, weight and mitral stenosis, varicose veins and stature, are thus bound together, and variability of the dimension in the population with the disease is low.

Thus, not only the mean dimensions associated with any disease, but also their variability, are of importance in judging the cause and effect of any disease or defect on the human proportions.

SECTION I.

PHYSICAL MEASUREMENTS.

A. THE IMPORTANCE OF ANTHROPOMETRY IN THE ARMY.

An army is made up chiefly of men and their machines. The men deserve first attention. Their mental qualities and their behavior are of importance, but of no less obvious importance is their physique. The significance of the physique of the soldier to the army is everywhere recognized and much effort is expended to select the physically fit. A soldier must have a good nervous system, heart and vessels without serious defect, good feet, strong inguinal muscles and fascia, strong bones and ligaments, and well-functioning joints, keen sense organs, and freedom from organic diseases.

Not only must the soldier be healthy, but he must fall within certain limits of size. In the British army the lower limit of stature during the World War was 60 inches (152 centimeters); in the French army, 154 centimeters (60.6 inches); in the Italian army, 150 centimeters (59.05 inches).³ It may be interesting to consider the following comparative data taken from Baxter¹ (Vol. I, pp. IX-XXXVII). In France the lower limit of stature in the year 1701 was 162 centimeters (63.9 inches); in the year 1804, 154 centimeters (60.6 inches); after Napoleon's return from the fatal invasion of Russia all limitation of the height of conscripts was practically abolished; in the year 1818, 157 centimeters (61.8 inches); in 1830, 154 centimeters (60.6 inches); in 1832, 156 centimeters (61.4 inches); in 1868, 155 centimeters (61 inches); in 1872, 154 centimeters (60.6 inches). Great Britain, in the year 1872, adopted these standards: Cavalry, 66 inches (167.6 centimeters) to 71 inches (180.3 centimeters); Infantry, 165.1 centimeters (64.5 inches) upward. Belgium, in the year 1871: Infantry, 155 centimeters (61 inches); Switzerland, in the year 1857, about 154.9 centimeters (61 inches); Prussia, in the year 1875, 157 centimeters (61.8 inches); Austria, Infantry, in the year 1869, 155.45 centimeters (61.2 inches). In the United States the regulations for the year 1802 placed the minimum height at 66 inches (167.6 centimeters). In 1846 the minimum was placed at 63 inches (160 centimeters); in 1861 at 63 inches and in 1864 at 60 inches (152.4 centimeters). These minimum measurements in 1861 and 1864 were for the Regular Army only. Baxter (Vol. I, p. 22) states that the minimum height authorized by the War Department at the outbreak of the Civil War was 63 inches, and continued to be the regulation height until 1864. However, the enrollment law expressly declared that no exemption should be made on account of height. Gould² (p. 90) also says that no limit of stature appears to have been established for

volunteer troops of the Civil War, and the rule of the Board of Enrollment was that: "The matter of stature shall be considered only in the general examination as to the physical fitness of the men for military service." In 1867 the minimum was placed at 62 inches; in 1874 at 64 inches (Baxter,¹ Vol. I, XLIX). During a period of years preceding 1917 it was 64 inches (162.6 centimeters). In the regulations governing physical examinations under the selective service act, 1917 (P. M. G. O., Form No. 11), the minimum height was placed at 61 inches (154.9 centimeters) and the maximum at 78 inches (198.1 centimeters), and it was stated: "To be acceptable men below 64 inches in height must be of good physique, well developed, and muscular." Also it was stated that "unless exceptionally well proportioned, men above 6 feet 6 inches in height should be rejected." In January, 1918, the minimum height was lowered to 60 inches (152.4 centimeters) (P. M. G. O. Changes No. 3). Special Regulations No. 65, authorized June 5, 1918 (but which came into general use some weeks later), set the minimum stature at 63 inches (160 centimeters); but this was again soon lowered, by an order of the War Department, to 60 inches. Consequently, the minimum height was 61 inches (154.9 centimeters) for the period June, 1917, to February, 1918, and 60 inches (152.4 centimeters) thereafter. Military men urge that soldiers shorter than 60 inches (152.4 centimeters) are not capable of carrying the weight of the prescribed equipment.

The stature of the recruits is of military importance in other respects than as an index of their ability to carry weight. The Division of Food and Nutrition, Office of the Surgeon General of the Army, was interested in the size of soldiers in relation to the standard ration, since this would vary with the size of the body. The 77th Division (containing a large proportion of South Italians and Polish Jews from New York City) required a smaller average ration than the men of the 88th Division, mobilized at Camp Dodge and containing a large proportion of Scandinavians and Germans. The knowledge of the size of the body is also important for making standards for uniforms.

Stature is correlated with length of leg, and length of leg is important from a military standpoint. Prof. Manouvrier,⁴ of Paris, has pointed out that the marching capacity of a company is determined more by the length of leg than by total stature. Hence, soldiers in ranks or platoons, should be sorted on the basis of leg length (crotch height or pubic height) rather than by total stature.

A knowledge of the size of body is important because it varies markedly with the race. Thus, among the races represented in the United States, the average stature of the male is distributed as shown in Table I.

TABLE 1.—*Approximate average stature of principal races represented in the United States, arranged in order of size (from Martin,⁵ pp. 213-217).*

Race.	Mean stature (centimeters).	Inches.
Cochin Chinese.....	158	62.2
Japanese.....	159	62.6
Ruthenians.....	160	63.0
Polish Jews.....	161	63.4
South Italians.....	162	63.8
Roumanians from Hungary.....	164	64.6
French.....	164	64.6
Great Russians.....	164	64.6
Poles from Galicia.....	165	65.0
Roumanians.....	165	65.0
South Russian Jews.....	165	65.0
White Russians.....	165	65.0
Belgians.....	166	65.4
Bavarians.....	166	65.4
Finn.....	167	65.7
Dutch from Holland.....	168	66.1
Danes.....	170	66.9
Serbs.....	170	66.9
Negroes of various origins.....	160-170	63.0-66.9
American Indians.....	162-173	63.8-68.1
Little Russians.....	170	66.9
Letts.....	171	67.3
Swedes.....	171	67.3
Norwegians.....	172	67.7
English (middle class).....	173	68.1
Scotch.....	175	68.9

Thus, between the Cochin Chinese, with a mean stature of 158 centimeters (62.2 inches), and the Scotch, with a stature of 175 centimeters (68.9 inches), is a range in the means of 17 centimeters, or over half a foot.

This diversity of race size has an important bearing on the clothing of the Army. The tariffs of sizes to be supplied to any distribution zone for a draft army will depend on the racial constitution of the population living in that zone. This racial constitution can be approximately known by consulting the most recent census report, which gives for each State the desired information as to country of birth of residents and of their parents.

Another point of contact that the Army has with the race is in forming regiments or companies of particular races. Two divisions (the 92d and 93d) were comprised wholly of Negro troops. The question whether a given person had Negro blood must often have arisen.

On July 31, 1918, the War Department, by General Orders No. 70, issued regulations to govern the raising of troops for a Slavic legion which should be composed of Jugo-Slavs, Czecho-Slovaks, and Ruthenians (Ukrainians). It was ordered that: "Companies will, if practicable, be composed of members of the same race, i. e., Jugo-Slavs, Czecho-Slovaks, or Ruthenians. So far as practicable, Italian regiments will also be organized on this basis. All officers, except field officers of these regiments, will be, so far as practicable, of those races of which the units are composed." It is clear that many cases might arise of doubtful classification, and the special knowledge of anthropology would in such cases be of value in helping Army officials to classify. Actually, on account of the practical cessation of mobilization in the autumn of 1918, the plans for raising such military units composed of European races did not progress far. The incident serves, nevertheless, to illustrate the need in the Army of special knowledge of anthropology.

Again, there is the importance to the Medical Department of a knowledge of the physical dimensions of soldiers individually and in the aggregate or on the average. Thus, despite all other medical methods for diagnosing pulmonary tuberculosis, loss of weight remains one of value. Hence, weight at induction needs to be known accurately. As weight in relation to stature is more important than absolute weight, stature needs to be known accurately. Chest circumference is important for the same reason as weight. The average weight is important in relation to the size of the mess ration as indicated above. Moreover, a knowledge of the proportions of man in relation to certain diseases will direct the wise physician to exercise a special care over the health of men of aberrant proportions, such as narrow or flat chest, extremely long or extremely short legs, a large neck circumference, etc. Special reference will be made in a later section, under the different measurements, of the military bearing of each.

There is still another class of work of an anthropological sort that has to be done in raising and maintaining an army, and that is making and classifying finger prints and other means of identification.

One of the lessons taught by the experience of raising an army in 1917-1918 is that, at the outset, there should be appointed among the officers of the Medical Department a broadly trained anthropologist to whom should be assigned the following tasks: (1) Collaboration in drawing up schedules of the physical examinations; (2) consultation on the taking of the standard measurements and observations on recruits throughout the country and especially at military camps and posts; (3) general supervision of the service of taking identification data; and (4) organization of the service of answering questions that may arise about the racial classification and racial differences of individuals.

B. HISTORY OF THE ANTHROPOLOGICAL WORK IN CONNECTION WITH THE ARMY, 1917-1919.

I. ANTHROPOMETRIC WORK IN CONNECTION WITH THE DRAFT RECRUITS.

On April 6, 1917, Congress declared war against Germany, and on May 17 the selective service act became a law. In accordance with the provisions of this act, 9,925,751 males between the ages of 21 and 30 were registered between June 5, 1917, and September 11, 1918. In addition to this number, between the date of the first registrations, June 5, 1917, and August 24, 1918, 912,564 young men who had in the meantime reached the age of 21 registered. On September 12, 1918, 13,395,706 men between the ages of 18 and 20 and 31 and 45 were also registered. The total number for the three registrations for the United States without the Territories then amounted to 23,908,576.⁶ Out of the approximately 10,000,000 males registered on June 5, 1917, 2,510,706⁷ were measured and examined physically by local boards prior to December 15, 1917. Of this number, 516,212⁸ were entrained for camps. After December 15, 1917, due to the reclassification, upon economic grounds, of all registrants who had not entrained for camps, 3,247,888⁹ men were placed in Class 1. This number included such of the men examined prior to December 15, who were subsequently classified in Class 1, as had not already (prior to Dec. 15, 1917) entrained for camps.

The records of the physical examinations of all the selective service men who had entrained prior to December 15, 1917, and of such of the Class 1 men as were sent to mobilization camps subsequent to that date, was forwarded to the Office of the Adjutant General of the Army.

In October, 1917, Major Albert G. Love was assigned to duty¹⁰ as officer in charge of the Medical Record Section of the Sanitation Division, Surgeon General's Office. Lieutenant (later major) Robert H. Delafield,^a was assigned to duty¹⁰ as assistant to the officer in charge. Steps were immediately taken to reorganize the section for its war work. This work consisted, in brief, of the receipt of all records of sickness or injuries of any character that occurred among the United States soldiers; the examination, care, and preservation of these records; the furnishing of information from them to authorized authorities requesting it; the compiling of statistical material from them for use in the Annual Report of the Surgeon General and in the Medical and Surgical History of the War; and the preparation of the statistical section of the Surgeon General's Report, with the editing of the whole.

Prior to that time the statistics for the report had been compiled by hand method. A punch-card system was at once installed; a code book prepared and published; and Hollerith tabulating and sorting machines installed. It

^a Major Delafield went overseas at his own request in March, 1918, to assist in installing a Hollerith punch-card system in the office of the Chief Surgeon, A. E. F., par. 14, S. O. No. 54, W. D., 1918.

was soon apparent that the work of the section would be incomplete without a thorough statistical study of the reports of the physical examination of the draft recruits. It was also apparent that this work could be done more economically in this section than elsewhere, as it was engaged in similar work with the records of the sick and injured in the military service.

The office of the Provost Marshal General, as well as the Surgeon General's Office, recognized that the data recorded on the reports of the physical examination were of great importance, not only on account of the records of the physical defects noted thereon, but also on account of the anthropological information. Consequently, on December 9, 1917, the Provost Marshal General and the Surgeon General signed a joint communication¹¹ to the Adjutant General requesting that the Surgeon General's Office be allowed to take, under proper safeguard, to the building where the Medical Record Section of the Surgeon General's Office was located, a limited number of these records of physical examination from day to day, that the statistical data might be extracted on Hollerith cards from a sufficient number of them. The Adjutant General, recognizing the desirability of this statistical study, approved the request.¹¹

Instructions were subsequently issued by the Provost Marshal General to the local boards directing them to send to the Office of the Surgeon General one copy of the report of the physical examination of all Class 1 men who had been examined and found by them to be totally disqualified, mentally or physically, for all military service. As the result of this order 549,099 records were received. A Hollerith statistical card was immediately drawn up for this work and a compilation of the statistical data was begun and carried on as opportunity permitted.

In April, 1918, Dr. Charles B. Davenport, of the Carnegie Institute of Washington, became associated with the Section of the Medical Records, where he served in civilian capacity until commissioned major in the Sanitary Corps in July, 1918. A subsection of anthropology was also authorized as a part of the Medical Record Section. The specific purpose of the organization of this subsection at that time was defined as follows:¹²

To secure the highest quality of the measurement of recruits and of identification records as done by the Surgeon General's Office for the purposes of the War Department; to assist, as called upon, in the analysis and synthesis of the statistics compiled from medical records; * * * and to assist the War Department in all questions about racial dimensions and differences.

First Lieutenants E. H. Hawkes and Wilson D. Wallis and Second Lieutenant Louis R. Sullivan were appointed in the Sanitary Corps for anthropological work, with special reference to supervising the finger-print identification work and the recording of the physical examination data at some of the larger camps.¹³

As the result of the statistical study of the draft records, "Physical Examination of the First Million Draft Recruits, Methods and Results," was published in Bulletin No. 11 of the Office of the Surgeon General, March, 1919. This dealt with the varying physical standards and their application at mobilization camps and the distribution of physical defects by States and also by urban and rural districts. Subsequently the complete study of the records of the physical examination of 1,961,692 of these selective service men who were inducted and sent to military camps, and of 549,099 who were rejected by the local boards as totally,

physically or mentally, unfit for military service, was completed and published in "Defects Found in Drafted Men."⁹ In this publication the distribution of the defects is given not only for States and urban and rural districts, but also for 156 population sections of the country separately and grouped into an occupational series, a physiographic series, and a racial series.

Many of the defects and diseases whose distribution is described in these reports are of great anthropological interest, especially the distribution in the racial series of grouped "sections." Some of the findings are that sections containing many French Canadians are characterized by defective appendages (but not an excessive amount of flat-foot), of defective physical development, deficient chest measurements, underweight, underheight, malnutrition, monorchism, cryptorchism, cleft palate, tuberculosis, nervous and mental defects, defective vision, otitis media, defects of the heart, valvular heart disorders, and bad teeth. They form the poorest of the groups from a military standpoint ("Defects Found in Drafted Men,"⁹ p. 299).

The sections containing a large proportion of Scandinavians are characterized by little tuberculosis, venereal diseases, alcoholism, and drug addiction, and by a large excess of goiter and a slight excess of curvature of the spine.

Sections containing a large percentage of "Germans and Austrians" are characterized by relatively little tuberculosis, venereal disease, cancer, arthritis, and obesity, but more than the average of goiter, alcoholism, and drug addiction. Epilepsy, hysteria, mental deficiencies, and defective speech are less common than the average, also teeth defects and hernia. But varicose veins, varicocele, and flat-foot are in excess.

Sections containing a large proportion of Finns have relatively high ratios for multiple sclerosis, monoplegia, disorders of heart action, chorea, defective teeth, and cleft palate.

Sections containing 10 per cent or more of agricultural Russians have high ratios for errors of refraction, diseases of the cornea and retina, otitis media, valvular diseases of the heart, varicose veins, foot defects, and muscular atrophy.

Sections containing many Indians showed a prevalence of well-developed men, except for the congenital defect of cleft palate and harelip.

Sections of the black belt of the South gave an excess of venereal disease, benign tumors, arthritis, mental deficiency, hysteria, dementia praecox, psychoneuroses, manic-depressive psychoses, valvular disease of the heart (especially endocarditis, cardiac hypertrophy, tachycardia), and arteriosclerosis. The following are less than normally common among negroes: Curvature of spine, obesity, the minor paralyses, ear and eye defects, diseases of the throat, varicocele, varicose veins, cardiac arrhythmia, pes planus, cryptorchidism, hypospadias, cleft palate, and harelip.

Measurements of draft recruits.—It has long been recognized that in the Army recruit service the following dimensions should be taken of all recruits: Stature and chest circumference (at expiration and inspiration), and since the Civil War the weight. These measurements were actually taken for all selective service recruits. The regulations issued to the local boards and to the camp examining boards prescribed that all of them be taken with the recruits stripped.

was soon apparent that a thorough statistical study of the draft recruits. It was economically in this way with the records of the

The office of the Office, recognized that the nation were of great physical defects and information. Colonel and the Surgeon General requesting that safeguard, to the General's Office examination from Hollerith card recognizing that

Instruction the local board one copy of had been physically were received this work as opposed

In April 1917, the local board served July, 1917 the M. C. C. subsequent

To the done upon and

F ant wor the

in: lis: Th tic an ex m.

demobilization, and recognizing the importance of anthropometry to the Army, to the Nation, and to science, an effort had been made since the summer of 1917 by the National Academy of Science to secure authorization for special measurements. A special committee was appointed, which met and rendered a report recommending special anthropological measurement. In the stress of the preparation for warfare such authorization was not deemed desirable by the military authorities, nor was such work considered advisable during the period of active hostilities. However, in the latter half of 1919 an order was issued by the Secretary of War to have special measurements of 100,000 men taken upon demobilization, to secure data for dimensions for uniforms.

A telegram was sent by the Surgeon General to Major Davenport, who had been discharged at his own request in January, 1919, as major in the Sanitary Corps (though continuing to serve in the Medical Record Section as a civilian three days a week until about June 1, 1919), requesting him to supervise the measurements to be taken. In accordance therewith he reported to the Surgeon General of the Army on July 7, 1919.

1. ORDERS ISSUED RELATIVE TO SPECIAL UNIFORM MEASUREMENTS.

Orders authorizing special measurements.—On June 9, 1919, the following order was issued by the Acting Director of Operations, General Staff, to The Adjutant General of the Army:

Subject: Sizes of clothing.

11. The Secretary of War further directs that 105,000 data cards be printed by The Adjutant General and turned over to the Surgeon General of the Army to be used in recording data ordered in Section 1. These cards must show the exact places measurements are to be taken in language sufficiently technical to insure accuracy by Medical Department personnel who are to do the work. In addition to the written descriptions of the locations where measurements are to be taken, the data cards should have outlined figures of the body showing front view, with the exact places measurements are to be taken indicated on them, so that they will be readily understood by the persons employed to make the manikins from the measurements. A sample of the outline figures to be shown in the data card will be furnished to The Adjutant General to turn over to the Surgeon General when completed. The measurements and other information to be indicated on the data cards will include the following:

- (1) Name, (2) Home State, (3) Born of native white parents? (4) Born of parents of African descent? (5) Nationality, if born in a foreign country, or of parents who were born in a foreign country, (6) Height (taken standing), (7) Height (taken sitting), (8) Measurement from finger tip to finger tip with arms extended horizontally, (9) Distance from spinous process of vertebra at level of spine of scapulæ laterally back of shoulder and behind elbow (arm held horizontally with elbow bent) to level of tip of styloid process of ulna, (10) Distance, when standing, from floor to presternal notch, (11) Height from floor to superior border of pubis, (12) Transverse diameter of shoulders at level of acromion processes, (13) Transverse diameter of chest just under the arm; that is, at level of articulation of humeri with scapulæ, (14) Transverse diameter of hips level of anterior-superior spines, (15) Anterior-posterior diameter of chest level of junction of ensiform with gladiolus, (16) Circumference of chest, level of nipples, (17) Circumference of waist, level of umbilicus, (18) Circumference of thigh below crotch, (19) Circumference of leg just above patella, (20) Circumference of knee, level of patella, (21) Circumference of calf (at largest part), (22) Circumference of leg just below level of tuberosity of tibia, (23) Inside length of leg from crotch to tip of internal malleolus of tibia, (24) Circum-

ference of neck, level of larynx, (25) If soldier has been fitted by Resco shoe-fitting system under supervision of an officer, state size of shoe worn,

NOTE.—Tape used in measurements should be drawn snug without looseness or compression. Calipers should be used in taking diameter measurements. All measurements will be given in the metric system.

On June 25, 1919, The Adjutant General of the Army sent the following to the Surgeon General:

Subject: Measurement for sizes of clothing.

1. You are directed to have measurements of 100,000 men made. When measured, men should be naked, except for breechcloth, and should have had at least four months of military training. Measurements shall be taken as follows:

Zone 1, 6,000; zone 2, 24,000; zone 4, 3,500; zone 5, 10,500; zone 7, 26,000; zone 8, 10,500; zone 9, 3,500; zone 10, 4,000; zone 11, 4,500; zone 12, 1,500; zone 13, 6,000.

When men about to be demobilized are measured, the taking of the measurements shall not be permitted to interfere in any way with demobilization. The personnel used in taking these measurements should receive such uniform instruction as will insure correctness and uniformity in data.

2. In zones 5, 9, and 10, 35 per cent, 30 per cent, and 25 per cent, respectively, of the men measured should be of African descent. Data cards will be furnished by The Adjutant General, as per memorandum herewith, and when completed should be transmitted to the Equipment Branch, General Staff. These measurements will be used in making manikins from which a pattern for each size can be made.

Haste was essential, since demobilization was being rapidly completed, and at times it was feared that it would be impossible to complete the quota before demobilization had come to an end. This state of mind reflected in some of the orders cited below.

(a) *Detailed directions for measurement.*—On July 23 the following letter was issued to camp commanders by The Adjutant General:

Subject: Measurements for sizes of clothing.

1. The Secretary of War has directed the Surgeon General to have measurements taken of 100,000 soldiers in various camps and stations in the United States, to be used in the construction of manikins of various sizes with the aim of affording better-fitting uniforms for the Army. Your camp has been designated for taking the measurements of

2. An expert anthropologist will be sent to your camp by the Surgeon General to supervise the measuring of the requisite number of men. He should be directed to report to, and to consult with, the camp surgeon, under whose general direction it is intended that the work shall be conducted. To enable him to satisfactorily perform this work the following enlisted personnel is required, which should be furnished by you from whatever source you may see fit. In view of the great scarcity of Medical Department enlisted personnel now on duty in camps it is not contemplated that the number required be drawn from this source alone, but from other staff and line troops as well.

One assistant measurer for every 80 men measured per eight-hour day.

These men should be selected with a view toward accuracy and reliability, noncommissioned officers if practicable.

One enlisted recorder for every assistant measurer.

One enlisted recorder for every 90 men measured per hour, for the purpose of recording descriptive data (name, age, birthplace, etc.) on the face of the blank forms.

One enlisted weigher and one recorder for each 90 men weighed per hour.

One enlisted orderly for every four assistant measurers.

3. In addition each measurer will require about 25 square feet of working space, which should be well lighted, inclosed, and sufficiently quiet so as not to interfere with the proper recording of the data; sufficient furniture, stationery, etc., to enable the work to be expeditiously performed will also be necessary. Blank forms for recording measurements will be furnished by The Adjutant General. The expert anthropologist will bring with him the necessary measuring apparatus.

4. It is directed that the measurements be taken while the men are stripped, and in the case of men who are about to be demobilized who are measured the procedure should not be permitted to interfere in any way with the demobilization. It is believed that this can be accomplished by having these measurements taken as a final step in the physical examination prior to demobilization.

5. As this work is of great importance, you are directed to afford the expert anthropologist every facility possible, both in personnel and material, for performing the duties with which he is charged.

6. You will assign to this work only men of the Regular service. Their work will be so arranged and coordinated by the demobilization officer as not to materially lengthen at any time the period of retention of men sent to your camp for discharge. During periods when the men sent for discharge are not sufficient to keep the measurers busy, men belonging to permanent camp organizations should be sent for measurement. During rush periods when daily discharges exceed the quota which can be measured per day, the excess will not be detained solely for the purpose of being measured.

7. No emergency man, who could otherwise be spared from camp organizations and discharged, will be retained due to the work of the measuring board.

(b) *Instructions issued by Surgeon General.*—On the following days additional instructions and memoranda were issued by the Surgeon General:

JULY 24, 1919.

Subject: Measurement for sizes of clothing.

1. The Surgeon General has received the following instructions from the Secretary of War in a letter dated June 25, 1919:

You are directed to have measurements of 100,000 men made. When measured, men should be naked, except for breechcloth, and should have had at least four months of military training. Measurements should be taken as follows:

Zone 1, 6,000; zone 2, 24,000; zone 4, 3,500; zone 5, 10,500; zone 7, 26,000; zone 8, 10,500; zone 9, 3,500; zone 10, 4,000; zone 11, 4,500; zone 12, 1,500; zone 13, 6,000.

When men about to be demobilized are measured the taking of the measurements shall not be permitted to interfere in any way with demobilization. The personnel used in taking these measurements should receive such uniform instructions as will insure correctness and uniformity in data.

2. Authority has been obtained for the employment of a group of expert anthropologists to undertake this work in the various camps under the general direction of Dr. Charles B. Davenport, now employed in this office. This personnel has already been selected and is now being given instructions by Dr. Davenport relative to methods and procedure in taking measurements in camp. Blank forms have been printed and the necessary apparatus accumulated, and it is proposed to begin this work within the next few days. Necessary instructions have been sent to the commanding general of camps in which measurements are to be made. Your camp has been designated for the measurement of men.

3. As the Surgeon General is charged with carrying out this work, it is desired that the post surgeon act as his representative in camp and give the necessary support and cooperation to the expert anthropologist in immediate charge of the work. As the anthropologist is a civilian and unfamiliar with Army procedure, he will need assistance and guidance from you in order to accomplish successfully his task. The time element is important, as these men are employed under special authority under a limited allotment of funds, and the work in each camp must be pushed with all possible expedition, in order to bring it to a conclusion with our present allotment. It is desired that the post surgeon assume the administrative responsibility for the expeditious handling of the work. The responsibility for the technical features of the work will rest on the expert anthropologist.

(c) *Daily reports.*—On July 25 a letter of instructions was issued relative to the subject and daily reports by telegraph from the anthropologists were called for. From the daily telegraphic reports a table was made up showing the progress of the work day by day.

2. SUPERVISING PERSONNEL AND CAMPS WHERE MEASUREMENTS WERE TAKEN.

(a) *Supervising personnel.*—Personnel to take charge of the measurements at camps had to be assembled and given instruction, and this was accompanied with some difficulties, owing to the fact that most anthropologists had scattered to their summer homes or were working in the West among Indians under the United States Bureau of Ethnology. Eventually the services of the following anthropologists, anatomists, and Army officers were secured to supervise the taking of the measurements at the designated camps. When two or more are named for one camp, the first in order was chiefly responsible for the work. The one or two others were assistants or continued the work after it was well organized:

Dr. Chas. H. Danforth, associate professor of anatomy, Washington University, St. Louis, Mo.
Camp Dix, N. J.

Mr. Frank J. Kelley, biologist, United States Department of Agriculture. Camp Dix, N. J.
First Lieut. Samuel H. Miller, Medical Corps. Camp Dix, N. J.

Mr. Geo. A. Miller, assistant anthropologist, National Museum. Camp Dix, N. J.

Dr. Geo. G. MacCurdy, professor of anthropology, Yale University. Camp Devens, Mass.

Second Lieut. W. B. Davis, Thirty-sixth Infantry. Camp Devens, Mass.

Dr. Robert B. Bean, professor of anatomy, University of Virginia. Camp Lee, Va., and Camp Gordon, Ga.

Dr. E. A. Hooton, instructor in anthropology, Harvard University. Camp Grant, Ill.

Dr. J. A. Mason, anthropologist, Field Museum of Natural History, Chicago. Camp Dodge, Iowa.
and Fort D. A. Russell, Wyo.

Capt. Fred. P. Nevius, Medical Corps. Fort D. A. Russell, Wyo.

Dr. J. R. Terry, professor of anatomy, Medical School, Washington University, St. Louis, Mo.
Camp Sherman, Ohio, and Camp Taylor, Ky.

Maj. Chas. P. Martin, Medical Corps. Camp Sherman, Ohio.

Maj. R. C. Chitting, Medical Corps. Camp Taylor, Ky.

Dr. Daniel Folkmar, anthropologist and statistician, Washington, D. C. Fort D. A. Russell, Wyo., and Camp Lewis, Wash.

Dr. Wm. Howard Griffith, assistant in physical education, University of Pennsylvania. Camp Pike, Ark.

Maj. R. D. Milner, Sanitary Corps. Camp Shelby, Miss., and Camps Travis and Bliss, Tex.

Maj. Samuel Clifford Cox, Medical Corps. Camp Meade, Md., and Camp Holabird, Md.

Capt. Richard M. Alley, Sanitary Corps. Camp Meade, Md.

Capt. Phil. Russell Pope. Camp Shelby, Miss.

To secure uniformity in the measurements taken, the anthropologists, anatomists, and officers who were to be in charge were ordered to Washington, D. C., for special instructions. The offer of Dr. Hrdlicka, curator, Division of Anthropology, United States National Museum, Washington, D. C., to instruct them was accepted. Models were furnished by the Adjutant, Army Medical School, and to each anthropologist, singly or in groups, as the case might be, instructions were given in the prescribed measurements and in the method of taking them. Dr. Hrdlicka also consented to assist as an inspector of the work that was being done in some of the camps. He was consequently appointed on temporary duty in August, 1919, to visit Camp Dix and Camp Devens and to give any assistance that might be possible and to further make a report of the conditions as he found them in those camps.

(b) *Camps, number of men measured.*—The following number of men were measured at the various camps:

Camp Bliss.....	1,509	Camp Meade.....	6,001
Camp Devens.....	6,111	Camp Pike.....	10,500
Camp Dix.....	24,040	Camp D. A. Russell.....	136
Camp Dodge.....	5,046	Camp Shelby.....	3,504
Camp Gordon.....	9,724	Camp Sherman.....	6,981
Camp Grant.....	8,500	Camp Taylor.....	7,014
Camp Holabird.....	1,505	Camp Travis.....	6,005
Camp Lee.....	3,508		
Camp Lewis.....	3,825	Total.....	103,909

3. APPARATUS USED.

It was decided to use the following apparatus:

1. The Seaver measuring rod: Wooden sliding calipers having a 3-foot rod metrically divided, as made by the Narragansett Machine Company.

2. The cloth tape, metrically graduated, made by the same company: These tapes wore out rapidly and had to be replaced. The graduation marks became rapidly obliterated on that part of the tape held by the fingers. For a time steel tapes were used but these occasionally cut the skin and frequently broke if kinked, so that experience proved they were inferior to the cloth tapes. In practice a single tape proved to be good for the measurement of only about 500 men.

3. Graduated paper metric scales furnished by the United States Bureau of Standards: These paper scales were less accurate than metal scales, being subject to alteration in length according to the amount of moisture in the air. Wooden scales would have been better and these were sometimes ruled on the studding of the building by the anthropologist in charge.

4. A plumb line and sinker to measure height of sternal notch from floor, subject standing. Instructions were to use a horizontal arm at the notch from which the line would depend; a pencil or a "tongue depressor" was employed.

4. DIRECTIONS FOR TAKING AND RECORDING MEASUREMENTS.

The following directions for general arrangements at camps for taking measurements, and for recording descriptive matter on the forms that were prepared were issued to the anthropologists in charge:

DIRECTIONS FOR TAKING MEASUREMENTS.

(a) *Stature (W.)*.—Each soldier is to stand against a wall upon which the metric scale has been fastened, accurately calibrated from the floor. The subject stands, heels together and in contact with the wall by buttocks and shoulders, and head in the "front" position, looking straight forward. The squared block is to be placed vertically in contact both with the scale and with the vertex of the head until the resistance of the skull is felt. Standing on the subject's left side, read from the under side of the block while subject is still standing in position.

(b) *Span (W.)* is to be taken standing, the subject touching a fixed strip with the longest finger tip of one hand and reaching out over a graduated scale with the finger tip of the opposite hand, stretching to a maximum. The thumb nail of the operator may be placed in contact with the movable finger tip of the hand which lies upon the scale and the scale read from the maximum position of the thumb nail of the operator.

(c) *Height sitting (W.)*.—A strong box or bench, with a perfectly flat top, is to be placed in contact with the wall, underneath the metric scale on which is to be measured the height of the vertex. The position of the scale should be carefully calibrated, the zero point being at the level of the top

of the box. The subject should sit with the buttocks, shoulder, and head in contact with the wall, unless contact of the head should require the soldier to look upward.

(d) *Height of knee joint (C.)*.—While subject is sitting, with *under* side of movable arm of sliding calipers get height of top of patella from floor.

(e) *Height of sternal notch (L.)*.—This is to be secured by dropping the plumb line and sinker from a short strip of wood ("tongue depressor") held horizontally, subject standing. The plumb line should be held fast by the thumb when the sinker touches the floor and the length of the line plus sinker are to be measured on the scale attached to the wall. The purpose of the horizontal arm is to bring the plumb line in front of the protuberance, if any, of the stomach. The measurement should give the vertical distance of the bottom of the depression of the sternal notch above the floor on which the subject is standing.

(f) *Height of pubis (C.)*.—Use wooden sliding calipers. Standing in front of subject, bring top of sliding arm to level of superior border of the pubis at symphysis. The rod is to be kept horizontal.

(g) *Transverse diameter of shoulders at level of heads of humeri (C.)*.—Use sliding calipers. These are to be in contact horizontally with the skin over the heads of the humeri, the arms of the subject being held at the sides of the body in the attitude of attention. The skin is to be compressed only sufficiently to permit the arms of the calipers to be brought in full contact with the skin, immediately over the head of the humerus. As the contour of the arm at this point is usually not directly vertical, there will be something of a compression of the skin at the lower edge of the arm of the calipers.

(h) *Transverse diameter of pelvis at level of the crests of ilium (C.)*.—The calipers, held horizontally, are to be placed in contact with and pressing upon the skin over the widest part of the ilium, until bone resistance is felt.

(i) *Transverse diameter of chest at level of nipples (C.)*.—The subject stands erect with arms slightly raised in a relaxed position. One arm of the sliding calipers is held fixed against the chest at the level of the nipples. The rod is applied to the chest in front. The movable arm is adjusted by the thumb until brought into contact with the wall of the chest. A series of contacts is made and a mental note made of the readings. This is to allow for changes in form of the chest during respiration. The middle position of the readings is to be recorded. The arms of the calipers will be held somewhat oblique, perpendicular to the axis of the trunk at this level.

(j) *Anterio-posterior diameter of chest (C.)*.—The subject stands in the same position as in (i). The fixed arm of the calipers is applied to the front of the chest at the level of the nipples, the plane of measurement is perpendicular to the axis of the trunk, the movable arm of the calipers is brought in contact with the back or vertebræ. The movable arm of the calipers is brought repeatedly in contact with the back at different phases of inspiration and expiration. The median position of the movable arm in these contacts is recorded.

(k) *Second dorsal vertebra to styloid process of right ulna (T.)*.—Stand behind and to the right of the subject, whose right humerus is raised to a horizontal position; forearm flexed, extending forward at right angles to the humerus. Measure with the tape from the spinous process on the same level with the humerus, along the length of the arm and forearm to the apex of the styloid process of ulna.

(l) *Circumference of neck, level of laryngeal prominence (T.)*.—This measurement is made with the tape from the front. Feel the apex of the laryngeal prominence and pass the tape from the back of the neck slightly down around this prominence perpendicular to the axis of the neck. In measuring with the tape, hold the zero end with the fingers of the left hand in contact with the skin and hold the movable part of the tape with the right hand, guiding that part which comes in contact with the zero end of the tape by means of the forefinger of the right hand. In case of the measurement of a circumference which, like that of the chest, undergoes changes with respiration, read the maximum and minimum and take a strictly intermediate dimension for record.

(m) *Circumference of chest, level of nipples (T.)*.—Arms in the position of (i). The tape is to be placed around the chest and gradually by sliding movements depressed to the required position, which is perpendicular to the axis of the trunk. Make the reading from in front, the tape passing over the nipples.

(n) *Circumference of waist, level of umbilicus (T.)*.—The tape is held in a nearly horizontal position at what is, in "spare" persons, the minimum circumference of the trunk. Read as before.

(o) *Circumference of thigh, maximum (T).*—The measurer kneels at the right side of the subject. The tape is placed around the upper portion of the thigh and passed slowly upward by sliding movements until it reaches the level of the gluteal fold. Legs of the subject slightly spread.

(p) *Circumference of leg just above patella (T).*—The tape is to be passed around the leg and held horizontally, being brought to the desired position, just above the patella.

(q) *Circumference of knee, level of patella (T).*—The tape is to be placed horizontally around the leg and at the middle of the patella in front.

(r) *Circumference of leg just below level of tuberosity of tibia (T).*—The tape is to be brought into the horizontal position, as before, just below the tuberosity of the tibia which lies in the median position in front.

(s) *Circumference of calf, maximum (T).*—The tape is to be brought into a position slightly above the thickest part of the calf, then gradually worked down the leg with repeated readings until the maximum circumference is determined. This is recorded.

(t) *Inside length of leg from the gluteal fold to tip of internal malleolus of tibia (T).*—This is to be measured by the tape from the gluteal fold downward to the apex of the internal malleolus.

(u) The weight of all soldiers measured should be recorded.

In general: Measurements are to be taken so that tape is in close contact with the skin without indenting or depressing it.

Abbreviations: (C.), Calipers; (L.), line and sinker; (T.), tape; (W.), wall.

5. DIRECTIONS FOR USE OF RECORD ON "DESCRIPTIVE" FACE OF FORM.

Write legibly; surname to be printed with pen in capital letters.

1. Under "color," check appropriate square. Judge fraction of Negro blood by estimate of skin color. The mulatto is $\frac{1}{2}$ black, clear brown or dark *café au lait*. If skin color is darker than clear brown, mark $\frac{3}{4}$ black; if light brownish yellow or lighter (and clearly of African descent), mark $\frac{1}{4}$ black. In case of a person of probable Indian, Chinese, or Japanese descent, ask: "Of what race?"

(a) *Hair color.*—There are two series—not-red and red. The not-red series is of four grades. Distinguish clear red and red more or less concealed by brown.

(b) *Eye color.*—Soldier should face light. If no brown pigment on iris, check "clear blue." If some brown pigment but blue field not covered, check "blue with brown spots." If whole iris covered with brown check light, medium, and dark according to degree.

6. SPECIFICATIONS FOR ARRANGEMENTS REQUIRED AT CAMP AND FOR TAKING MEASUREMENTS THERE.

In the building where the physical examinations are taken have erected at the corner of the building nearest the end of the examination line a sufficient number of vertical partitions running perpendicular to the long side of the building to permit of the simultaneous measurement of the number of men specified for each camp. Thus, for the maximum number of 12 sets of apparatus, permitting of the measurement of 12 men simultaneously, there will be required 12 wall spaces at least 6 feet 6 inches wide. These can be secured by using the short end of the room for the measurement of two men and by erecting five additional partitions parallel to the short end of the room against each of which can be measured two men by using the two sides of the partition. The partitions should be not less than 5 feet apart. Adequate lighting by electricity or otherwise is essential and must be secured.

Each partition is to have at the extreme edge a vertical strip of wood about 1 inch wide and $\frac{1}{2}$ inch thick, extending from between 3 and 6 feet from the floor. Midway in the partition are to be affixed to the partition the metric ruled strips or scales provided in the set of apparatus. The scale is printed in 50-centimeter strips. Place two strips vertically, one immediately above the other, the bottom of the lowest strip being precisely 100 centimeters above the floor and the top of the uppermost strip 200 centimeters above the floor. Place two of the 50-centimeter scales in a horizontal position one above the other, so that the ends of the scales nearest to the vertical strip of wood, described above, shall be 150 centimeters therefrom. The bottom of the lower scale is to be 125 centimeters from the floor and the top of the upper scale is to be 165 centimeters from the floor. On the wall rule vertical lines a centimeter apart, connecting these two scales.

Secure a stout box about 50 centimeters high, 50 centimeters long, and 30 centimeters wide, upon which the subject will sit in measuring sitting height. A specially made bench is to be pre-

ferred to a box if such can be made by the camp carpenter. This bench is to be placed at one side of the middle of the partition wall. Immediately over the middle line of the bench is to be affixed to the wall in a strictly vertical position a 50-centimeter section of the scale. The bottom of this scale to be exactly 60 centimeters above the upper surface of the box or bench. The zero end of the scale should then be changed to 60 centimeters; the 10-centimeter mark to 70 centimeters, and so on, the upper limit of the scale then reading 110 centimeters in place of 50 centimeters.

A recorder for each measurer should be seated at a desk in the interspace between every two partitions, or any other convenient position, to record the measurements called off to him by the measurer.

The details of the arrangements of partitions and the direction of passage of the examination line will have to be adjusted to meet the conditions found at the different examination rooms.

Omission of measurements.—The weight was omitted at Camp Gordon, Camp Lee, and Camp Devens. The knee height was omitted at Camp Devens, Camp Sherman, and Camp Taylor. The measurement from the styloid process of the ulna to the elbow was omitted from Camp Sherman and Camp Taylor.

7. STATISTICAL TREATMENT OF DATA.

(a) *System used.*—The taking of the measurements was completed in October, 1919. The data were then transferred to Hollerith punch cards by the use of a prearranged code. This coding and the subsequent handling of the data was all done in the Medical Record Section of the Surgeon General's Office.

(b) *Nationality.*—To determine the nationality of the soldiers measured the following rules were observed:

1. The nationality of all, except Hebrews, who were born in a foreign country, were credited to that country. Hebrews were counted as such without regard to country of birth.

2. Where neither parent was born in the United States, and both were born in the same foreign country, the soldier's nationality was credited to that country; if both parents were not born in the same foreign country the soldier was entered as of mixed origin.

3. If the soldier and both parents were born in the United States, but if three or four grandparents were born in the same foreign country, the soldier's nationality was credited to that country. If three grandparents were not born in the same foreign country, the soldier was classified as of mixed origin.

4. If only one parent was born in the United States and three or four grandparents were born in the same foreign country, the soldier was counted as of that country; otherwise as of mixed origin.

5. When the data furnished were insufficient to determine the nationality, the name was used to determine it, provided the evidence was sufficiently clear.

6. To further determine the nationality the religion was used in such countries as Ireland, where the races are mixed. For example, where both parents were born in Ireland and of the Catholic religion, the nationality was credited to the Irish, but where they were both born in Ireland and of the Protestant religion, the nationality was credited to the Scotch.

Provisions were also made for determining mixed nationalities, but it was decided that it was not advisable to attempt to tabulate statistics for the mixed races.

MEASUREMENT CARD FOR CLOTHING PATTERNS DEMOBILIZATION-1919

Name <u>John Doe</u>		Army Serial No. <u>278659</u>		Home State <u>New York</u>	
Organization <u>Hdqs Co., 313 Infantry</u>		Age <u>22</u>	Color <input checked="" type="checkbox"/> White <input type="checkbox"/> Black <input type="checkbox"/> Indian <input type="checkbox"/> (Check in square.) <input type="checkbox"/> Negro <input type="checkbox"/> Chinese <input type="checkbox"/> Japanese <input type="checkbox"/> Other <input type="checkbox"/>		
Place of observation <u>Camp Dix</u>		Date of observation <u>Sept. 10, 1918</u>		Initials of officer in charge <u>A.M.S.</u>	
Place of birth of—		Country.	State or Province.	City or Town.	
Self		<u>United States</u>	<u>New York</u>	<u>New York City</u>	
Father		<u>United States</u>	<u>New York</u>	<u>New York City</u>	
Mother		<u>United States</u>	<u>New York</u>	<u>New York City</u>	
Nationality of father's father <u>American</u>		Nationality of mother's father <u>American</u>			
Nationality of father's mother <u>American</u>		Nationality of mother's mother <u>American</u>			
Native language of mother <u>English</u>					
Religion of father <u>Protestant</u>		Hair, color— (Check in square.)		<input type="checkbox"/> Flaxen <input type="checkbox"/> Dark brown <input type="checkbox"/> <input type="checkbox"/> Light brown <input type="checkbox"/> Clear red <input type="checkbox"/> <input type="checkbox"/> Medium brown <input type="checkbox"/> Red and black <input type="checkbox"/>	
Other noteworthy racial traits		Eye, color— (Check in square.)		<input type="checkbox"/> Clear blue <input type="checkbox"/> Light brown <input type="checkbox"/> <input type="checkbox"/> Blue with brown spots <input type="checkbox"/> Dark brown <input type="checkbox"/>	

2-404

MEASUREMENT CARD FOR CLOTHING PATTERNS

MEASUREMENTS—ALL METRIC	
0. Weight	145.0
1. Height, standing (stature)	180.1
2. Span (maximum, between finger and tip of outstretched arm)	161.0
3. Height, sitting	86.0
4. Height of sternal notch	132.0
5. Height of pelvis	75.8
6. Transverse diam. of shoulders at level of head of humeri	42.7
7. Transverse diam. of hips, level of crease of thigh	29.8
8. Transverse diam. of chest at level of nipples, arms elevated and flexed	28.8
9. Ant.-post. diam. chest; level of nipples	21.0
10. Greatest dorsal vertebra to styloid process of ulna (elbow bent, horizontal)	68.0
11. Circumference of neck, level of laryngeal prominence, perpendicular to side of neck	58.0
12. Circumference of chest, level of nipples	86.7
13. Circumference of waist, level of umbilicus	85.0
14. Circumference of thigh (maximum)	55.2
15. Circumference of leg, just above patella	37.0
16. Circumference of knee, level of patella	35.0
17. Circumference of leg, just below level of tuberosity of tibia	31.9
18. Circumference of calf (maximum)	34.7
19. Inside length of leg, from gluteal fold to tip of lateral malleolus of tibia	84.0
20. Size of shoe worn if fitted above July 14, 1918, under par. 14, R. R. 28	8-C
Number of measurement <u>6</u>	
21. Height of knee	45.4
22. Length of forearm	28.0

DOTTED LINES INDICATE DIAMETERS
 FULL LINES INDICATE CIRCUMFERENCE
 NOS. 1, 2, AND 19 NOT SHOWN ON FIGURE

Statistics were tabulated for the following nationalities, which were determined as follows:

Irish.—Soldier, both parents, or three or four of the grandparents, all of Catholic religion, born in Ireland. If the data are not clear as to nationality, if the name begins with Mc or O', and if the mother's language is English and the religion is Catholic, he is classified as Irish.

Italian.—Soldier, both parents, or three or four of the grandparents born in any part of Italy other than the northern provinces. If the data are not clear, and if the name ends in a vowel (not Irish or French), with the religion Catholic, classify as Italian.

Hebrews.—All soldiers included in this race were of Jewish or Hebrew religion, whether born in this country or in any of the foreign countries.

English.—All soldiers were classified as English whenever either they, both of their parents, or three or four of their grandparents were born in England, Canada (French Canada excepted), Australia, or New Zealand.

Scotch.—All soldiers were classified as Scotch whenever either they, both of their parents, or three or four of their grandparents were born in Scotland or in Ireland and were of the Protestant religion.

German.—All soldiers were classified as Germans whenever they, both of their parents, or three or four of their grandparents were born in either of the following countries: Germany and Switzerland (mother's language German).

French.—Soldiers were classified as French where either they, both parents, or three or four of their grandparents were born in any of the following countries: France, Switzerland (mother's language French, and religion Catholic), and French Canada (Quebec, Catholic).

Polish.—Soldiers were classified as Polish whenever either they, both of their parents, or three or four of their grandparents were born in Poland (Hebrews excepted).

STATISTICAL PERFORATED CARDS

FIG. 1

Sacramento #2 Name Doe, John L.

Board	5	Sr. No.	00249	X	05	1	62	65	34	37	4	4	5	326	2	18	40	X	11
Occup.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Nat.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Age	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
Bl. No.	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3
Wt.	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4
Ht.	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5
Phy. Det.	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6
Do. Yr.	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7
Man No.	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8
Man No.	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9

WAR DEPT., SURGEON-GENERAL'S OFFICE

FIG. 2

Sacramento #2 Name Doe, John L.

Board	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Occup.	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
Nat.	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3
Age	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4
Bl. No.	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5
Wt.	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6
Ht.	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7
Phy. Det.	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8
Do. Yr.	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
Man No.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Man No.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1

WAR DEPT., SURGEON-GENERAL'S OFFICE

FIG. 3

CAMP Dix

MAN NO.	278659	278659	278659	278659	278659	278659	278659	278659	278659	278659	278659	278659	278659	278659	278659	278659	278659	278659	278659
Occup.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Nat.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Age	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
Bl. No.	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3
Wt.	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4
Ht.	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5
Phy. Det.	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6
Do. Yr.	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7
Man No.	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8
Man No.	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9

WAR DEPT., SURGEON-GENERAL'S OFFICE

PLATE III.

Fig. 1. Statistical card used for tabulating the statistics of the first million draft recruits (P_1).Fig. 2. The same for the second million draft recruits (P_2).

Fig. 3. The same for the special measurements of one hundred thousand veterans, 1919.

C. RESULTS OF THE STANDARD ARMY PHYSICAL MEASUREMENTS.

I. AGE OF RECRUITS.

Table 2, prepared from material published in Gould ^a and from material furnished by the War Risk Bureau, gives the relative frequency of the various ages of officers and men serving in the Civil and World Wars. It is apparent that the great majority of the men measured for the data in this book were between the ages of 18 and 31, inclusive. (See Plate IV.)

TABLE 2.—*Ages of soldiers (officers and men) serving in the Civil ^a and World Wars.^b*

Age.	Civil War.		World War.	
	Number.	Proportion per 1,000.	Number.	Proportion per 1,000.
13.....	127	0.12
14.....	330	.31	16	0.00
15.....	771	.71	140	.04
16.....	2,763	2.63	935	.25
17.....	6,430	6.13	12,846	3.50
18.....	133,653	127.35	62,849	17.11
19.....	90,624	86.36	122,977	33.48
20.....	71,745	68.36	152,635	41.56
21.....	98,766	94.11	293,161	79.81
22.....	75,230	71.69	506,426	137.87
23.....	61,818	61.76	440,581	119.94
24.....	54,329	51.76	381,321	103.81
25.....	48,787	46.49	328,185	89.34
26.....	42,357	40.36	283,276	77.11
27.....	36,254	31.55	235,901	64.22
28.....	37,383	35.62	214,133	58.30
29.....	26,269	25.03	187,010	50.91
30.....	30,196	28.78	160,735	43.76
31.....	19,383	18.47	117,316	31.94
32.....	23,580	22.47	47,890	13.04
33.....	19,401	18.49	20,967	5.71
34.....	17,064	16.26	16,407	4.47
35.....	20,414	19.45	13,318	3.63
36.....	15,278	14.56	10,992	2.99
37.....	12,851	12.25	9,356	2.55
38.....	14,379	13.70	9,086	2.47
39.....	10,409	9.92	8,039	2.19
40.....	14,869	14.17	6,747	1.84
41.....	7,992	7.62	5,165	1.41
42.....	11,585	11.01	4,067	1.11
43.....	10,825	10.32	3,438	.94
44.....	16,668	15.88	3,077	.81
45.....	7,490	7.14	2,560	.70
46.....	1,184	1.13	2,050	.56
47.....	896	.85	1,680	.46
48.....	874	.83	1,543	.42
49.....	590	.56	1,237	.34
50 and over.....	2,889	2.75	5,038	1.37
Total.....	1,019,456	1,000.01	3,673,133	999.99

Average age, Civil War, Gould's figures, 25.54, volunteer officers and enlisted men.

Average age, Civil War, Baxter's figures, 27.307 (Baxter, p. 51), drafted recruits, substitutes, and late volunteers.


Average age, World War, 24.89 for all officers and enlisted men. Other data in this study for draft, enlisted men only.

^a Gould, 1899, pp. 31 and 57.

^b Estimated from ages furnished by 3,683,134 applicants for War Risk Insurance.

AGE DISTRIBUTION CIVIL WAR VOLS., AND WORLD WAR TROOPS, OFFICERS AND ENLISTED MEN

RATIOS PER 1000



AVERAGE AGE, CIVIL AND WORLD WAR

25.84

27.307

24.89

MIL. TROOPS

PLATE IV

Gould stated that apparently many who were under 18 or 21 gave their ages as such that they might be able to enlist at the minimum age of 18 (with consent) or at the minimum legal age of 21.

Baxter's drafted recruits included all troops raised during the draft period, that is draftees, substitutes, and late volunteers

II. STATURE.

1. GENERAL DISCUSSION.

The distance from the sole of the foot to the vertex of the head is one of the most striking of human dimensions and one of the most easily secured. It is used in anthropology as the basal dimension with which minor dimensions are compared in forming the so-called *relative* lengths. Such relative lengths are obtained by dividing the minor dimensions by the stature.

Despite its striking nature, the ease of taking it and its universal use (often as standard of comparison), stature is not altogether satisfactory as a fundamental dimension. The principal objection to it is that it is a complex of dimensions of varied significance, the length of the trunk (in many ways the most significant single measure, but difficult to take), the length of the neck, the height of the head, and the length of the inferior (posterior) appendages of the trunk. Actually, the dimensions of the trunk and legs overlap. In many ways the best standard for human dimensions would be the distance of the sternal notch from the buttocks, that is, the sitting height of the sternal notch. This may readily be taken. The *relative* dimensions of this paper, however, will have for their basis the total stature.

The military reason for laying much stress on stature lies partly in its convenience as a fundamental measure and partly in military history. The potentates of Europe from early time prided themselves on their tall soldiers; they rejected the poorly developed as fit only to stay at home to cultivate the land and to reproduce their kind. It is customary, also, in many army formations to keep together men of about the same height, partly to enable the ranks to keep step better. The latter purpose is imperfectly met, in so far as keeping step depends rather upon similarity of leg length than of total stature; and the two dimensions are not very closely correlated. The military importance of stature is emphasized by the fact that total stature of recruits is taken at practically all recruit stations in all countries. Thus, armies may be compared in respect to average height of their soldiers. Differences in sizes of men of military age between various countries may be quantitatively expressed.

Stature is of great medico-military importance, as it is the basis by which may be judged the build or robustness of the man. Experience has shown that a certain chest circumference and a certain weight are essential for the successful soldier. These measurements are, however, to be judged in relation to stature and not absolutely. This will appear directly in the section relating to the standards of height, weight, and chest circumference. The importance of stature in relation to weight and chest circumference depends on the fact that it gives a warning for tuberculosis, hook-worm, and other diseases.

The method of measuring stature is a simple one. There is affixed to the wall a bit of metric (or English) scale, preferably of wood and accurately calibrated so that it records the vertical distance from the floor. For military purposes the range of the scale need be only from 150 to 200 centimeters, or 59 to 79 inches. To measure, the subject's shoes must be removed and the subject made to stand with his back to the wall at the point of the scale. For a vertical

arm, by which the height of the vertex is secured, one can not do better than to follow the directions given in the Standards of Physical Examination of the P. M. G. O., Form 75 (second edition), page 79, which read as follows:

Directions for taking height.—Use a board at least 2 inches wide by 80 inches long, placed vertically, and carefully graduated to one-quarter inch between 58 inches from the floor and the top end. Obtain the height by placing vertically in firm contact with the top of the head and against the measuring rod an accurately squared board of about 6 by 6 by 2 inches, best permanently attached to graduated board by a long cord. The registrant should stand erect with back to the graduated board, eyes straight to the front.

It remains only to state that the subject should be cautioned to stand in the "front" position, heels close together, buttocks (and shoulders) in contact with the wall.

2. MEAN STATURE.

The mean stature of the 868,445 recruits of whom the weight was also secured is, as shown in Table 11 (based on Table I), 67.49 inches, or 171.4 centimeters. The mean, in English units, is easily remembered as very near to $67\frac{1}{2}$ inches, or 5 feet $7\frac{1}{2}$ inches; also the metric height is almost exactly $1\frac{1}{4}$ meters. This number is probably close to the average for the entire male population of the ages of 21 to 30 years, inclusive, since the 873,000 men were drawn from all States of the Union in about the proportion of the population and without any obvious selection. It includes thus a great mixture of races whose height is known to be very variable.

The mean stature of 102,304 men, measured at demobilization and including both white and colored, is 67.72 inches, or 1,720 millimeters (Table 14 based on Table CXXXIII). This shows an increase in mean stature of men measured at demobilization over men measured at draft of 0.23 inch, or 6 millimeters. The increase in stature may possibly be due in part to the fact that the men at demobilization averaged more than a year older than at mobilization; in part that they were straighter, in part that some of the shortest divisions were not included in the measurements made at demobilization, and in part that some of the shorter men were excluded at the mobilization examination and hence not included in the demobilization measurements.

3. COMPARISON OF MEAN STATURE WITH CIVIL WAR RECORDS.

This mean stature of 67.49 inches may be compared with the statures obtained from recruits during the Civil War as given by Baxter¹ and Gould.² The average stature given by Baxter (Vol. I, p. 23) for 501,068 recruits of all nationalities measured by the Provost Marshal General's Bureau of Civil War times is 67.30 inches (1,709 millimeters). This is an average of stature obtained probably by the same method as that employed in measuring the drafted men of 1917-1918. Our measurements show an increase of 5 millimeters over the Civil War data. Our data alone exclude men rejected by the State or local boards. There was no minimum height for the Civil War draft, it being stated that no exemptions should be made on account of stature (Baxter, Vol. I, p. 22); at the beginning of the draft in 1917 it was 61 inches. In the Civil War draft the manhood of the Northern States had been much depleted by volunteer

enlistment prior to the draft. For Gould's (p. 105) data for 1,104,841 white volunteer soldiers, probably very crudely measured and recorded at the beginning of the Civil War, when the minimum height of 63 inches was prescribed but probably not adhered to, the average height was 67.64, or 171.8 centimeters. This is 0.15 inch greater than our average, which was in turn 0.19 inch greater than Baxter's average. The weighted average for the two groups combined was 67.502 inches, practically the same as our own.

It might be concluded, then, that the mean stature of men of military age has changed little in the United States in the last 50 years, and that our population, so far as stature goes, is placed in the same category as the Scandinavians and below the English middle class. But this conclusion would be hasty. The men of 1917-1918 were taken from all parts of the United States, while those of 1864-1865 largely excluded the Southern States; and since the men of these States are exceptionally tall, their inclusion probably tends to raise the mean stature. A more careful consideration has shown that the mean stature of American males 21 to 30 years has probably diminished since Civil War days about one-half inch. This is chiefly the result of the immigrants during the past half century of short races.

4. COMPARISON OF MEAN STATURE IN VARIOUS COUNTRIES.

It may be instructive to compare the mean height of other countries with the 1,714 millimeters which constitutes the mean height of the young males of the United States (21 to 30 years of age). This average places the United States in the group of nations characterized by a high average stature. This average is almost the same as that of Scandinavian males, 1,710 millimeters. It is about 30 millimeters less than the average of Scotch, 1,746 millimeters, and about 80 millimeters less than the agricultural Scotch of Galway, who, according to Deniker¹⁴ (p. 584), have an average stature of 1,792 millimeters. This average, however, is based on only 75 subjects, and thus may be influenced by accidental inclusion of a few exceptionally tall men. The following table gives the stature of various European races as listed by Martin⁵ (pp. 213-217):

TABLE 3.—Average statures of European males of various countries.

Group.	Stature (milli- meters).	Group.	Stature (milli- meters).
Laplanders from Scandinavia.....	1,523	Turks from Balkans.....	1,660
Jews of Russian Poland.....	1,612	Venetians.....	1,666
Magyars from west Hungary (conscrip- ts).....	1,619	Finns.....	1,666
Corsicans.....	1,633	Thuringians of Saxony (conscrip- ts).....	1,667
Austrian Jews of Hungary.....	1,633	Ukrainians.....	1,669
Roumanians of Hungary.....	1,635	Dutch in general.....	1,675
Portuguese.....	1,637	Poles in general.....	1,679
Hungarians (conscrip- ts).....	1,637	Swedes of Kalmar (conscrip- ts).....	1,681
Bulgarians of western Bulgaria.....	1,638	Danes.....	1,691
Lithuanians of Russian Poland.....	1,639	Welsh.....	1,685
Italians in general.....	1,640	Swedes in general (soldiers).....	1,705
French (conscrip- ts) (Rapillault, 1902).....	1,641	Serbs (conscrip- ts).....	1,709
Esthonians.....	1,642	Bosnian-Herzegovinians (soldiers).....	1,710
Lithuanians of Lithuania (conscrip- ts).....	1,643	Inhabitants of United Kingdom of Great Britain and Ireland.....	1,719
Spaniards.....	1,645	Norwegians (soldiers).....	1,720
Conscrip- ts of French Switzerland.....	1,646	Laplanders.....	1,736
Roumanians (conscrip- ts).....	1,650	Scotch in general.....	1,746
South Russian Jews (Weissenberg, 1895).....	1,651	Scotch of the north, Ayrshire, etc.....	1,786
Greeks.....	1,651	75 Scotch, agriculturists of Galway.....	1,792
White Russians.....	1,652		
Dutch of the Province of Zeeland (conscrip- ts).....	1,655		

TABLE 4.—*Stature, its mean, standard deviation, and coefficient of variation for men (and in part for women also) for certain especially studied groups (Harris and Benedict,¹ pp. 53-54).*

Series.	Men.			Women.		
	Mean.	Standard deviation.	Coefficient of variation.	Mean.	Standard deviation.	Coefficient of variation.
	<i>Centimeters.</i>	<i>Centimeters.</i>	<i>Per cent.</i>	<i>Centimeters.</i>	<i>Centimeters.</i>	<i>Per cent.</i>
American:						
Harvard students.....	175.34	6.58	3.76			
Army recruits.....	170.94	6.56	3.84			
English:						
Oxford students.....	176.50	6.61	3.74			
Cambridge students, Pearson.....	174.91	6.41	3.66	162.26	6.00	3.70
Cambridge students, MacDonell.....	174.88	6.46	3.70			
Pearson's second generation.....	174.37	6.88	3.95	162.23	6.63	4.09
Pearson's family records.....	172.81	7.04	4.07	159.90	6.44	4.03
Pearson's parental generation.....	171.91	6.86	3.99	158.70	6.07	3.83
New South Wales criminals.....	169.87	6.58	3.87	158.09	6.15	3.89
Scottish students.....	171.70	5.94	3.46			
MacDonell's convicts.....	166.46	6.45	3.88			
Goring's convicts.....	166.29	6.76	4.06			
Swedes.....	169.79	6.81	4.01	158.71	6.72	4.23
Hessians.....	167.36	7.19	4.30	156.18	6.90	4.40
French.....	166.80	6.47	3.88	156.10	6.79	4.35
Bavarians, Pearl.....	166.55	6.39	3.84	154.71	6.21	4.02
Bavarians, Pearson.....	165.93	6.68	4.02	163.85	6.55	4.26

TABLE 5.—*Average stature of adult males of various nationalities in the United States in the Civil War period (from Baxter,¹ Vol. I, p. 32).*

Nativity.	Number of men.	Mean height.	
		<i>Inches.</i>	<i>Centimeters.</i>
United States, Indians.....	121	67.934	172.55
United States, whites.....	315,620	67.672	171.89
Norway.....	2,290	67.467	171.37
Scotland.....	3,476	67.066	170.35
British America.....	21,645	67.014	170.22
Sweden.....	1,190	66.896	169.92
Ireland.....	50,537	66.741	169.52
Denmark.....	383	66.648	169.29
Holland.....	989	66.637	169.26
Hungary.....	89	66.584	169.12
England.....	16,196	66.577	169.11
Germany.....	64,944	66.536	169.00
United States, colored.....	25,828	66.531	168.99
Wales.....	1,104	66.418	168.70
Russia.....	122	66.393	168.64
Switzerland.....	1,802	66.381	168.61
West Indies.....	580	66.307	168.42
France.....	3,243	66.277	168.34
Poland.....	171	66.211	168.18
Mexico.....	91	66.110	167.92
Italy.....	339	66.000	167.64
South America.....	79	65.899	167.38
Spain.....	148	65.635	166.71
Portugal.....	81	65.432	166.20
Total.....	501,068		
Total frequency and mean of.....		67.300	170.94

TABLE 6.—*Frequency distribution of stature by classes at mobilization and demobilization (white and Negro troops), 1917-1919.*

A. First million draft recruits. ¹		B. 103,410 troops at demobilization. ²		
Inches.	Per 1,000.	Centimeters.	Inches.	Per 1,000.
59.....	3.534	148-149	58.3-58.7	0.22
60.....	3.354	150-151	59.1-59.4	.55
61.....	8.672	152-153	59.8-60.2	1.47
62.....	18.150	154-155	60.6-61.0	3.88
63.....	35.740	156-157	61.4-61.8	7.41
64.....	60.611	158-159	62.2-62.5	14.27
65.....	94.400	160-161	63.0-63.4	27.51
66.....	126.914	162-163	63.8-64.2	44.04
67.....	146.927	164-165	64.6-65.0	65.27
68.....	149.599	166-167	65.4-65.7	85.09
69.....	127.265	168-169	66.1-66.5	103.63
70.....	96.065	170-171	66.9-67.3	120.25
71.....	62.542	172-173	67.7-68.1	119.27
72.....	36.102	174-175	68.5-68.9	112.33
73.....	17.504	176-177	69.3-69.7	94.80
74.....	7.342	178-179	70.1-70.5	74.49
75.....	3.001	180-181	70.9-71.3	49.72
76.....	1.237	182-183	71.7-72.0	32.01
77.....	0.413	184-185	72.4-72.8	19.65
78.....	0.293	186-187	73.2-73.6	11.75
79.....	0.341	188-189	74.0-74.4	6.67
		190-191	74.8-75.2	2.94
		192-193	75.6-76.0	1.54
		194-195	76.4-76.8	.57
		196-197	77.2-77.6	.38
		198-199	78.0-78.3	.22
		200-201	78.7-79.1	.06
		202-203	79.5-79.9	.02
		204-205	80.3-80.7	.11
		206-207	81.1-81.5	.00
		208-209	81.9-82.3	.00

¹ From Table I.² From Table LXXV.TABLE 7.—*Stature of Army conscripts and recruits, in inches, as determined by Laplace-Charlier frequency curves (by Arne Fisher, from Hoffman,¹⁰ p. 53).*

[Ratio per 1,000.]

Inches of stature.	United States Army recruits, 1906-1915.	Norwegian conscripts, 1913.	Swedish conscripts, 1914.	Danish conscripts, 1916.	Wurttemberg conscripts, 1911.	Japanese conscripts, 1916.
56.....						4.7
57.....						12.5
58.....						31.6
59.....			1.3	2.9	1.2	64.0
60.....			2.2	6.7	7.0	106.5
61.....		2.1	5.4	15.0	22.7	148.6
62.....	3.8	9.9	12.9	30.2	53.2	173.0
63.....	19.2	29.4	27.5	54.0	96.9	169.8
64.....	53.8	60.1	53.4	92.6	141.7	132.8
65.....	105.5	100.3	88.6	130.5	167.5	83.9
66.....	155.7	137.6	127.8	157.7	164.8	44.2
67.....	182.2	165.2	155.3	160.0	137.2	20.8
68.....	169.5	163.1	159.7	136.0	97.0	6.4
69.....	129.4	132.8	138.5	96.9	59.2	1.2
70.....	86.8	96.4	102.0	59.4	30.9	
71.....	51.0	58.9	63.4	31.7	13.6	
72.....	26.3	28.7	34.6	15.2	5.0	
73.....	11.4	11.3	16.4	6.7	1.5	
74.....	4.2	3.2	7.0	2.8	.3	
75.....	1.2	.5	2.7	1.1		
76.....		.3	1.3	.4		
77.....		.2		.2		

TABLE 8.—*Calculated frequency distributions of statures of men of United States Civil War period, France, Belgium, and Italy (Baxter,¹ Vol. I, p. LXXXI, and Livi,¹⁷ Anthropometria Militare).*

[Ratio per 1,000.]

Stature.		United (States B. A. Gould).	France (D'Hagen- villers).	Belgium (Quetelet).	Italy (Livi).
Centimeters.	Inches.				
133	52.5			0.1	0.3
136	53.5		0.5	.3	.5
139	54.7		1.6	1	1
142	55.9		4.5	3	2
145	57.1		11	7	4
149	58.3	1	24	14	9
151	59.4	4	44	28	24
154	60.6	11	73	53	59
157	61.8	24	106	107	106
160	63.0	45	132	136	122
162	63.8	75	145	150	150
165	65.0	109	140	150	167
168	66.1	137	118	136	137
170	67.0	150	87	107	96
173	68.1	142	55	53	70
176	69.3	117	32	28	35
179	70.5	84	16	14	15
181	71.3	52	7	7	4
184	72.4	28	3	3	2
187	73.6	13	1	1	1
190	74.8	5	.3	.3	.2
192	75.0	2		.1	
194	76.4	1			
		1,000	1,000	1,000	1,000

5. FREQUENCY DISTRIBUTION.

While the mean is probably the best single measure of the stature of the country as a whole, still the relative frequency of the different statures (inches) will be highly instructive. This is shown in the second column of Table 6, which gives the proportion of drafted men of 1917-1918 of each stature from 59 inches (strictly, 59 inches and below) up to 79 inches (strictly, 79 inches and above). One sees that the statures below 62 inches are relatively uncommon; but this is in part due to the fact that, during a brief period of the draft, men with a stature below 63 inches were rejected, so that some such men were excluded. The sudden diminution of the number of men below 63 inches is thus in part due to a process of selective elimination of the short men. The effects of this selection are still more marked in the case of men 59 inches and under. No men of this stature were supposed to be accepted for military service. Their inclusion, therefore, is partly accidental, and partly due to the intentional acceptance, in spite of their short stature, of men of exceptionally good build. Instead of less than 4 men per 1,000 of our population being 59 inches or under, it is probable that the inclusion of all cases would give 10 per 1,000 or more.

As the distribution in Table 6 shows, the commonest stature at mobilization was 68 (67.5 to 68.4) inches—a stature found in about 15 per cent of our young men.^a About 10 per cent measured 70 inches in height, less than 4 per cent 72 inches in height, and above that stature to that of 78 inches the proportional numbers fall rapidly.

^a In Danish conscripts the mode is 67 inches, found in 160 per 1,000 men. For conscripts from Wurttemberg the modal stature is 65 inches, found in 168 per 1,000 men. For Japanese conscripts the mode is 62 inches, found in 173 per 1,000 men. (See Table 7.)

For comparison with Table 6 there are printed Tables 7 and 8, which give for various countries the findings as to frequency distribution of statures. The frequency distribution of stature of 103,410 men at demobilization is given in 2-centimeter classes in Table 6, extracted from Table LXXV. The total range in stature is from 148 to 209 centimeters and above. This tends to raise the class 208-209 above the class 206-207 centimeters, because the former class really has a much more inclusive range than the latter. The total range is from 58.3 to 82.3 inches. There are only seven cases above 200 centimeters, or 79 inches, and it is probable that some of these are due to errors in recording.

Table 7, taken from Hoffman¹⁶ (p. 33), and Table 8 give the comparative distribution in statures of conscripts of different countries, Civil War volunteers, per 1,000. We note that for the United States Army recruits, 1906-1915, the commonest or modal height is 67 inches, a class that contained 182 per 1,000 men. For Norwegian conscripts the mode is also 67 inches, with 165 per 1,000 men. For Swedish conscripts the mode is 68 inches, found in about 160 per 1,000 men.

The accompanying Table 9 gives the direct comparison of the distribution of statures of recruits of 1917-1918 (Table I) with that of Civil War recruits as given by Baxter¹ (Vol. II, Table 3) for 501,068 Civil War draft recruits of all countries of origin.

TABLE 9.—*Comparison of frequency distribution of statures, United States recruits, Civil War and World War.*

Classes (inches).	Civil War.	1917-1918	Classes (inches).	Civil War.	1917-1918.
Under 61.....	7,478	6,923	67-68.9.....	258,683	296,882
61-62.9.....	41,587	26,624	69-70.9.....	177,205	223,630
63-64.9.....	141,773	96,127	71-72.9.....	64,488	98,714
65-66.9.....	263,611	220,932	73 and over.....	15,174	30,164

This table shows that there were slightly fewer (per mille) recruits under 61 inches chosen in 1917 than 1864. There were nearly twice as many men 73 inches and over chosen in the latter as in the earlier period, and 50 per cent more men of 71.3 inches. The great deficiency in the later series is in men of mediocre size, namely 63-66.9 inches. This, again, is in accordance with the history of immigration, since within the last 50 years the United States has experienced a great immigration of Scandinavians on the one hand and of south Italians and Polish Jews on the other. However, as pointed out above, the great excess of relatively tall men in the later series is due to the inclusion therein of many tall white men from the Southern States.

The data supplied by the draft boards gave no indication of age; therefore it is impossible to make comparison with the statistics of Gould,² in which the statures are carefully distributed by age of recruits. From Hoffman's¹⁶ (p. 37) paper based on stature of the United States Army recruits, 1906-1915, it appears that the mean stature increases preceptibly up to 22 to 24 years and then diminishes at greater ages.

TABLE 10.—*Mean stature at each age, 18 to 25 years, United States Army recruits, 1906-1915 (Hoffman,¹⁰ p. 37).*

Age.	Mean stature.	
	Inches.	Centimeters.
18.....	66.900	169.83
19.....	66.965	170.09
20.....	67.024	170.24
21.....	67.329	171.02
22.....	67.341	171.05
23.....	67.329	171.01
24.....	67.367	171.16
25 and over.....	67.325	171.01

6. STANDARD DEVIATION.

The standard deviation of stature for the first million recruits, 1917-1918, is 2.71 inches (6.88 centimeters). (See Table I.) The standard deviation of the English upper middle class, with a stature of 69.22 inches, is 2.59 inches, and for Cambridge University students, with 68.86 inches of stature, the standard deviation is 2.52 inches. Since variability is measured by standard deviation, and since it tends to increase with the mean, it is more usual to make comparison with the standard deviation divided by the mean, the so-called coefficient of variation. The coefficient of variation thus obtained is, for the United States recruits, 4.02 per cent; for the English middle class, 3.74; for Cambridge University students, 3.66. The relatively large size of the coefficient of variation for United States recruits signifies that the population is much more variable in stature than even the population of the English middle class. It is indeed about 8 per cent more variable. We can understand this high variability of the mean stature for the United States recruits in view of the heterogeneous composition of the population of the United States.

The standard deviation of 501,068 Civil War recruits, using Baxter's figures, (Vol, II, Table 3), is 2.664 ± 0.002 . Of recruits of 1917-1918 the standard deviation of stature is 2.71 inches. Thus it appears that the standard deviation of the military population of the United States in 1917-1918 has increased slightly from that of 1865. Similarly the coefficient of variation has increased from 3.96 to 4.02. The difference is clearly to be explained by the inclusion in the 1917-1918 figures of many Scandinavians on the one hand and representatives of the south Italian and Jewish races on the other. It is also influenced by this inclusion of tall southern recruits in the later series.

The standard deviation of mean stature for white troops at demobilization is 6.66 centimeters (1.69 inches), with a probable error of ± 0.01 ; for Negro troops at demobilization, 6.91 ± 0.04 (1.76 inches). Negro soldiers are more variable than white.

TABLE 11.—*Distribution of stature and weight, draft recruits of 1917-1918.*

Classes of stature (inches).	Classes of weight (pounds).						Total.
	Under 100	100-119	120-139	140-159	160-179	180 and over.	
Under 61.....	0.038	2.021	2.627	1.611	0.500	0.126	6.923
61-62.9.....	.078	10.556	12.335	2.863	.586	.206	26.624
63-64.9.....	.061	24.605	52.594	15.586	2.663	.618	96.127
65-66.9.....	.029	28.185	120.903	58.668	10.849	2.298	220.932
67-68.9.....	.005	14.333	134.539	115.311	26.780	5.914	296.882
69-70.9.....	.001	3.349	68.364	107.105	36.330	8.481	223.630
71-72.9.....		.507	15.782	49.064	26.321	7.080	98.714
73 and over.....		.137	1.961	11.883	11.623	4.560	30.164
Total.....	.212	83.693	409.085	362.091	115.652	29.263	999.996

Mean stature, 67.49 inches. Standard deviation, 2.714 inches. Coefficient of variation, 4.021 inches.

TABLE 12.—*Distribution of stature and weight in 6,359 American born Civil War draft recruits (Baxter,¹ Vol. II, p. 300).*

Classes of stature (inches).	Classes of weight (pounds).						Total.
	Under 100	100-119	120-139	140-159	160-179	180 and over.	
Under 61.....	0.315	0.629	0.472	0.157			1.573
61-62.9.....	.629	13.996	8.177	1.887	0.157		24.847
63-64.9.....	.315	60.230	67.621	9.435	1.258		138.858
65-66.9.....	.629	58.500	170.939	57.242	4.403	1.258	292.971
67-68.9.....	.315	17.927	161.661	116.056	13.681	1.887	311.527
69-70.9.....		4.560	50.951	84.604	27.363	3.145	170.624
71-72.9.....			6.133	24.375	16.355	2.359	49.222
73 and over.....			.786	4.089	3.931	1.573	10.379
Total.....	2.202	155.842	466.740	297.846	67.149	10.222	1,000.000

Mean stature, 67.30 inches. Standard deviation, 2.3956 inches. Coefficient of variation, 3.560 inches.

7. MEAN STATURE FROM DIFFERENT STATES.

(a) *Recruits*.—The mean stature of 67.49 inches for recruits is obtained by lumping the statures of recruits from all States. It will be of interest to compare the stature of men from the different States. This comparison is made in Table 13, which gives the mean stature both in inches and centimeters for the different States, arranged in order of standing, the State with the highest stature being placed first. This table shows that the men of Texas have approximately an inch greater stature, on the average, than those of the entire United States, while men from Rhode Island have a stature an inch below the mean of the United States. The great stature of men from Texas is partly due to the fact that there has been to that State a very small immigration of men with the shorter statures characteristic of southeastern Europe. As shown in Table 17 probably in Texas under 1 per cent of the population is Italian, while Germans and Austrians are relatively common; native whites of native parentage comprise nearly 50 per cent, while nearly 25 per cent are Negroes. On the other hand, in Rhode Island 8 per cent of the population is Italian, 11 per cent French Canadian, and only 2 per cent German; 33 per cent were foreign-born whites. An examination of the table shows that the Southern States, Texas, Oklahoma, Mississippi, Tennessee, and Arkansas stand

at the head of the list, while the States of the Northeast, especially those engaged in manufacturing, lie at the bottom of the list (Rhode Island, Connecticut, Pennsylvania, New York, Massachusetts, and New Jersey). The high stature of the men of the Southern States is due, as indicated, in part to the absence of recent immigration from southeastern Europe, and also in part to the average tall stature of Negroes. The short stature of the population of the manufacturing and maritime States of the Northeast is due in part to the presence in them of members of the shortest European races. In the upper half of the table one finds also States like Kansas, Idaho, Oregon, Nebraska, South Dakota, Iowa, and Minnesota, which are populated largely by Nordics.

TABLE 13.—*Mean stature by States, first million draft recruits; States arranged in order of standing with proportional weight and chest circumference at (expiration) for each inch of stature.*

State.	Number of men measured.	Mean height.		Mean weight.	Mean chest.
		Inches.	Centimeters.	Pounds.	Inches.
Texas.....	34,531	68.40	173.74	2.079	0.483
Oklahoma.....	19,429	68.28	173.43	2.084	.485
Mississippi.....	8,543	68.27	173.41	2.10	.486
Tennessee.....	14,426	68.27	173.41	2.052	.483
Arkansas.....	10,111	68.20	173.23	2.071	.486
Kansas.....	9,571	68.20	173.23	2.107	.487
Alaska.....	106	68.15	173.10	2.208	.493
Colorado.....	6,635	68.15	173.10	2.069	.485
North Carolina.....	14,668	68.15	173.10	2.076	.487
Arizona.....	3,850	68.13	173.05	2.099	.488
Idaho.....	4,031	68.10	172.97	2.133	.496
Oregon.....	2,748	68.09	172.95	2.150	.492
Nebraska.....	10,774	68.08	172.92	2.126	.488
South Dakota.....	3,892	68.05	172.85	2.159	.493
Iowa.....	19,537	68.04	172.82	2.126	.491
Minnesota.....	27,341	68.04	172.82	2.15	.494
Kentucky.....	15,502	68.02	172.77	2.058	.484
Alabama.....	15,988	68.01	172.75	2.077	.485
Montana.....	11,648	68.01	172.75	2.151	.492
Georgia.....	20,305	67.99	172.69	2.071	.488
Washington.....	13,316	67.96	172.62	2.140	.492
Missouri.....	24,964	67.95	172.59	2.081	.486
North Dakota.....	6,444	67.92	172.52	2.163	.497
West Virginia.....	12,367	67.87	172.39	2.086	.490
Utah.....	4,568	67.85	172.34	2.109	.488
Nevada.....	1,441	67.83	172.29	2.143	.497
Virginia.....	17,616	67.80	172.21	2.070	.489
Wyoming.....	1,927	67.79	172.19	2.13	.492
Indiana.....	23,194	67.75	172.09	2.090	.489
California.....	35,461	67.67	171.88	2.127	.493
South Carolina.....	9,343	67.64	171.81	2.077	.489
District of Columbia.....	4,486	67.63	171.78	2.077	.482
Louisiana.....	12,356	67.60	171.70	2.065	.489
Wisconsin.....	18,433	67.60	171.70	2.137	.496
Florida.....	5,895	67.58	171.65	2.061	.489
New Mexico.....	2,690	67.50	171.45	2.051	.491
Illinois.....	69,491	67.40	171.20	2.103	.493
Ohio.....	52,814	67.38	171.15	2.098	.491
Maine.....	3,315	67.28	170.89	2.10	.497
Michigan.....	41,872	67.23	170.76	2.11	.496
Delaware.....	1,891	67.19	170.66	2.075	.492
Vermont.....	2,077	67.12	170.48	2.091	.498
Maryland.....	9,192	67.08	170.38	2.09	.494
New Hampshire.....	2,240	66.97	170.10	2.085	.495
New Jersey.....	29,958	66.77	169.60	2.079	.498
Massachusetts.....	29,534	66.76	169.57	2.07	.496
New York.....	87,818	66.72	169.47	2.091	.497
Pennsylvania.....	77,186	66.72	169.47	2.094	.496
Connecticut.....	13,585	66.71	169.44	2.095	.501
Rhode Island.....	3,928	66.40	168.66	2.06	.494

(b) *Demobilized men.*—Table 14 gives the distribution of mean stature of men at demobilization, by States. In this table the States are arranged in order of mean stature of men, the States with the tallest men being placed at the top of the table.

TABLE 14.—Mean stature, by States, of soldiers at demobilization (1919).

State.	Number of men measured.	Mean stature.	
		Inches.	Centimeters.
United States.....	102,304	67.72	172.09
Alaska.....	13	69.43	176.35
Mississippi.....	2,099	68.61	174.28
Tennessee.....	2,807	68.61	174.28
Texas.....	4,361	68.60	174.24
Alabama.....	1,930	68.57	174.16
Georgia.....	3,397	68.51	174.01
Oklahoma.....	2,310	68.44	173.84
Nebraska.....	819	68.44	173.84
Kansas.....	1,012	68.43	173.82
Arkansas.....	2,576	68.41	173.76
South Dakota.....	416	68.39	173.70
Oregon.....	1,069	68.38	173.68
Washington.....	2,025	68.38	173.67
Montana.....	264	68.35	173.60
Arizona.....	130	68.33	173.55
South Carolina.....	828	68.32	173.54
Minnesota.....	1,950	68.31	173.51
Iowa.....	1,609	68.28	173.42
Idaho.....	161	68.26	173.39
Florida.....	1,022	68.22	173.28
North Carolina.....	1,815	68.22	173.27
West Virginia.....	1,686	68.20	173.24
Utah.....	104	68.19	173.21
Wyoming.....	80	68.16	173.13
Kentucky.....	2,921	68.13	173.05
Colorado.....	225	68.12	173.02
Virginia.....	1,920	68.01	172.75
Missouri.....	2,836	67.98	172.66
North Dakota.....	358	67.96	172.61
Nevada.....	18	67.91	172.50
California.....	481	67.91	172.49
Louisiana.....	2,070	67.86	172.36
New Mexico.....	229	67.82	172.27
Wisconsin.....	2,675	67.79	172.18
Indiana.....	3,994	67.73	172.08
Illinois.....	6,687	67.65	171.83
District of Columbia.....	231	67.60	171.70
Ohio.....	7,076	67.48	171.39
Michigan.....	3,715	67.32	170.99
Delaware.....	300	67.26	170.83
Maryland.....	1,138	67.20	170.70
Vermont.....	446	67.19	170.67
Maine.....	693	67.17	170.60
Connecticut.....	996	67.08	170.38
Pennsylvania.....	10,874	67.01	170.21
New Jersey.....	3,180	66.93	169.99
New York.....	9,207	66.92	169.98
New Hampshire.....	413	66.80	169.67
Massachusetts.....	4,782	66.77	169.60
Rhode Island.....	403	66.54	169.00

TABLE 15.—Increase in stature of soldiers at demobilization over stature of recruits, 1917-1919 (inches).

State.	Increase (inches).	State.	Increase (inches).
United States.....	0.23	Virginia.....	0.21
Alaska.....	1.28	New York.....	.20
South Carolina.....	.68	Arizona.....	.20
Florida.....	.64	Texas.....	.20
Alabama.....	.56	Wisconsin.....	.19
Georgia.....	.52	Idaho.....	.16
Washington.....	.41	New Jersey.....	.16
Connecticut.....	.37	Oklahoma.....	.16
Wyoming.....	.37	Rhode Island.....	.14
Nebraska.....	.36	Maryland.....	.12
Tennessee.....	.34	Kentucky.....	.11
Utah.....	.34	Ohio.....	.10
Mississippi.....	.34	Michigan.....	.09
Montana.....	.34	Nevada.....	.08
South Dakota.....	.34	North Carolina.....	.07
West Virginia.....	.33	Vermont.....	.07
New Mexico.....	.32	Delaware.....	.07
Oregon.....	.29	North Dakota.....	.04
Pennsylvania.....	.29	Missouri.....	.03
Minnesota.....	.27	Massachusetts.....	.01
Louisiana.....	.26	Indiana.....	-.02
Illinois.....	.25	Colorado.....	-.03
California.....	.24	District of Columbia.....	-.03
Iowa.....	.24	Maine.....	-.11
Kansas.....	.23	New Hampshire.....	-.17
Arkansas.....	.21		

8. COMPARISON OF STATURE OF RECRUITS AND VETERANS, BY STATES.

A comparison of Tables 13 and 14 and reference to Table 15 bring out many interesting differences in the stature of recruits and veterans. The increase in stature for the troops measured for the United States as a whole is about 0.23 inch. The State that showed the greatest increase in stature at demobilization as compared with mobilization is Alaska. The increase amounts to about 1.28 inches, but since this difference is based on only 13 men measured at demobilization, little stress is to be laid on it. The next on the list are the four Southern States of South Carolina, Florida, Alabama, and Georgia, in which the increase is from 0.68 to 0.52 inch. From these States there came many Negroes and also many white men of exceptionally tall stature. The end result of increase in stature is probably due to a combination of circumstances. Many of the Negroes assume a lax posture which the Army training would do much to correct and straighten. Similarly, many of the tall Southerners, as is well known, early acquire a stoop. Probably the mean for the recruits at induction was lowered to a certain extent by the inclusion of the measurement of some men subsequently rejected by the camp boards for underweight, defective physical development, etc. Finally, the men have acquired between one and two years additional age and, in the case of the younger troops who are still growing, this would mean an addition in stature, and this addition would be absolutely the greatest in the case of the tallest population, and this tallest population comes from just those Southern States. In the Southern States there are found in the upper half of the table the States which have acquired an increase of 0.25 inch or over, West Virginia, Tennessee, Mississippi, and Louisiana. Only the Southern States of Arkansas, Virginia, Texas, Kentucky, and North Carolina show an increase of less than 0.25 inch.

The increase of stature affected different States differently, so that the order in which they stand is changed in the two periods. Thus, Mississippi, which stood third in stature of recruits, is second in the stature of demobilized troops. Tennessee and Texas changed places. Alabama and Georgia are placed relatively much higher in the order of States at demobilization than at mobilization. On the other hand, farmers from Kansas increased only slightly in stature and consequently stand relatively low in the demobilized list.

In general, the Southern States show greater improvement in stature than the Northern States, and, as indicated above, there was greater room for improvement. Part of the improvement is doubtless to be attributed to the greatly bettered sanitation in the Army over that which they experienced at home. With the elimination of the hookworm infections and the "straightening up" resulting from the setting-up exercises of military drill, muscular weakness was relieved and the back strengthened. Consequently, 1 centimeter or more was added to the stature.

Among Northern States which showed a considerable increase in stature are: Washington, 0.42 inch; Connecticut, 0.37; Nebraska, 0.36; and Utah, 0.34. The States of the Northwest for the most part lie in the upper part of the table, and this is because they contain so many tall men who showed the greatest absolute increment in stature even if they are not proportionately increased over the shorter men.

While Rhode Island retains her position at the bottom of the list, her men made greater improvement in stature than those from some other States. At the bottom of the table of increase stand New Hampshire, the District of Columbia, and Indiana, in which there has been an average decrease in height at demobilization. Why there should have been a decrease of 0.17 inch in the case of New Hampshire troops is hard to say. Perhaps it is because the number of men examined is only 94 and the diminution is due to the accident of small numbers. Men from the District of Columbia remained practically unchanged in stature and this is probably because the District is a city made up, so far as white population goes, of men who are used to holding themselves well, assuming a good posture, for it is well known that the standing posture of men in cities is, on the whole, superior to that of rural districts. Similarly, the men of Massachusetts (largely urban in its population) have changed little in stature. In the lower half of the table, showing an increase of less than 0.20 inch, lie certain States of the Central West, such as Indiana, Missouri, North Dakota, Michigan, Ohio, and Oklahoma; also certain Eastern States, such as Delaware, Vermont, Maryland, Rhode Island, and New Jersey, States for the most part not marked by extremely tall stature, in which, therefore, any increase in size with age will be less marked than in the case of States containing tall men.

9. COMPARISON OF STATURE OF RECRUITS FROM THE VARIOUS STATES, 1863-1864 AND 1917-1918.

A natural inquiry is: How does the stature of draft recruits of 1917-1918 compare with that of recruits of the Civil War, 1861-1864, 55 years earlier? The mean stature of 1,104,841 white volunteer recruits in the first years of the Civil War was, according to Gould² (p. 105), 67.64; for 501,068 draft recruits (Baxter,¹ Vol. I, p. 23) it was 67.30.^a The weighted average for the two groups was 67.502. To conclude that the average of our male population has diminished 0.15 inch, has increased 0.19 inch, or has remained practically stationary with a decrease of only 0.01 inch, would probably not be justified, for the population measured in 1861 is not strictly comparable with that measured in 1917-1918. For, first, the population of the Civil War recruits largely excluded the Southern States, which were in secession, while that of the World War included them. It is these Southern States that in 1917-1918 showed the tallest average stature; and the inclusion in the later data (and not in former) of several States above the average probably tends unduly to raise the 1917-1918 mean stature as compared with that of the Civil War. Second, in the Civil War there was a larger percentage of men below the ages of 21 and 24 than in the World War. In the Civil War 292 per 1,000 were below the age of 21 and 519.56 were below the age of 24, while in the World War only 95.94 were below the age of 21 and 433.56 below the age of 24.^b Since many men under 21 have not reached their full stature and some not even until the age of 24, the exclusion of a number of men of the younger ages tends to raise the average for the World War.

A more just basis of comparison of mean stature in the two epochs is that between individual States. Table 16 has been drawn up from Gould's Table I,

^a The "Draft recruits," considered by Baxter here, as well as elsewhere, include also draft substitutes and late volunteers, all raised during the "Draft" period.

^b However, the present statistics deal only with men of ages 21 to 30, inclusive. The younger and older men included in the age compilation (see Table 2, Gould,² pp. 69, 34, and 57) were volunteers, officers, and enlisted men.

Chapter V, page 94. It should be recalled that this table includes only volunteers from the unculled part of the northern population during the first years of the Civil War. This table reveals a certain measure of stability in the order of average male statures in the different States, even during the course of half a century. In both the earlier and the later series Iowa, Kentucky, Missouri, and West Virginia stand near the top of the list (of these Northern States) and Connecticut and Rhode Island at the very bottom. New York, Pennsylvania, and Massachusetts stand low in both series—the effect of the immigration of South Irish and South German stock was already evident in 1861–1864. On examining the different columns it appears that there is an increase in mean stature in Minnesota, due to recent Scandinavian immigration thither; Wisconsin shows little change in mean stature because the increase of Scandinavians has counterbalanced the effect of the shorter immigrants. New Jersey's increase is probably largely due to its large commuting population, the overflow of the best of the metropolis which has attracted great numbers of men of exceptionally fine physique. The following States show a decrease: Illinois, 0.57 inch; Ohio, 0.46 inch; Rhode Island and Connecticut about 0.69 and 0.38, respectively; New York, 0.37; Indiana, 0.31; Michigan, 0.31; Massachusetts, 0.29. These are the States which have received most of the recent immigration of the Mediterraneans, Polish Jews, and Balkanese. The great reductions in Maine, New Hampshire, and Vermont are due chiefly to the immigration of the French Canadians into these States.

It is reasonable to suppose that, since this country has received a very large number of immigrants of prevaillingly low statures from southern Europe during the last 50 years, the average stature of the population of the country should show a decrease. Such is, however, very difficult to demonstrate mathematically, since the methods used in the recruiting of the two armies at the two periods differed so materially.

TABLE 16.—*Comparison of stature (in inches) of native and foreign born white and colored draft recruits, United States, 1917–1918, and white recruits of the Civil War (Gould,² Table I, Chap. V), by States in order of 1917–1918 average statures (Louisiana omitted on account of scanty data in Gould's Table).*

State.	Stature (inches).		Difference.
	1917–1918	1861–1864	
Minnesota.....	68.04	67.63	+0.41
Iowa.....	68.04	68.13	— .09
Kentucky.....	68.02	68.16	— .14
Missouri.....	67.95	68.03	— .08
West Virginia.....	67.87	68.43	— .56
Indiana.....	67.75	68.06	— .31
Wisconsin.....	67.60	67.65	— .05
Illinois.....	67.40	67.97	— .57
Ohio.....	67.38	67.84	— .46
Maine.....	67.28	68.12	— .84
Michigan.....	67.23	67.62	— .39
Vermont.....	67.12	67.61	— .49
Maryland.....	67.08	67.31	— .23
New Hampshire.....	66.97	67.40	— .43
New Jersey.....	66.77	66.58	+ .19
Massachusetts.....	66.76	67.05	— .29
New York.....	66.72	67.09	— .37
Pennsylvania.....	66.72	67.14	— .42
Connecticut.....	66.71	a 67.09	— .38
Rhode Island.....	66.40	a 67.09	— .69

a Data for Rhode Island and Connecticut consolidated, 67.09.

TABLE 17.—*Characteristics and composition of the popu-*

State No.	Designation of section.	Characteristics.	Total population.	Density per square mile.	Cities of 25,000 or over.	Per cent urban.
1	Alabama 1.....	Mining and manufacturing area.	760,740	49.0	Birmingham.....	26.9
	Alabama 2.....	Large Negro population.....	563,441	44.0	Montgomery.....	14.8
	Alabama 3.....	Large native white population..	577,827	35.0		5.3
	Alabama 4.....	Large Negro population.....	127,817	31.0		
	Alabama 5.....	Urban and suburban areas.....	85,308	41.0	Mobile.....	54.0
2	Arizona 1.....	Large Indian population, sparsely settled.	57,653	.9		12.2
	Arizona 2.....	Chiefly white population.....	146,371	2.9		38.4
3	Arkansas 1.....	Negro, Mississippi bottoms.....	641,940	38.0	Little Rock.....	17.6
	Arkansas 2.....	Large native white population, hill country.	212,005	19.0		3.4
	Arkansas 3.....	Large native white population..	720,504	30.0		11.3
4	California 1.....	Chiefly agricultural area.....	1,433,865	16.1	(Oakland..... Sacramento.....)	46.9
	California 2.....	Mining area.....	63,226	4.4		10.5
	California 3.....	Sparsely populated.....	114,318	2.5		43.9
	California 4.....	Urban area.....	319,198		Los Angeles.....	100.0
	California 5.....	do.....	416,912	9,689.0	San Francisco.....	100.0
5	Colorado 1.....	Large native white population..	108,622	3.4		18.1
	Colorado 2.....	Russian population.....	89,813	8.0		28.8
	Colorado 3.....	English population.....	78,716	10.0		29.4
	Colorado 4.....	Prevaillingly agricultural.....	139,574	5.0	Colorado Springs.....	33.5
	Colorado 5.....	Urban population.....	213,381	3,679.0	Denver.....	100.0
	Colorado 6.....	Austrian and Italian population.	159,918	8.0	Pueblo.....	46.5
6	Connecticut 1.....	Prevaillingly agricultural and near metropolitan.	400,100	114.7	Norwich.....	78.2
	Connecticut 2.....	Manufacturing area.....	714,656	536.5	(New Haven..... Bridgeport.....)	96.1
7	Delaware.....	State undivided.....	202,322	103.0	Wilmington.....	48.0
8	District of Columbia.	District undivided.....	331,089	5,518.0	Washington.....	100.0
9	Florida 1.....	More white and maritime.....	248,836	18.7	Jacksonville.....	35.0
	Florida 2.....	More Negro and rural population	220,302	21.0		14.1
	Florida 3.....	(Cuban, Spanish, West Indian population.)	21,563	19.0		92.5
	Florida 4.....	Peninsular.....	261,918	8.7	Tampa.....	30.9
10	Georgia 1.....	Mixed population, native white predominating.	1,334,222	43.0	Atlanta.....	19.4
	Georgia 2.....	Large Negro population.....	1,274,899	45.0	(Savannah..... Augusta.....)	21.9
11	Idaho.....	State undivided.....	325,594	3.9		21.5
12	Illinois 1.....	Densely populated.....	434,972	192.5	(Joliet..... Aurora.....)	53.5
	Illinois 2.....	(Mixed native and foreign population.)	753,575	68.2	(Peoria..... Rockford.....)	43.9
	Illinois 3.....	Agricultural area, native.....	995,129	51.0	(Springfield..... Decatur.....)	24.3
	Illinois 4.....	Largely German population.....	344,621	80.0	East St. Louis..	45.3
	Illinois 5.....	Urban area.....	2,185,283	11,812.0	Chicago.....	100.0
	Illinois 6.....	Negro population (Egypt).....	52,591	80.0		41.9
	Illinois 7.....	Agricultural area.....	805,587	49.0	Bloomington.....	31.1
	Illinois 8.....	Agriculture and manufacturing area.	266,833	45.3		28.2
13	Indiana 1.....	Manufacturing.....	282,521	117.0	South Bend.....	69.3
	Indiana 2.....	Agricultural, considerable German.	128,679	37.0		18.6
	Indiana 3.....	Agricultural area, native stock..	2,299,676	76.0	(Indianapolis.. Evansville.....)	40.4
14	Iowa 1.....	(Foreign white, German and Scandinavian.)	1,442,410	38.0	(Sioux City..... Davenport.....)	29.6

* Indian.

* Chinese.

* Japanese.

* Russian.

lation of the various sections of the United States.

Native white.		Foreign-born white.					Austrian and Russian.	Scandinavian.	Italian.	Canadian, French.				
Native part-entage.	Foreign part-entage.													
71.5	2.2	1.1	25.6											
28.5	.6	.3	70.6											
67.6	.6	.6	31.0											
28.9	.2	.1	72.8											
46.3	7.5	3.2	42.8		1.7	1.3								
34.2	13.1	15.4	.6	30.6	1.5	2.2								
42.8	23.6	25.9	1.1	6.6	2.2	1.9	2.5			1.2				
41.7	2.0	.9	55.3											
96.9	1.4	.5	.7											
83.9	2.9	1.3	12.0											
49.6	25.2	20.0	.6	1.7	5.7	3.7	2.8		2.6	3.8				
				11.5										
47.2	27.3	19.9	.2	12.1										
				12.6										
				1.6	4.3	4.0	6.0	1.1	1.5	6.9				
				11.0				2.0						
57.6	17.5	17.8	1.1	13.6	3.4	1.7	2.5							
				1.6										
				1.7										
33.2	23.4	19.0	2.4	1.6	5.9	2.8	3.5	2.2	1.7	1.9				
				1.3										
27.7	36.9	31.4	.4	2.5	11.7	12.1	3.5	1.5	4.4	6.4				
				1.1				1.6						
73.9	15.7	8.6	.4	1.0	8.5	1.6	1.1	1.0	1.2	1.2				
64.3	19.8	14.5	.2	1.6	4.9	1.2	1.8	8.3	3.3					
54.3	27.1	17.6	.7	5.3	2.8	6.1	6.1	1.9	4.8	2.0				
								1.2						
69.5	18.2	10.7	1.4	4.8	1.9	1.8	1.8	1.2	2.2	1.1				
								1.7						
50.1	28.7	18.2	2.5	7.6	4.8	3.2	3.2	1.3	3.7	2.3				
								4.4						
52.4	22.9	22.4	1.9	3.6	2.9	2.8	2.8	8.0	2.0	8.6				
44.3	29.5	24.9	1.2	5.1	10.5	3.0	3.0	2.9	2.5	5.1	5.5			1.0
								4.5						1.0
30.6	35.9	32.0	1.4	6.4	15.2	3.4	3.4	3.8	3.2	9.6	1.8	2.1		
								9.0						
63.2	12.8	8.6	15.4	3.0	6.2	1.3	1.3	2.6	2.1					
50.4	13.6	7.4	28.5	4.0	4.2	1.2	1.2	1.7	1.2					
54.2	3.3	2.0	40.5						2					
40.5	1.0	.9	57.8											
16.8	24.3	27.1			1.0	2.5								
55.9	7.1	8.4	28.5											
68.6	.9	.6	29.9											
37.3	1.1	.6	61.0											
62.6	23.1	12.4	.2	1.1	3.6	1.4	3.4	3.3	5.7		1.9			
				1.3										
				1.4										
34.6	38.2	23.9	1.0	21.2	3.5	2.8	2.8	4.3	6.2	2.2	1.3	1.0		
								3.4						
52.3	29.8	16.9	1.0	10.9	3.5	2.4	2.4	1.3	7.9	2.2				1.0
								1.4						
83.2	10.6	4.4	1.6	4.3	1.3	1.1	1.1							
52.9	29.6	13.5	3.8	17.4	2.1	1.6	1.6	2.8				1.1		
								1.5						
20.4	41.8	35.8	2.0	19.5	7.5	2.0	2.0	8.3	6.9	3.3	1.5	1.7		
60.4	7.4	2.1	30.5	3.5	1.0									
71.0	20.2	8.2	1.0	10.2	2.6	1.4	1.4		1.0					
54.1	31.5	14.2	.2	15.2	4.1	2.4	2.4		5.4					
50.6	27.0	21.8	.6	17.2	1.5			4.1	2.4	1.0		5.3		
								2.7						
76.2	16.8	6.4	.4	8.1	2.0									
82.5	11.0	3.0	2.5	5.6	1.3									
50.7	34.3	14.8	.2	15.9	2.5	1.5	1.5	1.9	8.0					

*Austrian.

*Cuban.

*West Indian.

TABLE 17.—*Characteristics and composition of the population*

State No.	Designation of section.	Characteristics.	Total population.	Density per square mile.	Cities of 25,000 or over.	Per cent urban.
14	Iowa 2.....	Native White.....	782,361	44.0	Des Moines.....	32.3
15	Kansas 1.....	Russian population.....	196,998	12.0	16.3
	Kansas 2.....	Native and German population.....	1,491,951	23.0	{Kansas City... Wichita.....}	30.9
16	Kentucky 1.....	Mountainous area, native white.....	599,797	44.0	4.5
	Kentucky 2.....	Agricultural area.....	1,720,108	63.0	{Louisville..... Covington.....}	30.8
17	Louisiana 1.....	Mississippi bottoms and upland, large Negro population.....	599,548	36.8	Shreveport.....	13.3
	Louisiana 2.....	Urban area.....	339,075	1,695.0	New Orleans.....	100.0
	Louisiana 3.....	Rural, chiefly white population.....	717,765	24.8	10.8
18	Maine 1.....	English Canadian.....	222,741	13.0	41.2
	Maine 2.....	Native white stock, maritime.....	124,720	37.0	28.7
	Maine 3.....	French Canadian population.....	394,901	37.0	{Portland..... Lewiston.....}	64.2
19	Maryland 1.....	Urban area.....	680,834	1,001.0	Baltimore.....	82.0
	Maryland 2.....	Peninsular area.....	176,412	65.0	12.7
	Maryland 3.....	Large white population.....	400,364	77.0	19.3
	Maryland 4.....	Large Negro population.....	43,741	41.0
20	Massachusetts 1.....	Mountainous area.....	148,850	89.0	Pittsfield.....	67.3
	Massachusetts 2.....	Manufacturing center.....	2,306,884	454.0	{Worcester..... Fall River.....}	93.3
	Massachusetts 3.....	Peninsular region.....	179,345	144.0	Brockton.....	73.7
	Massachusetts 4.....	Urban area.....	731,888	14,341.0	{Boston..... Chelsea.....}	100.0
21	Michigan 1.....	Finnish population.....	206,943	21.0	40.3
	Michigan 2.....	{Prevalingly native white popu- lation.....}	1,158,767	34.0	{Grand Rapids..... Kalamazoo.....}	33.6
	Michigan 3.....	Foreign population.....	613,048	65.9	{Bay..... Saginaw.....}	33.6
	Michigan 4.....	Urban area.....	465,766	Detroit.....	100.0
	Michigan 5.....	Dutch and other foreign popu- lation.....	256,078	65.6	27.2
22	Minnesota 1.....	Scandinavian population.....	558,953	12.0	10.6
	Minnesota 2.....	German and Scandinavian popu- lation.....	752,212	31.0	18.7
	Minnesota 3.....	Scandinavians and Finns.....	207,388	15.0	Duluth.....	61.7
	Minnesota 4.....	Urban area, "Twin Cities".....	557,155	766.0	{Minneapolis..... St. Paul.....}	93.8
23	Mississippi 1.....	Rural area, large Negro popu- lation.....	1,029,399	45.0	10.7
	Mississippi 2.....	Rural area, large native white population.....	714,715	32.0	12.7
24	Missouri 1.....	Native white, agricultural.....	1,636,845	41.0	{Kansas City.. St. Joseph.....}	30.0
	Missouri 2.....	Mississippi bottoms, consider- able Negro population.....	510,181	38.0	24.2
	Missouri 3.....	Native white, Ozark region.....	159,280	24.0	4.4
	Missouri 4.....	Urban area.....	687,029	11,263.0	St. Louis.....	100.0
25	Montana 1.....	Mining area, foreign population.....	228,098	5.6	Butte.....	49.6
	Montana 2.....	Sparsely settled, mountainous area.....	150,955	1.4	14.4
26	Nebraska 1.....	German and Irish, foreign stocks.....	776,717	13.0	{Omaha..... Lincoln.....}	32.6
	Nebraska 2.....	{German, Austrian, and Russian stocks.....}	413,497	23.0	13.9
27	Nevada 1.....	State undivided, sparse popu- lation.....	81,875	.7	16.3
28	New Hampshire 1.....	Mountainous area.....	88,721	19.0	35.5
	New Hampshire 2.....	Manufacturing area.....	341,851	75.0	{Manchester..... Nashua.....}	65.4
29	New Jersey 1.....	Densely populated.....	1,514,588	2,145.0	{Newark..... Jersey City.....}	89.6
	New Jersey 2.....	Plains section, rural.....	733,624	177.6	{Trenton..... Camden.....}	56.0
	New Jersey 3.....	{Mountainous area plus Atlantic County.....}	288,955	107.9	Atlantic City.....	68.1

* Austrian.

* Russian.

* Japanese

of the various sections of the United States—Continued.

Native white.		Foreign-born white.	Negro.	Indian, Chinese, Japanese.	German.	Irish.	English.	Austrian and Russian.	Scandinavian.	Italian.	Canadian, French.	Canadian, other.	Hungarian.	Mexican.	Scotch.
Native percentage.	Foreign percentage.														
73.1	17.7	7.6	1.4	4.9	2.1	1.6	2.9
60.3	25.7	12.7	1.3	6.1	1.0	{1.8 13.1}	3.3
72.9	16.1	7.4	3.4	* 2	5.4	1.5	1.3	1.2	1.5
96.4	.7	.3	2.5
76.4	6.9	2.2	14.4	3.8	1.2
31.8	3.1	2.0	63.0	2.2
43.5	21.9	8.2	26.3	6.5	3.1	4.8
61.0	2.7	1.7	34.4	1.3
59.3	23.9	16.3	.2	1.9	1.0	5.0	15.3
86.1	7.8	5.8	.2	1.29	2.9
64.7	18.2	16.9	.2	3.5	1.6	* 1.3	12.5	5.1
49.6	23.1	12.9	14.3	13.3	3.4	{1.7 6.4}	1.1
65.6	1.7	1.1	31.6
73.4	7.9	3.4	14.8	2.7	1.1
50.0	1.3	.8	47.8
46.7	30.7	21.7	.8	4.1	10.4	2.6	{3.4 3.3}	3.2	7.6	1.4	1.3
23.3	34.7	31.2	.8	1.6	14.9	4.7	{1.9 3.8}	2.0	2.8	9.7	6.1	1.4
51.6	25.2	20.9	2.0	1.0	9.6	2.3	* 3.4	3.1	3.3	2.7	5.0
23.9	38.2	35.8	1.9	2.7	22.0	2.9	* 10.2	1.5	7.1	9.7	1.1
11.6	48.1	39.8	.1	* 3	5.2	2.8	8.5	{5.5 2.4}	23.1	4.4	6.6	3.2	1.1
55.6	29.4	14.5	.4	* 4	7.2	1.5	2.1	2.5	2.1	6.9
42.9	37.5	19.0	.5	16.3	2.1	2.5	* 1.5	1.8	10.7
24.7	40.4	33.6	1.2	24.5	3.1	2.8	{4.3 5.6}	1.7	1.4	10.1	1.5
51.9	31.3	16.0	.6	* 1	8.7	1.4	1.6	* 1.2	2.4	2.5
23.3	49.0	26.2	* 1.2	10.3	1.3	* 1.4	37.4	1.3	2.1
31.9	47.8	20.1	22.3	2.6	* 2.9	16.8	1.1
15.7	38.3	44.9	.8	5.5	1.8	1.9	{8.6 2.7}	31.1	2.9	5.4
20.9	40.8	27.2	1.0	12.2	4.2	1.5	* 3.3 2.7	22.0	1.2	2.4
27.3	.9	.5	71.2
64.5	.3	.6	33.4
81.4	10.9	4.4	3.2	4.7	1.2
76.6	10.4	3.1	9.9	5.8
94.4	3.9	1.4	.3	1.5
39.3	35.9	18.3	6.4	20.0	6.0	1.3	{2.5 3.5}	1.6	1.4
37.5	31.4	28.5	.6	* 1.9	5.5	8.1	5.0	* 4.5	7.7	2.4	4.1	1.2
51.5	23.9	18.1	.3	* 6.1	5.2	2.6	2.2	* 1.5	5.9	1.6	3.7	1.5
54.3	29.3	15.0	.9	* 5	12.2	2.5	1.4	{3.9 1.9}	6.4
52.9	39.5	14.3	.1	* 1	13.5	1.2	{5.5 2.2}	6.5
33.1	25.6	22.0	.6	{8.4 1.1 1.1}	4.9	5.4	4.0	* 1.2	2.6	4.6	2.5
60.8	21.6	17.4	1.6	1.0	12.9	9.6
51.6	24.5	23.7	.1	1.0	6.9	1.9	* 1.5	17.3	4.7
28.7	37.5	31.5	2.2	14.0	10.0	3.5	{4.4 7.1}	8.9	2.2	1.4
54.7	21.7	18.1	5.6	6.6	5.6	2.7	{2.1 4.4}	4.4	3.4
60.4	17.6	16.7	5.2	4.4	4.9	2.4	{1.4 2.0}	6.7	3.2

* Indian.

* Chinese.

TABLE 17.—*Characteristics and composition of the population*

State No.	Designation of section.	Characteristics.	Total population.	Density per square mile.	Cities of 25,000 or over.	Per cent urban.
80	New Mexico 1.....	Indian population.....	59,970	2.0
	New Mexico 2.....	Native white population.....	212,657	3.0	18.5
	New Mexico 3.....	Noteworthy Mexican element.....	54,614	1.7	13.1
31	New York 1.....	Suburban territory.....	565,449	210.0	Yonkers..... Mount Vernon.....	57.9
	New York 2.....	Urban area, densely populated.....	4,766,883	16,667.0	New York City.....	100.0
	New York 3.....	Eastern manufacturing region.....	658,978	85.0	Albany..... Schenectady.....	56.3
	New York 4.....	Western manufacturing region.....	1,361,257	141.0	Rochester..... Syracuse.....	61.7
	New York 5.....	Mountainous Catskill region.....	284,857	101.0	Newburgh..... Kingston.....	39.9
	New York 6.....	Urban area.....	423,715	Buffalo.....	100.0
	New York 7.....	Agricultural and dairying.....	774,620	62.0	Binghamton..... Elmira.....	37.7
	New York 8.....	Mountainous Adirondack area.....	277,855	25.0	26.5
32	North Carolina 1.....	Sparsely populated mountainous area.....	375,905	38.0	7.4
	North Carolina 2.....	Intermediate.....	657,162	62.0	Charlotte.....	21.7
	North Carolina 3.....	Native white of Scotch origin.....	296,425	40.0	3.8
	North Carolina 4.....	Large Negro population.....	651,669	51.0	16.1
	North Carolina 5.....	Island and peninsular area.....	55,975	19.0	0.0
	North Carolina 6.....	Remainder of State.....	133,408	29.0	Wilmington.....	19.3
33	North Dakota 1.....	Scandinavian and Canadian population.....	113,603	12.0	10.9
	North Dakota 2.....	Scandinavian population.....	262,681	8.0	12.8
	North Dakota 3.....	Russian population.....	200,772	6.0	8.6
34	Ohio 1.....	Dense foreign population.....	989,804	478.0	Cleveland..... Toledo.....	85.3
	Ohio 2.....	Intermediate.....	919,823	114.0	Youngstown..... Akron.....	51.3
	Ohio 3.....	Agricultural area.....	2,493,883	81.0	Columbus..... Dayton.....	38.2
	Ohio 4.....	Urban area.....	363,591	7,279.0	Cincinnati.....	100.0
35	Oklahoma 1.....	Marked Indian and Negro population.....	615,973	24.0	Muscooke.....	17.2
	Oklahoma 2.....	Chiefly white population.....	1,041,182	23.0	Oklahoma City.....	20.6
36	Oregon 1.....	Fairly densely populated.....	445,464	29.5	Portland.....	56.9
	Oregon 2.....	{Columbia River Valley and coastal dry plain, sparsely populated.....}	227,301	2.8	23.4
37	Pennsylvania 1.....	Urban area.....	1,549,008	11,647.0	Philadelphia.....	100.0
	Pennsylvania 2.....	Rural area, native stock.....	1,877,385	132.0	Reading..... Harrisburg.....	42.5
	Pennsylvania 3.....	Mining area.....	1,067,487	245.0	Scranton..... Wilkes-Barre.....	66.7
	Pennsylvania 4.....	Coal mining.....	357,356	118.5	33.7
	Pennsylvania 5.....	Manufacturing.....	750,892	182.0	Johnstown..... Altoona.....	37.7
	Pennsylvania 6.....	Rural area.....	892,495	74.0	Erie..... New Castle.....	40.5
	Pennsylvania 7.....	{Allegheny County plus a small rural area.....}	1,363,333	181.0	Pittsburgh..... McKeesport.....	70.4
38	Rhode Island.....	State undivided.....	542,610	508.0	Providence..... Pawtucket.....	96.7
39	South Carolina 1.....	Native white.....	300,348	77.0	16.9
	South Carolina 2.....	Large Negro population.....	638,941	50.0	Columbia.....	12.8
	South Carolina 3.....	Peninsular and rural areas.....	576,111	41.0	Charleston.....	16.1
40	South Dakota 1.....	Dry farming area.....	480,230	9.0	15.2
	South Dakota 2.....	Large Russian population.....	87,826	8.0	4.2
	South Dakota 3.....	Indian population.....	15,832	1.0
41	Tennessee 1.....	Negroes, Mississippi bottoms.....	352,510	57.5	9.5
	Tennessee 2.....	Agricultural region.....	1,148,013	51.5	Memphis..... Nashville.....	27.8
	Tennessee 3.....	Mountainous region.....	683,266	51.5	Chattanooga.....	12.9

		Negro.		Indian.		Chinese.		Japanese.	
61.1	3.8	6.2	.1	23.1	1.4			1.1	1.6
86.9	6.2	5.0	.6	11.2	1.2				1.5
61.5	19.8	15.8	.6	11.3	1.2				14.3
44.7	27.6	24.6	2.9		7.3	10.6	2.8	8.3	1.5
19.3	38.1	40.4	1.9		12.7	11.7		11.1	2.3
59.4	24.2	15.7	.7		6.6	8.2	2.4	3.7	1.8
47.8	30.9	20.6	.5		10.8	7.6	3.6	4.7	3.4
60.4	20.0	16.0	2.6		5.8	7.6	2.1	5.4	
28.2	43.2	28.0	.4		27.9	6.1	2.4	4.4	4.7
70.8		10.5	.6		4.9	5.3	1.8	2.4	2.4
62.5	24.7	12.0	.2		1.1	5.4		1.0	6.7
90.8	.5	.2	8.3						5.1
74.7	.4	.3	24.7						
00.9	.4	.2	38.1						
51.9	.3	.2	47.3						
69.6	.3	.1	29.8						
57.1	.9	.7	41.2						
21.2	47.7	28.6		2.4	6.2	1.5	1.2	2.2	24.0
31.5	43.3	24.2		1.6	8.0	1.5		1.4	30.6
27.3	41.4	20.9		1.2	8.3	1.2	7.3	20.7	13.9
33.1	37.1	28.4	1.3		15.9	4.2	3.1	78.5	
64.7	20.0	15.0	1.1		6.3	2.3	3.9	78.5	
78.7	13.7	4.8	2.8		7.1	1.6		72.8	
42.6	36.4	15.6	5.4		24.3	5.3		71.2	
72.6	2.9	1.5	13.7	10.2					
82.7	7.2	3.7	5.0	11.7	2.4			1.2	
65.5	23.1	18.5	.3	1.4	7.4	1.9	2.2	1.3	5.2
74.5	14.3	9.1	.1	1.5				1.8	1.2
37.7	32.1	24.7	5.4	1.4	3.2	1.4	1.4		2.7
79.5	9.8	7.9	2.6	1.3					1.7
42.5	32.8	23.8	.2	1.3	9.7	12.8	3.7	2.9	1.1
61.3	18.1	16.4	2.2	1.5	3.2	2.5	1.0	1.6	1.2
56.8	19.7	22.2	1.3	1.0	5.6	6.7	6.4	1.0	2.3
64.1	30.1	14.8	.6	1.1	1.6	1.7	2.3	1.1	3.5
45.6	29.3	22.3	2.7	1.4	4.5	1.5	2.0	1.4	5.5
29.4	35.9	32.8	1.8	1.3	5.4	2.7	1.8	1.4	1.7
87.8	.4	.3	31.4	1.7	10.7	5.8	2.9	1.7	
39.5	.5	.3	59.9	1.7	13.5	7.8		3.3	
34.7	1.2	.6		1.6				2.3	7.8
44.7	37.3	16.8		1.1	10.7	2.4	1.6	11.4	1.9
33.5	43.6	22.3			10.3	1.2	1.0		
8.1	1.6	2.5		187.2				7.5	
54.5	.8	.4	44.2						
74.1	2.4	1.1	22.0						
89.6	1.1	.6	9.3						

TABLE 17.—*Characteristics and composition of the population*

State No.	Designation of section.	Characteristics.	Total population.	Density per square mile.	Cities of 25,000 or over.	Per cent urban.
42	Texas 1.....	Large Mexican population.....	606,641	8.0	{San Antonio.....	33.8
					{El Paso.....	
	Texas 2.....	Sparsely settled, white.....	2,023,848	16.7	{Dallas.....	22.3
					{Houston.....	22.8
	Texas 3.....	German and Negro population..	199,787	32.5	{Austin.....	31.5
	Texas 4.....	Coastal native population.....	268,413	17.5	{Galveston.....	6.3
	Texas 5.....	Large Negro population.....	157,353	24.0		
43	Utah 1.....	Sparsely populated.....	88,753	1.3		17.2
	Utah 2.....	More densely populated.....	254,504	44.0	{Salt Lake City..	60.6
					{Ogden.....	11.4
	Utah.....	Mining area.....	30,094	3.0		47.5
44	Vermont.....	State undivided.....	255,956	39.0		
45	Virginia 1.....	Peninsular region and east shore.	324,242	130.0	{Norfolk.....	38.9
					{Portsmouth.....	27.9
	Virginia 2.....	Large Negro population.....	604,358	50.0	{Richmond.....	16.6
	Virginia 3.....	Native rural region.....	498,840	44.0	{Lynchburg.....	15.6
	Virginia 4.....	Mountain, white.....	640,172	43.0	{Roanoke.....	43.4
46	Washington 1.....	{Coastal region plus eastern coun- ties.....	436,342	14.0	{Spokane.....	68.1
	Washington 2.....	Puget Sound, foreign white.....	569,055	54.0	{Seattle.....	17.7
					{Tacoma.....	
	Washington 3.....	Mountainous area.....	136,283	6.0		
47	West Virginia 1.....	do.....	186,238	29.0	{Wheeling.....	19.6
	West Virginia.....	Agricultural region.....	1,034,881	59.0	{Huntington.....	26.4
48	Wisconsin 1.....	{Scandinavian and German popu- lation.....	496,265	24.0	{La Crosse.....	30.4
					{Oshkosh.....	90.9
	Wisconsin 2.....	German population.....	1,053,772	35.0	{Green Bay.....	45.5
	Wisconsin 3.....	Urban and foreign stock.....	433,187	1,881.0	{Milwaukee.....	29.6
	Wisconsin 4.....	Lake counties.....	350,636	84.0	{Superior.....	
					{Racine.....	
49	Wyoming.....	{State undivided, sparsely popu- lated.....	145,965	1.5		

of the various sections of the United States—Continued.

Native white.		Foreign-born white.	Negro.	Indian, Chinese, Japanese.	German.	Irish.	English.	Austrian and Russian.	Scandinavian.	Italian.	Canadian, French.	Canadian, other.	Hungarian.	Mexican.	Scotch.
Native percentage.	Foreign percentage.														
44.1	25.0	21.2	9.6	5.5	17.1
77.6	4.1	2.3	15.9	1.6
33.9	26.5	11.4	28.1	7.1	¹ 5.4	1.7
52.3	13.1	7.7	26.8	4.7	.9	¹ 1.3	1.5	1.2
37.3	7.6	4.0	51.1	30.0	¹ 3.0	1.5
53.6	31.4	11.7	.1	² 3.7	8.3	10.4	1.0	1.2
43.5	36.8	18.6	.4	² 7	2.4	1.1	13.2	10.5	1.8
44.5	33.7	20.2	² 1.6	1.6	13.2	3.4	3.7	2.0
64.4	21.1	14.0	.5	4.1	1.0	² 1.0	1.8	7.8	4.4	1.1
49.5	3.6	2.8	44.0
46.6	2.4	1.4	49.6
64.8	1.3	.7	33.2
88.0	.9	.8	10.2
57.6	22.9	17.7	.4	² 7 ² 4.3 ² 4.4 ² 3.6 ² 4.2 ² 1.8	6.2	2.0	1.9	² 1.3 ² 2.0	6.7	1.0	3.3	
44.5	27.2	25.1	.7	² 3.3 ² 4.1 ² 4.4	5.7	2.4	3.2	² 1.5 ² 1.1	13.5	1.6	5.0	1.2
59.4	20.6	15.6	.5	² 3.3 ² 4.1 ² 4.4	4.6	1.6	2.4	² 2.2 ² 1.4	3.3	1.8	3.3	1.0
86.8	3.7	4.8	4.5	² 1.1	2.4
85.7	4.9	4.6	5.4	1.6	1.6
31.8	44.2	23.1	.1	² 8	13.6	1.7	² 3.2 ² 1.3 ² 1.8 ² 1.0	22.3	1.5	2.5
38.0	43.2	18.2	.1	² 5	26.3	2.9	1.7	² 4.5 ² 4.1 ² 4.3 ² 2.5	10.2
21.6	48.3	29.8	.2	43.9	2.1	1.1	² 4.5 ² 4.1 ² 4.3 ² 2.5	1.1	1.1	1.9
31.7	45.9	21.9	.1	² 3	27.1	2.1	1.3	² 4.5 ² 4.1 ² 4.3 ² 2.5	4.6
55.3	22.3	18.6	1.5	² 1.0 ² 1.1	4.3	2.5	3.8	² 3.6	2.7	1.6	1.2	2.0
Austrian.		Indian.		Russian.		Chinese.		Japanese.							

¹ Austrian.² Indian.³ Russian.⁴ Chinese.⁵ Japanese.

TABLE 18.—*List of counties comprised in each "section."***ALABAMA.**

Section I: Blount, Cherokee, Colbert, Cullman, De Kalb, Etowah, Fayette, Franklin, Jackson, Jefferson, Lamar, Lauderdale, Lawrence, Limestone, Madison, Marion, Marshall, Morgan, Tuscaloosa, Walker, Winston.

Section II: Autauga, Barbour, Bullock, Butler, Chambers, Clarke, Dallas, Lee, Lowndes, Macon, Marengo, Monroe, Montgomery, Perry, Russell, Wilcox.

Section III: Baldwin, Bibb, Calhoun, Chilton, Clay, Cleburne, Coffee, Conecuh, Coosa, Covington, Crenshaw, Dale, Elmore, Escambia, Geneva, Henry, Houston, Pike, Randolph, St. Clair, Shelby, Talladega, Tallapoosa.

Section IV: Choctaw, Greene, Hale, Pickens, Sumter.

Section V: Mobile and Washington.

ARIZONA.

Section I: Apache, Coconino, Gila, Mohave, Navajo, Pinal.

Section II: Cochise, Graham, Greenlee, Maricopa, Pima, Santa Cruz, Yavapai, Yuma.

ARKANSAS.

Section I: Ashley, Chicot, Columbia, Crittenden, Cross, Desha, Drew, Hempstead, Jackson, Jefferson, Lafayette, Lee, Lincoln, Little River, Lonoke, Miller, Mississippi, Monroe, Ouachita, Phillips, Pulaski, St. Francis, Union, Woodruff.

Section II: Baxter, Boone, Carroll, Cleburne, Fulton, Izard, Madison, Marion, Montgomery, Newton, Polk, Searcy, Scott, Sharp, Stone, Van Buren.

Section III: Arkansas, Benton, Bradley, Calhoun, Clark, Clay, Cleveland, Conway, Craighead, Crawford, Dallas, Faulkner, Franklin, Garland, Grant, Greene, Hot Spring, Howard, Independence, Johnson, Lawrence, Logan, Nevada, Perry, Pike, Polk, Pope, Prairie, Randolph, Saline, Sebastian, Sevier, Washington, White, Yell.

CALIFORNIA.

Section I: Alameda, Butte, Colusa, Contra Costa, Del Norte, Fresno, Glenn, Humboldt, Kern, Kings, Lake, Los Angeles, Madera, Marin, Mendocino, Merced, Monterey, Napa, Orange, Sacramento, San Benito, San Diego, San Joaquin, San Luis Obispo, San Mateo, Santa Barbara, Santa Clara, Santa Cruz, Shasta, Siskiyou, Solano, Sonoma, Stanislaus, Sutter, Tehama, Trinity, Tulare, Ventura, Yolo, Yuba.

Section II: Amador, Calaveras, Eldorado, Lassen, Mariposa, Modoc, Nevada, Placer, Plumas, Sierra, Tuolumne.

Section III: Alpine, Imperial, Inyo, Mono, Riverside, San Bernardino.

Section IV: Includes city of Los Angeles.

Section V: Includes city of San Francisco.

COLORADO.

Section I: Alamosa, Archuleta, Conejos, Costilla, Delta, Garfield, Grand, Hinsdale, Jackson, La Plata, Mesa, Mineral, Moffat, Montezuma, Montrose, Rio Blanco, Rio Grande, Routt, Saguache.

Section II: Larimer, Logan, Morgan, Phillips, Sedgwick, Weld.

Section III: Boulder, Clear Creek, Eagle, Douglas, Gilpin, Jefferson, Park, Summit, Teller.

Section IV: Adams, Arapahoe, Baca, Bent, Cheyenne, Crowley, Elbert, El Paso, Kiowa, Kit Carson, Lincoln, Otero, Prowers, Washington, Yuma.

Section V: Includes city and county of Denver.

Section VI: Chaffee, Custer, Dolores, Fremont, Gunnison, Huerfano, Lake, Las Animas, Ouray, Pitkin, Pueblo, San Juan, San Miguel.

CONNECTICUT.

Section I: Fairfield, Litchfield, Middlesex, New London, Tolland, Windham.

Section II: Hartford, New Haven.

Cities not included in counties, Bridgeport and Stamford.

DELAWARE.

Section I: Includes entire State.

DISTRICT OF COLUMBIA.

Section I: Includes entire District.

FLORIDA.

Section I: Bay, Calhoun, Duval, Escambia, Franklin, Holmes, Jackson, Lafayette, Liberty, Nassau, Okaloosa, Santa Rosa, Taylor, Wakulla, Walton, Washington.

Section II: Alachua, Citrus, Columbia, Gadsden, Hamilton, Hernando, Jefferson, Leon, Levy, Madison, Marion, Putnam, Suwanee.

Section III: Includes county of Monroe.

Section IV: Baker, Bradford, Brevard, Broward, Clay, Dade, De Soto, Hillsboro, Lake, Lee, Manatee, Orange, Osceola, Palm Beach, Pasco, Pinellas, Polk, St. Johns, St. Lucie, Seminole, Sumter, Volusia.

GEORGIA.

Section I: Appling, Bacon, Banks, Barrow, Bartow, Ben Hill, Berrien, Brooks, Bullock, Campbell, Candler, Carroll, Catoosa, Charlton, Chattooga, Cherokee, Clayton, Clinch, Cobb, Coffee, Colquitt, Dade, Dawson, Dekalb, Dodge, Douglas, Echols, Effingham, Emanuel, Evans, Fannin, Fayette, Floyd, Forsyth, Franklin, Fulton, Gilmer, Gordon, Grady, Gwinnett, Hall, Habersham, Haralson, Hart, Heard, Irwin, Jackson, Jeff Davis, Johnson, Laurens, Lowndes, Madison, Lumpkin, Milton, Montgomery, Murray, Oconee, Paulding, Pickens, Pierce, Polk, Rabun, Rockdale, Stephens, Tattnall, Telfair, Thomas, Tift, Toombs, Towns, Turner, Union, Walker, Walton, Ware, Wayne, Wheeler, White, Whitfield, Wilcox, and Worth.

Section II: Baker, Baldwin, Bibb, Bleckley, Bryan, Burke, Butts, Calhoun, Camden, Chatham, Chattahoochee, Clarke, Clay, Columbia, Coweta, Crawford, Crisp, Decatur, Dooly, Dougherty, Early, Elbert, Glasscock, Glynn, Greene, Hancock, Harris, Henry, Houston, Jasper, Jefferson, Jenkins, Jones, Lee, Liberty, Lincoln, McDuffie, McIntosh, Macon, Marion, Meriwether, Miller, Mitchell, Monroe, Morgan, Muscogee, Newton, Oglethorpe, Pike, Pulaski, Putnam, Quitman, Randolph, Richmond, Schley, Screven, Spalding, Stewart, Sumter, Talbot, Taliaferro, Taylor, Terrell, Troup, Twiggs, Upson, Warren, Washington, Webster, Wilkes, and Wilkinson.

IDAHO.

Section I: Includes entire State.

ILLINOIS.

Section I: Cook (except city of Chicago), Dupage, Kane, Lake.

Cities not included in counties, Joliet.

Section II: Adams, Bureau, Fulton, Grundy, Hancock, Henderson, Henry, Kendall, Knox, La Salle, Marshall, Mercer, Peoria, Putnam, Rock Island, Stark, Warren, Will.

Cities not included in counties, Rockford.

Section III: Bond, Calhoun, Christian, Clark, Clay, Coles, Crawford, Cumberland, Douglas, Edgar, Edwards, Effingham, Fayette, Franklin, Gallatin, Greene, Hamilton, Hardin, Jackson, Jasper, Jefferson, Jersey, Johnson, Lawrence, Marlon, Montgomery, Morgan, Moultrie, Perry, Pike, Pope, Richland, Saline, Sangamon, Scott, Shelby, Union, Wabash, Wayne, White, Williamson.

Cities not included in counties, Decatur and Danville.

Section IV: Clinton, Macoupin, Madison, Monroe, Randolph, St. Clair, Washington.

Section V: Includes city of Chicago.

Section VI: Alexander, Massac, Pulaski.

Section VII: Brown, Cass, Champaign, Dewitt, Ford, Iroquois, Kankakee, Livingston, Logan, McDonough, McLean, Macon, Mason, Menard, Piatt, Schuyler, Tazewell, Vermillion, Woodford.

Section VIII: Boone, Carroll, Dekalb, Jo Daviess, Lee, McHenry, Ogle, Stephenson, Whiteside, Winnebago.

INDIANA.

Section I: Elkhart, Lake, Laporte, Porter, St. Joseph.

Section II: Benton, Jasper, Newton, Pulaski, Starke, Tippecanoe, Warren, White.

Section III: Adams, Allen, Bartholomew, Blackford, Boone, Brown, Carroll, Cass, Clark, Clay, Clinton, Crawford, Daviess, Dearborn, Decatur, Dekalb, Delaware, Dubois, Fayette, Floyd, Fountain, Franklin, Fulton, Gibson, Grant, Greene, Hamilton, Hancock, Harrison, Hendricks, Henry, Howard, Huntington, Jackson, Jay, Jefferson, Jennings, Johnson, Knox, Kosciusko, Lagrange, Lawrence, Madison, Marion, Marshall, Martin, Miami, Monroe, Montgomery, Morgan, Noble, Ohio, Orange, Owen, Parke, Perry, Pike, Posey, Putnam, Randolph, Ripley, Rush, Scott, Shelby, Spencer, Steuben, Sullivan, Switzerland, Tipton, Union, Vanderburg, Vermillion, Vigo, Wabash, Warrick, Washington, Wayne, Wells, Whitley.

IOWA.

Section I: Allamakee, Audubon, Benton, Blackhawk, Boone, Bremer, Buchanan, Buena Vista, Butler, Calhoun, Carroll, Cass, Cedar, Cerro Gordo, Cherokee, Chickasaw, Clay, Clayton, Clinton, Crawford, Delaware, Dickinson, Dubuque, Emmett, Fayette, Floyd, Franklin, Grundy, Hamilton, Hancock, Hardin, Harrison, Howard, Humboldt, Ida, Iowa, Jackson, Johnson, Jones, Linn, Lyon, Marshall, Mitchell, Monona, Muscatine, O'Brien, Osceola, Palo Alto, Plymouth, Pocahontas, Pottawattamie, Sac, Scott, Shelby, Sioux, Story, Tama, Webster, Winnebago, Winneshiek, Woodbury, Worth, Wright.

Section II: Adair, Adams, Appanoose, Clarke, Dallas, Davis, Decatur, Des Moines, Fremont, Greene, Guthrie, Henry, Jasper, Jefferson, Keokuk, Lee, Louisa, Lucas, Madison, Mahaska, Marion, Mills, Monroe, Montgomery, Page, Polk, Poweshiek, Ringgold, Taylor, Union, Van Buren, Wapello, Warren, Washington, Wayne.

KANSAS.

Section I: Barton, Ellis, Gove, Greeley, Hamilton, Harvey, Kearny, Logan, McPherson, Marlon, Ness, Reno, Rice, Rush, Russell, Trego, Wallace, Wichita.

Section II: Allen, Anderson, Atchison, Barber, Bourbon, Brown, Butler, Chase, Chautauqua, Cherokee, Cheyenne, Clark, Clay, Cloud, Coffey, Comanche, Cowley, Crawford, Decatur, Dickinson, Doniphan, Douglas, Edwards, Elk, Ellsworth, Finney, Ford, Franklin, Geary, Graham, Grant, Gray, Greenwood, Harper, Haskell, Hodgeman, Jackson, Jefferson, Jewell, Johnson, Kingman, Kiowa, Labette, Lane, Leavenworth, Lincoln, Linn, Lyon, Marshall, Meade, Miami, Mitchell, Montgomery, Morris, Morton, Nemaha, Neosho, Norton, Osage, Osborne, Ottawa, Pawnee, Phillips, Pottawatomie, Pratt, Rawlins, Republic, Riley, Rooks, Saline, Scott, Sedgwick, Seward, Shawnee, Sheridan, Sherman, Smith, Stafford, Stanton, Stevens, Sumner, Thomas, Wabaunsee, Washington, Wilson, Woodson, Wyandotte.

KENTUCKY.

Section I: Bell, Boyd, Breathitt, Carter, Clay, Clinton, Cumberland, Elliott, Floyd, Greenup, Harlan, Jackson, Johnson, Knott, Knox, Laurel, Lawrence, Lee, Leslie, Letcher, Lewis, Magoffin, Martin, Menifee, Monroe, Morgan, Owsley, Perry, Pike, Pulaski, Rockcastle, Rowan, Russell, Wayne, Whitley, Wolfe.

Section II: Adair, Allen, Anderson, Ballard, Barren, Bath, Boone, Bourbon, Boyle, Bracken, Breckinridge, Bullitt, Butler, Caldwell, Calloway, Campbell, Carlisle, Carroll, Casey, Christian, Clark, Crittenden, Daviess, Edmonson, Estill, Fayette, Fleming, Franklin, Fulton, Gallatin, Garrard, Grant, Graves, Grayson, Green, Hancock, Hardin, Harrison, Hart, Henderson, Henry, Hickman, Hopkins, Jefferson, Jessamine, Kenton, Larue, Lincoln, Livingston, Logan, Lyon, McCracken, McLean, Madison, Marion, Marshall, Mason, Meade, Mercer, Metcalfe, Montgomery, Muhlenberg, Nelson, Nicholas, Ohio, Oldham, Owen, Pendleton, Powell, Robertson, Scott, Shelby, Simpson, Spencer, Taylor, Todd, Trigg, Trimble, Union, Warren, Washington, Webster, Woodford.

LOUISIANA.

Section I (parishes): Ascension Bossier, Caddo, Claiborne, Concordia, De Soto, East Baton Rouge, East Carroll, East Feliciana, Iberville, Jefferson, Madison, Morehouse, Natchitoches, Ouachita, Plaquemines, Pointe Coupee, Red River, Richland, St. Charles, St. James, St. John the Baptist, St. Mary, Tensas, Webster, West Baton Rouge, West Carroll, West Feliciana.

Section II: Includes parish of Orleans:

Section III (parishes): Acadia, Allen, Assumption, Avoyelles, Beauregard, Blenville, Calcasieu, Caldwell, Cameron, Catahoula, Evangeline, Franklin, Grant, Iberia, Jackson, Jefferson Davis, Lafayette, Lafourche, La Salle, Lincoln, Livingston, Rapides, Sabine, St. Bernard, St. Helena, St. Landry, St. Martin, St. Tammany, Tangipahoa, Terrebonne, Union, Vermillion, Vernon, Washington, Winn.

MAINE.

Section I: Aroostook, Penobscot, Piscataquis, Washington.

Section II: Hancock, Knox, Lincoln, Sagadahoc, Waldo.

Section III: Androscoggin, Cumberland, Franklin, Kennebec, Oxford, Somerset, York.

MARYLAND.

Section I: Includes county and city of Baltimore.

Section II: Caroline, Dorchester, Kent, Queen Anne, Somerset, Talbot, Wicomico, Worcester.

Section III: Allegany, Anne Arundel, Carroll, Cecil, Frederick, Garrett, Harford, Howard, Montgomery, Prince Georges, Washington.

Section IV: Calvert, Charles, St. Marys.

MASSACHUSETTS.

Section I: Berkshire, Franklin.

Section II: Bristol, Essex, Hampden, Hampshire, Middlesex, Norfolk, Worcester.

Section III: Barnstable, Dukes, Nantucket, Plymouth.

Section IV: Suffolk.

MICHIGAN.

Section I: Alger, Baraga, Gogebic, Houghton, Iron, Keweenaw, Luce, Marquette, Ontonagon.

Section II: Alcona, Alpena, Antrim, Arenac, Barry, Benzie, Branch, Calhoun, Cass, Charlevoix, Cheboygan, Chippewa, Clare, Clinton, Crawford, Delta, Dickinson, Eaton, Emmet, Genesee, Gladwin, Grand Traverse, Gratiot, Hillsdale, Ingham, Ionia, Iosco, Isabella, Jackson, Kalamazoo, Kalkaska, Lake, Leelanau, Livingston, Mackinac, Manistee, Mason, Mecosta, Menominee, Midland, Missaukee, Montcalm, Montmorency, Newaygo, Oceana, Ogemaw, Osceola, Oscoda, Otsego, Presque Isle, Roscommon, St. Joseph, Schoolcraft, Shiawassee, Wexford.

City not included in counties, Grand Rapids.

Section III: Bay, Huron, Lapeer, Lenawee, Macomb, Monroe, Oakland, Saginaw, St. Clair, Sanilac, Tuscola, Washtenaw, Wayne (except for city of Detroit).

Section IV: Includes city of Detroit.

Section V: Allegan, Berrien, Kent, Muskegon, Ottawa, Van Buren.

MINNESOTA.

Section I: Aitkin, Anoka, Becker, Beltrami, Big Stone, Cass, Chippewa, Chisago, Clay, Clearwater, Crow Wing, Douglas, Grant, Hubbard, Isanti, Kanabec, Kandiyohi, Kittson, Koochiching, Lac qui Parle, Mahnomen, Marshall, Meeker, Mille Lacs, Norman, Otter Tail, Pennington, Pine, Polk, Pope, Red Lake, Renville, Roseau, Sherburne, Stevens, Swift, Todd, Traverse, Wadena, Wilkin.

Section II: Benton, Blue Earth, Brown, Carver, Cottonwood, Dakota, Dodge, Faribault, Fillmore, Freeborn, Goodhue, Hennepin, Houston, Jackson, Le Sueur, Lincoln, Lyon, McLeod, Martin, Morrison, Mower, Murray, Nicollet, Nobles, Olm-

stead, Pipestone, Ramsay, Redwood, Rice, Rock, Scott, Sibley, Stearns, Steele, Wabasha, Waseca, Washington, Watonwan, Winona, Wright, Yellow Medicine.

Section III: Carlton, Cook, Itasca, Lake, St. Louis.

Section IV: Includes cities of Minneapolis and St. Paul.

MISSISSIPPI.

Section I: Adams, Amite, Attala, Benton, Bolivar, Carroll, Chickasaw, Claiborne, Clay, Coahoma, Copiah, De Soto, Grenada, Hinds, Holmes, Issaquena, Jefferson, Jefferson Davis, Kemper, Lafayette, Leflore, Lowndes, Madison, Marshall, Monroe, Montgomery, Noxubee, Panola, Oktibbeha, Rankin, Sharkey, Sunflower, Tallahatchie, Tate, Tunica, Warren, Washington, Wilkinson, Yazoo.

Section II: Alcorn, Calhoun, Choctaw, Clarke, Covington, Forrest, Franklin, George, Greene, Hancock, Harrison, Itawamba, Jackson, Jasper, Jones, Lamar, Lauderdale, Lawrence, Leake, Lee, Lincoln, Marion, Neshoba, Newton, Pearl River, Perry, Pike, Pontotoc, Prentiss, Scott, Simpson, Smith, Stone, Tippah, Tishomingo, Union, Walthall, Wayne, Webster, Winston.

MISSOURI.

Section I: Adair, Andrew, Atchison, Barton, Bates, Benton, Bollinger, Buchanan, Butler, Caldwell, Camden, Carroll, Carter, Cass, Cedar, Clark, Clay, Clinton, Cole, Crawford, Dade, Dallas, Daviess, Dekalb, Dent, Dunklin, Franklin, Gasconade, Gentry, Greene, Grundy, Harrison, Henry, Hickory, Holt, Iron, Jasper, Jefferson, Johnson, Knox, Laclede, Lawrence, Lewis, Linn, Livingston, Macon, Madison, Maries, Mercer, Miller, Moniteau, Morgan, Newton, Nodaway, Oregon, Osage, Perry, Phelps, Platte, Polk, Pulaski, Putnam, Ray, Reynolds, Ripley, St. Clair, St. Francois, St. Louis, Ste. Genevieve, Schuyler, Scotland, Shannon, Shelby, Stoddard, Sullivan, Texas, Vernon, Washington, Wayne, Worth.

City not included in counties, Kansas City.

Section II: Audrain, Boone, Callaway, Cape Girardeau, Chariton, Cooper, Howard, Jackson, Lafayette, Lincoln, Marion, Mississippi, Monroe, Montgomery, New Madrid, Pemiscot, Pettis, Pike, Ralls, Randolph, St. Charles, Saline, Scott, Warren.

Section III: Barry, Christian, Douglas, Howell, McDonald, Ozark, Stone, Taney, Webster, Wright.

Section IV: Includes city of St. Louis.

MONTANA.

Section I: Broadwater, Carbon, Cascade, Deer Lodge, Flathead, Granite, Jefferson, Lewis and Clark, Lincoln, Mineral, Missoula, Powell, Sanders, Silver Bow, Stillwater, Yellowstone.

Section II: Beaverhead, Bighorn, Blaine, Carter, Chouteau, Custer, Dawson, Fallon, Fergus, Gallatin, Hill, Madison, Meagher, Musselshell, Park, Phillips, Prairie, Ravalli, Richland, Rosebud, Sheridan, Sweetgrass, Teton, Toole, Valley, Wheatland, Wibaux.

NEBRASKA.

Section I: Antelope, Banner, Blaine, Boxbutte, Boyd, Brown, Burt, Cass, Cedar, Chase, Cherry, Cheyenne, Cuming, Custer, Dakota, Dawes, Dawson, Deuel, Dixon, Dodge, Douglas, Dundy, Frontier, Gage, Garden, Garfield, Gosper, Grant, Greeley, Hayes, Holt, Hooker, Johnson, Keith, Keyapaha, Kimball, Knox, Lancaster, Lincoln, Logan, Loup, McPherson, Morrill, Nemaha, Otoe, Pawnee, Perkins, Pierce, Richardson, Rock, Sarpy, Saunders, Scotts Bluff, Sheridan, Sherman, Sioux, Thomas, Thurston, Valley, Washington, Wayne, Wheeler.

Section II: Adams, Boone, Buffalo, Butler, Clay, Colfax, Fillmore, Franklin, Furnas, Hall, Hamilton, Harlan, Hitchcock, Howard, Jefferson, Kearney, Madison, Merrick, Nance, Nuckolls, Phelps, Platte, Polk, Redwillow, Saline, Seward, Stanton, Thayer, Webster, York.

NEVADA.

Section I: Includes entire State.

NEW HAMPSHIRE.

Section I: Carroll, Coos, Grafton.
 Section II: Belknap, Cheshire, Hillsborough, Merrimack, Rockingham, Strafford, Sullivan.

NEW JERSEY.

Section I: Bergen, Essex, Hudson, Passaic, Union.
 Section II: Burlington, Camden, Cape May, Cumberland, Gloucester, Mercer, Middlesex, Monmouth, Ocean, Salem.
 City not included in counties, Orange.
 Section III: Atlantic, Hunterdon, Morris, Somerset, Sussex, Warren.

NEW MEXICO.

Section I: McKinley, Rio Arriba, Sandoval, San Juan, Valencia.
 Section II: Bernalillo, Chaves, Colfax, Curry, De Baca, Guadalupe, Lea, Lincoln, Mora, Quay, Roosevelt, San Miguel, Santa Fe, Socorro, Taos, Torrance, Union.
 Section III: Dona Ana, Eddy, Grant, Lea (one-half), Luna, Otero, Sierra.

NEW YORK.

Section I: Dutchess, Nassau, Putnam, Suffolk, Westchester.
 Section II: Kings, New York, Queens, Richmond.
 Section III: Albany, Columbia, Fulton, Herkimer, Montgomery, Otsego, Rensselaer, Saratoga, Schenectady, Schoharie, Washington.
 Section IV: Cayuga, Erie (except city of Buffalo), Genesee, Jefferson, Monroe, Niagara, Oneida, Onondaga, Ontario, Orleans, Oswego, Seneca, Wayne.
 Cities not included in counties, Amsterdam, Niagara Falls, Troy.
 Section V: Greene, Orange, Rockland, Ulster.
 Section VI: Includes city of Buffalo.
 Section VII: Allegany, Broome, Cattaragus, Chautauqua, Chemung, Chenango, Cortland, Delaware, Livingston, Madison, Schuyler, Steuben, Sullivan, Tioga, Tompkins, Wyoming, Yates.
 Section VIII: Clinton, Essex, Franklin, Hamilton, Lewis, St. Lawrence, Warren.

NORTH CAROLINA.

Section I: Ashe, Alleghany, Alexander, Avery, Buncombe, Burke, Caldwell, Cherokee, Clay, Haywood, Graham, Henderson, Jackson, McDowell, Macon, Madison, Mitchell, Polk, Rutherford, Swain, Transylvania, Watauga, Wilkes, Yancey.
 Section II: Alamance, Cabarrus, Caswell, Catawba, Chatham, Cleveland, Davidson, Davie, Forsyth, Gaston, Guilford, Iredell, Lincoln, Mecklenburg, Orange, Person, Randolph, Rockingham, Rowan, Stokes, Surry, and Yadkin.
 Section III: Anson, Cumberland, Harnett, Hoke, Lee, Montgomery, Moore, Richmond, Robeson, Sampson, Scotland, Stanley, Union.
 Section IV: Beaufort, Bertie, Chowan, Craven, Durham, Edgecombe, Franklin, Gates, Granville, Greene, Halifax, Hertford, Jones, Johnston, Lenoir, Martin, Nash, Northhampton, Onslow, Pasquotank, Perquimans, Pitt, Vance, Wake, Warren, Washington, Wayne, Wilson.
 Section V: Camden, Carteret, Currituck, Dare, Hyde, Pamlico, Tyrrell.
 Section VI: Bladen, Brunswick, Columbus, Duplin, New Hanover, Pender.

NORTH DAKOTA.

Section I: Bottineau, Cavaller, Golden Valley, Grand Forks, Pembina, Rquette, Towner, Walsh.
 Section II: Adams, Barnes, Benson, Billings, Bowman, Burke, Cass, Divide, Eddy, Foster, Griggs, McKenzie, Mountrall, Nelson, Ramsay, Ranson, Renville, Richland, Sargent, Slope, Steele, Trall, Ward, Williams.
 Section III: Burleigh, Dickey, Dunn, Emmons, Grant, Hettinger, Kidder, Lamoure, Logan, McHenry, McIntosh, McLean, Mercer, Morton, Oliver, Pierce, Sheridan, Sioux, Stark, Stutsman, Wells.

OHIO.

Section I: Cuyahoga, Erie, Lake, Lorain, Lucas, Ottawa.

Section II: Ashtabula, Belmont, Carroll, Columbiana, Geauga, Guernsey, Harrison, Jefferson, Mahoning, Medina, Portage, Stark, Summit, Trumbull, Tuscarawas, Wayne.

Section III: Adams, Allen, Ashland, Athens, Auglaize, Brown, Butler, Champaign, Clark, Clermont, Clinton, Coshocton, Crawford, Darke, Defiance, Delaware, Fairfield, Fayette, Franklin, Fulton, Gallia, Greene, Hamilton, Hancock, Hardin, Henry, Highland, Hocking, Holmes, Huron, Jackson, Knox, Lawrence, Licking, Logan, Madison, Marion, Meigs, Mercer, Miami, Monroe, Montgomery, Morgan, Morrow, Muskingum, Noble, Paulding, Perry, Pickaway, Pike, Preble, Putnam, Richland, Ross, Sandusky, Scioto, Seneca, Shelby, Union, Van Wert, Vinton, Warren, Washington, Williams, Wood, Wyandot.

Section IV: City of Cincinnati.

OKLAHOMA.

Section I: Adair, Atoka, Bryan, Cherokee, Choctaw, Craig, Delaware, Haskell, Hughes, Johnston, Latimer, Le Flore, McCurtain, McIntosh, Mayes, Muskogee, Nowata, Okfuskee, Okmulgee, Osage, Ottawa, Pittsburg, Pushmataha, Rogers, Seminole, Sequoyah, Tulsa, Wagoner, Washington.

Section II: Alfalfa, Beaver, Beckham, Blaine, Caddo, Canadian, Carter, Cimarron, Cleveland, Coal, Comanche, Cotton, Creek, Custer, Dewey, Ellis, Garfield, Garvin, Grady, Grant, Greer, Harmon, Harper, Jackson, Jefferson, Kay, Kingfisher, Kiowa, Lincoln, Logan, Love, McClain, Major, Marshall, Murray, Noble, Oklahoma, Pawnee, Payne, Pontotoc, Pottawatomie, Roger Mills, Stephens, Texas, Tillman, Washita, Woods, Woodward.

OREGON.

Section I: Benton, Clackamas, Clatsop, Columbia, Hood River, Lincoln, Linn, Marion, Multnomah, Polk, Tillamook, Wasco, Washington, Yamhill.

Section II: Baker, Coos, Crook, Curry, Douglas, Gilliam, Grant, Harney, Jackson, Josephine, Klamath, Lake, Lane, Malheur, Morrow, Sherman, Umatilla, Union, Wallowa, Wheeler.

PENNSYLVANIA.

Section I: Philadelphia.

Section II: Adams, Bedford, Berks, Bucks, Chester, Cumberland, Dauphin, Delaware, Franklin, Fulton, Huntington, Juniata, Lancaster, Lebanon, Lehigh, Mifflin, Monroe, Montgomery, Northampton, Perry, Pike, Snyder, Union, York.

Section III: Carbon, Columbia, Lackawanna, Luzerne, Montour, Northumberland, Schuylkill, Wayne.

Section IV: Beaver, Butler, Greene, Lawrence, Washington.

Section V: Blair, Cambria, Fayette, Somerset, Westmoreland.

City not included in counties, Altoona.

Section VI: Armstrong, Cameron, Clarion, Clearfield, Crawford, Elk, Erie, Forest, Indiana, Jefferson, McKean, Mercer, Potter, Venango, Warren, Wyoming. Cities not included in counties, Williamsport and New Castle.

Section VII: Allegheny, Bradford, Center, Clinton, Lycoming, Sullivan, Susquehanna, Tioga.

City not included in counties, McKeesport.

RHODE ISLAND.

Section I: Includes entire State.

SOUTH CAROLINA.

Section I: Anderson, Cherokee, Greenville, Oconee, Pickens, Spartanburg.

Section II: Abbeville, Aiken, Bamberg, Barnwell, Calhoun, Chester, Edgefield, Fairfield, Greenwood, Kershaw, Lancaster, Laurens, Lexington, McCormick, Newberry, Orangeburg, Richland, Saluda, Union and York.

Section III: Beaufort, Berkley, Charlestown, Chesterfield, Clarendon, Colleton, Darlington, Dillon, Dorchester, Florence, Georgetown, Hampton, Horry, Jasper, Lee, Marion, Marlboro, Sumter, Williamsburg.

SOUTH DAKOTA.

Section I: Aurora, Beadle, Brookings, Brown, Brule, Buffalo, Butte, Charles Mix, Clark, Clay, Codrington, Custer, Davison, Day, Deuel, Douglas, Fall River, Faulk, Grant, Gregory, Hamlin, Hand, Harding, Hyde, Jerauld, Kingsbury, Lake, Lawrence, Lincoln, Lyman, McCook, Marshall, Meade, Miner, Minnehaha, Moody, Pennington, Perkins, Roberts, Sanborn, Spink, Stanley, Union, Yankton.

Section II: Bonhomme, Campbell, Edmunds, Hanson, Hughes, Hutchinson, McPherson, Potter, Sully, Turner, Walworth.

Section III: Armstrong, Bennett, Corson, Dewey, Mellette, Shannon, Todd, Washabaugh, Washington, Ziebach.

TENNESSEE.

Section I: Crockett, Dyer, Fayette, Gibson, Hardeman, Haywood, Lake, Lauderdale, Madison, Obion, Shelby, Tipton.

Section II: Bedford, Benton, Cannon, Carroll, Cheatham, Chester, Clay, Coffee, Davidson, Decatur, Dekalb, Dickson, Fentress, Franklin, Giles, Grundy, Hardin, Henderson, Henry, Hickman, Houston, Humphreys, Jackson, Lawrence, Lewis, Lincoln, McNairy, Macon, Marshall, Maury, Montgomery, Moore, Overton, Perry, Pickett, Putnam, Robertson, Rutherford, Scott, Smith, Stewart, Sumner, Trousdale, Van Buren, Warren, Wayne, Weakley, White, Williamson, Wilson.

Cities not included in counties, Memphis and Knoxville.

Section III: Anderson, Bledsoe, Blount, Bradley, Campbell, Carter, Claiborne, Cocke, Cumberland, Grainger, Greene, Hamblen, Hamilton, Hancock, Hawkins, James, Jefferson, Johnson, Knox, Loudon, McMinn, Marion, Meigs, Monroe, Morgan, Polk, Rhea, Reane, Sequatchie, Sevier, Sullivan, Unico, Union, Washington.

TEXAS.

Section I: Atacosa, Bastrop, Bee, Bexar, Brewster, Brooks, Caldwell, Cameron, Comal, Culberson, Dimmit, Duval, El Paso, Eril, Goliad, Guadalupe, Hays, Hidalgo, Hudspeth, Jeff Davis, Jim Hogg, Jim Wells, Karnes, Kinney, Kleberg, La Salle, Live Oak, McMullen, Maverick, Medina, Nueces, Pecos, Presidio, Reeves, San Patricio, Starr, Terrell, Travis, Uvalde, Valverde, Webb, Willacy, Williamson, Wilson, Zapata, Zavalla.

Section II: Anderson, Andrews, Angelina, Archer, Armstrong, Bailey, Bandera, Baylor, Bell, Blanco, Borden, Bosque, Bowie, Briscoe, Brown, Burnet, Callahan, Camp, Carson, Cass, Castro, Cherokee, Childress, Clay, Cochran, Coke, Coleman, Collin, Collingsworth, Comanche, Concho, Cooke, Coryell, Cottle, Crane, Crockett, Crosby, Dallam, Dallas, Dawson, Deaf Smith, Delta, Denton, Dickens, Donley, Eastland, Ector, Edwards, Ellis, Erath, Falls, Fannin, Fisher, Floyd, Foard, Franklin, Freestone, Gaines, Garza, Gillespie, Glasscock, Gray, Grayson, Gregg, Hale, Hall, Hamilton, Hansford, Hardeman, Harrison, Hartley, Haskell, Hemphill, Henderson, Hill, Hockley, Hood, Hopkins, Houston, Howard, Hunt, Hutchinson, Irion, Jack, Jones, Kaufman, Kendall, Kerr, Kimble, King, Knox, Lamar, Lamb, Lampasas, Lee, Leon, Limestone, Lipscomb, Llano, Loving, Lubbock, Lynn, McCulloch, McLennan, Madison, Marion, Martin, Mason, Menard, Midland, Milam, Mills, Mitchell, Montague, Moore, Morris, Motley, Nacogdoches, Navarro, Nolan, Ochiltree, Oldham, Palo Pinto, Panola, Parker, Parmer, Polk, Potter, Rains, Randall, Reagan, Real, Red River, Roberts, Rockwall, Runnels, Rusk, San Augustine, San Saba, Schleicher, Scurry, Shackelford, Shelby, Sherman, Smith, Somervell, Stephens, Sterling, Stonewall, Sutton, Swisher, Tarrant, Taylor, Terry, Throckmorton, Titus, Tom Green, Trinity, Tyler, Upshur, Upton, Van Zandt, Ward, Wheeler, Wichita, Wilbarger, Winkler, Wise, Wood, Yoakum, Young.

City not included in counties, Houston.

Section III: Austin, Colorado, De Witt, Fayette, Gonzales, Lavaca, Washington.

City not included in counties, Austin.

Section IV: Aransas, Brazoria, Calhoun, Chambers, Galveston, Hardin, Harris, Jackson, Jasper, Jefferson, Liberty, Matagorda, Newton, Orange, Refugio, Sabine, Victoria, Wharton.

Section V: Brazos, Burleson, Ford Bend, Grimes, Montgomery, Robertson, San Jacinto, Walker, Waller.

UTAH.

Section I: Beaver, Box Elder, Emery, Garfield, Grand, Iron, Juab, Kane, Millard, Piute, San Juan, Sevier, Tooele, Uinta, Washington, Wayne.

Section II: Cache, Davis, Salt Lake, Sanpete, Utah, Weber.

Section III: Carbon, Duchesne, Morgan, Rich, Summit, Wasatch.

VERMONT.

Section I: Undivided.

VIRGINIA.

Section I: Accomac, Elizabeth City, Gloucester, Lancaster, Mathews, Middlesex, Norfolk, Northampton, Northumberland, Princess Anne, Warwick, York.

Section II: Amelia, Brunswick, Caroline, Charlotte, Charles City, Chesterfield, Cumberland, Dinwiddie, Essex, Goochland, Greenville, Halifax, Hanover, Henrico, Isle of Wight, James City, King and Queen, King George, King William, Lunenburg, Macklenburg, Nansemond, New Kent, Nottoway, Powhatan, Prince Edward, Prince George, Richmond, Surry, Sussex, Southampton, Westmoreland.

Section III: Albemarle, Alexandria, Amherst, Appomattox, Bedford, Buckingham, Campbell, Culpeper, Fairfax, Fauquier, Fluvanna, Franklin, Greene, Henry, Loudoun, Louisa, Madison, Nelson, Orange, Pittsylvania, Prince William, Rappahannock, Spotsylvania, Stafford.

Section IV: Alleghany, Augusta, Bath, Bland, Botetourt, Buchanan, Carroll, Clarke, Craig, Dickenson, Floyd, Frederick, Giles, Grayson, Highland, Lee, Montgomery, Page, Patrick, Pulaski, Roanoke, Rockbridge, Rockingham, Russell, Scott, Shenandoah, Smythe, Tazewell, Warren, Washington, Wise, Wythe.

WASHINGTON.

Section I: Adams, Asotin, Benton, Clallam, Clarke, Columbia, Cowlitz, Franklin, Garfield, Grays Harbor, Jefferson, Klickitat, Lewis, Lincoln, Mason, Pacific, Skamania, Spokane, Thurston, Wahkiakum, Walla Walla, Whitman.

Section II: Island, King, Kitsap, Pierce, San Juan, Skagit, Snohomish, Whatcom.

Section III: Chelan, Douglas, Ferry, Grant, Kittitas, Okanogan, Pend Oreille, Stevens, Yakima.

WEST VIRGINIA.

Section I: Berkeley, Grant, Hampshire, Hardy, Jefferson, Mineral, Morgan, Pendleton, Pocahontas, Preston, Randolph, Tucker.

Section II: Barbour, Boone, Braxton, Brooke, Cabell, Calhoun, Clay, Doddridge, Fayette, Gilmer, Greenbrier, Hancock, Harrison, Jackson, Kanawha, Lewis, Lincoln, Logan, McDowell, Marion, Marshall, Mason, Mercer, Mingo, Monongalia, Monroe, Nicholas, Ohio, Pleasants, Putnam, Raleigh, Ritchie, Roane, Summers, Taylor, Tyler, Upshur, Wayne, Webster, Wetzel, Wirt, Wood, Wyoming.

WISCONSIN.

Section I: Ashland, Barron, Bayfield, Buffalo, Burnett, Chippewa, Crawford, Douglas, Dunn, Eau Claire, Iron, Jackson, La Crosse, Pepin, Pierce, Polk, Price, Rusk, St. Croix, Sawyer, Taylor, Trempealeau, Vernon, Washburn.

Section II: Adams, Clark, Columbia, Dane, Dodge, Florence, Fond du Lac, Forest, Grant, Green, Green Lake, Iowa, Jefferson, Juneau, Lafayette, Langlade, Lincoln, Marathon, Marinette, Marquette, Monroe, Oconto, Oneida, Outagamie, Portage, Richland, Rock, Sauk, Shawano, Vilas, Walworth, Washington, Waukesha, Wampaca, Waushara, Winnebago, Wood.

City not included in counties, Green Bay.

Section III: Milwaukee.

Section IV: Brown, Calumet, Door, Kenosha, Kewaunee, Manitowoc, Ozaukee, Racine, Sheboygan.

City not included in counties, Superior.

WYOMING.

Section I: Includes entire State.




PLATE V.

TABLE 19.—*Consolidation of similar sections; the series and their constituent groups.*

Series I. The occupational series:

- Group 1. Agricultural, North, native white, 73 per cent.
- Group 2. Agricultural, North and West, mixed foreign and native white.
- Group 3. Agricultural, South, native white.
- Group 4. Agricultural, South, Negro, 45 per cent plus.
- Group 5. Eastern manufacturing.
- Group 6. Commuter.
- Group 7. Mining.

Series II. The physiographic series:

- Group 8. Sparsely settled, not more than 3 per square mile.
- Group 9. Desert.
- Group 10. Maritime.
- Group 11. Mountain.

Series III. The racial series:

- Group 12. Mountain whites.
- Group 13. Indian, sparsely settled.
- Group 14. Mexican, sparsely settled.
- Group 15. Native whites of Scotch origin.
- Group 16. Russian, 10 per cent plus.
- Group 17. Scandinavian, 10 per cent plus.
- Group 18. Finn, 10 per cent plus.
- Group 19. French Canadian, 10 per cent plus.
- Group 20. German and Scandinavian, 10 per cent plus.
- Group 21. German and Austrian, 20 per cent plus.
- Group 22. German and Austrian, 15 per cent plus.

TABLE 20.—*Consolidation of similar sections; the groups and their composition out of sections.*

Group 1. Agricultural, North, native white, 73 per cent:

Illinois.....	3	Ohio.....	3
Indiana.....	3	Pennsylvania.....	2
Iowa.....	2		

Group 2. Agricultural, North and West, mixed foreign and native white:

Colorado.....	4	New York.....	7
Illinois.....	8	Ohio.....	2
Indiana.....	2	Pennsylvania.....	6
Iowa.....	1	South Dakota.....	1
Kansas.....	2	Vermont.....	1
Michigan.....	2	Washington.....	1
Nebraska.....	2	Wisconsin.....	2
New Jersey.....	2		

Group 3. Agricultural, South, native white:

Alabama.....	3	North Carolina.....	2, 3, 6
Arkansas.....	2, 3	Oklahoma.....	1, 2
Kentucky.....	2	Tennessee.....	2
Louisiana.....	3	Texas.....	2, 4
Maryland.....	3	Virginia.....	3
Mississippi.....	2	West Virginia.....	2
Missouri.....	1, 3		

Group 4. Agricultural, South, Negro, 45 per cent plus:

Alabama.....	2, 4	North Carolina.....	4
Arkansas.....	1	South Carolina.....	2, 3
Georgia.....	2	Tennessee.....	1
Louisiana.....	1	Texas.....	5
Mississippi.....	1	Virginia.....	2

TABLE 23.—Consolidation of similar sections; the groups and their composition out of sections—Con.

Group 5. Eastern manufacturing:			
Connecticut.....	2	New York.....	3
Massachusetts.....	2	Ohio.....	1
New Hampshire.....	2	Pennsylvania.....	5
New Jersey.....	1	Rhode Island.....	1
Group 6. Commuter:			
Illinois.....	1	New York.....	1
New Jersey.....	1		
Group 7. Mining:			
Alabama.....	1	Montana.....	1
California.....	2	Nevada.....	1
Colorado.....	1, 3, 6	Pennsylvania.....	3, 4
Idaho.....	1	Utah.....	3
Group 8. Sparsely settled, not more than 3 per square mile:			
California.....	3	Oregon.....	2
Montana.....	2	Utah.....	1
Nevada.....	1	Wyoming.....	1
New Mexico.....	2		
Group 9. Desert:			
Arizona.....	2	New Mexico.....	2
Nevada.....	1		
Group 10. Maritime:			
Maine.....	2	North Carolina.....	5
Maryland.....	2, 4	Virginia.....	1
Massachusetts.....	3		
Group 11. Mountain:			
Arkansas.....	2	New Hampshire.....	1
Massachusetts.....	1	New York.....	5, 8
Missouri.....	3	Washington.....	3
Montana.....	1	Wyoming.....	1
Group 12. Mountain whites:			
Kentucky.....	1	Tennessee.....	3
North Carolina.....	1	Virginia.....	4
South Carolina.....	1	West Virginia.....	1
Group 13. Indian, sparsely settled:			
Arizona.....	1	Oklahoma.....	1
New Mexico.....	1	South Dakota.....	3
Group 14. Mexican, sparsely settled:			
Arizona.....	1, 2	Texas.....	1
New Mexico.....	3		
Group 15. Native whites of Scotch origin:			
Kentucky.....	2	North Carolina.....	3
Group 16. Russian, 10 per cent plus:			
Colorado.....	2	Pennsylvania.....	3
Kansas.....	1	South Dakota.....	2
North Dakota.....	3		
Group 17. Scandinavian, 10 per cent plus:			
Michigan.....	1	Utah.....	1, 2
Minnesota.....	1, 2, 3	Washington.....	2
North Dakota.....	1, 2, 3	Wisconsin.....	1, 2
South Dakota.....	1		

TABLE 20.—*Consolidation of similar sections; the groups and their composition out of sections—Con.*

Group 18. Finn, 10 per cent plus:

Michigan.....	1	Minnesota.....	3
---------------	---	----------------	---

Group 19. French Canadian, 10 per cent plus:

Maine.....	3	New Hampshire.....	1, 2
Massachusetts.....	2	Rhode Island.....	1

Group 20. German and Scandinavian, 10 per cent plus:

Minnesota.....	1, 2	Wisconsin.....	1, 2
South Dakota.....	1		

Group 21. German and Austrian, 20 per cent plus:

Illinois.....	1, 4	Minnesota.....	2
Indiana.....	1	Ohio.....	1

Group 22. German and Austrian, 15 per cent plus:

Illinois.....	1, 4	New Jersey.....	1
Indiana.....	1	Ohio.....	1
Iowa.....	1	Pennsylvania.....	3, 5, 7
Minnesota.....	2	Wisconsin.....	1, 2, 4
Nebraska.....	1, 2		

10. AVERAGE STATURE OF RECRUITS FROM DIFFERENT SECTIONS.

For various purposes the country has been divided into 156 sections, on the basis of population. Table 21 gives the average stature of recruits from the different sections arranged in order of this stature. At the head of this table stands Section 1 of North Carolina, the sparsely populated mountainous area of that State. Here the stature is 68.67 inches (174.42 centimeters), being 1.18 inches above the average of the United States. This tall stature is practically the same as that given for 1,304 Scotch in general, namely, 174.6 centimeters. The reason for the exceptionally great stature of men from Section 1 of North Carolina is primarily that many are of Scotch origin. As is well known, North Carolina, especially the Cape Fear region, was settled by Scotch Presbyterians in the middle of the seventeenth century. Their descendants have penetrated to the higher regions of the Cape Fear River in Scotland County and many of them have settled in the mountain region of western North Carolina. It is probable that there has been something of a selection of the largest and hardiest of these Scotch to settle the mountain region. It appears also that in Section 2, comprising the intermediate part of North Carolina, the stature is very great, 68.26 inches. In Section 3, comprising a large proportion of native whites of Scotch origin, the stature is 68.24 inches, while in those parts of North Carolina which lie near the sea coast the population is only slightly above the average for the United States. Unfortunately, it is not possible to say what was the stature of men of North Carolina at the time of the Civil War because this State was one of those in secession and its statistics are not included in those of recruits of the northern Army. During the Civil War the greatest average stature was found in men from Kentucky and Tennessee. In the present table Section 1 of Kentucky (mountainous area, native whites), gives an average stature of 68.21 inches, which is 0.72 above the average of the whole United States, and Section 2 of Kentucky (agricultural area of the central and western part) has an average stature of 67.95, or nearly 0.5 inch

above the mean of the whole United States. The mean stature for Kentucky, 68.02, is less than that given by Gould² (p. 95) for men from Kentucky and Tennessee—namely, 68.16.

TABLE 21.—Mean height, by sections; sections arranged in order of standing, with proportional weight and chest circumference (expiration) for each inch of height; also standard deviation for each height, first million draft recruits.

Alabama.....	3	Large native white population.....	2,670	68.21	2.74	2.063	.484
Washington.....	3	Mountainous area.....	1,539	68.19	2.66	2.142	.493
Texas.....	1	Large Mexican population.....	6,676	68.19	2.70	2.080	.487
South Carolina.....	1	Native white.....	1,564	68.19	2.53	2.060	.484
Kansas.....	2	Native and German population.....	8,505	68.18	2.54	2.105	.488
Arizona.....	2	Chiefly white population.....	2,823	68.17	2.61	2.066	.487
Montana.....	2	Sparsely settled, mountainous area.....	6,531	68.17	2.57	2.150	.493
Illinois.....	6	Negro population (Egypt).....	409	68.16	2.38	2.043	.482
Oklahoma.....	1	Marked Indian and Negro population.....	8,471	68.16	2.59	2.078	.485
Utah.....	1	Sparsely populated.....	1,224	68.16	2.64	2.114	.492
Alabama.....	4	Large Negro population.....	669	68.16	2.61	2.115	.486
Alaska.....	All	Undivided.....	106	68.15	2.30	2.208	.493
Mississippi.....	1	Rural area, large Negro population.....	5,149	68.15	2.67	2.120	.488
Minnesota.....	2	German and Scandinavian population.....	7,601	68.14	2.63	2.170	.497
Virginia.....	4	Mountain, white.....	5,512	68.14	2.54	2.055	.489
Oregon.....	2	Columbia River Valley and coastal dry plain, sparsely populated.....	1,077	68.13	2.52	2.140	.490
South Dakota.....	3	Indian population.....	247	68.13	2.41	2.180	.496
Wisconsin.....	1	Scandinavian and German population.....	3,297	68.13	2.66	2.130	.494
Colorado.....	3	English population.....	381	68.12	2.66	2.086	.487
Indiana.....	2	Agricultural, considerable German.....	837	68.12	2.48	2.120	.491
Virginia.....	3	Native rural region.....	3,866	68.12	2.73	2.066	.489
Idaho.....	All	State undivided.....	4,034	68.10	2.57	2.133	.495
Missouri.....	2	Mississippi bottoms, considerable Negro population.....	3,448	68.10	2.63	2.090	.486
Iowa.....	1	Foreign white, German and Scandinavian.....	12,139	68.09	2.56	2.139	.492
Missouri.....	1	Native white, agricultural.....	13,588	68.09	2.59	2.080	.486
Texas.....	4	Coastal native population.....	2,722	68.09	2.70	2.090	.487
Georgia.....	1	Mixed population, native white predominating.....	10,248	68.08	2.63	2.064	.486
Oregon.....	1	Fairly densely populated.....	2,748	68.08	2.61	2.133	.492
South Dakota.....	1	Dry farming area.....	3,051	68.07	2.68	2.160	.492
Tennessee.....	1	Negroes, Mississippi bottoms.....	2,218	68.07	2.59	2.090	.483
Colorado.....	1	Large native white population.....	1,056	68.08	2.79	2.061	.489
Arkansas.....	1	Negro, Mississippi bottoms.....	4,945	68.06	2.68	2.063	.487
Colorado.....	4	Prevalently agricultural.....	1,227	68.05	2.70	2.067	.486
North Dakota.....	2	Scandinavian population.....	3,307	68.03	2.48	2.159	.497
Arizona.....	1	Large Indian population, sparsely settled.....	1,027	68.02	2.73	2.108	.489
Nebraska.....	1	German and Irish, foreign stocks.....	7,629	68.02	2.69	2.120	.488
Washington.....	1	Coastal region plus eastern counties.....	5,176	68.01	2.60	2.139	.492
West Virginia.....	1	Mountainous area.....	1,507	67.98	2.71	2.072	.488
Alabama.....	1	Mining and manufacturing area.....	2,641	67.97	2.67	2.071	.484
Iowa.....	2	Native white.....	7,404	67.96	2.61	2.106	.488
Alabama.....	2	Large Negro population.....	3,327	67.95	2.71	2.096	.489
Kentucky.....	2	Agricultural area.....	11,469	67.95	2.62	2.060	.484
South Dakota.....	2	Large Russian population.....	564	67.92	2.53	2.170	.495

TABLE 21.—Mean height, by sections; sections arranged in order of standing, with proportional weight and chest circumference (expiration) for each inch of height; also standard deviation for each height; first million draft recruits—Continued.

States.	Section.	Characteristics of sections.	Number of men measured.	Mean height.	Standard deviation (height).	Mean weight. Mean height.	Mean chest. Mean height.
				Inches.	Inches.	Pounds.	Inch.
Georgia.	2	Large Negro population.	10,078	67.91	2.66	2.077	0.490
Louisiana.	3	Rural, chiefly white population.	5,235	67.89	2.69	2.064	.488
New Mexico.	3	Noteworthy Mexican element.	540	67.89	2.73	2.048	.480
North Dakota.	3	Russian population.	2,005	67.87	2.61	2.172	.496
Washington.	2	Puget Sound, foreign white.	6,601	67.87	2.70	2.140	.492
Illinois.	7	Agricultural area.	5,442	67.86	2.58	2.092	.490
Do.	3	Agricultural area, native.	8,928	67.86	2.59	2.094	.487
West Virginia.	2	Agricultural region.	10,860	67.85	2.70	2.087	.491
Indiana.	3	Agricultural area, native stock.	18,743	67.84	2.56	2.083	.487
Colorado.	2	Russian population.	1,105	67.83	2.67	2.094	.490
Minnesota.	4	Urban area, "Twin Cities."	9,759	67.83	2.63	2.130	.489
Nevada.	All.	State undivided.	1,441	67.83	2.69	2.143	.497
Montana.	1	Mining area, foreign population.	5,117	67.82	2.65	2.150	.491
Wisconsin.	2	German population.	7,685	67.82	2.58	2.140	.495
Alabama.	5	Urban and suburban area.	481	67.81	2.56	2.086	.483
North Carolina.	4	Large Negro population.	4,570	67.79	2.72	2.087	.489
Wyoming.	All.	State undivided.	1,927	67.79	2.63	2.130	.492
Illinois.	8	Agricultural and manufacturing area.	2,451	67.77	2.63	2.110	.493
California.	1	Chiefly agricultural area.	17,793	67.75	2.63	2.137	.494
Ohio.	3	Agricultural area.	17,606	67.75	2.59	2.085	.489
Utah.	2	More densely populated.	2,781	67.75	2.56	2.105	.485
Louisiana.	1	Mississippi bottoms and upland, large Negro population.	4,074	67.73	2.63	2.073	.491
North Carolina.	6	Remainder of State.	744	67.73	2.63	2.076	.489
South Carolina.	2	Large Negro population.	3,975	67.72	2.77	2.100	.490
California.	4	Urban area.	7,428	67.71	2.64	2.069	.487
Do.	2	Mining area.	943	67.69	2.64	2.154	.499
Florida.	1	More white and maritime.	2,486	67.69	2.67	2.050	.486
North Carolina.	5	Island and peninsular area.	254	67.69	2.61	2.087	.491
Florida.	2	More Negro and rural population.	996	67.69	2.63	2.070	.490
Colorado.	5	Urban population.	1,644	67.68	2.69	2.070	.485
North Dakota.	1	Scandinavian and Canadian population.	1,132	67.67	2.56	2.159	.496
Colorado.	6	Austrian and Italian population.	1,222	67.65	2.71	2.060	.484
Minnesota.	3	Scandinavians and Finns.	3,520	67.65	2.66	2.170	.502
Utah.	3	Mining area.	563	67.65	2.78	2.127	.494
Michigan.	2	Prevalently native white population.	12,567	67.63	2.55	2.100	.493
District of Columbia.	All.	Undivided.	4,493	67.63	2.65	2.077	.482
Maine.	2	Native white stock, maritime.	828	67.60	2.59	2.091	.497
Illinois.	4	Largely German population.	4,238	67.59	2.64	2.115	.494
Do.	2	Mixed native and foreign population.	7,803	67.59	2.60	2.114	.494
Michigan.	5	Dutch and other foreign population.	2,892	67.51	2.65	2.080	.491
Missouri.	4	Urban area.	6,789	67.49	2.63	2.080	.488
Virginia.	2	Large Negro population.	5,352	67.46	2.72	2.077	.490
New York.	7	Agricultural and dairying.	6,465	67.45	2.64	2.098	.496
Florida.	4	Peninsular.	2,340	67.44	2.57	2.069	.491
Illinois.	1	Densely populated.	6,303	67.43	2.67	2.123	.495
New Mexico.	2	Native white population.	1,857	67.43	2.85	2.049	.493
Michigan.	3	Foreign population.	6,298	67.40	2.62	2.110	.497
Wisconsin.	4	Lake counties.	2,890	67.39	2.57	2.140	.500
Ohio.	4	Urban area.	3,557	67.39	2.90	2.104	.489
Maryland.	2	Peninsular area.	1,068	67.37	2.69	2.080	.490
Pennsylvania.	6	Rural area.	8,616	67.37	2.90	2.099	.494
Virginia.	1	Peninsular region and east shore.	2,896	67.34	2.73	2.091	.487
South Carolina.	3	Peninsular and rural areas.	3,804	67.33	2.64	2.060	.491
Ohio.	2	Intermediate.	14,443	67.31	2.74	2.096	.491
California.	5	Urban area.	7,189	67.28	2.61	2.137	.495
Maine.	1	English Canadian.	1,240	67.28	2.59	2.110	.497
Maryland.	3	Large white population.	2,683	67.26	2.48	2.080	.496
New Mexico.	1	Indian population.	293	67.26	2.90	2.068	.494
New Hampshire.	1	Mountainous area.	665	67.25	2.54	2.106	.501
Indiana.	1	Manufacturing.	3,614	67.22	2.64	2.113	.497
Florida.	3	Cuban, Spanish, West Indian population.	84	67.21	2.60	2.026	.487
Delaware.	All.	State undivided.	1,894	67.19	2.61	2.075	.492
New York.	5	Mountainous, Catskill region.	795	67.16	2.69	2.074	.493
Vermont.	All.	State undivided.	2,077	67.12	2.52	2.091	.496
Michigan.	1	Finnish population.	2,344	67.10	2.61	2.160	.501
Illinois.	5	Urban area.	33,919	67.09	2.67	2.099	.495
Maine.	3	French Canadian population.	1,247	67.07	2.58	2.080	.496
New York.	8	Mountainous, Adirondack area.	2,990	67.06	2.64	2.090	.497
Ohio.	1	Dense foreign population.	17,208	67.06	2.67	2.111	.495
New York.	4	Western manufacturing region.	14,222	67.01	2.67	2.096	.495
Wisconsin.	3	Urban and foreign stock.	4,527	66.99	2.56	2.100	.497
New York.	6	Urban area.	6,344	66.95	2.66	2.126	.498
Massachusetts.	4	do.	8,387	66.94	2.64	2.090	.494
Louisiana.	2	do.	3,047	66.93	2.66	2.056	.487
Maryland.	1	do.	5,411	66.93	2.69	2.100	.493
Massachusetts.	3	Peninsular region.	1,127	66.90	2.70	2.070	.491

TABLE 21.—*Mean height, by sections; sections arranged in order of standing, with proportional weight and chest circumference (expiration) for each inch of height; also standard deviation for each height; first million draft recruits—Continued.*

States.	Section.	Characteristics of sections.	Number of men measured.	Mean height.	Standard deviation (height).	Mean weight. Mean height.	Mean chest. Mean height.
				<i>Inches.</i>	<i>Inches.</i>	<i>Pounds.</i>	<i>Inch.</i>
New York.....	3	Eastern manufacturing region.....	5,150	66.67	2.66	2.092	0.498
New Hampshire.....	2	Manufacturing area.....	1,581	66.86	2.61	2.081	.493
Massachusetts.....	1	Mountainous area.....	1,373	66.85	2.67	2.070	.492
Michigan.....	4	Urban area.....	17,771	66.84	2.61	2.110	.490
New Jersey.....	3	Mountainous area plus Atlantic County.....	3,196	66.84	2.76	2.082	.501
Do.....	2	Plains section, rural.....	8,985	66.83	2.70	2.078	.499
Pennsylvania.....	4	Coal mining.....	4,827	66.80	2.69	2.109	.496
Connecticut.....	2	Manufacturing area.....	8,708	66.73	2.73	2.096	.499
Pennsylvania.....	2	Rural area, native stock.....	14,218	66.73	2.62	2.095	.497
New Jersey.....	1	Densely populated.....	17,795	66.72	2.74	2.078	.497
Connecticut.....	1	Prevaillingly agricultural and near metropolitan.....	4,877	66.67	2.68	2.094	.503
Massachusetts.....	2	Manufacturing center.....	18,447	66.67	2.67	2.070	.497
Pennsylvania.....	7	Allegheny County plus a small rural area.....	17,243	66.67	2.65	2.093	.495
Do.....	5	Manufacturing.....	8,892	66.66	2.69	2.116	.497
New York.....	1	Suburban territory.....	4,934	66.65	2.76	2.091	.497
Pennsylvania.....	1	Urban area.....	16,085	66.62	2.65	2.065	.494
Do.....	3	Mining area.....	7,305	66.55	2.57	2.105	.500
New York.....	2	Urban area densely populated.....	46,718	66.46	2.77	2.084	.498
Rhode Island.....	All.	State undivided.....	3,928	66.40	2.61	2.060	.494

To return to Table 21, the second entry from the top is Arkansas, Section 2. This section comprises about 97 per cent "native whites of native parentage"; that is, the old American southern white stock that lives in the hill country of northwestern Arkansas. The third section in order is Missouri 3, which included native whites of the Ozark region, a region practically contiguous with Arkansas 2, and composed of the same sort of men. In this section about 95 per cent of the population are of old white American stock, and fewer than 3 per thousand are Negroes. As has often been remarked, there is great resemblance in the general constitution of the population of the Ozark region in Missouri to that of the mountains of Kentucky and Tennessee.

The next three sections are in Texas, and two of these contain a considerable Negro population. As already pointed out, the proportion of immigrants from southeastern Europe in Texas is negligible. The State was settled chiefly by the tall southern stock. Next on the table comes Minnesota, Section 1. This comprises the northern counties, with prevaillingly Scandinavian population. We have already seen from the table of statures, page 68, that the Scandinavians are among the tallest races of Europe. This characteristic they have carried with them into Minnesota and have transmitted to their sons.

In the upper part of the table one finds certain other sections of interest, such as the mountain region of Tennessee (Section 3), the State of Oklahoma in general, recently populated by a selected lot of whites; Arkansas in general, including sections with a prevaillingly white population; Kansas, both sections, with the prevaillingly native, German, and Mennonite Russian population; Section 6 in the State of Illinois, so-called "Egypt," with a prevaillingly Negro population; and in general, those sections of the Southern States which have a large Negro population.

The bottom of the table is occupied by Rhode Island. The reason for this has already been pointed out. It is the presence of short races, Italian,

French Canadians, and Portuguese. Next to the bottom comes Section 2 of New York, comprising Greater New York, the most densely populated part of the Western Hemisphere. Here the mean stature is 66.46, or approximately 1 inch below the average for the United States. This low average stature of Greater New York is associated with a very high standard deviation, namely, 2.77. This indicates, as common observation confirms, that the stature of the population is exceptionally variable, comprising tall elements, selected from the most vigorous representatives of the northwestern and western races of Europe, including many of German and British stock, and, on the other hand, a very large proportion of representatives of the shortest races of Europe: Polish Jews, South Italians, and Greeks. The preponderance of the short races has resulted in bringing the average stature well toward the bottom of the list. The third section from the bottom is Pennsylvania 3. This comprises certain mining counties in the eastern part of the State. In the census of 1910 these included 4 per cent Italians, 21 per cent Austrians and Russians, 2.3 per cent Hungarians, and 42.5 per cent native whites of native parentage. The whites of native parentage were, however, in turn largely descended from the short races. Fourth from the bottom lies Pennsylvania 1, Philadelphia. This city comprises over 10 per cent Austrians and Russians (largely Jews), 5 per cent Italians, and only 37 per cent native whites of native parentage. Philadelphia approximates New York City in its possession of a large mixture of southeastern and eastern Europeans, and hence tends to fall near the bottom of the list. The next section is that of New York 1, which includes territory surrounding Greater New York, and whose population naturally is largely influenced by conditions in the great city. Then come certain manufacturing and mining populations. Next comes Massachusetts 2, a manufacturing center of that State outside of Boston. Reasons similar to those cited above account for the low position in the table of Section 1 of Connecticut and Section 1 of New Jersey (being densely populated portions of the State adjacent to Greater New York), and all other sections in Connecticut, Pennsylvania, and New Jersey. Michigan 4 comprises Detroit, and Section 2 of New Hampshire includes the manufacturing area of that State along the Merrimac River. The remaining sections of the table are those in which the population is less strikingly selected for great or small height or in which no great mixture of statures occurs.

In examining the table more generally, we find that there are very few sections with a large Negro population in which the stature is below the average. In fact, Virginia 2 is the only instance of this kind. On the other hand, there are relatively few mining areas in which the population is markedly above the average. The most striking of these are Alabama 1, the population tributary to Birmingham, which consists almost exclusively of native whites, 72 per cent, and Negroes, 26 per cent. Another instance is Montana 1 (67.82 inches), in which the foreign population is largely Irish and Scandinavian. California 2, with an average stature of 67.69 inches, has a high proportion of native whites of native parentage (47 per cent) and many English, Irish, and German, together with some Italians. In Utah 3, with a mean stature of 67.65 inches, the mining population included a large proportion of English. These have

doubtless migrated into the mining region from the more densely populated part of the State which has attracted to itself, through Mormon proselytizing, many representatives of the English and Scandinavian peoples. Those sections that include a large proportion of Germans and Scandinavians naturally lie in the upper part of the table. The great cities lie prevailing in the lower part of the table, not because city life tends to stunt growth but because cities attract the people from southeastern Europe, who remain in them instead of going upon the farms. On the other hand, the agricultural areas are occupied by persons of tall stature because the small races of southeastern Europe do not go to them in large numbers, whereas Scandinavians and many of the Germans do. Some of these conclusions will be strengthened and new ones will be gained by a study of the groups of similar sections shown in Table 22.

11. HIGH AND LOW STANDARD DEVIATIONS IN THE DIFFERENT SECTIONS.

Table 21 gives the standard deviation in stature for each section. For the United States as a whole the standard deviation in stature is 2.71 inches. Some of the highest standard deviations are: Ohio 4 (Cincinnati), 2.90; Pennsylvania 6 (a rural area in the northwestern part of the State), 2.90; New Mexico 1 (including many tuberculous whites, and also Indians and Mexicans), 2.90; New Mexico 2 (with more whites, but also Mexicans and Indians), 2.85; South Carolina 1 (mountain whites, but also a large colored population), 2.83. High variability is found in many large cities and suburban areas, for the reason suggested above; e. g., New York 2 (New York City), 2.77; New Jersey 1 (suburban), 2.74; New York 1 (suburban), 2.76. Low variabilities are found in Alaska, 2.30; Illinois 6 (31 per cent Negro), 2.38; South Dakota 3 (87 per cent Indian), 2.41; Missouri 3 (the Ozark region, with 94 per cent whites, prevailing tall), 2.48. Low variability implies homogeneity in the population; high variability, heterogeneity.

12. MEAN STATURE, BY GROUPS OF SIMILAR SECTIONS.

In Table 22 and Table IV the different sections are grouped so as to bring together those which have certain points of similarity. The mean stature and standard deviation have been worked out for these groups. The groups are arranged in order of the average stature. At the top of the list lies the group of mountain whites (group 12), including sections from the States of Kentucky, North and South Carolina, Tennessee, Virginia, and West Virginia. The average stature of men from these sections is 68.29, which is 0.8 inch above the average for the whole United States. Since these sections, except South Carolina, have a small proportion (less than 10 per cent) of Negroes, their exceptionally high average stature depends upon the physique of the mountain whites. These mountain whites, as pointed out, are, in the case of North Carolina and Kentucky, largely of Scotch origin. In the other States it is probable that there is a large mixture of Scotch and also some of the best physically developed of the stock that originally settled Virginia. The group is characterized by small variability, indicating that the population is fairly homogeneous in origin. The largest variability is found in South Carolina 1, in which the Negro element constitutes 31 per cent. The smallest variability (2.51) is found in the moun-

tain whites of Kentucky, comprising the smallest proportion of Negroes, 2.5 per cent.

The second group (group 3) in rank is that of the agricultural areas of the South that comprise a rather small proportion of Negroes. The proportion varies, however, in the different sections from 0.7 to 47.3 per cent. The average stature of this group is 68.18 inches, and all but one representative of this group are markedly above the average for the whole United States. The exception is Maryland 3, in the western part of the State, including nearly 75 per cent native whites and almost entirely native-born Americans. The variability of the group is low, namely, 2.64, as contrasted with 2.71 for the whole United States. The other sections obviously comprise exceptionally tall white men, and this is because of the racial stock which settled Alabama, Arkansas, North Carolina, Virginia, Kentucky, Tennessee, and Texas. They seem to have been a taller lot than settled New England. This can not be inferred from present day statistics, because of the recent immigrants, but from the statistics of the Civil War. According to Gould² (p. 125), the stature of native-born volunteers from New Hampshire was only 67.93; Vermont, 67.88; Rhode Island and Connecticut, 67.43; New York, 67.42; and Massachusetts, 67.41. To northern eyes, even at the time of the Civil War, southern whites appeared tall and lank.

The third section (group 14) in order includes four sparsely settled sections near the Mexican border. One of these includes 17 per cent Mexicans, another 14 per cent, the others less. The highest average stature is found in Texas 1, which comprises 17 per cent Mexicans. These are largely of Indian stock and the tall stature is no doubt due to the infusion of Indian blood. This appears also in the next group (group 13) of sections selected because of their large Indian population. In South Dakota 3, with 87 per cent of Indians, the average stature is 68.13, or 0.64 inch above the average.

The next group (group 20) in point of stature includes certain agricultural areas of the North, with a large German and Scandinavian population. The average stature of this group is 68.11, or 0.62 inch above the average for the whole United States. The tallest men are found in Minnesota 1, which includes 37 per cent of Scandinavians.

The next group (group 8) includes seven sparsely settled sections, mostly of the Southwest, excepting two sections of Wyoming and Oregon. In this group the average stature is 68.01, or 0.52 inch above the average. These sections include a large sprinkling of Indians and a very small percentage of recent immigrants.

The next group (group 15) includes two sections of native white persons of Scotch origin. In this group the mean height is about 0.5 inch above the average for the United States. Next comes a group (group 17) which includes a number of sections characterized by having 10 per cent or more Scandinavians. In this the mean height is 67.96, or 0.47 inch above the average. The tallest section is Minnesota 1, already referred to in another connection, with its mean stature of 68.44 inches. The next tallest section is Utah 1, including over 10 per cent Scandinavians and 8 per cent English, with a mean height of 68.16. The shortest people of this group are found in Michigan 1, 67.10 inches,

which includes a large Finnish population, and this helps to pull down the average.

The next group (group 9) includes three desert sections whose population includes many white people from other sections who suffer from tuberculosis. The average for the whole group is 67.86 inches, which is 0.37 inch above the average for the whole United States. Arizona 2, which includes Tucson, gives the tallest men of this group, 68.17 inches.

Passing now to the bottom of the table, we find that those sections in which the French Canadians (group 19) constitute 10 per cent or more of the population form the group with the least mean height, 66.67, or 0.82 inch below the average for the whole United States. Of these sections, Rhode Island, with 11.4 per cent French Canadians and a large number of Portuguese, is the shortest.

The next taller group (group 5) is the eastern manufacturing group, in which the mean height is 66.77, or 0.72 inch below the average. The sections of this group are characterized by a large proportion of the short races of southeastern Europe.

Next comes the group (group 6) including commuters. The sections of this group lie adjacent to the large manufacturing cities of the East and partake of many characteristics of their population.

The next taller group (group 16) is that which contains sections made up of about 10 per cent or more Russians. These are largely Russian Mennonites, chiefly engaged in mining. The section with the shortest stature is that of Pennsylvania 3, including a large mining population, while the tallest is Kansas 1, with 13 per cent Russians, engaged in agricultural pursuits. The differences in the stature of these populations are due chiefly to the difference in stature of the associated peoples.

Next above comes the group (group 22) in which the German and Austrian part of the population constitutes more than 15 per cent of the whole. Here the average stature is 67.27 inches, or only 0.22 inch below the average of all. When we select just those sections in which the Germans and Austrians constitute 20 per cent or more, the average stature, 67.41 inches, approaches still more closely the average stature of the whole country.

The mining group (group 7) comprises a population with just exactly the average stature of the whole United States and with a variability the same as that of the whole United States. The mining sections are for the most part regions of great admixture of various foreign nationalities.

It is noteworthy that those agricultural areas of the South which comprise 45 per cent or more of Negroes (group 4) have a shorter average stature than those agricultural areas of the South in which the proportion of Negroes is less. Since there is little difference in the average stature of white and colored, this result is to be ascribed to the fact that in the sections inhabited by the taller white man, there are fewer Negroes than in other sections of the South.

TABLE 22.—*Mean height, by groups of sections; groups arranged in order of standing, with proportional weight and chest circumference (expiration) for each inch of height; also the standard deviation for each height; first million draft recruits.*

[From Table IV, p. 427.]

[Height and chest in inches, and weight in pounds.]

Group No.	Description.	Number of men measured.	Mean height.	Standard deviation (height).	Mean weight. Mean height.	Mean chest. Mean height.
			Inches.	Inches.	Pounds.	Inch.
	Average for the United States.....	868,445	67.49	2.01	2.097	0.492
12	Mountain whites.....	21,254	68.29	2.57	2.05	.4862
3	Agricultural, native white, South.....	117,890	68.18	2.61	2.07	.4854
14	Mexican, sparsely settled.....	11,064	68.16	2.69	2.09	.4874
13	Indian, sparsely settled.....	10,038	68.12	2.61	2.08	.4864
20	German and Scandinavian, over 10 per cent.....	28,095	68.11	2.62	2.15	.4951
8	Sparsely settled, not more than 3 per square mile.....	16,165	68.01	2.63	2.13	.4929
15	Native whites of Scotch origin.....	13,522	68.00	2.61	2.06	.4844
17	Scandinavian, 10 per cent.....	51,009	67.96	2.63	2.15	.4952
9	Desert.....	6,121	67.87	2.72	2.09	.4917
4	Agricultural Negroes, 45 per cent plus.....	49,506	67.82	2.68	2.09	.4894
11	Mountain.....	17,101	67.72	2.68	2.11	.4921
2	Agricultural, mixed, foreign—native white.....	97,340	67.62	2.66	2.11	.4934
1	Agricultural, native white, North, native white over 75 per cent, North.....	66,885	67.60	2.63	2.09	.4900
7	Mining.....	35,730	67.49	2.72	2.11	.4929
18	Finn, 10 per cent.....	5,864	67.43	2.65	2.16	.5016
21	German and Austrian, over 20 per cent.....	38,962	67.41	2.69	2.13	.4955
10	Maritime.....	6,161	67.31	2.70	2.09	.4903
22	German and Austrian, over 15 per cent.....	126,994	67.27	2.72	2.12	.4954
16	Russian, 10 per cent plus.....	12,076	67.11	2.68	2.12	.4976
6	Commuter.....	29,032	66.86	2.75	2.09	.4970
5	Eastern manufacturing.....	81,718	66.77	2.70	2.09	.4970
19	French Canadian, 10 per cent.....	25,862	66.67	2.65	2.07	.4966

STATURE.

TABLE 23.—*Height distribution shown by groups of sections, first million draft recruits.*

SECTION A: ABSOLUTE NUMBERS.

Group No.	Description.	Num-ber meas-ured.	Height, in inches.																			79 and over.	
			59 and under.	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77		78
1	Agricultural, North, native white over 75 per cent.	66,885	245	178	415	975	2,062	3,705	5,990	8,415	10,181	10,444	8,956	6,709	4,254	2,456	1,100	472	154	79	22	15	26
2	Sweden and native white.	97,338	313	297	667	1,420	3,024	5,524	8,090	12,255	14,468	15,014	12,802	9,828	6,446	3,569	1,737	718	204	114	38	29	45
3	South.	117,548	316	235	510	997	2,311	4,436	8,243	12,605	16,212	18,685	17,469	14,170	9,857	6,007	3,100	1,358	688	247	75	59	59
4	per cent plus.	46,503	163	132	294	632	1,311	2,440	4,133	5,951	7,063	7,726	6,840	5,226	3,573	2,100	1,103	440	193	69	31	21	33
5	Commuters	81,718	395	425	1,161	2,548	4,541	6,997	9,681	11,721	12,171	11,047	8,613	5,779	3,452	1,797	794	336	118	51	13	15	19
6	Mining.	29,032	173	164	421	868	1,570	2,292	3,338	4,070	4,182	4,075	3,118	2,158	1,314	720	327	143	59	15	7	3	6
7	Sparsely settled, not more than 3 per square mile.	35,780	123	121	319	615	1,256	2,146	3,394	4,733	5,203	5,205	4,504	3,409	2,307	1,254	620	256	120	49	21	14	12
8	Desert.	16,166	36	28	66	185	352	718	1,277	1,796	2,312	2,553	2,281	1,849	1,318	724	355	164	82	25	10	7	7
9	Maritime.	6,121	22	12	35	91	159	302	505	716	888	894	681	548	449	288	141	66	25	12	2	3	2
10	Mountain.	17,090	55	47	119	260	477	858	1,537	2,071	2,577	2,654	2,271	1,763	1,224	703	326	135	59	19	10	8	6
11	Mountain.	21,264	55	34	60	135	340	777	1,394	2,087	3,014	3,480	3,165	2,990	1,909	1,117	579	238	108	44	16	5	7
12	Mountain.	10,098	29	11	40	92	210	396	712	1,103	1,427	1,629	1,411	1,225	845	462	239	104	52	16	3	7	5
13	Mountain.	10,779	30	17	48	110	241	456	799	1,126	1,473	1,647	1,402	1,113	890	619	293	119	61	22	7	1	10
14	Mountain.	13,522	42	41	69	138	295	597	1,019	1,495	1,947	2,151	1,962	1,532	1,045	628	299	126	53	27	7	1	10
15	Mountain.	12,076	61	55	136	263	527	863	1,313	1,769	2,181	2,317	2,073	1,596	1,095	632	317	157	63	25	6	8	6
16	Mountain.	51,009	171	95	296	528	1,174	2,250	3,992	5,941	7,441	8,163	7,273	5,755	3,809	2,254	1,126	526	217	70	30	19	24
17	Mountain.	5,864	18	23	46	103	218	383	567	769	874	898	768	547	381	180	83	41	23	3	1	1	1
18	Mountain.	25,862	101	112	378	845	1,532	2,321	3,210	3,840	4,383	4,478	3,581	2,732	1,036	508	217	78	31	16	4	6	7
19	German and Scandinavians, 10 per cent plus.	28,088	109	83	198	214	549	1,104	1,967	3,141	4,034	4,590	4,116	3,311	2,215	1,372	674	318	126	46	21	14	19
20	German and Austrians, 20 per cent plus.	38,963	197	110	332	695	1,416	2,358	3,808	5,138	6,954	8,065	6,949	5,209	3,359	2,090	1,309	609	261	104	38	16	11
21	German and Austrians, 15 per cent plus.	126,994	633	531	1,308	2,649	5,143	8,463	12,938	16,913	19,021	19,637	15,275	11,184	7,045	4,041	1,858	821	314	104	54	32	42
22	Total.	987,755	3,224	2,749	6,820	14,464	28,976	49,726	79,025	108,553	127,114	131,454	112,708	83,561	52,744	32,744	15,859	6,822	2,825	1,077	393	291	354

SECTION B. RATIOS PER 1,000

13. THE FREQUENCY DISTRIBUTION OF STATURES IN THE GROUPS OF SECTIONS.

The average is quite an inadequate method of indicating the composition of a population with reference to stature, for two populations which are very different in composition may have the same average. Thus one locality may have a large proportion of its men of average stature and another may be composed of nearly equal proportion of very short and very tall men. The average for the two populations might be alike. In Table 23 is given the distribution of statures for men from the different groups in ratios per 1,000 men of a given group. If we compare the ratios of men 61 inches tall in the different groups, we find that there is a large proportion of such exceedingly short men in those sections where French Canadians constitute 10 per cent or more of the population. Next in order come the commuter and eastern manufacturing groups with a large proportion of south Italians and Polish Jews. Then come the sections containing 10 per cent or more Russians, and after them the maritime groups.

The smallest proportion of 61-inch men is found among the mountain whites, the sections containing a large proportion of Indians, the districts characterized by 10 per cent or more Germans and Scandinavians, the southern white agricultural districts, and those sparsely settled areas which contain a good many Indians.

If, now, we turn to the very tall men, say 74 inches, we find that they are commonest in the southern white agricultural groups. Next come the groups of Germans and Scandinavians 10 per cent, then the mountain whites, the desert districts, and those containing a large proportion of Indians on reservations and elsewhere. The smallest proportion of these tall men is found in those sections occupied by 10 per cent or more French Canadians. Next come the eastern manufacturing and commuter sections, and next the group containing 10 per cent or more Russians. It is significant to note that, though the commuter group contains a slightly larger proportion of 61-inch men than the eastern manufacturing group, it contains proportionately very many more men of 72, 73, 74, 75 inches than does the eastern manufacturing group. This indicates that the commuter groups contain not only representatives of the races of eastern and southeastern Europe, who crowd the factories, but also men of the Nordic race, who are more largely leaders in affairs of the cities. In other words, the commuter groups are characterized by a deficiency of men of mediocre stature, 64-67 inches, as compared with the eastern manufacturing group.

A comparison of the southern white agricultural groups with agricultural groups containing 45 per cent or more Negroes shows in the latter a relative excess of short statures, 66 inches and less, and a relative deficiency of statures over 72 inches. This is partly associated with an inferiority in stature of Negroes over the average southern whites (Pl. XVIII). A comparison of the northern agricultural areas, one with over 73 per cent native whites and the other with larger admixture of foreigners, reveals an excess of men under 62 inches in the latter group and also an excess of men 69 inches and over. This shows that the agricultural areas containing a mixture of foreign and native whites are, as might be expected, much more variable in stature, just on account of the

variety of races present. The consequence is that the foreign and native groups have a smaller proportion of men of mediocre stature, 67-69 inches, than have the northern native agricultural groups.

If we compare the mining groups with the average of the whole country, we find they are characterized by an excess of short men, 66 inches and under. They are also characterized by a slight deficiency in very tall men, 71 inches or taller. A comparison of the mountain whites of the Alleghenies and the inhabitants of the mountain in other districts shows that the mountain whites have a relative deficiency in men under 67 inches and a marked excess of men with a stature over 69 inches.

Table 22 gives for the different groups of sections the mean height of the drafted men. This is a summary table of Table IV already discussed. In this table there are given the averages and standard deviations. A study of the standard deviations is significant, since this is the measure of variability.

The groups are arranged in descending order of the mean height. This brings out clearly, what has been indicated before, that the mountain whites and southern agriculturists, the Indians, and the Mexicans constitute the tallest part of our population and the groups containing many French Canadians, eastern manufacturing and commuter groups include the shortest of our population. The average height for the United States is, as we have repeatedly seen before, 67.49 inches, and the standard deviation is 2.71, which means that this is the center of gravity, as it were, of the variation above and below the average. When the variation above and below the average is slight, the standard deviation is small; when it is great, the standard deviation is large. Referring to Table 22, we find that the smallest standard deviation applies to the group of the mountain whites, this despite the fact that they are the tallest men, and in the tallest men one would expect a greater variability than in the shorter men, just because there are more inches of height to vary. The fact that the standard deviation is so small, 2.57 inches, indicates that we have to do here with a very homogeneous population. As a matter of fact, this group contains relatively few colored men; it is made up of the old British stock descended from the immigrants of colonial days. At the other extreme, the greatest variability is found in the commuter group. This, of course, is not a biological group at all, but a mixture of successful business men of the Nordic strain together with great numbers of recent immigrants who tend to settle in the seaboard cities and in their suburbs. The latter include, of course, the short races; the combination is the reason for the high standard deviation. Among other small standard deviations is that of the Indian group, 2.61, again containing a fairly homogeneous population. The Scandinavian and German-Scandinavian groups have likewise standard deviations less than the average. The same is true of the northern agricultural groups with their 73 per cent of native whites. The "sparsely settled group" has the same standard deviation. The groups of native whites of Scotch origin and the southern agricultural native white groups are others with small standard deviations. The groups with 45 per cent Negroes or more have a greater variability, owing to the mixture of races. Groups which have a variability above the average for the whole country are, the mining group, to which all kinds of men resort;

the desert group, which includes orientals and tall tuberculous cases from the other sections, and the German and Austrian group, 15 per cent.

Plates VI and VII show for each one of the groups of sections the distribution of the frequency of heights. In each of the charts the average for the whole United States is given so that the departure from this average in each of the different classes may be seen at a glance. It appears that the sections containing 10 per cent Finns have a distribution of height agreeing most closely with that of the United States as a whole. The group containing French Canadians shows the greatest departure from the United States as a whole, owing to the short stature of the people of this section. The Mexican group has a peculiar form, including a more mediocre and a taller subgroup. The taller subgroup is possibly due to the persons affected with pulmonary tuberculosis who are above the average in stature, together with tall Indians.

14. COMPARISON OF STATURE IN EIGHT EUROPEAN RACES OF MEN AT DEMOBILIZATION.


For the sake of completion there are added here the results of measurements of stature at demobilization (1919), in the case of eight European races. Table 25 gives the proportional distribution of different classes of stature. The order is given in the following table:

TABLE 24.—*Mean stature and standard deviation of each of the eight European races.*

Race.	Number exam- ined.	Mean stature.		Standard deviation.	
		Centi- meters.	Inches.	Centi- meters.	Inches.
Scotch.....	2,074	172.54	67.93	6.39	2.52
English.....	4,204	172.08	67.75	6.62	2.61
German.....	7,077	172.04	67.73	6.61	2.60
Irish.....	6,164	171.36	67.46	6.31	2.48
Polish.....	2,408	169.41	66.70	6.12	2.41
French.....	1,457	168.59	66.37	6.50	2.56
Hebrews.....	1,692	168.91	65.71	6.20	2.44
Italians.....	3,519	165.18	65.03	6.06	2.39

The standard deviation in stature is least in the Italians (probably because they are shortest) and greatest in the English (6.62 centimeters), indicating a great admixture of race statures in that people. Other high standard deviations are German, 6.61; French, 6.50. Next to the Italians (6.06) in stature-variability stand the Polish with a standard deviation of 6.12, and the Hebrews with 6.20. The Irish have a standard deviation of 6.31, and the Scotch of 6.39.




HEIGHT DISTRIBUTION BY GROUPS OF SECTIONS (P.)



FINE LINE CURVE DENOTES AVERAGE FOR U. S.

HEIGHT DISTRIBUTION BY GROUPS OF SECTIONS (P)

RATIOS PER 1000



STATURE.

FINE LINE CURVE DENOTES AVERAGE FOR U.S.
PLATE VII.

Corresponding to their tall stature, we find among the Scotch a larger proportion of men of stature class 172–173 centimeters than among any other people. Indeed, this constitutes the modal class for the Scotch. For the English, 170–171 centimeters is the modal class and the same holds for the German, Irish, Polish, and French. For Hebrews and Italians, however, the modal class is 164–165 centimeters. Under the English system, the modal stature of the Scotch is about 68 inches (172.72 centimeters), of the Italians 65 inches (165.10 centimeters).

TABLE 26.—*The mean stature in five color races, demobilization, 1919.*^a

Race.	Number of men examined.	Mean stature.	
		Inches.	Centimeters.
White.....	96,596	67.71	171.99
Negro.....	6,454	67.70	171.97
Indian.....	107	67.52	171.51
Chinese.....	23	67.37	171.11
Japanese.....	32	67.30	170.94

^a It will be noted, from examination of Tables 74, 75, 83, 84, 85, 86, 87, 88, 89, 90, and 91, that the average stature varies slightly for the white and Negro troops, the variation depending upon the number of men measured.

15. COMPARISON OF WHITE AND COLOR RACES.

A comparison of stature of white and color races is afforded by Table 26 taken from Tables 107 and LXXXIV and LXXXIX. It gives for the different color races the mean stature (in centimeters and inches). It appears that the stature of the white troops exceeds that of the negro by only 0.02 centimeter, or 0.01 inch. As Table LXXXIX shows, the colored troops are markedly more variable, having a standard deviation of 6.908 ± 0.041 , while that of the white troops is only 6.660 ± 0.010 (Table LXXXIV). As the difference is about six times the probable error, it is doubtless significant. The remaining three races are of decidedly shorter stature, and of them the Japanese are the shortest with a stature of 170.94 centimeters (67.30 inches). This figure is far greater than the average for Japanese given by Miwa as 159.3 centimeters (62.72 inches). We conclude, therefore, that the 32 Japanese included in our measurements were exceptionally tall representatives of that short race.

III. WEIGHT.

1. GENERAL DISCUSSION.

This measurement is of great importance in itself and of still greater importance in relation to stature. The varying relation of weight to stature is a measure of build. Build is of importance as an index of physical robustness and general health. Just how weight should be expressed in relation to stature has been much discussed and will be further elaborated in the fifth section, dealing with build. Different races differ in size and average build. In judging weight as an index of health one must, accordingly, take into account the

racial constitution of the individual and not apply the same absolute standard to Scotch, French, and Polish Jews.

The medical significance of weight is indicated by its deviation from the normal in various diseases. Table I gives the normal distribution of weights, as determined from 868,445 drafted men. This normal distribution for each stature is shown in Plates XI and XII. The mean weight of the whole population is 141.54 pounds (Table I). If, now, there be selected a group of men having a special disease, it is found that their mean weight frequently varies markedly from this average of all. Thus, it is evident at a glance that men affected with tuberculosis (Plate XXXIV) have a low weight, while men with varicose veins (Plate XXXV) and flat feet have a weight that is above the normal. Abnormal weight may, therefore, be symptomatic of these and other diseases.

Weight is of medico-military importance since a marked progressive change of weight under fairly uniform conditions of nutrition and exercise is an important diagnostic feature. Loss of weight under these circumstances suggests need of careful examination. Increase of weight requires careful consideration of possible endocrine glandular disturbance.

2. METHOD.

The method of measuring weight is fairly simple. When practicable, the subject should be weighed without clothing, since the weight of the latter and contents of the pockets can not be judged accurately enough for "practical" purposes. For recording in times of peace, any good beam scale, in which the weight has to be adjusted, may suffice; but for rapid work in mobilization examination, an automatic, springless scale (like that known under the trade name of "Toledo") has advantages over the beam scale, both for speed and elimination of error in reading the somewhat obscure markings on the beam scale. Care, of course, will be exercised that the subject is standing directly on the platform of the scale and free from contact with anything else.

Mean weight without relation to stature is of only limited significance; still it must be considered in army statistics, since the food requirements of a body of men are better indicated by weight than by any other single measure. The absolute weight of adults varies, of course, with stature. In the very careful measurements made at the Nutrition Laboratory of the Carnegie Institution of Washington (Harris and Benedict¹⁵ pp. 53, 57), the absolute weight of a series of men of which the average stature was also found is given (Table 27).

TABLE 27.—*Weights associated with statures with the standard deviations, and the coefficient of variation for each, in various classes of American males (Harris and Benedict¹⁵).*

Series.	N.	Average stature (centimeters).	Standard deviation.	Coefficient of variation.
Original series:				
Athletes.....	16	177.44±1.57	9.33±1.11	5.26±0.63
Others.....	62	171.82±0.58	6.79±0.41	3.95±.24
Whole series.....	89	172.45±.56	7.80±.39	4.53±.23
Gephart and Dubois selection.....	72	172.75±.56	6.98±.39	4.04±.23
First supplementary series.....	28	174.61±1.04	8.17±.74	4.68±.42
Original and first supplementary series.....	117	172.97±.50	7.94±.35	4.59±.20
Second supplementary series.....	19	172.95±.75	4.83±.53	2.79±.31
Other than Gephart and Dubois series.....	64	173.20±.69	8.21±.49	4.74±.28
All men of three series.....	136	172.96±.44	7.59±.44	4.39±.18

TABLE 27.—Weights associated with statures with the standard deviations, and the coefficient of variation for each, in various classes of American males (Harris and Benedict¹⁵)—Continued.

Series.	N	Average weight (kilograms).	Standard deviation.	Coefficient of variation.
Original series:				
Athletes.....	16	73.82±2.17	12.87±1.53	17.43±2.14
Others.....	62	63.03±0.77	9.92±0.55	14.32±0.88
Whole series.....	89	64.33±.77	10.73±.54	16.68±.87
Gephart and Dubois selection.....	72	63.35±.67	8.37±.47	13.22±.76
First supplementary series.....	28	62.69±1.34	10.48±.94	16.72±1.55
Original and first supplementary series.....	117	63.84±.67	10.69±.47	16.73±.76
Second supplementary series.....	19	65.06±1.13	7.30±.80	11.22±1.24
Other than Gephart and Dubois selection.....	64	64.96±1.02	12.04±.72	18.54±1.14
All men of three series.....	136	64.10±.60	10.30±.42	16.06±.67

3. MEAN WEIGHT.

The mean weight of the population of 868,445 accepted recruits of 1917–1918 considered in this paper is 141.54 pounds, or 64.26 kilograms.

Baxter¹ (Vol. I, pp. 51 and 52) states:

While the annals of recruiting contain copious details as to stature, the amount of information furnished upon the subject of weight is, for the most part, extremely meager. A principal reason for this is to be found in the fact that weight is not a regulated quality in any code of laws governing the enlistment of recruits. The circumference of chest thought to be indispensable as an accompanist to certain degrees of stature, is carefully laid down in the English regulations, but weight is not even mentioned. It is to be presumed that the matter is left to the discretion of the examining surgeon, with whom the decision as to the other qualities named might, it is thought, be also left with advantage. A due proportion in the weight is quite as essential in the soldier as a well-formed chest, and is of greater importance than lofty stature. In former times, when it was necessary to make use of a ramrod in loading a musket, men of a certain height were absolutely necessary for the service; but in these days of breech-loading arms, a man from 5 feet to 5 feet 4 inches in stature, and well proportioned in build and weight is, *ceteris paribus*, as serviceable a soldier as can be desired.

The instructions delivered to enrolling surgeons during the War of the Rebellion contained no injunctions as to weight. As a matter of course, it was duly considered in the estimate of "physical fitness" of the conscript; but, unfortunately for the purpose of the present investigation, it was not an obligatory process, and a large part of the returns contain no entry upon the subject. Some energetic officers, however, saw fit to make their work complete by adding the particulars of weight of the other details given and for their records the tables in which the weight is a component, were completed. It is reasonable to assume, as the information was voluntarily furnished, that it was procured with due accuracy. The men weighed were invariably quite naked.

However, the mean weight of recruits of 1917–1918 may be compared with such information as is given by Baxter for recruits of the Civil War. This is, for 6,359 white Americans, 136.05 pounds (61.77 kilograms), and for 377 colored natives, 141.67 pounds (64.32 kilograms). The weight of recruits of British, American, English, Irish, and German origins averaged somewhat under 137 pounds. This smaller weight of Civil War recruits is associated with their shorter mean stature and lower mean age.

Men at demobilization, 1919 (white and colored combined), weighed 144.89 pounds, an increase of 3.35 pounds over weight of recruits. The whites alone, at demobilization, 1919, weighed 144.67 pounds, whereas the whites at demobilization, Civil War, weighed 141.38 pounds. Here again appears the superiority of weight of the troops of 1919 as compared with those of 1865.

The position of males of the United States in relation to those of other countries is indicated by the accompanying table (Table 28) of average weights of adult males of different nationalities:

TABLE 28.—Average weights of adult males of various nationalities (*Martin*,⁵ p. 258).

Race.	Weight.	
	Kilo-grams.	Pounds.
Tribes of Central Africa.....	53.5	118
Japanese.....	54.5	120
Polish Jews.....	55.0	121
Roumanians.....	58.4	127
South Russian Jews.....	61.3	135
Upper Bavarians.....	63.2	139
French.....	64.9	143
Belgians.....	65.0	143
East Friesians.....	65.1	144
American mulattoes (Gould).....	65.8	145
Norwegians.....	66.0	146

4. THE FREQUENCY DISTRIBUTION OF WEIGHT.

(a) *Recruits 1917-1918*.—Table I (page 421) gives the absolute and relative frequencies of each of the different classes of weight into which the 868,445 recruits of 1917-1918 fall. Each of the classes has a range of 5 pounds. The modal class is seen to be 137 pounds, and this class includes 123 per 1,000 men. The frequency is very little less in classes 132 pounds and 142, but below and above these limits the frequency rapidly diminishes to 97 pounds on the one hand and to 202 pounds on the other. Below the lower limit of 97 pounds it is clear that there are proportionately few individuals, but at 202 pounds the upper limit is by no means reached, inasmuch as the class of 202 pounds and over comprises 5.4 per 1,000 persons.

(b) *At demobilization*.—Table 29 gives the relative frequency of the different classes of weight as found at demobilization in 1919. The weights are here taken in classes with a range of 10 pounds. For comparison, weights from Table I are given in the first column. The comparison reveals the fact that in veterans as compared with recruits, the mode shifts from 130-139 to 140-149. Of the veterans there were less than half as many of the weight 100-109 and there were also fewer of them of the weight 190-199. As a result of military training and warfare, either the lightest and heaviest men had been weeded out or else the light men had become heavier and the overweight men had lost weight; there was a tendency for the men to become more nearly uniform. However, the frequency of the modal class has not increased, but has fallen slightly, from 238 to 236. The average weight increased from 141.54 to 144.89 pounds.

TABLE 29.—*Frequency distribution of the various classes of weight (per mille) at mobilization, 1917-1918, and demobilization, 1919.*

Class range.	Distribution of weight per Mille.	
	At mobilization.	At demobilization.
90-99 pounds.....	0.21
100-109 pounds.....	11.27	5.206
110-119 pounds.....	72.42	41.602
120-129 pounds.....	170.76	132.605
130-139 pounds.....	238.32	222.553
140-149 pounds.....	217.25	235.943
150-159 pounds.....	144.85	177.611
160-169 pounds.....	79.29	104.061
170-179 pounds.....	36.37	48.003
180-189 pounds.....	15.96	20.587
190-199 pounds.....	7.92	7.246
200 pounds and over.....	5.40	4.561

Weight—

Mobilization; mean 141.54; standard deviation 17.42 pounds.

Demobilization; mean 144.89; standard deviation 17.06 pounds.

Plates XI and XII show, for the first million men, the relation of weight to stature. This is done by a series of 12 graphs, one for each class of stature from 62 to 73 inches, inclusive. On each graph is drawn in a faint line the normal distribution of weight for the entire population. This is for comparison with the curve drawn in heavier line showing the relative frequency of the different classes of weight for men of the respective stature. One learns from these graphs that, as is to be expected, the distribution of frequencies of weights in men 67 inches tall accords most closely with that of the whole population, although the weights of men with a stature of 67 inches are less variable than the weights of the entire population. As the stature diminishes from 67 inches the modal weight departs toward the lighter end of the series and as the stature increases from 67 inches, the modal weight departs toward the heavier end of the series.

5. THE STANDARD DEVIATION OF THE WEIGHT SERIES.

The standard deviation of the weight variability of the 873,159 recruits was 17.42 pounds, or 7,908 grams. The standard-deviation of weight of men at demobilization was 17.06 pounds. This means that the demobilized men were 2 per cent less variable in weight than the recruits. This result is doubtless due in part to the cutting off of the extremes by discharge for disability and in part by the equalizing effect of an approximately uniform good environment.

6. MEAN WEIGHT FOR THE DIFFERENT STATES.

Table 30 shows, by States arranged in order of size, the average weight at mobilization and, for comparison, at demobilization. From this table is compiled the next Table 31, in which the States are arranged in order of the differences of average weight of recruits and veterans.

TABLE 30.—Average weight, by States, at mobilization, 1917-18, and demobilization, 1919 (in pounds); States arranged in order of standing, with proportional weight for each inch of height, and chest circumference (expiration) for each pound of weight, for the first million draft recruits.

State.	Number of men measured.	Mean weight at demobilization.	Mean weight. Mean height.	Mean chest. Mean weight.	Mean weight. Mean chest.	Demobilization. (average weight).	Difference.
		<i>Pounds.</i>	<i>Pounds.</i>	<i>Inch.</i>	<i>Pounds.</i>	<i>Pounds.</i>	
Alaska.....	106	150.49	2.208	0.223	4.472	162.00	11.51
South Dakota.....	3,892	146.96	2.159	.228	4.382	152.19	5.23
North Dakota.....	6,444	146.95	2.163	.229	4.353	150.89	3.94
Minnesota.....	27,341	146.41	2.150	.230	4.354	151.37	4.96
Oregon.....	2,748	146.38	2.150	.228	4.368	148.32	1.94
Montana.....	11,648	146.32	2.151	.228	4.372	151.11	4.79
Washington.....	13,316	145.44	2.140	.230	4.347	148.39	2.95
Nevada.....	1,441	145.35	2.143	.232	4.307	149.50	4.15
Idaho.....	4,031	145.31	2.133	.232	4.307	150.97	5.66
Nebraska.....	10,774	144.74	2.126	.229	4.354	151.23	6.49
Iowa.....	19,537	144.72	2.126	.230	4.332	150.05	5.33
Wyoming.....	1,927	144.61	2.130	.231	4.332	148.44	3.83
Wisconsin.....	18,433	144.50	2.137	.232	4.307	147.87	3.37
California.....	35,461	143.98	2.127	.231	4.312	145.37	1.39
Kansas.....	9,671	143.72	2.107	.231	4.319	150.14	6.42
Mississippi.....	8,543	143.23	2.100	.231	4.330	147.54	4.31
Utah.....	4,568	143.13	2.109	.231	4.319	149.25	6.12
Arizona.....	3,850	143.04	2.099	.232	4.301	148.34	5.30
Oklahoma.....	19,429	142.35	2.084	.232	4.293	148.47	6.12
Texas.....	34,531	142.22	2.079	.232	4.307	147.36	5.14
Michigan.....	41,872	141.99	2.110	.235	4.258	145.07	3.08
Illinois.....	69,491	141.77	2.103	.234	4.260	145.42	3.65
Indiana.....	23,194	141.64	2.090	.233	4.274	144.78	3.14
West Virginia.....	12,367	141.53	2.085	.235	4.251	146.60	5.07
North Carolina.....	14,668	141.49	2.076	.235	4.255	146.17	4.68
Missouri.....	24,964	141.43	2.081	.233	4.275	145.70	4.27
Ohio.....	52,814	141.38	2.098	.234	4.268	144.45	3.07
Alabama.....	15,988	141.28	2.077	.233	4.277	144.79	3.51
Arkansas.....	10,111	141.28	2.071	.234	4.259	146.83	5.55
Colorado.....	6,635	141.06	2.069	.234	4.265	147.38	6.32
Maine.....	3,315	141.03	2.100	.237	4.221	142.97	1.94
Georgia.....	20,305	140.82	2.071	.235	4.241	143.94	3.12
District of Columbia.....	4,486	140.53	2.077	.232	4.303	140.80	.27
South Carolina.....	9,343	140.49	2.077	.235	4.244	144.89	4.40
Maryland.....	9,192	140.40	2.090	.236	4.240	141.81	1.41
Virginia.....	17,616	140.34	2.070	.236	4.230	146.05	5.71
Vermont.....	2,077	140.33	2.091	.238	4.198	136.95	-3.38
New Hampshire.....	2,240	140.33	2.095	.236	4.227	142.67	2.34
Tennessee.....	14,426	140.10	2.052	.235	4.249	145.54	5.44
Kentucky.....	15,502	140.00	2.058	.235	4.245	144.50	4.50
Connecticut.....	13,585	139.82	2.095	.239	4.182	141.05	1.23
Pennsylvania.....	77,186	139.72	2.094	.236	4.221	142.46	2.74
Louisiana.....	12,356	139.62	2.065	.236	4.221	146.41	6.79
New York.....	87,818	139.53	2.091	.238	4.200	140.43	.90
Delaware.....	1,491	139.45	2.075	.237	4.212	142.22	2.77
Florida.....	5,895	139.32	2.061	.237	4.214	141.50	2.18
New Jersey.....	29,958	138.81	2.079	.239	4.170	140.29	1.48
New Mexico.....	2,690	138.47	2.051	.239	4.178	144.00	5.53
Massachusetts.....	29,534	138.40	2.070	.239	4.181	139.74	1.34
Rhode Island.....	3,928	136.44	2.060	.241	4.156	140.19	3.75

Here, again, the numbers at demobilization from certain of the States and Territories, like Alaska, Nevada, and Wyoming, are so small that no stress must be laid upon the average that they show.

7. INCREASE IN WEIGHT AT DEMOBILIZATION OVER MOBILIZATION (TABLE 31).

For the United States as a whole, the troops show an increase in weight of 3.35 pounds. The greatest increase was found in men from Alaska, 11.51 pounds, where the number weighed was too small to furnish reliable averages. In the upper half of the list, showing an increase of 4 pounds or more, we find certain Southern States, such as Louisiana, with an average increase of 6.8 pounds; Oklahoma, 6.1 pounds; Virginia, 5.7; Arkansas, 5.6; Tennessee, 5.4; Texas, 5.1; West Virginia, 5.1; North Carolina, 4.7; Kentucky, 4.5; South Carolina, 4.4; and Mississippi, 4.3. On the other hand, the only Southern

States in which the troops showed an increase of less than 4 pounds were Alabama, 3.5; Georgia, 3.1; and Florida, 2.2. Evidently the tall and slender men were most improved in absolute weight by army life, partly because there was the greatest room for improvement. A striking increase in weight was shown also by troops from Nebraska, Kansas, Colorado, Utah, Iowa, and South Dakota, a group which (with the exception of Colorado) contains prevailingly agricultural States.

At the other end of the table stands last New Hampshire, with a decrease of over 3 pounds on the average in her troops at demobilization as compared with mobilization. As pointed out above, the numbers were small, and it is possible that the troops at demobilization were a specially selected lot. Next from the bottom stand men from the District of Columbia with practically no change. Then come men from New York, Connecticut, Massachusetts, Maryland, New Jersey, all States containing large cities in which the population is probably well nourished and free from parasitic diseases such as keep the weight of the southern men down. Consequently they show the least change as a result of the medical treatment and sanitary conditions in the Army.

TABLE 31.—States arranged in order of difference of weight at mobilization, 1917-1918, and demobilization, 1919.

State.	Difference.	State.	Difference.
	Pounds.		Pounds.
United States.....	3.35	North Dakota.....	3.94
Alaska.....	11.51	Wyoming.....	3.83
Louisiana.....	6.79	Illinois.....	3.65
Nebraska.....	6.49	Rhode Island.....	3.65
Kansas.....	6.42	Alabama.....	3.51
Colorado.....	6.32	Wisconsin.....	3.37
Utah.....	6.12	Indiana.....	3.14
Oklahoma.....	6.12	Georgia.....	3.12
Virginia.....	5.71	Michigan.....	3.08
Idaho.....	5.66	Ohio.....	3.07
Arkansas.....	5.55	Washington.....	2.95
New Mexico.....	5.53	Delaware.....	2.77
Tennessee.....	5.44	Pennsylvania.....	2.74
Iowa.....	5.33	Vermont.....	2.34
Arizona.....	5.30	Florida.....	2.18
South Dakota.....	5.23	Maine.....	1.94
Texas.....	5.14	Oregon.....	1.94
West Virginia.....	5.07	New Jersey.....	1.48
Minnesota.....	4.96	Maryland.....	1.41
Montana.....	4.79	California.....	1.39
North Carolina.....	4.68	Massachusetts.....	1.34
Kentucky.....	1.50	Connecticut.....	1.23
South Carolina.....	4.40	New York.....	.90
Mississippi.....	4.31	District of Columbia.....	.27
Missouri.....	4.27	New Hampshire.....	-3.38
Nevada.....	4.15		

TABLE 32.—Comparative view of mean height and mean weight of men from different States: (a) First million draft recruits (white and colored), 1917 and 1918; (b) 100,000 demobilized troops (white and colored), 1919; and (c) Civil War volunteer recruits (Gould), 1869, pp. 104 and 105.

States.	First million draft recruits (white and colored), 1917 and 1918.			100,000 demobilized troops (white and colored), 1919.				Civil War volunteer recruits (Gould), 1869, pp. 104 and 105.	
	Number of men measured.	Mean height.	Mean weight.	Number of men measured.	Mean weight.	Number of men measured.	Mean height.	Number of men measured.	Mean height.
		Inches.	Pounds.		Pounds.		Inches.		Inches.
Average for United States.	866,445	67.49	141.54	83,585	144.89	102,304	67.72	1,104,841	67.64
Alabama.....	15,988	68.01	141.28	383	144.79	1,930	68.57		
Alaska.....	106	68.15	150.49	12	162.00	13	69.43		
Arizona.....	3,850	68.13	143.04	125	148.34	130	68.33		
Arkansas.....	10,111	68.20	141.28	2,538	146.83	2,576	68.41		
California.....	35,461	67.67	143.98	414	145.37	481	67.91		
Colorado.....	6,635	68.15	141.06	208	147.38	225	68.12		
Connecticut.....	13,584	66.71	139.82	550	141.05	996	67.08		
Delaware.....	1,891	67.19	139.45	189	142.22	300	67.26		
District of Columbia.....	4,486	67.63	140.53	184	140.80	231	67.60		
Florida.....	5,895	67.58	139.32	140	141.50	1,022	68.22		
Georgia.....	20,305	67.99	140.82	446	143.94	3,397	68.51		
Idaho.....	4,031	68.10	145.31	153	150.97	164	68.26		
Illinois.....	604,491	67.40	141.77	6,462	145.42	6,687	67.65	188,507	67.97
Indiana.....	23,194	67.75	141.64	3,804	144.78	3,944	67.73	118,251	68.06
Iowa.....	19,537	68.04	144.72	1,543	150.05	1,609	68.28	29,604	68.13
Kansas.....	9,571	68.20	143.72	978	150.14	1,012	68.43		
Kentucky.....	15,502	68.02	140.00	2,753	144.50	2,921	68.13	23,993	68.16
Louisiana.....	12,356	67.60	139.62	1,726	146.41	2,070	67.96	2,582	66.83
Maine.....	3,315	67.28	141.03	209	142.97	663	67.17	52,314	64.12
Maryland.....	9,192	67.08	140.40	983	141.81	1,138	67.20	7,333	67.31
Massachusetts.....	29,534	66.76	138.40	1,320	139.74	4,782	66.77	40,855	67.05
Michigan.....	41,872	67.23	141.99	3,618	145.07	3,715	67.32	23,322	67.62
Minnesota.....	27,341	68.04	146.41	1,882	151.37	1,950	68.31	6,697	67.63
Mississippi.....	8,543	68.27	143.23	1,566	147.54	2,099	68.61		
Missouri.....	24,964	67.95	141.43	2,752	145.70	2,836	67.98	57,494	68.03
Montana.....	11,648	68.01	146.32	245	151.11	264	68.35		
Nebraska.....	10,774	68.06	144.74	791	151.23	819	68.44		
Nevada.....	1,441	67.83	145.35	16	149.50	18	67.91		
New Hampshire.....	2,240	66.97	140.33	94	136.95	413	66.80	26,821	67.40
New Jersey.....	29,958	66.77	138.81	3,103	140.29	3,180	66.93	18,875	66.58
New Mexico.....	2,690	67.50	138.47	221	144.00	229	67.82		
New York.....	87,818	66.72	139.53	8,965	140.43	9,207	66.92	188,008	67.09
North Carolina.....	14,668	68.15	141.49	570	146.17	1,815	68.22		
North Dakota.....	6,444	67.92	146.95	335	150.89	358	67.96		
Ohio.....	52,814	67.38	141.38	6,900	144.45	7,076	67.48	106,288	67.84
Oklahoma.....	19,429	68.28	142.35	2,274	148.47	2,310	68.44		
Oregon.....	3,825	68.09	146.38	1,049	148.32	1,069	68.38		
Pennsylvania.....	77,186	66.72	139.72	10,408	142.46	10,874	67.01	77,761	67.14
Rhode Island.....	3,928	68.40	136.44	209	140.19	403	66.54	41,305	67.09
South Carolina.....	9,343	67.64	140.49	205	144.89	828	68.32		
South Dakota.....	3,892	68.05	146.96	399	152.19	416	68.39		
Tennessee.....	14,426	68.27	140.10	781	145.54	2,807	68.61		
Texas.....	34,531	68.40	142.22	4,282	147.36	4,361	68.60		
Utah.....	4,568	67.85	143.13	99	149.25	104	68.19		
Vermont.....	2,077	67.12	140.33	93	142.67	446	67.19	24,062	67.61
Virginia.....	17,616	67.80	140.34	1,421	146.05	1,920	68.01		
Washington.....	13,316	67.96	145.44	1,984	148.39	2,025	68.38		
West Virginia.....	12,367	67.87	141.53	1,516	146.60	1,686	68.20	17,563	68.43
Wisconsin.....	18,433	67.60	144.50	2,616	147.87	2,675	67.79	51,302	67.65
Wyoming.....	1,927	67.79	144.61	71	148.44	80	68.16		

8. MEAN WEIGHT OF RECRUITS FROM THE DIFFERENT SECTIONS.

From the mean weight of 141.54 pounds for recruits from the United States at large, that of the various sections showed considerable deviation (see Table 33). Thus, excepting Alaska, the greatest average weight is found in South Dakota 3 (148.3 pounds), whose population is largely Indian. Next comes Minnesota 1, with a prevalingly Scandinavian population. Other high mean weights (of 147 or more) are found in Minnesota 2, North Dakota 3, and South Dakota 2. These contain (besides Scandinavians) Germans and Russian Mennonites. Sections with mean weights between 146 and 147 pounds are:

Montana 2, South Dakota 1, Oregon 1, Minnesota 3, North Dakota 1, and Washington 3.

The foregoing is a strikingly different list of sections from that standing at the top of Table 13, of mean stature; those were all southern sections. These comprise heavy men of only slightly greater stature than the average; those are tall and lank. The first southern section to come in as we proceed downward on Table 33 is Texas 5, with a large Negro population, mean weight 144.7 pounds.

At the bottom of the table of mean weights is Florida 3 (Key West), with a population that is prevailing Cuban, Spanish, and West Indian, racially small and living under insanitary conditions, with a mean weight of only 136.2 pounds. Next comes Rhode Island and then Philadelphia (137.6 pounds). New Orleans, with its numerous French, comes next highest; then the manufacturing section of northeast Massachusetts; then the part of New Mexico where many tuberculous patients dwell; and then, New York City with a mean weight of 138.5 pounds. Above lie numerous sections of the Middle and New England States—homes of men of small races. Relatively few southern sections are found in the lowest 10 per cent of the table; another of Florida's sections, however, is found here, possibly a consequence of hookworm and malaria. Chicago stands a little below the middle of the table (mean weight 140.9 pounds). Minneapolis and St. Paul stand in the upper third (144.2 pounds). Many other points of interest will be revealed from a study of the table.

TABLE 33.—Mean weight by sections; sections arranged in order of standing with proportional weight for each inch of height and chest circumference (expiration) for each pound of weight; also standard deviation for each weight; first million draft recruits.

State.	Section.	Characteristics of sections					
Average for United States.							
Alaska.....	All.	Undivided.....	106	150.49	14.95	2.208	.223
South Dakota....	3	Indian population.....	247	148.30	16.77	2.180	.228
Minnesota.....	1	Scandinavian population.....	6,461	148.28	16.61	2.170	.228
Minnesota.....	2	German and Scandinavian population..	7,601	147.64	17.31	2.170	.229
North Dakota....	3	Russian population.....	2,005	147.48	16.63	2.172	.228
South Dakota....	2	Large Russian population.....	594	147.22	16.15	2.170	.228
North Dakota....	2	Scandinavian population.....	3,307	146.93	16.23	2.150	.230
Montana.....	2	Sparsely settled, mountainous area.....	6,531	146.80	16.65	2.150	.229
South Dakota....	1	Dry farming area.....	3,051	146.80	18.54	2.100	.228
Oregon.....	1	Fairly densely populated.....	2,748	146.61	17.44	2.153	.228
Minnesota.....	3	Scandinavians and Finns.....	3,520	146.44	16.64	2.170	.232
North Dakota....	1	Scandinavian and Canadian population	1,132	146.10	16.20	2.150	.230
Washington.....	3	Mountainous area.....	1,537	146.07	16.29	2.142	.230
California.....	2	Mining area.....	943	145.94	16.85	2.134	.231
Oregon.....	2	Columbia River Valley and coastal dry plain, sparsely populated.....	1,077	145.82	16.64	2.140	.229
Montana.....	1	Mining area, foreign population.....	5,117	145.70	16.65	2.150	.229
Nebraska.....	2	German, Austrian, and Russian stocks.	3,145	145.70	17.73	2.136	.228
Iowa.....	1	Foreign white, German and Scandinavian.	12,136	145.67	17.10	2.139	.230
Washington.....	1	Coastal region plus eastern counties..	5,176	145.50	17.10	2.136	.230
Nevada.....	All.	State undivided, sparse population....	1,441	145.35	17.11	2.143	.232
Idaho.....	All.	State undivided.....	4,034	145.31	16.29	2.133	.232
Washington.....	2	Puget Sound, foreign white.....	6,601	145.25	17.28	2.140	.230
Wisconsin.....	1	Scandinavian and German population..	3,297	145.13	16.93	2.130	.232
Kansas.....	1	Russian population.....	1,057	144.95	17.44	2.122	.229
Wisconsin.....	2	German population.....	7,685	144.94	17.13	2.140	.232

TABLE 33.—*Mean weight by sections; sections arranged in order of standing with proportional weight for each inch of height and chest circumference (expiration) for each pound of weight; also standard deviation for each weight; first million draft recruits—Continued.*

State.	Section.	Characteristics of sections.	Number of men measured.	Mean weight.	Standard deviation (weight).	Mean weight. Mean height.	Mean chest. Mean weight.
				<i>Pounds.</i>	<i>Pounds.</i>	<i>Pounds.</i>	<i>Inch.</i>
Virginia.....	3	Native rural region.....	3,866	140.77	16.28	2.066	0.237
Maryland.....	3	Large white population.....	2,683	140.76	16.48	2.090	.237
Alabama.....	3	Large native white population.....	2,670	140.71	15.84	2.062	.235
North Carolina.....	3	Remainder of State.....	743	140.63	16.35	2.076	.235
Wisconsin.....	6	Urban and foreign stock.....	4,527	140.62	18.04	2.100	.237
Georgia.....	1	Mixed population, native white predominating.....	10,248	140.55	16.71	2.064	.235
District of Columbia.....	All.	District undivided.....	4,493	140.53	18.03	2.077	.232
Louisiana.....	1	Mississippi bottoms and upland, large Negro population.....	4,074	140.47	16.55	2.073	.236
New York.....	4	Western manufacturing region.....	14,220	140.46	17.49	2.096	.236
Missouri.....	4	Urban area.....	6,789	140.44	18.40	2.080	.235
South Carolina.....	1	Native white.....	1,564	140.42	16.72	2.060	.235
Vermont.....	All.	State undivided.....	2,073	140.33	16.43	2.091	.238
Maryland.....	1	Urban area.....	5,441	140.29	17.49	2.100	.235
New York.....	8	Mountainous, Adirondack area.....	2,990	140.21	16.71	2.090	.237
Alabama.....	5	Urban and suburban area.....	481	140.16	16.61	2.066	.234
Colorado.....	5	Urban population.....	1,644	140.16	16.26	2.070	.234
Florida.....	2	More Negro and rural population.....	996	140.14	17.53	2.070	.236
Louisiana.....	3	Rural, chiefly white population.....	5,235	140.13	16.22	2.064	.236
Pennsylvania.....	3	Mining area.....	7,305	140.10	17.17	2.105	.237
Virginia.....	2	Large Negro population.....	5,352	140.10	16.43	2.077	.236
Kentucky.....	2	Agricultural area.....	11,469	140.02	16.76	2.060	.234
Tennessee.....	3	Mountainous region.....	5,900	140.02	16.43	2.050	.235
Virginia.....	4	Mountain, white.....	5,512	140.02	15.94	2.055	.238
Maryland.....	2	Peninsular area.....	1,068	140.01	16.56	2.080	.236
New York.....	3	Eastern manufacturing region.....	5,150	139.94	17.50	2.062	.238
Connecticut.....	2	Manufacturing area.....	8,708	139.92	18.20	2.096	.238
Kentucky.....	1	Mountainous area, native white.....	4,033	139.92	15.26	2.051	.237
Pennsylvania.....	2	Rural area, native stock.....	14,207	139.83	17.06	2.065	.237
Maine.....	3	French Canadian population.....	1,247	139.71	17.21	2.080	.238
Connecticut.....	1	Prevalently agricultural and near metropolitan.....	4,876	139.65	17.73	2.094	.240
Florida.....	4	Peninsular.....	2,340	139.60	16.85	2.099	.237
Massachusetts.....	4	Urban area.....	8,587	139.59	17.65	2.090	.237
Pennsylvania.....	7	Allegheny County plus a small rural area.....	17,243	139.55	17.56	2.093	.236
Tennessee.....	2	Agricultural region.....	6,308	139.50	16.33	2.040	.237
Delaware.....	All.	State undivided.....	1,894	139.45	17.06	2.075	.237
Colorado.....	6	Austrian and Italian population.....	1,222	139.40	16.10	2.060	.235
New York.....	1	Suburban territory.....	4,934	139.39	17.09	2.091	.238
Do.....	5	Mountainous, Catskill region.....	795	139.30	16.74	2.074	.238
Illinois.....	6	Negro population (Egypt).....	409	139.27	16.39	2.043	.236
New Jersey.....	3	Mountainous area plus Atlantic County.....	3,195	139.18	16.13	2.062	.240
New Hampshire.....	2	Manufacturing area.....	1,575	139.13	17.55	2.081	.237
New Mexico.....	1	Indian population.....	293	139.12	18.49	2.068	.239
Do.....	3	Noteworthy Mexican element.....	540	139.01	17.36	2.048	.234
New Jersey.....	2	Plains section, rural.....	8,968	138.92	17.34	2.078	.240
South Carolina.....	3	Peninsular and rural areas.....	3,804	138.90	15.70	2.060	.238
Florida.....	1	More white and maritime.....	2,846	138.83	16.46	2.050	.237
Massachusetts.....	3	Peninsular region.....	1,127	138.70	16.76	2.070	.237
New Jersey.....	1	Densely populated.....	17,795	138.69	17.59	2.078	.239
Massachusetts.....	1	Mountainous area.....	1,373	138.52	17.13	2.070	.237
New York.....	2	Urban area, densely populated.....	46,718	138.50	18.29	2.094	.239
New Mexico.....	2	Native white population.....	1,857	138.20	16.42	2.049	.240
Massachusetts.....	2	Manufacturing center.....	18,447	137.82	17.25	2.070	.241
Louisiana.....	2	Urban area.....	3,047	137.62	16.55	2.056	.237
Pennsylvania.....	1	Urban area.....	16,065	137.61	17.48	2.065	.239
Rhode Island.....	All.	State undivided.....	3,928	136.44	17.69	2.060	.241
Florida.....	3	Cuban, Spanish, West Indian population.....	84	136.23	16.98	2.026	.240

9. MEAN WEIGHT FOR THE DIFFERENT GROUPS.

Tables 35, Section A, gives the absolute distribution of frequency of weights of men found in the 22 groups. The ratios per 1,000 are given in Table 35, Section B. The tables show that the lowest average weights are found in those sections containing 10 per cent or more of French Canadians (group 19) and in the eastern manufacturing group (group 5) and commuter group (group 6).

The higher weights, on the other hand, of 180 pounds or more, are found especially in the group (group 20) containing 10 per cent or more of Germans and Scandinavians, in group 17 containing 10 per cent or more of Scandinavians alone, in group 18, containing 10 per cent or more of Finns, in the sparsely settled and Mexican groups (group 8 and group 14), and in those containing 20 per cent or more of Germans and Austrians (group 21). The largest proportion of extremely heavy men is found in the sections with 10 per cent or more of Germans and Scandinavians and 20 per cent or more of Germans and Austrians. If we compare now the southern white agricultural and Negro agricultural groups, we find relatively little difference except that the white group contains proportionately fewer men under 115 pounds and over 140 pounds. Of these men, however, there is an excess in the white agricultural groups with a weight of 185 pounds and over. Apparently obese Negroes are less common than obese whites.

If we compare the northern native white agricultural groups with those of mixed population, we find an excess of underweight or low weight in the former and a slight excess of heavy weights in the latter. However, of extremely obese men, 190 pounds or over, there is an excess in the native white group.

Comparing the eastern manufacturing with the commuter groups we find an excess of thin men in the former and of men of 155 pounds and more in the latter. There is, however, a very slight excess of extremely obese men in the eastern manufacturing over the commuter groups. Comparing the mountain whites with inhabitants of other mountainous areas, there is an excess of thin men in the mountain whites and a deficiency of heavy men. The native whites of Scotch origin show a slight excess of low-weight men, and a corresponding deficiency of heavy men. And the French Canadian group, as might be expected, shows a very large excess of slight men and a corresponding deficiency of heavy men.

Table 34 gives the mean weights and standard deviations for the groups as well as relative stature and chest between them. It may be worth while to consider the significance of certain extremes in the standard deviations. Thus in weight, we find the highest standard deviation, or the greatest variability, in the sections containing 20 per cent or more of Germans and Austrians. Such sections are characterized by a mixture of strains dissimilar in weight. The smallest standard deviation in weight is that of the mountain whites, obviously a homogeneous people. Other high standard deviations, 17.70 or over, are found in the eastern manufacturing group and in the commuter group, of which the significance has already been discussed; also in the group containing Germans and Austrians, 15 per cent. Of groups with small standard deviations, 16.90 or under, we have the sections occupied by 10 per cent or more of Finns, mountain populations aside from the southern Alleghenies, the mining sections, the southern white agricultural sections, the maritime sections, and the Negro agricultural sections. These are more homogeneous in their racial characteristics than the other groups.

The relation between the distribution of weights in the populations of the different groups, or sections, as compared with their distribution in recruits in general is shown in the graphs of Plates VIII and IX. A study of these

curves reveals the following facts: Groups containing over 10 per cent of Scandinavians have a population of men strikingly heavier than recruits at large. Thus there is a deficiency of men under 140 pounds and an excess of men over 140 pounds in weight. The modal weight of Scandinavian groups is 5 pounds above that of recruits in general. This is, of course, associated with the excess height of Scandinavians.

The groups of sections having 10 per cent or more of Finns reveals a population that is much heavier than the average. There is a deficiency of men under 135 pounds and an excess of men weighing 140 pounds or more, and this despite the fact that in these same sections the distribution of statures is essentially that of the whole population of recruits. This shows then that in those sections which are characterized by an excess of Finns we have men of exceptionally robust build, and it is well known from other sources of information that the Finns, like most races of the extreme north, tend to put on weight and are of heavy build.

On the other hand, the groups containing 10 per cent or more of French Canadians are characterized by a great excess of men with a weight under 135 pounds and a deficiency of men above 135 pounds. The mode is indeed shifted from 137 pounds to about 132 pounds. This low weight of the groups with a large proportion of French Canadians is associated with the small stature of the population of these groups. These groups therefore contain an excess of population of small size.

The populations of the groups containing native whites of Scotch origin are peculiar in this, that they have an excess of men under average weight, while at the same time they have an excess of men over average stature. Thus, as the graphs in Plate IX show, the modal weight is clearly below that of the population of recruits in general and the group is less variable than that of recruits in general, which suggests that we have to do here with a racial characteristic. We may say then that, from the evidence of these graphs, the Scotch groups are characterized by an excess of tall, gaunt men.

The remaining groups show less striking deviations from the average of all recruits. The groups with an excess of Austrians and Germans are somewhat heavier than the average and the same is true of the groups containing 10 per cent or more of Russians. The groups containing nearly half Negroes are slightly above the average in weight, much more than the southern agricultural groups containing a larger proportion of native whites. Thus the Negro groups appear better nourished than those groups that contain an excess of native whites. This is possibly due to the greater resistance on the part of the Negroes to those parasites that tend to keep down the weight.

WEIGHT DISTRIBUTION BY GROUPS OF SECTIONS (P.)

RATIOS PER 1000

1 2

M

3
ADMC NATIVE WHITE SOUTH4
ADMC. NEGROES. 45 % +

PM

5
MEDIAN6
MEDIAN

PM

FINE LINE CURVE DENOTES AVERAGE FOR U. S.
PLATE VIII.

WEIGHT DISTRIBUTION BY GROUPS OF SECTIONS (P)

RATIOS PER 1000

1 2

3

4

5

FRENCH CANADIAN 10 %

6

WEIGHT.

FINE LINE CURVE DENOTES AVERAGE FOR U. S.
PLATE IX.

TABLE 34.—Mean weight by groups of sections; groups arranged in order of standing with proportional weight for each inch of height and chest circumference (expiration), for each pound of weight; also the standard deviation for each weight; first million draft recruits.

[From Table V, p. 434.]

Group No.	Description.	Number of men measured.	Mean weight.	Standard deviation (weight).	Mean weight. Mean height.	Mean chest. Mean weight.	Mean weight. Mean chest.
	Average for the United States.....	868,445	Pounds. 141.54	Pounds. 17.42	Pounds. 2.007	Inch. 0.234	Pounds. 4.200
20	German and Scandinavian, over 10 per cent.....	28,095	146.66	17.00	2.15	.230	4.350
17	Scandinavian, 10 per cent.....	51,009	146.13	16.99	2.15	.230	4.343
18	Finn, 10 per cent.....	5,864	145.76	16.86	2.16	.232	4.311
8	Sparsely settled, not more than 3 per square mile.....	16,165	144.84	16.93	2.13	.232	4.320
21	German and Austrian, over 20 per cent.....	38,962	143.27	18.05	2.13	.233	4.287
11	Mountain.....	17,103	142.97	16.76	2.11	.233	4.290
2	Agricultural, mixed foreign, native white.....	97,340	142.79	17.28	2.11	.234	4.277
22	German and Austrian, over 15 per cent.....	126,994	142.31	17.73	2.12	.234	4.271
16	Russian, 10 per cent plus.....	12,076	142.30	17.21	2.12	.235	4.264
14	Mexican, sparsely settled.....	10,779	142.18	17.36	2.09	.234	4.283
7	Mining.....	35,730	142.25	16.86	2.11	.234	4.282
9	Desert.....	6,121	142.08	17.23	2.09	.235	4.256
13	Indian, sparsely settled.....	10,088	141.89	16.91	2.08	.234	4.283
4	Agricultural Negroes, 45 per cent plus.....	49,503	141.61	16.64	2.09	.234	4.266
3	Agricultural, native white, South.....	117,548	141.44	16.83	2.07	.234	4.274
1	Agricultural, North, native white over 73 per cent.....	66,885	141.32	17.45	2.09	.234	4.270
10	Maritime.....	6,161	140.38	16.86	2.09	.235	4.255
12	Mountain whites.....	21,254	140.24	16.05	2.05	.237	4.225
15	Native whites of Scotch origin.....	13,522	140.26	16.77	2.06	.236	4.260
6	Commuter.....	29,082	139.79	17.66	2.09	.238	4.205
5	Eastern manufacturing.....	81,718	139.48	17.71	2.09	.238	4.204
19	French Canadian, 10 per cent.....	25,862	137.88	17.38	2.07	.240	4.172

TABLE 35.—Weight distribution shown by groups of sections, first million draft recruits.

SECTION A: ABSOLUTE NUMBERS.

Group No.	Description.	Num- ber meas- ured.	Weight, in pounds.																200 and over					
			95- 99	100- 104	105- 109	110- 114	115- 119	120- 124	125- 129	130- 134	135- 139	140- 144	145- 149	150- 154	155- 159	160- 164	165- 169	170- 174		175- 179	180- 184	185- 189	190- 194	195- 199
1	Agricultural, North, native white over 73 per cent.	66,885	144	506	1,648	8,212	4,863	6,093	7,979	8,538	7,789	6,634	5,315	3,934	2,927	2,156	1,437	925	615	426	357	246	307	
2	Agricultural, foreign and native white over 73 per cent.	97,840	12	207	660	1,951	3,899	6,234	8,803	10,718	12,104	11,542	10,504	8,728	6,455	4,916	3,549	2,333	1,556	962	655	444	350	
3	Agricultural, native white, South.	117,546	18	259	841	2,508	5,149	8,371	11,547	14,190	15,232	14,171	12,373	9,894	7,207	5,175	3,657	2,352	1,549	947	688	459	340	
4	Agricultural, Negroes, 45 per cent plus	40,568	13	125	363	1,119	2,144	3,371	4,833	5,618	6,109	6,077	5,287	4,444	3,128	2,356	1,670	1,022	647	436	251	145	105	
5	Eastern manufacturing	81,718	14	325	1,028	2,736	5,188	7,060	8,810	9,792	9,816	9,926	7,486	5,898	4,233	3,169	2,346	1,544	1,078	703	504	375	312	
6	Communities	29,032	1	107	356	982	1,768	2,452	3,033	3,464	3,509	3,179	2,706	2,080	1,590	1,179	889	555	393	249	187	140	115	
7	Mining	85,720	5	86	256	762	1,486	2,397	3,223	4,004	4,359	4,380	3,624	3,226	2,868	1,722	1,334	786	528	371	183	131	96	
8	Sparsely settled, not more than 3 per square mile.	16,165	21	72	205	462	850	1,271	1,710	1,934	2,003	1,872	1,638	1,506	1,271	906	722	436	311	196	106	82	57	
9	Desert.	6,121	19	56	133	235	423	569	722	764	794	747	607	454	307	201	173	115	92	63	34	27	13	
10	Maritime.	6,161	4	17	58	147	307	508	617	755	797	747	607	454	307	201	173	115	92	63	34	27	13	
11	Mountain.	17,099	25	80	282	674	1,090	1,569	1,938	2,141	2,060	1,854	1,560	1,197	847	547	404	268	181	97	64	45	19	
12	Mountain whites.	21,254	6	30	133	470	1,012	1,563	2,313	2,702	2,760	2,525	2,273	1,718	1,245	851	574	351	233	137	85	53	49	
13	Mountain whites.	10,038	1	43	52	186	405	727	913	1,206	1,296	1,268	1,065	769	632	478	335	238	121	77	52	39	37	
14	Mountain whites.	10,779	1	33	97	257	464	738	978	1,245	1,296	1,268	1,065	769	632	478	335	238	121	77	52	39	37	
15	Mountain whites.	13,522	1	24	116	318	601	1,096	1,450	1,718	1,758	1,525	1,306	1,033	751	546	384	259	161	107	81	53	43	
16	Mountain whites.	12,076	1	30	94	271	538	799	1,091	1,399	1,423	1,425	1,265	1,057	826	622	436	274	193	130	90	65	34	
17	Mountain whites.	51,008	4	44	160	541	1,264	2,388	3,626	4,917	6,078	6,262	6,003	5,271	4,148	3,266	2,461	1,606	1,006	691	447	271	211	
18	Mountain whites.	5,864	2	20	50	145	283	453	578	727	708	686	594	471	344	240	168	108	74	49	33	18	8	
19	Mountain whites.	25,892	5	123	349	1,044	1,888	2,628	3,404	3,181	3,085	2,618	2,215	1,724	1,290	852	632	427	328	201	131	109	87	
20	German and Scandinavians, 10 per cent plus.	26,005	1	24	80	286	632	1,228	1,856	2,682	3,270	3,449	3,429	2,984	2,267	1,876	1,417	948	565	365	263	144	122	
21	German and Austrians, 30 per cent plus.	38,982	6	106	312	800	1,674	2,499	3,367	4,231	4,678	4,515	4,141	3,354	2,499	2,011	1,530	997	655	467	296	211	171	
22	German and Austrians, 15 per cent plus.	126,994	42	386	1,138	2,912	5,876	8,762	11,608	14,114	15,265	14,707	13,139	10,734	8,245	6,141	4,729	2,992	2,007	1,315	900	625	465	
	Total	967,757	141	2,130	6,869	19,583	39,227	60,364	81,738	94,933	107,001	101,890	90,442	72,816	55,363	41,327	30,533	19,636	12,973	8,505	5,605	3,870	2,961	4,749

10. COMPARISON OF WEIGHT IN EIGHT EUROPEAN RACES OF MEN AT DEMOBILIZATION.

For the sake of completion there are added here the results of weights taken at demobilization, 1919, in the case of eight European races. Table 37 gives the proportional distribution of different classes of weight. The order of weight is as follows:

TABLE 36.—Mean weight and standard deviation in each of eight European races.

Race.	Number measured.	Mean weight.		Standard deviation.	
		Kilos.	Pounds.	Kilos.	Pounds.
German.....	6,767	67.22	148.20	7.72	17.02
Polish.....	2,225	66.05	145.62	6.95	15.29
English.....	3,608	65.76	144.98	7.57	17.35
Scotch.....	1,821	65.74	144.93	7.90	17.41
Irish.....	4,907	64.84	142.96	7.75	17.08
French.....	746	64.48	142.16	7.27	16.03
Italian.....	3,075	62.59	137.99	7.03	15.49
Hebrew.....	1,531	62.53	137.85	7.27	16.03

TABLE 37.—Comparative frequency distribution of weight in each of eight races, demobilization.

SECTION A: ABSOLUTE NUMBERS.

Race.	Total.	Weight, in pounds.										
		100-109	110-119	120-129	130-139	140-149	150-159	160-169	170-179	180-189	190-199	200 and over.
English.....	3,608	24	158	538	790	808	618	377	178	74	31	12
Scotch.....	1,821	12	79	254	436	404	308	175	89	37	19	8
Irish.....	4,907	34	259	796	1,233	1,151	700	407	175	89	35	28
German.....	6,767	16	183	678	1,406	1,589	1,351	867	399	177	56	46
French.....	746	7	39	123	181	183	122	59	19	5	5	3
Italian.....	3,075	44	274	664	845	631	362	154	65	30	5	1
Polish.....	2,225	4	64	245	518	599	444	212	88	36	12	3
Hebrew.....	1,531	24	144	341	402	325	168	71	34	11	5	6
Number measured.	24,680	165	1,200	3,639	5,811	5,690	4,073	2,322	1,047	459	167	107
Not measured.	3,990											
Total.	28,670											

SECTION B: PROPORTIONAL RATIOS PER 1,000.

Race.	Total.	Weight, in pounds.											
		100-109	110-119	120-129	130-139	140-149	150-159	160-169	170-179	180-189	190-199	200and over.	Total.
English.....	3,608	6.65	43.80	149.11	218.96	223.94	171.29	104.49	49.34	20.51	8.59	3.33	1,000
Scotch.....	1,821	6.59	43.39	139.49	239.44	221.85	169.13	98.11	48.88	20.32	10.43	4.39	1,000
Irish.....	4,907	6.93	52.78	162.22	251.29	234.56	142.65	82.94	35.66	18.14	7.13	5.71	1,000
German.....	6,767	2.36	27.04	100.19	207.79	234.81	199.65	128.12	58.96	26.16	8.13	6.80	1,000
French.....	746	9.38	52.28	164.89	242.61	245.31	163.55	79.09	25.47	6.70	6.70	4.02	1,000
Italian.....	3,075	14.31	99.10	215.94	274.80	205.21	117.72	50.08	21.14	9.76	1.63	.33	1,000
Polish.....	2,225	1.80	28.76	110.12	232.81	289.21	199.56	95.28	39.55	16.18	5.39	1.35	1,000
Hebrew.....	1,531	15.68	94.06	222.72	262.57	212.28	109.73	46.37	22.21	7.19	3.27	3.92	1,000
Number measured.....	24,680	6.69	48.62	147.44	235.46	230.55	165.03	94.08	42.42	18.60	6.77	4.34	1,000
Not measured.....	3,990												
Total.....	28,670												

It is seen that the Germans have the highest mean weight, although they are not the tallest of the eight races. It appears also that the Hebrews are lighter in weight than the Italians, although slightly taller.

The standard deviation in weight is greatest in the Scotch, despite their average extreme stature. They show a fairly large proportion of men under 120 pounds, also they are exceeded by only two other races in the proportion of men weighing over 200 pounds. This large proportion of the extreme classes is responsible for the high standard deviation. The next highest standard deviation is seen in the English group and the third in the Irish group. The reason in the case of the Irish is fairly clear from the fact that this group contains in its composition two or more races, one of which, the Scotch-Irish, is tall and spare, and the other of which, the Celtic-Irish, is short and stocky. The least variability is found in the Polish group and next to the lowest in the Italian group. The Hebrew and French groups show the same variability despite the marked difference in average weight.

TABLE 38.—*Mean weight in five color races with the standard deviation for the white and Negro troops, demobilization, 1919.*

Race.	Number exam- ined.	Mean weight.		Standard deviation.	
		Kilos.	Pounds.	Kilos.	Pounds.
White.....	79,708	65.62	144.67	7.67±0.02	16.92±0.04
Negro.....	3,319	67.83	149.53	8.00±.07	17.53±.15
Indian.....	103	68.10	150.13		
Chinese.....	18	67.56	148.94		
Japanese.....	24	65.73	144.92		

11. COMPARISON OF THE WEIGHT OF THE COLOR RACES.

A comparison of mean weights of the five color races measured at demobilization is afforded by the accompanying Table 38 taken from Tables 103, 104, and 107. It gives for the different color races the mean weight in kilograms and pounds. It appears that, though the white and Negro troops have almost exactly the same average stature, the Negro troops exceed the white in weight by about 5 pounds, and the weight is slightly more variable in the Negro troops. The Indians are on the average still heavier than the Negro troops, although from the small numbers it is probable that they are a highly selected lot. The Chinese come next in weight and the Japanese lowest with a weight of almost 145 pounds, a trifle in excess of that of the white troops.

IV. CHEST CIRCUMFERENCE.

1. GENERAL DISCUSSION.

This dimension is of both military and anthropological importance. It is of medico-military importance, first, because it may be used to measure lung capacity and, second, because it is an index of certain diseases. It is used, as will be shown later, in obtaining the index of robustness, an index which is believed to give a fair measure of vital resistance.

The medical importance of chest circumference is indicated in Section II of this work, which discusses the relation of chest circumferences to the different diseases. For example, from the summaries given there, it appears that chest circumference is small in men with pulmonary tuberculosis, and also in persons with various heart disorders. It is exceptionally large in one group of asthmat-

ics, doubtless due to the exercise of the chest muscles in the forced breathing which is a symptom of this disease.

The Army has long laid stress upon the difference between chest circumference at expiration and full inflation. This difference is called mobility. The minimum mobility for Army purposes is usually set at 2 to 3 inches for men under 6 feet (180 centimeters) and 3 to 4 inches for men over 6 feet. (See Table 138, p. 297.) Into mobility there enters a large nervous and mental factor; not infrequently the examiners find that the subject is unable to expand the chest, not through small lung capacity but through an inability to exercise a voluntary control over the muscles of the chest. Such control may, however, usually be secured by practice. Dr. O. L. Williamson, of Mariana, Ark. (Hoffman,¹⁶ p. 5), stated at the Conference of Physical Examination under the Selective Service (held in Chicago, June 13, 1918): "Many physically fit Negroes have not a chest mobility of 2 inches and they do not know how to expand the chest."

The occupational and racial significance of chest circumferences must not be overlooked whenever an attempt is made to draw inferences from the measurements. A comparison of our soldiers before and after training indicated how responsive chest circumference is to such training, for it increases with severe exercise of the arms and chest and diminishes in the sedentary. Thus Livi finds (Martin,⁵ p. 278), that in Italy farmers have the largest chest circumference, and tailors, barbers, and students have the smallest. However, it must be recognized that natural feebleness of muscular development may be one of the causes that leads some men to abandon the farm and become barbers, students, etc.

The chest circumference is particularly important in relation to the general size of the individual, as measured by his stature. Relative chest circumference is, where possible, to be considered; i. e., chest circumference divided by total stature.

2. METHODS OF MEASUREMENT.

The measurement of chest circumference requires attention to a few technical details. The graduated tape is passed around the chest (subject's arms lifted) until it lies under armpits, over the nipples, and perpendicular to the axis of the trunk at this level. Since the axis of the trunk is rarely vertical, the tape will rarely lie horizontal. Pressure is not to be applied so as markedly to indent the flesh. The subject's arms are lowered to his sides and the reading is taken.

Differences in technique are used by different anthropometrists. The method recommended by Martin⁵, (pp. 149-150) may be translated as follows:

61. Circumference of the chest in quiet breathing (Brustumfang während der Atempause oder in sogenannter Normalstellung; périmètre ou circonférence thoracique; girth of chest):

The individual to be measured stands upright, holding his arms at first laterally up to the level of the shoulders. The tape is placed high in the axillæ at the level of the mesosternal (above the nipples), horizontally about the thorax, and the two ends, passing each other, are held firmly with the ends upon the chest wall. The arms are then dropped and lie quiet at the side of the body. It is necessary to take care that the tape lies horizontally everywhere, even at the back, in contact with the body, without cutting into the skin. The part of the back lying between the two scapulæ will usually not be in contact with the tape, but will be bridged over by it. It is usual in most individuals to pass over the lower angle of the scapulæ. One observes the change in the position

of the tape caused by the light breathing movements for about half a minute and notes the middle position.

In many examinations, among others the military, the tape is placed about the chest just below the nipples and the lower angle of the scapulæ. Other authors measure without regard to the meso-sternal and nipples, as high as possible in the axillæ. By others the level of the processus ensiformis is recommended.

3. MEAN CHEST CIRCUMFERENCE AT EXPIRATION.

The average circumference of the deflated chest for the whole United States for 873,159 recruits is 33.22 inches, or 84.38 centimeters. The mean circumference for the uninflated chest of 95,867 troops at demobilization is 34.94 inches, or 88.74 centimeters. This gives a difference of 1.72 inches, or 4.36 centimeters, in the two sets of measurements. In comparing the means for recruits and men at demobilization, it is to be kept in mind that recruits were encouraged to deflate the chest as much as possible, since there was sought not merely the chest circumference but also the chest mobility. In the measurements of men at demobilization, instructions were that the chest should be in a quiescent condition, that is, neither inflated nor uninflated, as far as possible. However, since the difference in circumference of the quiescent chest and that from which the air has been driven as far as possible is usually between $\frac{1}{2}$ and $1\frac{1}{2}$ inches and averages about $\frac{3}{4}$ inch, only about 1 inch of the added chest girth is to be ascribed to the intensive training which the troops have received. This tended on the one hand to develop the lung capacity and on the other to develop the muscles of the chest and particularly those attached to the scapulæ.

The foregoing measurements of chest circumference are absolute. One may reduce them to relative measurements by dividing the average chest circumference by the average stature, both for men at mobilization and at demobilization. The relative chest circumference obtained in this way gives for men (deflated chest) at mobilization 49.2 per cent, and for men at demobilization (quiescent chest) 51.6 per cent.

The relation between the distribution of chest circumference of men of different statures and that of the whole population of recruits is shown graphically in Plate XIII. As is to be expected, the chest circumference for the shorter statures is below the distributions for the statures 67–68 inches, which are close to the average. For statures above this they are clearly above the average. The curve of distribution of chest circumference of men 62 inches tall is seen to be highly unsymmetrical owing to the fact that chest circumferences which were 3 or more inches below the average in the case of short men were rejected, whereas chest circumferences 3 or more inches up to 8 inches above the average for any stature were accepted. This elimination of the extremes results in a high mode for men with short statures. They form a less variable group than the men with mediocre or taller statures.

4. COMPARISON WITH CIVIL WAR DATA.

The Civil War statistics, obtained by Gould² (p. 280), give a mean circumference of chest at expiration, for white soldiers, of 34.49 inches. The mean girth at expiration of chest of recruits, according to Baxter¹ (Vol. I, p. 32), was 33.53 inches, or 85.17 centimeters, a very great discrepancy, which is doubtless

due to the fact that Gould's measurements were made at demobilization, whereas Baxter's statistics were of 500,000 drafted men taken from a population greatly depleted by volunteers. Thus Baxter's and Gould's measurements largely stand to each other as do ours of recruits and men at demobilization. In both cases the increase of circumferences after training is about one inch.

Comparing the recruits of Civil War times and 55 years later, we see a decrease of .3 inches in the latter group. Comparing men at demobilization, there is an increase of about one-half inch in the latter group, which difference is accounted for by the measurement at rest, rather than at expiration. Chest circumference has probably not diminished as much as stature.

5. COMPARISON WITH OTHER COUNTRIES.

For comparing the chest circumferences of our recruits with those of other countries, the following measurements will be of interest, probably all taken on the chest at rest, mostly from Martin⁵ (p. 278): Russians, 81 centimeters; Serbs, 80 centimeters; Bulgarians, 81 centimeters; English, 88.9 centimeters; Chinese, 77.5 centimeters; French, 88.7 centimeters; Bavarians (Ammon,¹⁸ p. 247), 87 centimeters.

Thus the chest circumference of our troops at demobilization exceeds, with a single exception, all the averages of different races as given. For the other races the dimensions lie either between those of our recruits and those of our veterans or else below the circumference of the recruits.

The relative chest circumference is more important in its racial variation than the absolute chest circumference. The following relative chest circumferences are given by Martin⁵ (p. 279): Russian Jews, 49.7; Belgians, 52.8; French, 53.7; Letts, 56.

Thus in the series given of the relative chest circumferences of European races all (except one) exceed that of our recruits and are equal to those of our veterans.

TABLE 39.—*Frequency and proportional distribution of chest circumferences (expiration) at mobilization, 1917-1918, and of chest circumference (rest) at demobilization, 1919.*

Chest circumference, in inches.	Mobilization.		Demobilization.							
	Number of men measured.	Ratio per 1,000.	Whites only.				White and colored.			
			Chest circumference, in centimeters.	Chest circumference, in inches, approximate.	Number of men measured.	Ratio per 1,000.	Chest circumference, in centimeters.	Chest circumference, in inches.	Number of men measured.	Ratio per 1,000.
28 and under.....			68-73	27-28	195	2.04	68-77	26.77-30.32	731	8.80
29.....	18,093	20.74	74-75		29	1.72	78-81	30.71-31.89	4,399	52.99
30.....	49,090	56.22	76-77		30	4.84	82-85	32.28-33.46	16,383	197.32
31.....	103,294	118.30	78-79		31	1,354	86-89	33.86-35.04	26,745	322.13
32.....	159,379	182.54	80-81		31	3,746	90-93	35.43-36.61	30,898	251.71
33.....	175,858	201.42	82-83		32	7,259	94-97	37.01-38.19	9,767	117.64
34.....	152,663	174.85	84-85		33	11,688	98-101	38.58-39.76	3,131	37.71
35.....	103,414	118.42	86-87		34	14,576	102 and over.	40.16 and over.	971	11.70
			88-89		35	16,172				
36.....	59,015	67.60	90-91		35	13,702				
37.....	28,175	32.27	92-93		36	10,553				
38.....	13,151	15.06	94-95		37	7,057				
39.....	11,027	12.63	96-97		38	4,184				
40 and over.....			98-99		39	2,522				
			100-117	40 and over.	2,210	23.05				
Total measured.....	873,159				95,867				83,025	1,000.00

TABLE 39.—*Frequency and proportional distribution of chest circumferences (expiration) at mobilization, 1917-1918, and of chest circumference (rest) at demobilization, 1919—Continued.*

	Mean chest circumference—			
	At mobilization (deflated).		At demobilization (at rest).	
	Inches.	Centi- meters.	Inches.	Centi- meters.
Mean chest circumference, white and colored.....	33.22	84.38	34.94	88.74
Standard deviation:				
White and colored.....	2.01	5.11		
White.....			2.04	5.09
Negro.....			1.87	4.76

6. DISTRIBUTION OF FREQUENCIES OF VARIOUS CLASSES OF CHEST CIRCUMFERENCE.

Table 39 gives for recruits and veterans the distribution of frequencies of the different classes of chest circumference in inches or in centimeters. The frequency is given in absolute numbers of men measured and also in the ratio per thousand. It is to be recalled that about three-fourths of an inch has to be added to the chest circumference at mobilization to make the measurements comparable with those taken at demobilization. Even after making this correction the great superiority of veterans over recruits is strikingly apparent. The mode for white veterans is at 35 inches instead of 33 plus; 23 per mille were found at 40 and over instead of practically none at all. Only 5 per mille of white veterans had a chest circumference of 30 inches; while 20 per mille of recruits had a circumference of 29 inches.

Also the standard deviation of the recruits (deflated chest) was 2.01 inches, and that of the white veterans 5.093 centimeters, or 2.04 inches. The coefficients of variation are respectively 6.05 and 5.86. That is, the chest circumferences of the veterans were much less variable than those of the recruits—doubtless due to the greater uniformity of conditions under which they had been trained.

There is given for comparison, extracted from Table XCIX, the distribution of chest circumference for 95,867 white men measured at demobilization. In this case the classes are in centimeters and here also is given the nearest corresponding English measure.

7. THE FREQUENCY DISTRIBUTION OF CHEST CIRCUMFERENCE, BY STATES.

Table 40 gives the mean chest circumference for recruits from each of the States, arranged in descending order of size of chest. In this table, North Dakota stands at the top with a mean chest circumference of 33.76 inches, over half an inch above the average. This great size of chest is associated with a robustness which is higher for this State than for any other of the United States proper. Next on the list stands Nevada, a State which has a high, though not extremely high, relative chest circumference. This is followed by Idaho, of which the relative chest circumference falls at the bottom of the upper third.

People from these States are therefore not especially stout, but have an absolutely large chest circumference, which is due probably to a combination of muscular activity, especially of the arms, and the rarified air of these States of high altitude. The inhabitants of Nevada and Idaho are largely miners, and no doubt that part of the population which is engaged in mining has acquired especially large chest circumference. At the same time these men, especially of Idaho, are above the average in stature and consequently have a high absolute chest circumference. Among the other States and Territories at the top of the list are Alaska, 33.65; Minnesota, Wisconsin, and North Dakota, which include men of exceptional robustness. These are followed by other States of the Northwest—Oregon, Montana, and Washington. At the bottom of the list lies the District of Columbia, the most urban of all of the States and Territories listed. Indeed, the District falls in a class by itself. The small chest circumference is no doubt due largely to the comparative lack of use of the muscles of the chest by an urban population, especially one in which the males are so largely engaged in clerical occupations. Next above comes Rhode Island, the second most urban of all of the States and one which stands at the bottom both in height and weight of its drafted men. The chest circumference in relation to stature is not extremely low; the small chest circumference is therefore due primarily to the small size of the inhabitants. Next come the States of Tennessee and Kentucky, with tall men of low weight and of extraordinarily small chest circumference. In fact, at the bottom of the table one finds a group of Southern States, including Alabama, Florida, Louisiana, Mississippi, and Missouri, the inhabitants of which are characterized by lankiness of form, which shows itself also in their low average chest circumference. The question arises how far this small chest circumference is influenced by the Negro population. From a set of measurements made at demobilization, it appears that the Negro troops have indeed a smaller chest circumference than white troops, as 34.64 to 34.96. These averages are, to be sure, very much higher than those obtained by local boards, but this is due to the training which the returned soldiers had undergone in the preceding months. There is no reason for thinking that the Negro troops were less active than the whites, and yet their mean chest circumference is 0.32 inch less than that of the whites. We may conclude therefore that the Negro population has a lower chest circumference than the white population; and since, in the Southern States, the Negro forms a relatively large proportion of the population, the low average chest circumference of men from the Gulf States is to be partly attributed to the presence in them of small-chested colored men. Among the States occupying a relatively low position for chest circumference is Colorado, the State which stood near the top in the number of rejections for tuberculosis of the lungs. The figures suggest that the well-known small chest circumference of the tuberculous has been influential in reducing the average chest circumference of men from Colorado. The small chest circumference of men from Massachusetts is largely due to their small size, since the relative chest circumference is high in them.

TABLE 40.—*Mean chest circumference (expiration), by States; States arranged in order of standing, with proportional chest circumference at expiration in inches for each inch of height and each pound of weight; also the proportional weight in pounds for each inch of chest circumference; first million draft recruits.*

State.	Number of men measured.	Mean chest.	Mean chest.	Mean chest.	Mean weight.
		Mean height.	Mean weight.	Mean chest.	Mean chest.
		Inches.	Inch.	Inch.	Pounds.
North Dakota.....	6,444	32.76	0.497	0.230	4.353
Nevada.....	1,441	33.75	.497	.232	4.307
Idaho.....	4,031	33.74	.495	.232	4.307
Alaska.....	106	33.65	.493	.223	4.472
Minnesota.....	27,341	32.63	.494	.230	4.354
Wisconsin.....	18,433	33.55	.496	.232	4.307
South Dakota.....	3,892	33.54	.493	.228	4.382
Oregon.....	2,748	33.51	.492	.228	4.368
Montana.....	11,648	33.47	.492	.228	4.372
Washington.....	11,316	33.46	.492	.230	4.347
Connecticut.....	13,585	33.43	.501	.239	4.182
Vermont.....	2,077	33.43	.498	.238	4.198
Iowa.....	19,537	33.41	.491	.230	4.332
Maine.....	3,315	33.41	.497	.237	4.221
California.....	35,461	33.39	.493	.231	4.312
Wyoming.....	1,927	33.38	.492	.231	4.332
Michigan.....	41,872	33.35	.496	.235	4.258
New Jersey.....	29,958	33.29	.498	.239	4.170
West Virginia.....	12,367	33.29	.490	.235	4.251
Illinois.....	69,491	33.28	.493	.234	4.260
Kansas.....	9,571	33.28	.487	.231	4.319
Arizona.....	3,850	33.26	.488	.232	4.301
North Carolina.....	14,668	33.25	.487	.235	4.255
Nebraska.....	10,774	33.24	.488	.229	4.354
New York.....	87,818	33.22	.497	.238	4.200
Georgia.....	20,305	33.21	.488	.235	4.241
New Hampshire.....	2,240	33.20	.495	.236	4.227
Virginia.....	17,616	33.18	.489	.236	4.230
Arkansas.....	10,111	33.17	.486	.234	4.269
Oklahoma.....	19,429	33.16	.485	.232	4.293
Indiana.....	23,194	33.14	.489	.233	4.274
New Mexico.....	2,690	33.14	.491	.239	4.178
Utah.....	4,568	33.14	.488	.231	4.319
Ohio.....	52,814	33.12	.491	.234	4.268
Delaware.....	1,891	33.11	.492	.237	4.212
Maryland.....	9,192	33.11	.494	.236	4.240
Massachusetts.....	29,534	33.10	.496	.239	4.281
South Carolina.....	9,343	33.10	.489	.235	4.244
Pennsylvania.....	77,186	33.10	.496	.236	4.221
Missouri.....	24,964	33.08	.486	.233	4.275
Mississippi.....	8,543	33.08	.485	.231	4.330
Louisiana.....	12,356	33.08	.489	.236	4.221
Colorado.....	6,635	33.07	.485	.234	4.265
Florida.....	5,895	33.06	.489	.237	4.214
Alabama.....	15,988	33.03	.485	.233	4.277
Texas.....	34,531	33.02	.483	.232	4.307
Kentucky.....	15,502	32.98	.494	.235	4.245
Tennessee.....	14,426	32.97	.483	.235	4.249
Rhode Island.....	3,928	32.83	.494	.241	4.156
District of Columbia.....	4,486	32.66	.482	.232	4.303

TABLE 41.—*Chest circumference (expiration) of native American white draft recruits of Civil War.*

[From Baxter,¹ Vol. I, p. 32, rearranged.]

State.	Inches.	Centi- meters.	State.	Inches.	Centi- meters.
Nevada.....	34.38	87.33	Illinois.....	33.65	85.48
Delaware.....	34.25	86.98	New Hampshire.....	33.60	85.34
California.....	34.11	86.63	Wisconsin.....	33.51	85.10
Minnesota.....	34.02	86.41	Michigan.....	33.50	85.08
Kansas.....	33.99	86.34	Pennsylvania.....	33.49	85.07
Kentucky.....	33.98	86.30	Vermont.....	33.38	84.77
Missouri.....	33.90	86.11	West Virginia.....	33.07	83.99
Maryland.....	33.90	86.10	New York.....	32.91	83.59
Iowa.....	33.87	86.02	Connecticut.....	32.57	82.74
Maine.....	33.81	85.87	New Jersey.....	32.33	82.11
Indiana.....	33.70	85.89	Rhode Island.....	32.27	81.97
Ohio.....	33.66	85.50	Massachusetts.....	31.99	81.25
District of Columbia.....	33.66	85.49			

In Table 42 the different States are arranged in order of the relative chest circumference obtained by dividing the mean chest circumference of each State by the mean height of men from that State. In this table the State showing the highest ratio of chest circumference to mean height is Connecticut. This is partly due to the small stature of the men of Connecticut and partly to the large chest circumference they show. This large chest circumference is more striking for men of the agricultural part of Connecticut than of the manufacturing area. It appears that Connecticut stands at the top of the list for relative chest circumference because it contains so many small men who are engaged in agricultural occupations and others involving exercise of the upper appendages and upper trunk. Vermont comes second in the list, again an agricultural State, comprising many persons of small size. New Jersey and Maine come next and their position is to be explained in similar fashion. Next in order comes North Dakota. Here, despite the great average stature of the inhabitants, the chest circumference is relatively large, again associated with the agricultural activity of this magnificently proportioned population. This is followed by a mixture of mining and agricultural States in which the population is largely engaged in occupations involving use of the upper part of the body.

At the other extreme of the table stands first the District of Columbia for reasons already put forward in accounting for the small absolute chest circumference of its population. Next come certain States containing very tall men, such as Tennessee, Texas, Kentucky, in which the chest circumference has not increased in proportion to the great stature. The ratio is small, partly because it is very small in the mountain-white sections of these States. Possibly hookworm has an important influence in keeping down the relative chest circumference. In the lower part of the table lie also Alabama, Mississippi, Missouri, Arkansas, North Carolina, and other Southern States, probably largely because of the admixture of Negroes who, as we have seen, have a relatively smaller chest circumference and about the same average stature as the whites.

The relative small chest circumference of the draft recruits from the Southern States is due in part to the fact, as shown in Plate XIV, figure 1, that the proportion of the chest circumference (expiration) to the stature increases as the stature decreases.

TABLE 42.—*Relative chest circumference (mean chest circumference divided by mean stature), by States, arranged in order of standing, first million draft recruits.*

States.	Relative chest circumference.	States.	Relative chest circumference.
Connecticut.....	0.501	Iowa.....	0.491
Vermont.....	.498	New Mexico.....	.491
New Jersey.....	.498	Ohio.....	.491
Maine.....	.497	West Virginia.....	.490
North Dakota.....	.497	Virginia.....	.489
Nevada.....	.497	Indiana.....	.488
New York.....	.497	South Carolina.....	.488
Pennsylvania.....	.496	Louisiana.....	.488
Wisconsin.....	.496	Florida.....	.498
Michigan.....	.496	Arizona.....	.498
Massachusetts.....	.496	Nebraska.....	.488
Idaho.....	.495	Utah.....	.488
New Hampshire.....	.495	Georgia.....	.488
Minnesota.....	.494	Kansas.....	.487
Maryland.....	.494	North Carolina.....	.487
Rhode Island.....	.494	Arkansas.....	.486
Alaska.....	.493	Missouri.....	.486
South Dakota.....	.493	Oklahoma.....	.485
California.....	.493	Mississippi.....	.485
Illinois.....	.493	Colorado.....	.485
Oregon.....	.492	Alabama.....	.485
Montana.....	.492	Kentucky.....	.484
Washington.....	.492	Texas.....	.483
Wyoming.....	.492	Tennessee.....	.483
Delaware.....	.492	District of Columbia.....	.482

8. MEAN CHEST CIRCUMFERENCE BY SECTIONS.

Table 43 gives the chest circumference for each of the sections into which the country has been divided, arranged in order of size of chest circumference, the largest being placed first. The average for the whole United States is 33.22 inches. We find that more than half of the sections have a chest circumference above the mean. At the top of the table stand three rural districts of Minnesota, comprising a large proportion of Scandinavians. That Minnesota as a whole does not occupy the first position is due to the reduction in stature of men from her large cities. Next comes North Dakota 2, largely Scandinavians, and next the mining area of California 2. The mining States of Nevada 1 and Idaho 1, as already shown, have a high average chest girth, as has also South Dakota 3, containing a large proportion of Indians. Next comes Wisconsin 4, containing a large proportion of Germans. The mountainous region of New Hampshire 1 comes next and this is followed by three sections containing Scandinavian and rural Russian population. The foregoing sections have a mean chest circumference about 0.5 inch above the average. These are followed by a number of sections among which the mountain areas are strikingly prevalent, followed by several agricultural areas more largely of native white population. In the middle of the list stand many sections with a large Negro population. At the very bottom of the list stands New Orleans (Louisiana 2), in which the chest circumference is 32.63—less than the men from the District of Columbia. The ratio of mean chest to stature, however, is greater than in the District of Columbia. Next to the bottom of the table lies New Mexico 3, with its noteworthy Mexican element, in which not only the stature but also the relative chest circumference is small. This is followed by the District of Columbia and

by the Key West Section (Florida 3), containing many Italians and Cubans. The district around Mobile (Alabama 5) affords a population with chest circumference of only 32.82, and indeed many southern sections, especially those containing few Negroes, are found in the lower part of the table. Rather striking is the position, toward the bottom, of Denver (Colorado 5), (associated with a large number of rejections for tuberculosis) and Philadelphia (Pennsylvania 1), Cincinnati (Ohio 4), St. Louis (Missouri 4), Baltimore (Maryland 1), Los Angeles (California 4), Boston (Massachusetts 4), and even New York city (New York 2), (mean chest girth, 33.14). It is clear that the inhabitants of cities tend to have reduced chest girth, possibly due to a smaller amount of exercise of the upper appendages and to the small races that congregate in them. This is illustrated by comparing the twin cities of Minnesota with the rest of the State. The men of the former have a chest circumference about 0.75 inch less than the latter.

TABLE 43.—Mean chest circumference (expiration) by sections; sections arranged in order of standing with proportional chest circumference (expiration) in inches for each inch of height and each pound of weight; also standard deviation for each chest circumference; first million draft recruits.

State.	Section.	Characteristics of sections.	Number of men measured.	Mean chest.	Standard deviation. (chest).	Mean chest. Mean height.	Mean chest. Mean weight.
				Inches.	Inches.	Inch.	Inch.
Average for United States.			873, 159	33. 22	2. 01	0. 492	0. 234
Minnesota.	3	Scandinavians and Finns.	3, 515	33. 95	1. 98	.502	.232
Do.	2	German and Scandinavian population.	7, 585	33. 89	1. 93	.497	.229
Do.	1	Scandinavian population.	6, 448	33. 86	1. 86	.495	.228
North Dakota.	2	do.	3, 305	33. 82	1. 88	.497	.230
California.	2	Mining area.	942	33. 81	1. 87	.499	.231
Nevada.	1	State undivided, sparse population.	1, 438	33. 75	2. 08	.497	.232
Idaho.	1	State undivided.	4, 031	33. 74	2. 04	.495	.232
South Dakota.	3	Indian population.	247	33. 74	1. 74	.495	.228
Wisconsin.	4	Lake counties.	2, 883	33. 73	2. 01	.500	.234
New Hampshire.	1	Mountainous area.	667	33. 72	2. 09	.501	.236
North Dakota.	3	Scandinavian and Canadian population.	1, 131	33. 72	1. 91	.498	.230
Do.	3	Russian population.	2, 005	33. 70	1. 86	.496	.228
Wisconsin.	1	Scandinavian and German population.	3, 290	33. 68	1. 89	.494	.232
Alaska.	All.	Undivided.	106	33. 65	1. 94	.493	.223
Maine.	2	Native white stock, maritime.	828	33. 64	1. 95	.497	.237
North Carolina.	1	Sparsely populated mountainous area.	2, 738	33. 64	1. 82	.489	.238
Michigan.	1	Finnish population.	2, 340	33. 63	1. 96	.501	.232
Washington.	3	Mountainous area.	1, 539	33. 62	1. 83	.493	.230
South Dakota.	2	Large Russian population.	694	33. 61	1. 87	.495	.228
Montana.	2	Sparsely settled, mountainous area.	6, 621	33. 60	1. 85	.493	.229
Utah.	1	Sparsely populated.	1, 224	33. 59	1. 82	.492	.233
Connecticut.	1	Prevalently agricultural and near metropolitan.	4, 877	33. 58	2. 10	.503	.240
Wisconsin.	2	German population.	7, 678	33. 56	1. 97	.495	.232
Iowa.	1	Foreign white, German and Scandinavian.	12, 139	33. 54	1. 93	.492	.230
Oregon.	1	Fairly densely populated.	2, 747	33. 54	2. 10	.492	.228
New Jersey.	3	Mountainous area plus Atlantic County.	3, 196	33. 52	2. 02	.501	.240
California.	1	Chiefly agricultural area.	11, 712	33. 52	2. 08	.494	.231
South Dakota.	1	Dry farming area.	3, 050	33. 51	1. 95	.492	.228
Indiana.	2	Agricultural, considerable German.	835	33. 49	2. 01	.491	.231
Michigan.	3	Foreign population.	6, 280	33. 49	2. 09	.497	.235
California.	3	Sparsely populated.	2, 106	33. 48	1. 97	.490	.231
New York.	7	Agricultural and dairying.	6, 465	33. 48	2. 06	.496	.236
Washington.	1	Coastal region plus eastern counties.	5, 174	33. 47	2. 02	.492	.230
Maine.	1	English Canadian.	1, 238	33. 46	1. 96	.497	.235
Indiana.	1	Manufacturing.	3, 609	33. 45	2. 12	.497	.235
Utah.	3	Mining area.	562	33. 44	1. 77	.494	.232
Vermont.	1	State undivided.	2, 079	33. 43	1. 90	.498	.238
Michigan.	8	Prevalently native white population.	12, 560	33. 42	1. 98	.493	.235
Illinois.	2	Agricultural and manufacturing area.	2, 451	33. 42	1. 96	.493	.233
Oregon.	2	Columbia River Valley and coastal dry plain, sparsely populated.	1, 076	33. 42	1. 91	.490	.229
Nebraska.	2	German, Austrian, and Russian stocks.	3, 138	33. 41	1. 95	.489	.229
Washington.	2	Puget Sound, foreign white.	6, 599	33. 41	1. 96	.492	.230
Illinois.	4	Largely German population.	4, 238	33. 40	2. 03	.494	.233

TABLE 43.—Mean chest circumference (expiration) by sections; sections arranged in order of standing with proportional chest circumference (expiration) in inches for each inch of height and each pound of weight; also standard deviation for each chest circumference; first million draft recruits—Contd.

State.	Sec- tion.	Characteristics of sections.	Number of men meas- ured.	Mean chest.	Stand- ard devia- tion (chest).	Mean chest. Mean height.	Mean chest. Mean weight.
				<i>Inches.</i>	<i>Inches.</i>	<i>Inch.</i>	<i>Inch.</i>
Illinois.	2	Mixed native and foreign population.	7,803	33.40	2.03	0.494	0.233
New Jersey.	2	Plains section, rural.	8,985	33.40	2.12	.499	.240
Texas.	3	German and Negro population.	1,415	33.40	2.09	.488	.231
Maryland.	3	Large white population.	2,675	33.39	1.99	.496	.237
Illinois.	1	Densely populated.	7,803	33.38	2.02	.495	.233
Wyoming.	1	State undivided, sparsely populated.	1,927	33.38	1.89	.492	.231
Texas.	5	Large Negro population.	1,346	33.36	2.05	.487	.231
Virginia.	3	Native rural region.	3,866	33.36	1.94	.489	.237
New York.	6	Urban area.	6,544	33.35	2.08	.498	.234
Connecticut.	2	Manufacturing area.	8,708	33.34	2.20	.499	.238
New York.	8	Mountainous, Adirondack area.	2,986	33.34	2.00	.497	.237
Virginia.	4	Mountain, white.	5,499	33.33	1.87	.489	.238
Georgia.	2	Large Negro population.	10,070	33.33	1.91	.490	.236
Illinois.	7	Agricultural area.	5,442	33.33	1.98	.491	.234
Wisconsin.	3	Urban and foreign stock.	4,513	33.33	2.11	.497	.237
New York.	3	Eastern manufacturing region.	5,131	33.32	2.07	.498	.238
Pennsylvania.	3	Mining area.	7,293	33.32	2.10	.500	.236
California.	5	Urban area.	7,189	33.32	2.09	.495	.231
Colorado.	1	Large native white population.	1,053	33.32	1.77	.489	.235
Montana.	1	Mining area, foreign population.	5,117	33.31	1.93	.491	.229
Colorado.	2	Russian population.	1,099	33.30	1.75	.490	.234
Missouri.	3	Native white, Ozark region.	1,138	33.30	1.76	.485	.234
Arkansas.	2	Large native white population, hill country.	1,559	33.29	1.80	.484	.236
Louisiana.	1	Mississippi bottoms and upland, large Negro population.	4,072	33.29	1.97	.491	.236
Pennsylvania.	6	Rural area.	8,616	33.29	1.98	.494	.235
Kansas.	2	Native and German population.	8,505	33.28	1.99	.488	.231
Arizona.	1	Large Indian population, sparsely settled.	1,027	33.28	1.91	.489	.232
Alabama.	2	Large Negro population.	3,327	33.27	1.90	.489	.233
New Mexico.	2	Native white population.	1,851	33.26	1.84	.493	.240
Arizona.	2	Chiefly white population.	2,821	33.25	1.99	.487	.232
Illinois.	5	Urban area.	33,905	33.25	2.12	.495	.236
New Mexico.	1	Indian population.	290	33.25	1.84	.494	.239
Texas.	1	Large Mexican population.	6,676	33.24	1.98	.487	.234
Kansas.	1	Russian population.	1,066	33.24	2.68	.486	.229
North Carolina.	5	Island and peninsular area.	254	33.24	1.84	.491	.235
Mississippi.	1	Rural area, large Negro population.	5,149	33.24	1.88	.488	.231
New York.	4	Western manufacturing region.	14,222	33.23	2.13	.495	.236
Oklahoma.	2	Chiefly white population.	10,958	33.22	1.95	.485	.232
Maine.	3	French Canadian population.	1,247	33.22	1.93	.495	.238
Colorado.	3	English population.	380	33.21	1.86	.487	.233
South Carolina.	2	Large Negro population.	3,976	33.20	1.85	.490	.234
Iowa.	2	Native white.	7,404	33.20	1.92	.488	.231
Ohio.	1	Dense foreign population.	17,208	33.20	2.04	.495	.234
West Virginia.	1	Mountainous area.	1,506	33.20	1.87	.488	.236
Kentucky.	1	Mountainous area, native white.	4,029	33.19	1.80	.486	.237
New Jersey.	1	Densely populated.	17,772	33.19	2.12	.497	.239
Arkansas.	1	Negro, Mississippi bottoms.	4,933	33.18	1.95	.487	.233
Florida.	4	Peninsular.	2,339	33.18	1.95	.491	.237
Do.	2	Negro and rural population.	995	33.18	2.02	.490	.236
Michigan.	4	Urban area.	17,751	33.18	2.08	.496	.235
Minnesota.	4	Urban area, "Twin Cities."	9,757	33.18	2.01	.489	.230
North Carolina.	2	Intermediate.	4,309	33.18	1.90	.486	.235
Pennsylvania.	2	Rural area, native stock.	14,218	33.18	2.02	.497	.237
Nebraska.	1	German and Irish, foreign stocks.	7,621	33.17	1.93	.488	.230
New York.	5	Mountainous, Catskill region.	795	33.17	2.01	.493	.238
Louisiana.	3	Rural, chiefly white population.	5,227	33.17	1.87	.486	.236
Texas.	4	Coastal native population.	2,722	33.16	1.99	.487	.233
Alabama.	1	Large Negro population.	665	33.16	1.84	.486	.229
New York.	1	Suburban territory.	4,919	33.16	2.08	.497	.238
North Carolina.	3	Native white of Scotch origin.	2,050	33.16	1.82	.485	.234
Do.	6	Remainder of State.	744	33.16	1.85	.489	.235
Massachusetts.	2	Manufacturing center.	18,352	33.15	2.04	.497	.241
North Carolina.	4	Large Negro population.	4,358	33.15	1.91	.489	.233
Pennsylvania.	5	Coal mining.	4,813	33.15	2.00	.496	.235
Do.	5	Manufacturing.	8,892	33.15	1.98	.497	.235
Michigan.	3	Dutch and other foreign population.	2,899	33.14	1.96	.491	.235
New York.	2	Urban area, densely populated.	46,651	33.14	2.15	.498	.239
Colorado.	3	Prevailing agricultural.	1,222	33.14	1.88	.486	.233
Ohio.	3	Agricultural area.	17,548	33.13	2.00	.489	.234
Missouri.	1	Native white, agricultural.	13,571	33.11	1.90	.486	.234
Do.	2	Mississippi bottoms, considerable Negro population.	3,448	33.11	1.89	.486	.233
Delaware.	1	State undivided.	1,891	33.11	1.97	.492	.237
Georgia.	1	Mixed population, native white pre- dominating.	10,235	33.10	1.88	.486	.235

TABLE 43.—*Mean chest circumference (expiration) by sections; sections arranged in order of standing with proportional chest circumference (expiration) in inches for each inch of height and each pound of weight; also standard deviation for each chest circumference; first million draft recruits—Contd.*

State.	Section.	Characteristics of sections.	Number of men measured.	Mean chest.	Standard deviation. (chest).	Mean chest. Mean height.	Mean chest. Mean weight.
				<i>Inches.</i>	<i>Inches.</i>	<i>Inch.</i>	<i>Inch.</i>
Arkansas.....	3	Large native white population.....	3,589	33.10	1.78	0.485	0.235
Oklahoma.....	1	Marked Indian and Negro population.....	8,471	33.09	1.87	.485	.233
Virginia.....	2	Large Negro population.....	5,339	33.07	1.89	.490	.236
Alabama.....	3	Large native white population.....	2,666	33.07	1.80	.484	.235
Illinois.....	3	Agricultural area, native.....	8,900	33.07	1.94	.487	.232
Indiana.....	3	Agricultural area, native stock.....	18,725	33.06	2.00	.487	.233
Ohio.....	2	Intermediate.....	14,443	33.06	1.98	.491	.234
Massachusetts.....	4	Urban area.....	8,553	33.06	2.14	.494	.237
South Carolina.....	3	Peninsular and rural areas.....	3,795	33.05	1.85	.491	.238
California.....	4	Urban area.....	7,428	33.04	2.02	.487	.234
Tennessee.....	2	Agricultural region.....	6,305	33.02	1.85	.484	.237
Pennsylvania.....	7	Allegheny County plus a small rural area.....	17,238	33.01	2.08	.495	.236
Maryland.....	2	Peninsular area.....	1,066	33.00	1.88	.490	.236
Do.....	1	Urban area.....	5,420	32.99	2.08	.493	.235
New Hampshire.....	2	Manufacturing area.....	1,581	32.98	2.00	.493	.237
South Carolina.....	1	Native white.....	1,563	32.97	1.83	.484	.235
Missouri.....	4	Urban area.....	6,784	32.96	2.07	.488	.235
Ohio.....	4	do.....	3,554	32.96	2.09	.489	.232
Tennessee.....	3	Mountainous region.....	5,896	32.93	1.85	.481	.231
Alabama.....	1	Mining and manufacturing area.....	8,633	32.93	1.84	.484	.233
Florida.....	1	Largely white and maritime.....	2,477	32.92	1.83	.486	.237
Pennsylvania.....	1	Urban area.....	16,053	32.91	2.02	.494	.239
Massachusetts.....	1	Mountainous area.....	1,373	32.90	2.09	.492	.237
Kentucky.....	2	Agricultural area.....	11,419	32.90	1.91	.484	.234
Tennessee.....	1	Negroes, Mississippi bottoms.....	2,217	32.90	1.84	.483	.232
Texas.....	2	Sparsely settled, white.....	22,372	32.90	1.85	.480	.231
Utah.....	3	More densely populated.....	2,781	32.89	1.88	.485	.230
Massachusetts.....	3	Peninsular region.....	1,123	32.88	2.12	.491	.237
Colorado.....	5	Urban population.....	1,640	32.88	1.83	.485	.234
Illinois.....	6	Negro population (Egypt).....	1,409	32.87	1.95	.482	.236
Virginia.....	1	Peninsular region and east shore.....	2,886	32.84	2.05	.487	.233
Mississippi.....	2	Rural area, large native white population.....	3,387	32.83	1.96	.490	.231
Rhode Island.....	1	State undivided.....	3,925	32.83	2.11	.494	.241
Alabama.....	5	Urban and suburban area.....	479	32.82	1.96	.485	.234
Colorado.....	6	Austrian and Italian population.....	1,224	32.79	1.89	.484	.235
Florida.....	3	Cuban, Spanish, West Indian population.....	84	32.74	1.99	.487	.240
Dis. of Columbia.....	1	District undivided.....	4,486	32.66	2.00	.482	.232
New Mexico.....	3	Noteworthy Mexican element.....	540	32.63	1.85	.480	.234
Louisiana.....	2	Urban area.....	3,040	32.63	2.09	.487	.237

9. STANDARD DEVIATIONS OF CHEST CIRCUMFERENCE BY SECTIONS.

Table 44 shows the variations in the standard deviations of chest circumference for the various sections. For the United States as a whole the standard deviation is close to 2 inches. In western Kansas it is 2.68 inches, a high variability associated with the mixture of Germans and large Scandinavians, on the one hand, and of smaller Russians on the other. In manufacturing Connecticut, in New York City, Boston, Chicago, suburban New Jersey, and Rhode Island; the standard deviation is also high. In general, the eastern cities attract both extremes in body size. Greater uniformity (smaller standard deviation) is found in the Southern States. Extremely low variability is found in South Dakota 3, with 87 per cent Indians; Colorado 2; and Missouri 3, the Ozark Mountains, 94 per cent native whites and mostly big men.

TABLE 44.—*The standard deviation of chest circumference (expiration), by sections, arranged in order of standing, first million draft recruits.*

State.	Section.	Standard deviation.	State.	Section.	Standard deviation.
United States		2.01	Florida.....	4	1.95
Kansas.....	1	2.68	Maine.....	2	1.95
Connecticut.....	2	2.20	South Dakota.....	1	1.95
New York.....	2	2.15	Oklahoma.....	2	1.95
Massachusetts.....	4	2.14	Texas.....	2	1.95
New York.....	4	2.13	Nebraska.....	2	1.95
Illinois.....	5	2.12	Illinois.....	6	1.95
New Jersey.....	2	2.12	Arkansas.....	1	1.95
Indiana.....	1	2.12	Illinois.....	3	1.94
New Jersey.....	1	2.12	Virginia.....	3	1.94
Massachusetts.....	3	2.12	Alaska.....	All.	1.94
Rhode Island.....	All.	2.11	Minnesota.....	2	1.93
Wisconsin.....	3	2.11	Iowa.....	1	1.93
Connecticut.....	1	2.10	Montana.....	1	1.93
Oregon.....	1	2.10	Maine.....	3	1.93
Pennsylvania.....	3	2.10	Nebraska.....	1	1.93
Louisiana.....	2	2.09	Iowa.....	2	1.92
Massachusetts.....	1	2.09	Kentucky.....	2	1.91
Ohio.....	4	2.09	North Carolina.....	4	1.91
New Hampshire.....	1	2.09	Arizona.....	1	1.91
Michigan.....	3	2.09	North Dakota.....	1	1.91
Texas.....	3	2.09	Oregon.....	2	1.91
California.....	5	2.09	Georgia.....	2	1.91
New York.....	6	2.08	Vermont.....	All.	1.90
Nevada.....	1	2.08	Alabama.....	2	1.90
Ohio.....	1	2.08	North Carolina.....	2	1.90
Michigan.....	4	2.08	Missouri.....	1	1.90
New York.....	1	2.08	Do.....	2	1.89
Pennsylvania.....	7	2.08	Virginia.....	2	1.89
Maryland.....	1	2.08	Wisconsin.....	1	1.89
Missouri.....	4	2.07	Wyoming.....	1	1.89
New York.....	3	2.07	Colorado.....	6	1.89
California.....	1	2.06	Georgia.....	1	1.88
New York.....	7	2.06	Colorado.....	4	1.88
Texas.....	5	2.05	Mississippi.....	1	1.88
Virginia.....	1	2.05	North Dakota.....	2	1.88
Massachusetts.....	2	2.04	Maryland.....	2	1.98
Idaho.....	1	2.04	Utah.....	2	1.88
Illinois.....	4	2.03	California.....	2	1.87
Do.....	1	2.03	South Dakota.....	2	1.87
Do.....	1	2.02	Oklahoma.....	1	1.87
Washington.....	1	2.02	Virginia.....	4	1.87
New Jersey.....	3	2.02	West Virginia.....	1	1.87
Pennsylvania.....	1	2.02	Louisiana.....	3	1.87
California.....	4	2.02	Maine.....	1	1.86
Pennsylvania.....	2	2.02	Minnesota.....	1	1.86
Florida.....	2	2.02	North Dakota.....	3	1.86
Wisconsin.....	4	2.01	Colorado.....	3	1.86
Indiana.....	2	2.01	Mississippi.....	2	1.86
Minnesota.....	4	2.01	South Carolina.....	2	1.85
New York.....	5	2.01	Do.....	3	1.85
Do.....	8	2.00	New Mexico.....	3	1.85
Pennsylvania.....	4	2.00	Tennessee.....	2	1.85
Ohio.....	3	2.00	Do.....	3	1.85
Indiana.....	3	2.00	North Carolina.....	6	1.85
New Hampshire.....	2	2.00	Montana.....	2	1.85
District of Columbia.....	All.	2.00	North Carolina.....	5	1.84
Kansas.....	2	1.99	Tennessee.....	1	1.84
Florida.....	3	1.99	Alabama.....	1	1.84
Arizona.....	2	1.99	Do.....	4	1.84
Texas.....	4	1.99	New Mexico.....	1	1.84
Maryland.....	3	1.99	Do.....	2	1.84
Pennsylvania.....	6	1.98	Colorado.....	5	1.83
Michigan.....	2	1.98	Florida.....	1	1.83
Minnesota.....	3	1.98	South Carolina.....	1	1.83
Ohio.....	2	1.98	Washington.....	3	1.83
Texas.....	1	1.98	North Carolina.....	1	1.82
Pennsylvania.....	5	1.98	Do.....	3	1.82
Illinois.....	8	1.98	Utah.....	1	1.82
Do.....	7	1.98	Alabama.....	3	1.90
Delaware.....	1	1.97	Arkansas.....	2	1.80
Wisconsin.....	2	1.97	Kentucky.....	1	1.80
California.....	3	1.97	Arkansas.....	3	1.78
Louisiana.....	1	1.97	Utah.....	3	1.77
Alabama.....	5	1.96	Colorado.....	1	1.77
Michigan.....	1	1.96	Missouri.....	3	1.76
Do.....	5	1.96	Colorado.....	2	1.75
Washington.....	2	1.96	South Dakota.....	3	1.74

10. MEAN CHEST CIRCUMFERENCE BY GROUPS OF SECTIONS.

Certain additional points are revealed in Table 45, giving the chest circumference by groups of sections. Of all the groups, group 18 (the two Finnish sections) show the highest absolute chest girth, namely, 33.82, or 0.60 inch above the average for the United States. Next come the German and Scandinavian sections, followed by the sparsely settled sections with a large sprinkling of Orientals, the German and Austrian, the Russian, the agricultural sections of mixed foreign and native white, and then desert sections, including many large men, among them many tuberculous patients. Men of the mountain sections have a chest circumference only slightly above the average. The groups of commuter sections, mining, sparsely settled Mexican, eastern manufacturing, and mountain whites are close to the average. At the bottom of the list are the native whites of Scotch origin, whose chest circumference shows up very small, both absolutely and relatively. Next above these are the maritime sections, southern agricultural sections, with a prevalence of whites; French-Canadian sections and agricultural sections, with 45 per cent or more of Negroes. That the Negro agricultural sections of the South have a larger chest circumference than the white agricultural sections, despite the smaller average chest circumference in Negroes, is doubtless due to the fact that in the latter there is a larger proportion of towns and cities in which the chest circumference tends to become reduced. The low chest circumference of French-Canadian sections is due to the small stature of the population in these sections, though relatively the chest girth stands rather high.

Table 46 shows that the sections with 10 per cent Finns, among the most northern of the sections of the United States, have the largest relative chest girth, and that for all other groups it is less than half the stature. According to the table of Martin ⁵ (p. 279) the measure of chest girth of Europeans gives for most races an excess of half the stature, and one is led to inquire if there has been a relative disuse of the arms and chest for severe manual labor in the United States, possibly due to replacement of manual by machine labor.

Next in order come the sections containing 10 per cent or more of agricultural Russians with a relative chest girth of 49.8 per cent. Sections containing a large proportion of French Canadians have a relative chest girth of 49.7. All these sections are engaged primarily in agriculture. Then come the eastern manufacturing and commuter groups, in which the high relative chest circumference must be largely ascribed to racial stock. These are followed by a series of northern, chiefly agricultural, areas, containing Austrians, Scandinavians, and Germans in large proportions. At the end of the series come the Scotch sections, with a chest relative circumference of 48.4, a result which is largely due to the excessive stature of the men from these sections, which is not completely equalized by the increased chest circumference.

TABLE 45.—*Mean chest circumference (expiration) by groups of sections; groups arranged in order of standing with proportional chest circumference (expiration) in inches for each inch of height and each pound of weight; also the standard deviation for each chest circumference; first million draft recruits.*

[From Table VI, p. 440.]

Group No.	Description.	Number of men measured.	Mean chest.	Standard deviation (chest).	Mean chest. Mean height.	Mean chest. Mean weight.	Mean weight. Mean chest.
				Inches.	Inch.	Inch.	Pounds.
	Average for the United States....	873,159	33.22	2.01	0.492	0.234	4.260
18	Finn, 10 per cent.....	5,855	33.82	1.99	.5016	.232	4.311
20	German and Scandinavian, over 10 per cent.....	28,056	33.72	1.95	.4951	.230	4.350
17	Scandinavian, 10 per cent.....	50,953	33.65	1.95	.4952	.230	4.343
8	Sparsely settled, not more than 3 per square mile.....	16,151	33.53	1.92	.4929	.232	4.320
21	German and Austrian, over 20 per cent.....	38,911	33.42	2.07	.4955	.233	4.287
16	Russian, 10 per cent plus.....	12,064	33.39	2.01	.4976	.235	4.264
2	Agricultural, mixed, foreign native white.....	97,319	33.38	2.00	.4934	.234	4.277
9	Desert.....	6,109	33.38	1.99	.4917	.235	4.256
22	German and Austrian, over 15 per cent.....	126,895	33.33	2.06	.4954	.234	4.271
11	Mountain.....	17,103	33.33	1.96	.4921	.233	4.290
6	Commuter.....	28,980	33.25	2.09	.4970	.238	4.205
7	Mining.....	35,691	33.23	1.97	.4929	.234	4.282
14	Mexican, sparsely settled.....	11,064	33.22	1.99	.4874	.234	4.283
5	Eastern manufacturing.....	81,598	33.20	2.08	.4970	.238	4.204
12	Mountain whites.....	21,254	33.20	1.87	.4862	.237	4.225
4	Agricultural Negroes, 45 per cent plus.....	49,465	33.19	1.91	.4894	.234	4.266
1	Agricultural, North, native white over 73 per cent.....	66,836	33.13	1.99	.4900	.234	4.270
13	Indian, sparsely settled.....	10,038	33.13	1.89	.4884	.234	4.283
19	French-Canadian, 10 per cent.....	25,787	33.11	2.07	.4966	.240	4.164
3	Agricultural, native white, South.....	117,800	33.09	1.91	.4854	.240	4.164
10	Maritime.....	6,157	33.00	2.04	.4903	.235	4.255
15	Native whites of Scotch origin.....	13,473	32.95	1.90	.4844	.235	4.260

Next above come the agricultural areas of the South with a prevailingly white population. The mountain whites have also a relatively low chest circumference. The southern agricultural sections with 45 per cent Negroes have a mean relative chest circumference of 48.9, slightly in excess of that of the agricultural areas of the South predominantly white, because the southern white man is lankier than the southern Negro.

TABLE 46.—*Relative chest circumference, by groups of sections (chest circumference divided by stature), first million draft recruits.¹⁰*

Group.	Relative chest circumference.	Group.	Relative chest circumference.
Finns.....	0.502	Mining.....	0.493
Russians, 10 per cent.....	.498	Desert.....	.492
French-Canadians.....	.497	Mountain.....	.492
Commuters.....	.497	Northern agricultural, native white.....	.490
Eastern manufacturing.....	.497	Maritime.....	.490
Germans and Austrians, 20 per cent plus.....	.496	Agricultural, Negro, 45 per cent plus.....	.489
Germans and Scandinavians, 10 per cent plus.....	.495	Mexican, sparsely settled.....	.487
Germans and Austrians, 15 per cent.....	.495	Mountain white.....	.486
Scandinavians, 10 per cent.....	.495	Indian.....	.486
Agricultural, mixed foreign and native white.....	.493	Agricultural, southern whites.....	.485
Sparsely settled.....	.493	Native whites, Scotch origin.....	.484

The relation between the distribution of chest circumference at expiration for each of the principal groups of sections and that of the whole population of recruits is shown graphically in Plate X. The inspection of these curves

shows that the groups containing 10 per cent or more of Finns have the greatest excess of chest girth. This is in accord with what we have already found regarding the robustness of the men of these sections.

Similarly the groups of sections characterized by having 10 per cent of Scandinavians are characterized by large chest girth and this is associated with what we have found in regard to the great stature and heavy build of men in this group of sections. Also the groups with 10 per cent or more of Russians are characterized by a slight excess of chest girth. On the other hand, the groups of sections containing a large proportion of men of Scotch origin are characterized by a deficiency of chest girth. This agrees with what we have already found concerning the lankness of form of the men of this group.

The graphs show, moreover, that the chest circumference of sections comprising half, or more, Negroes are on the average larger than those sections of the South containing a smaller proportion of Negroes. The sections containing 10 per cent or more of French Canadians are characterized by a deficiency of chest circumference.

TABLE 47.—*Distribution of chest circumference (expiration) shown by groups of sections, first million draft recruits.*

SECTION A: ABSOLUTE NUMBERS.

Group No.	Description.	Number measured.	Chest, in inches.										
			27	30	31	32	33	34	35	36	37	38	39
1	Agricultural, North, native white over 73 per cent.....	66,795	1,451	4,944	8,483	12,715	13,752	11,371	7,455	4,025	1,870	842	887
2	Foreign and native white.....	97,338	1,693	4,696	10,247	16,749	19,807	17,829	12,632	7,363	3,541	1,576	1,205
3	Agricultural, native white, South.....	117,890	2,421	6,776	14,684	22,478	25,217	21,006	13,163	6,924	2,884	1,324	1,013
4	Agricultural, Negroes, 45 per cent plus.....	49,447	853	2,591	5,673	9,162	10,378	9,227	5,917	3,287	1,381	638	340
5	Eastern manufacturing.....	81,569	2,047	5,018	10,234	14,920	15,799	13,552	9,209	5,438	2,725	1,371	1,256
6	Commuter.....	28,994	737	1,771	3,542	5,072	5,594	4,839	3,425	2,059	1,026	493	436
7	Mining.....	35,686	731	1,835	4,089	6,404	7,221	6,452	4,395	2,522	1,136	504	397
8	Sparsely settled, not more than 3 per square mile.....	16,143	157	606	1,456	2,704	3,348	3,228	2,325	1,316	579	249	175
9	Desert.....	6,110	99	265	644	1,097	1,279	1,098	784	449	221	94	80
10	Maritime.....	6,157	214	395	797	1,186	1,270	984	607	394	176	62	72
11	Mountain.....	17,101	259	800	1,849	3,080	3,541	3,234	2,127	1,251	575	226	179
12	Mountain whites.....	21,233	328	1,001	2,388	4,082	4,661	3,923	2,571	1,353	546	215	162
13	Indian, sparsely settled.....	10,035	178	468	1,260	1,979	2,144	1,829	1,151	616	201	129	80
14	Mexican, sparsely settled.....	11,064	221	586	1,271	2,029	2,373	1,911	1,327	718	347	159	122
15	Native whites of Scotch origin.....	13,469	303	826	1,855	2,811	2,904	2,213	1,315	730	270	129	113
16	Russian, 10 per cent plus.....	12,057	228	627	1,305	2,075	2,390	2,215	1,570	973	421	205	148
17	Scandinavian, 10 per cent.....	50,951	501	1,999	4,286	7,887	10,296	10,221	7,548	4,544	2,258	984	627
18	Finn, 10 per cent.....	5,855	49	163	421	815	1,145	1,123	869	614	322	149	85
19	French Canadian, 10 per cent.....	25,772	645	1,597	3,340	5,016	5,066	4,189	2,796	1,605	753	395	381
20	German and Scandinavian, 10 per cent and over.....	28,051	241	829	2,240	4,220	5,666	5,752	4,305	2,551	1,313	563	371
21	German and Austrian, over 20 per cent.....	38,943	723	1,934	4,147	6,582	7,662	7,112	4,924	3,026	1,499	703	631
22	German and Austrian, over 10 per cent.....	126,887	2,639	6,839	14,168	21,909	24,971	22,624	15,871	9,377	4,616	2,147	1,726
	Total.....	867,547	16,718	45,166	98,379	155,152	176,483	155,935	106,276	61,135	28,660	13,157	10,496

TABLE 47.—*Distribution of chest circumference (expiration) shown by groups of sections, first million draft recruits—Continued.*


SECTION B: RATIOS PER 1,000.

Group No.	Description.
1	Agricultural, North, native white over 78 per cent.....
2	Foreign and native white.....
3	Agricultural, native white, South.....
4	Agricultural, Negroes, 45 per cent plus.....
5	Eastern manufacturing.....
6	Commuter.....
7	Mining.....
8	Sparsely settled, not more than 3 per square mile.....
9	Desert.....
10	Maritime.....
11	Mountain.....
12	Mountain whites.....
13	Indian, sparsely settled.....
14	Mexican, sparsely settled.....
15	Native whites of Scotch origin.....
16	Russian, 10 per cent plus.....
17	Scandinavian, 10 per cent.....
18	Finn 10 per cent.....
19	French Canadian, 10 per cent.....
20	German and Scandinavian, 10 per cent and over.....
21	German and Austrian 20 per cent and over.....
22	German and Austrian over 10 per cent.....
	Total.....

11. MEAN CHEST CIRCUMFERENCE OF THE EIGHT EUROPEAN RACES OF MEN AT DEMOBILIZATION.

Table 49 gives the absolute and proportional frequencies of the different classes of chest circumference for men of the eight European races as taken at demobilization. These are summarized in Table 48. The second column of Table 48 gives the average chest circumference at rest. The greatest average chest circumference, 90.42, is found among the Poles, the next among the Germans, followed by Italians and Irish. The smallest chest circumference, 87.53, is found among the Hebrews; markedly above stand the English, followed by the French and Scotch. Our measurements correspond rather closely with Gould's. We may, therefore, compare his measurements (after reduction to centimeters) on page 280 "after expiration" with those of the present work. Thus, for the measurements in 1866 of the English, Gould gets 87.12 centimeters, about 1 centimeter less than the English troops measured half a century later. Gould's figures are: Chest circumference of the Scotch, 88.06, as contrasted with 88.57, 50 years later; of men from Ireland, 89.28, as contrasted with 88.67, which shows a reduction of 0.5 centimeter; of the French, etc., 87.12, as contrasted with our figure of 88.49, showing a marked increase; of the Germans 88.19, as contrasted with our average of 89.52, showing a marked increase. In general, excepting the Irish, the mean chest circumference for our races is greater than for those of Gould. This is largely due to the fact that in Gould's measurements, the circumference of the chest was taken at full inspiration, whereas in the present series it was taken of the chest at rest.

CHEST (EXP.) DISTRIBUTION BY GROUPS OF SECTIONS (N)



The middle column of Table 48 gives the standard deviation or index of variability for the chest circumference of the eight races. From this column it appears that the Irish are the most variable in respect to chest circumference, which may be due to the combination in that rubric of tall Scotch-Irish and the more thickset Celtic-Irish. Next in order come the Scotch, then the Hebrews and Germans. The lowest index of variability, 4.94, is found among the Italians, followed by the English, French, and Polish.

The second column from the right shows the proportion of chest circumference to total stature for each of the races. From this column it appears that in relation to stature the Italians have the largest chest circumference, followed by the Poles, French, and Hebrews. The English have the smallest relative chest circumference, 51.24, followed by the Scotch, Irish, and Germans. Thus it appears that the Mediterranean races, Poles, and Hebrews are relatively larger chested than the Nordics. Since chest circumference is not very closely correlated with stature, this difference in relative chest circumference is largely dependent upon the varying size of the divisors (stature) used in finding the quotients. Rather more to the point would be the relation of chest circumference to weight of the body and these quotients have been calculated and are given in the last column to the right of Table 48. According to the last column we find the greatest chest in relation to weight among the Italians, 0.644; next largest among the Hebrews and then the French and Polish. The smallest relation of chest to weight is found among the Germans, 0.604, next larger English, 0.608; Scotch, 0.611; and Irish, 0.620. This result runs somewhat parallel to the preceding column and justifies the general conclusion that whether in relation to the stature or in relation to weight the Mediterranean races and the Hebrews have a larger relative chest girth than the Nordic races.

TABLE 48.—*Absolute and relative chest circumference (rest) of eight European races, with standard deviation and the coefficient of variation for each, demobilization, 1919.*

Race.	Number measured.	Absolute chest circumference.	Standard deviation.	Coefficient of variation.	Relative chest circumference to stature.	Relative chest circumference to weight.
		<i>Centimeters.</i>	<i>Centimeters.</i>	<i>Per cent.</i>	<i>Per cent.</i>	<i>Per cent.</i>
English.....	4, 205	88.18	5.00	5.670	51.24	0.608
Scotch.....	2, 067	88.57	5.25	5.928	51.33	0.611
Irish.....	6, 142	88.67	5.31	5.989	51.74	0.620
German.....	7, 070	89.52	5.17	5.774	52.03	0.604
French.....	1, 450	88.49	5.08	5.741	52.49	0.623
Italian.....	3, 524	88.87	4.94	5.558	53.80	0.644
Polish.....	2, 409	90.42	5.11	5.651	53.37	0.621
Hebrew.....	1, 691	87.53	5.19	5.929	52.44	0.635

TABLE 49.—Comparative frequency distribution of chest circumference (real) in each of eight races, demobilization.

SECTION A: ABSOLUTE NUMBERS.																												
Race.	Chest circumference, in centimeters.																											
	Total.																											
	63-64	65-66	67-68	69-70	71-72	73-74	75-76	77-78	79-80	81-82	83-84	85-86	87-88	89-90	91-92	93-94	95-96	97-98	99-100	101-102	103-104	105-106	107-108	109-110	111-112	113-114	115-116	
English.....	4,206				3	7	14	44	125	268	508	654	755	821	471	348	182	111	73	33	11	3	4	2	2	1	1	
Scottish.....	2,067	1			1	5	9	28	56	128	232	283	382	377	254	183	92	80	44	13	9	5	2	2	1			
Irish.....	6,142	2			3	8	18	41	176	374	696	978	1,118	1,142	724	518	325	212	128	57	29	18	12	7	3	3		
German.....	7,070	2			1	6	17	49	133	310	612	829	1,118	1,142	724	518	497	329	156	76	39	25	12	4	2	1		
French.....	1,480				3	2	12	21	53	103	183	227	242	213	148	118	87	51	28	12	3	2	2	2	1			
Italian.....	3,428	1			2	6	14	27	78	171	333	463	594	692	493	318	211	124	68	22	9	7	2	1	2			
Polish.....	2,469	3			2	7	16	36	106	208	333	463	594	692	493	318	211	124	68	22	9	7	2	1	2			
Hebrew.....	1,681				1	3	4	39	70	137	269	385	484	540	143	114	60	43	20	12	9	4	2	1	1			
Number measured.....	28,368	8	2	8	11	15	36	81	242	706	1,542	2,843	3,945	4,411	3,642	2,638	1,679	1,076	587	250	118	70	31	22	13	5	1	
Not measured.....	102																											
Total.....	28,670																											

SECTION B: PROPORTIONAL RATIOS PER 1,000.

12. CHEST CIRCUMFERENCE OF MEN OF THE COLOR RACES.

The following table, derived from Tables 103 and 104, gives the means of comparing the two principal color races measured at demobilization. It will be recalled that no distinction of color races was made in the original schedules for recording measurements of drafted men.

TABLE 50.—*Mean and relative chest circumference (rest), white and Negro troops, demobilization, 1919.*

Race.	Number measured.	Mean in centimeters.	Relative chest circumference.
White.....	95,867	88.79	51.6
Colored (Negro).....	6,355	87.99	51.2

The table indicates that the chest circumference of the white troops exceeds that of the Negro troops by 8 millimeters. In relation to height the chest circumference of the Negro troops is slightly less than that of the white troops.

TABLE 51. — *Various heights, weights, and chest circumferences (expiration) shown for the United States, with ratio per 1,000 of each, first million draft recruits.*

Height to weight (Table I).					Weight to chest (Table II).					Weight to chest (Table III).					
Height.	Number of men measured.	Ratio per 1,000.	Weight.	Number of men measured.	Ratio per 1,000.	Height.	Number of men measured.	Ratio per 1,000.	Chest.	Number of men measured.	Ratio per 1,000.	Chest.	Number of men measured.	Ratio per 1,000.	
58 and under.	3,124	3.60	92	184	0.21	58 and under.	3,086	3.53	28 and under.	92	213	28 and under.	17,933	20.56	
59	2,857	3.32	97	2,356	0.21	59	2,921	3.35	29	102	2,313	29	49,056	56.23	
60	15,044	8.61	107	7,435	8.56	60	15,752	8.16	30	107	21,381	30	103,277	118.38	
61	30,935	35.62	117	21,938	24.63	61	31,848	18.15	31	112	41,665	31	199,456	182.77	
62	32,547	33.53	122	41,953	47.79	62	31,207	35.74	32	117	63,866	32	175,770	201.47	
63	81,904	94.31	127	63,567	73.20	63	82,923	60.61	33	122	85,072	33	152,555	174.86	
64	109,804	126.62	132	100,064	97.56	64	110,816	62.92	34	127	100,715	34	103,381	118.50	
65	127,864	147.21	137	100,869	115.24	65	128,291	64.90	35	132	107,129	35	58,967	67.48	
66	128,768	149.98	142	100,807	123.08	66	130,624	66.92	36	137	101,040	36	28,121	32.23	
67	110,768	127.55	147	111,523	127.08	67	131,523	67.87	37	142	86,316	37	15,065	14.98	
68	83,762	94.59	152	87,332	83.22	68	83,609	62.15	38	147	82,618	38	5,928	6.68	
69	37,370	42.59	157	39,731	41.33	69	34,699	32.10	39	152	78,948	39	5,110	5.86	
70	21,370	23.12	162	20,093	21.53	70	31,523	27.10	40 and over.	157	38,948	40 and over.			
71	15,636	17.36	167	13,536	15.37	71	18,411	17.34		162	28,151				
72	8,301	9.36	172	7,054	8.41	72	16,411	15.40		167	16,052				
73	6,301	7.36	177	5,654	6.41	73	13,620	12.54		172	13,062				
74	7,071	7.91	182	6,385	7.26	74	1,080	1.24		177	9,310				
75	3,861	4.30	187	3,482	3.94	75	1,361	1.54		182	5,546				
76	2,959	3.30	192	2,907	3.26	76	256	0.29		187	3,653				
77	2,968	3.32	197	2,963	3.32	77	268	0.30		192	2,967				
78	268	.34	202	4,688	5.40	78				197	5,432				
79						79				202					
80 and over.						80 and over.									
Total.	868,445	1,000.00		868,445	1,000.00		873,159	1,000.00			872,419	1,000.00		872,419	1,000.00

TABLE 52.—*Height and weight classes—Mean weight and the standard deviation for each height; also mean height and the standard deviation for each weight; derived from summation of sections (Table I); first million Draft Recruits.*

Height.	Number of men measured.	Mean weight.	Standard deviation.	Weight.	Number of men measured.	Mean height.	Standard deviation.
<i>Inches.</i>		<i>Pounds.</i>	<i>Pounds.</i>	<i>Pounds.</i>		<i>Inches.</i>	<i>Inches.</i>
59.....	3, 124	135.98	18.95	97.....	184	62.38	2.03
60.....	2, 887	128.11	20.30	102.....	2, 356	63.92	2.90
61.....	7, 477	124.80	19.29	107.....	7, 435	64.02	2.34
62.....	15, 644	125.24	17.66	112.....	21, 388	64.51	2.20
63.....	30, 935	127.49	16.46	117.....	41, 503	65.16	2.17
64.....	52, 547	130.24	15.41	122.....	63, 567	65.79	2.20
65.....	81, 904	133.11	14.87	127.....	84, 726	66.34	2.23
66.....	106, 964	136.24	14.72	132.....	100, 084	66.86	2.26
67.....	127, 844	139.46	15.18	137.....	106, 889	67.33	2.28
68.....	129, 987	142.82	16.13	142.....	100, 607	67.80	2.33
69.....	110, 508	146.25	17.63	147.....	88, 057	68.17	2.35
70.....	83, 702	149.49	19.28	152.....	72, 362	68.56	2.41
71.....	54, 357	153.26	21.55	157.....	53, 431	68.88	2.46
72.....	31, 370	156.64	23.84	162.....	39, 797	69.15	2.50
73.....	15, 198	160.40	26.52	167.....	29, 063	69.32	2.62
74.....	6, 391	163.90	29.14	172.....	18, 854	69.57	2.83
75.....	2, 620	166.85	31.66	177.....	12, 629	69.66	2.68
76.....	1, 071	167.30	32.48	182.....	8, 385	69.76	2.76
77.....	360	166.05	33.41	187.....	5, 467	69.70	2.73
78.....	259	161.89	31.08	192.....	3, 907	69.65	2.82
79 and over.....	296	158.05	30.58	197.....	2, 966	69.38	3.20
				212 and over.....	4, 688	70.16	2.78
Total.....	868, 445				868, 445		

Mean height: 67.49 inches; standard deviation, 2.71 inches.

Mean weight: 141.54 pounds; standard deviation, 17.42 pounds.

TABLE 53.—*Height and chest circumference (expiration) classes—Mean chest circumference (expiration) and the standard deviation for each height; also the standard deviation for each chest circumference; derived from summation of sections (Table II); first million Draft Recruits.*

Height.	Number of men measured.	Mean chest.	Standard deviation.	Chest.	Number of men measured.	Mean height.	Standard deviation.
<i>Inches.</i>		<i>Inches.</i>	<i>Inches.</i>	<i>Inches.</i>		<i>Inches.</i>	<i>Inches.</i>
59.....	3, 086	32.94	2.08	29.....	18, 063	63.91	2.55
60.....	2, 921	32.49	2.01	30.....	49, 090	66.29	2.54
61.....	7, 572	32.28	1.93	31.....	103, 294	66.71	2.56
62.....	15, 848	32.33	1.91	32.....	159, 379	67.12	2.60
63.....	31, 207	32.46	1.93	33.....	175, 858	67.51	2.63
64.....	52, 923	32.59	1.95	34.....	152, 063	67.85	2.66
65.....	82, 426	32.77	1.96	35.....	103, 414	68.11	2.70
66.....	110, 816	32.92	1.98	36.....	59, 015	68.36	2.73
67.....	128, 291	33.10	1.96	37.....	28, 175	68.47	2.76
68.....	130, 624	33.29	1.96	38.....	13, 151	68.51	2.75
69.....	111, 123	33.49	1.96	39 and over.....	11, 027	68.59	2.76
70.....	83, 880	33.68	1.95				
71.....	54, 609	33.86	1.95				
72.....	31, 523	34.06	1.96				
73.....	15, 294	34.28	1.93				
74.....	6, 411	34.46	1.94				
75.....	2, 620	34.65	1.96				
76.....	1, 080	34.57	1.98				
77.....	361	34.60	2.00				
78.....	256	34.48	1.98				
79 and over.....	296	34.30	2.21				
Total.....	873, 591				873, 591		

Height: Mean, 67.49 inches; standard deviation, 2.72 inches.

Chest circumference (expiration): Mean, 33.22 inches; standard deviation, 2.01 inches.

WEIGHT DISTRIBUTION BY HEIGHT (P.)

RATIOS PER 1000


2

PURE LINE CURVE DENOTES AVERAGE FOR U. S.
PLATE XII.

CHEST (EXP.) DISTRIBUTION BY HEIGHT (P.)
RATIOS PER 1000

3

1



FINE LINE CURVE DENOTES AVERAGE FOR U.S.
PLATE XIII.

TABLE 54.—*Weight and chest circumference (expiration) classes.—Mean chest circumference (expiration) and the standard deviation for each weight; also the mean weight and the standard deviation for each chest circumference; derived from summation of sections (Table III); first million draft recruits.*¹⁹

Weight.	Number of men measured.	Mean chest.	Stand-ard deviation.	Chest.	Number of men measured.	Mean weight.	Stand-ard deviation.
<i>Pounds.</i>		<i>Inches.</i>	<i>Inches.</i>	<i>Inches.</i>		<i>Pounds.</i>	<i>Pounds.</i>
97	213	29.61	1.15	29	17,933	119.53	12.92
102	2,313	30.59	2.14	30	49,056	123.64	10.98
107	7,391	30.56	1.71	31	103,277	128.56	10.97
112	21,382	30.82	1.47	32	159,456	134.02	11.29
117	41,665	31.20	1.43	33	175,770	139.66	11.89
122	63,866	31.64	1.40	34	152,555	145.64	12.58
127	85,072	32.06	1.40	35	103,381	151.69	13.37
132	100,715	32.47	1.40	36	58,867	158.33	14.48
137	107,129	32.88	1.41	37	28,121	165.53	16.09
142	101,040	33.28	1.43	38	13,065	172.95	17.85
147	88,316	33.69	1.44	39	5,828	178.10	21.69
152	72,618	34.08	1.46	40 and over	5,110	185.48	24.32
157	53,688	34.46	1.48				
162	39,998	34.85	1.52				
167	29,141	35.17	1.60				
172	19,052	35.61	1.63				
177	12,692	36.00	1.71				
182	8,210	36.44	1.68				
187	5,566	36.82	1.79				
192	3,853	37.14	1.91				
197	2,867	37.14	2.49				
202 and over	5,432	38.70	1.63				
	872,419				872,419		

Weight: Mean, 141.59 pounds; standard deviation, 17.49 pounds.

Chest circumference (expiration): Mean, 33.23 inches; standard deviation, 2.03 inches.

V. BUILD.

It is clear that the absolute weight and chest circumference are relatively unimportant in giving an idea of the build of man, unless we know something about his stature. It is customary, therefore, to consider not only these absolute measurements, but also these measurements in relation to stature. Weight considered in relation to stature gives us an index of build. A formula which will combine in proper fashion the weight, stature, and chest circumference will give us an index of robustness. The latter will be considered in another section.

1. IMPORTANCE OF THE INDEX OF BUILD.

Important as stature and weight are for military and medico-military purposes, they are hardly as important as the index of build, which tells us something about the physical constitution of a man, and, by implication and as a result of experience, also something about his ability to withstand the stress of warfare. The relativeness of weight to height has been long recognized in the Army, where the tables indicate the limitations of weight for men of respective height. Such is shown in Table 138. In fact, it is not too much to say that the principal reason for taking weight in connection with height is to secure a numerical statement of the build as a first means of deciding upon the acceptance or rejection of the recruit for military service.

2. METHOD OF DETERMINING.

The best method of expressing the index of build is not easily determined. The simplest method and that used by Army, Life Insurance examiners, etc., is that of dividing the weight by the stature, recognizing that in tall (large) persons the absolute increment per inch is greater than in short (small) persons. This method would be without objection if the body of men were cylinders of equal diameter but of varying height. In such a case the index would be constant, since the differences in weight would correspond to the differences in stature. It is clear, however, that the form of the body departs somewhat from this assumption.

If the body were a cube or sphere then body weight would vary as the cube of any one of the diameters, and the index of build would be most properly given by dividing the weight by the cube of any one of the diameters; but the body does not fulfill these conditions. Finally, it has been pointed out that inasmuch as the form of the body lies between the two hypothetical conditions just mentioned a more suitable index of build would be obtained by dividing the weight by the second power of the stature. Such a method was indeed discussed by Gould^a and it was shown by him to meet very satisfactorily the requirements of the index of build.

To decide between the foregoing methods of measuring the index of build, comparative tables have been made, Tables 55 and 56, giving the result of applying the three formulæ. That series must be regarded as the most satisfactory which gives a fairly constant quotient when applied to figures from different parts of the general correlation table of stature and weight on page 417. By comparing columns 3, 4, and 5, which give respectively the index obtained by the three methods described, it is to be noted that column 4 (weight in pounds $\times 1,000 \div$ by the square of the height) is the most constant, but that the index falls somewhat from the short stature of 61 inches to the tall stature of 74 inches. There is indeed some reason to believe that the weight of short men does not diminish pro rata with the stature and, therefore, this decrease in the size of the index obtained in column 4 agrees with the apparent facts. Column 3 tells a different story from column 4. It shows how sections of the body an inch thick weigh absolutely more in tall men than in short ones. The ratio of column 5 is of the same order as that of column 4, but shows a still more marked decrease in build, passing from 61 to 74 inches. The matter of choice between these three methods has been fully discussed elsewhere. Here may be given only the conclusion that in accordance with the findings of Gould and, before him, Quetelet, the ratio of weight divided by the second power of the height seems to be the most satisfactory index of build, and is one which we shall largely use in this section.

^a Subject first elaborated by Quetelet in 1835. See Baxter,¹ Vol. I, p. 52.

TABLE 55.—*Index of build calculated by three methods (based on Table 1, first million draft recruits).*

MODAL WEIGHT.

Height.	Modal weight.	Weight in pounds ($\times 10$).	Weight (\times 1,000).	Weight (\times 100,000).
		Height in inches.	Height (sq.).	Height (cubed).
<i>Inches.</i>	<i>Pounds.</i>			
61	117	19.18	31.44	51.55
62	117	18.87	30.44	49.09
63	122	19.37	30.74	48.79
64	127	19.84	31.01	48.45
65	127	19.54	30.06	46.24
66	132	20.00	30.30	45.91
67	137	20.45	30.52	45.55
68	137	20.15	29.63	43.57
69	142	20.58	29.83	43.23
70	147	21.00	30.00	42.84
71	153	21.41	30.15	42.47
72	152	21.11	29.32	40.72
73	157	21.51	29.46	40.36
74	157	21.22	28.67	38.74

TABLE 56.—*Index of build calculated by three methods (based on Table 1, first million draft recruits).*

AVERAGE WEIGHT.

Height.	Average weight.	Weight in pounds ($\times 10$).	Weight (\times 1,000).	Weight (\times 100,000).
		Height in inches.	Height (sq.).	Height (cubed).
<i>Inches.</i>	<i>Pounds.</i>			
61	135.96	22.29	36.54	59.91
62	128.11	20.66	33.33	53.76
63	124.80	19.81	31.44	49.91
64	125.24	19.57	30.58	47.77
65	127.49	19.61	30.17	46.42
66	130.24	19.74	29.90	45.31
67	133.11	19.87	29.65	44.26
68	136.24	20.04	29.46	43.33
69	139.46	20.21	29.29	42.45
70	142.82	20.40	29.15	41.64
71	146.25	20.60	29.01	40.86
72	149.49	20.76	28.84	40.06
73	153.26	20.99	28.76	39.40
74	156.64	21.17	28.61	38.66

3. INDEX OF BUILD FOR MEAN STATURE AND WEIGHT.

If we divide the mean weight ($\times 1,000$) of the whole population by the square of the mean height, we shall obtain by probably the most accurate method an average index of build of the whole population. The following brief table gives the average index of build thus obtained:

Recruits, World War.....	31.08
Men at demobilization, 1919.....	31.59
Earlier series of Gould (pp. 284-3).....	31.47
Later series, Gould (p. 403).....	31.35

4. THE INDEX OF BUILD OF CIVIL AND WORLD WAR VETERANS FOR EACH INCH OF STATURE.

Table 57 gives the index of build of veterans of the World War and Civil War. It appears that while men 70 inches tall or less were more robust in 1919, those from 71 to 75 inches were less robust in 1919 than in 1865. This is largely because the later figures contain many Southerners of slender build, who were absent from the earlier Civil War series. In the figures for the

World War veterans, the Negro troops are included. However, as the number of them was small they probably affect the average but slightly.

TABLE 57.—*Index of build of Civil War veterans^a (white troops) and World War veterans^b (white and Negro troops).*

Stature classes.	Civil War.		World War.	
	Weight.	Weight ($\times 1,000$). Height (sq.).	Weight.	Weight ($\times 1,000$). Height (sq.).
	<i>Pounds.</i>		<i>Pounds.</i>	
60.....	111.79	31.05	123.00	34.17
61.....	117.60	31.60	125.66	33.77
62.....	120.77	31.42	127.10	33.06
63.....	122.95	30.98	129.78	32.69
64.....	128.43	31.35	131.84	32.19
65.....	132.12	31.27	135.20	32.00
66.....	136.06	31.24	139.26	31.97
67.....	140.77	31.36	142.71	31.79
68.....	144.92	31.34	145.52	31.47
69.....	149.04	31.30	148.39	31.38
70.....	153.19	31.26	153.30	31.29
71.....	158.21	31.38	156.81	31.13
72.....	162.48	31.34	159.84	30.83
73.....	166.39	31.22	164.03	30.78
74.....	168.99	30.86	163.54	29.87
75.....	170.39	30.29	168.00	29.87

^a Calculated from Gould, p. 408, Table IX.

^b Calculated from Table No. LXXIV.

5. DISTRIBUTION OF INDEX OF BUILD, BY STATES.

Table 58 gives the distribution of the index of build at mobilization by States. In this table there are four columns. The first two give the index of build (mean weight divided by the square of the mean stature of recruits) and the last two columns give for the successive States another index of build obtained by dividing the mean weight by the first power of the mean stature. By squaring the stature, differences in stature are exaggerated and consequently the range of the first two columns seems more significant and the order of the States is, therefore, more important in this case. Of all States and Territories, Alaska stands first in robustness of its drafted men. This is followed by North Dakota, South Dakota, Montana, Minnesota, Wisconsin, Nevada, and Oregon. The men of the Northwest are tall men, but they are relatively so heavy that there is in those States a high index of build. In other words, they are large men. However, in the case of Wisconsin the high index of build is partly due to the relatively short stature (although above the average) of its drafted men. Examining now the bottom of the table, we find that, using the second power of stature as the divisor, the drafted men from Tennessee and Kentucky lie at the very bottom of the list. Men from these States have practically the same mean weight, but the men from Tennessee are taller. Accordingly, their index of build is much less than that of men from Kentucky. Indeed, they are the least robust of those of any State. The low rank of these States is due especially to mountain sections, although the men of Tennessee seem to be of the tall, slender type throughout the State. Next in order comes Colorado with an index of build of 30.37. The men from this State are not only tall but they are below the average in weight, a condition which is probably associated with the immigration of tuberculous patients to that State.

By any method of calculating build, the Southern States tend to lie toward the bottom of the list. Thus in column 1, Arkansas, Texas, Georgia, North Carolina, Florida, Virginia, Alabama, Louisiana, South Carolina, and Mississippi occupy relatively low positions. This low position is due both to the great stature of the men of these States and also to their relatively low mean weight. On account of the prevalence of malaria in these Southern States, as well as hookworm in many of them, it seems probable that the low index of build is due in part to the combination of these parasitic diseases. In addition, the low position of New Mexico is doubtless to be ascribed to the large amount of tuberculosis in the population. The low index of build of the men of Oklahoma is due to their great stature combined with only an average weight.

In the second list of States in Table 58, calculated by using as divisor the first power of the weight, the same general statement made above concerning the build of men from various parts of the country holds, though the order of the States is somewhat shifted.

TABLE 58.—*Index of build at mobilization, by States, 1917-1918.*

State.	Weight \times 1,000. (Height (sq.))	State.	Weight. Height.
United States.....	31.07	United States.....	2.097
Alaska.....	32.41	Alaska.....	2.208
North Dakota.....	31.85	North Dakota.....	2.163
South Dakota.....	31.73	South Dakota.....	2.159
Montana.....	31.64	Montana.....	2.151
Minnesota.....	31.63	Oregon.....	2.150
Wisconsin.....	31.62	Minnesota.....	2.150
Nevada.....	31.59	Nevada.....	2.143
Oregon.....	31.57	Washington.....	2.140
Washington.....	31.48	Wisconsin.....	2.137
Wyoming.....	31.47	Idaho.....	2.133
California.....	31.44	Wyoming.....	2.130
Connecticut.....	31.42	Nebraska.....	2.128
Michigan.....	31.41	Iowa.....	2.128
Pennsylvania.....	31.38	California.....	2.127
New York.....	31.34	Michigan.....	2.110
Idaho.....	31.33	Utah.....	2.109
New Hampshire.....	31.29	Kansas.....	2.107
Iowa.....	31.26	Illinois.....	2.103
Nebraska.....	31.22	Mississippi.....	2.100
Illinois.....	31.21	Maine.....	2.100
Maryland.....	31.20	Arizona.....	2.099
Vermont.....	31.16	Ohio.....	2.098
Maine.....	31.15	New Hampshire.....	2.095
Ohio.....	31.14	Connecticut.....	2.095
New Jersey.....	31.14	Pennsylvania.....	2.094
Utah.....	31.09	Vermont.....	2.091
Massachusetts.....	31.05	New York.....	2.091
Rhode Island.....	30.95	Indiana.....	2.090
Kansas.....	30.90	Maryland.....	2.090
Delaware.....	30.89	West Virginia.....	2.085
Indiana.....	30.88	Oklahoma.....	2.084
Arizona.....	30.81	Missouri.....	2.081
West Virginia.....	30.73	Texas.....	2.079
District of Columbia.....	30.72	New Jersey.....	2.079
Mississippi.....	30.72	Alabama.....	2.077
South Carolina.....	30.70	District of Columbia.....	2.077
Missouri.....	30.63	South Carolina.....	2.077
Louisiana.....	30.55	North Carolina.....	2.076
Alabama.....	30.55	Delaware.....	2.075
Virginia.....	30.53	Arkansas.....	2.071
Oklahoma.....	30.53	Georgia.....	2.071
Florida.....	30.51	Virginia.....	2.070
North Carolina.....	30.47	Massachusetts.....	2.070
Georgia.....	30.46	Colorado.....	2.069
Texas.....	30.40	Louisiana.....	2.065
New Mexico.....	30.39	Florida.....	2.061
Arkansas.....	30.37	Rhode Island.....	2.060
Colorado.....	30.37	Kentucky.....	2.058
Kentucky.....	30.26	Tennessee.....	2.052
Tennessee.....	30.06	New Mexico.....	2.061

TABLE 59.—*Index of build at demobilization, by States, 1919.*

State.	Weight \times 1,000. Stature (sq.).	State.	Weight. Stature.
Alaska.....	33.60	Alaska.....	2.333
North Dakota.....	32.67	South Dakota.....	2.225
South Dakota.....	32.54	North Dakota.....	2.220
Minnesota.....	32.44	Minnesota.....	2.216
Nevada.....	32.42	Idaho.....	2.212
Idaho.....	32.40	Montana.....	2.211
Montana.....	32.34	Nebraska.....	2.210
Nebraska.....	32.29	Nevada.....	2.201
Iowa.....	32.19	Iowa.....	2.198
Wisconsin.....	32.18	Kansas.....	2.194
Utah.....	32.10	Utah.....	2.189
Kansas.....	32.06	Wisconsin.....	2.181
Michigan.....	32.01	Wyoming.....	2.178
Wyoming.....	31.95	Arizona.....	2.171
Louisiana.....	31.79	Washington.....	2.170
Illinois.....	31.77	Oklahoma.....	2.169
Arizona.....	31.77	Oregon.....	2.169
Colorado.....	31.75	Colorado.....	2.163
Washington.....	31.73	Louisiana.....	2.158
Pennsylvania.....	31.73	Michigan.....	2.155
Oregon.....	31.72	Mississippi.....	2.150
Ohio.....	31.72	West Virginia.....	2.150
Oklahoma.....	31.70	Illinois.....	2.150
Maine.....	31.69	Texas.....	2.148
Rhode Island.....	31.66	Virginia.....	2.147
Vermont.....	31.61	Arkansas.....	2.146
Virginia.....	31.58	North Carolina.....	2.143
Indiana.....	31.56	Missouri.....	2.143
Missouri.....	31.53	Ohio.....	2.141
West Virginia.....	31.52	California.....	2.140
California.....	31.52	Indiana.....	2.138
Delaware.....	31.44	Maine.....	2.128
North Carolina.....	31.41	Pennsylvania.....	2.126
Maryland.....	31.40	Vermont.....	2.123
Arkansas.....	31.37	New Mexico.....	2.123
New York.....	31.36	Tennessee.....	2.121
Massachusetts.....	31.35	South Carolina.....	2.121
Connecticut.....	31.34	Kentucky.....	2.121
Mississippi.....	31.34	Delaware.....	2.114
Texas.....	31.31	Alabama.....	2.112
New Jersey.....	31.31	Maryland.....	2.110
New Mexico.....	31.30	Rhode Island.....	2.107
Kentucky.....	31.13	Connecticut.....	2.103
South Carolina.....	31.04	Georgia.....	2.101
Tennessee.....	30.92	New York.....	2.098
District of Columbia.....	30.81	New Jersey.....	2.096
Alabama.....	30.79	Massachusetts.....	2.093
New Hampshire.....	30.69	District of Columbia.....	2.083
Georgia.....	30.66	Florida.....	2.074
Florida.....	30.40	New Hampshire.....	2.050

ARMY AIRFIELD LOG

[illegible]

State	Weight Height	Weight Height	Percent of in- crease or decrease	State	Weight Height	Percent of in- crease or decrease
United States	1.71	1.74	United States	1.73	1.77	1.7
Alabama	1.79	1.7	Alabama	1.75	1.7	1.7
Alaska	1.79	1.7	Alaska	1.75	1.7	1.7
Arizona	1.79	1.7	Arizona	1.75	1.7	1.7
Arkansas	1.79	1.7	Arkansas	1.75	1.7	1.7
California	1.79	1.7	California	1.75	1.7	1.7
Colorado	1.79	1.7	Colorado	1.75	1.7	1.7
Connecticut	1.79	1.7	Connecticut	1.75	1.7	1.7
Delaware	1.79	1.7	Delaware	1.75	1.7	1.7
District of Columbia	1.79	1.7	District of Columbia	1.75	1.7	1.7
Florida	1.79	1.7	Florida	1.75	1.7	1.7
Georgia	1.79	1.7	Georgia	1.75	1.7	1.7
Hawaii	1.79	1.7	Hawaii	1.75	1.7	1.7
Idaho	1.79	1.7	Idaho	1.75	1.7	1.7
Illinois	1.79	1.7	Illinois	1.75	1.7	1.7
Indiana	1.79	1.7	Indiana	1.75	1.7	1.7
Iowa	1.79	1.7	Iowa	1.75	1.7	1.7
Kansas	1.79	1.7	Kansas	1.75	1.7	1.7
Kentucky	1.79	1.7	Kentucky	1.75	1.7	1.7
Louisiana	1.79	1.7	Louisiana	1.75	1.7	1.7
Maine	1.79	1.7	Maine	1.75	1.7	1.7
Maryland	1.79	1.7	Maryland	1.75	1.7	1.7
Massachusetts	1.79	1.7	Massachusetts	1.75	1.7	1.7
Michigan	1.79	1.7	Michigan	1.75	1.7	1.7
Minnesota	1.79	1.7	Minnesota	1.75	1.7	1.7
Mississippi	1.79	1.7	Mississippi	1.75	1.7	1.7
Missouri	1.79	1.7	Missouri	1.75	1.7	1.7
Montana	1.79	1.7	Montana	1.75	1.7	1.7
Nebraska	1.79	1.7	Nebraska	1.75	1.7	1.7
Nevada	1.79	1.7	Nevada	1.75	1.7	1.7
New Hampshire	1.79	1.7	New Hampshire	1.75	1.7	1.7
New Jersey	1.79	1.7	New Jersey	1.75	1.7	1.7
New Mexico	1.79	1.7	New Mexico	1.75	1.7	1.7
New York	1.79	1.7	New York	1.75	1.7	1.7
North Carolina	1.79	1.7	North Carolina	1.75	1.7	1.7
North Dakota	1.79	1.7	North Dakota	1.75	1.7	1.7
Ohio	1.79	1.7	Ohio	1.75	1.7	1.7
Oklahoma	1.79	1.7	Oklahoma	1.75	1.7	1.7
Oregon	1.79	1.7	Oregon	1.75	1.7	1.7
Pennsylvania	1.79	1.7	Pennsylvania	1.75	1.7	1.7
Rhode Island	1.79	1.7	Rhode Island	1.75	1.7	1.7
South Carolina	1.79	1.7	South Carolina	1.75	1.7	1.7
South Dakota	1.79	1.7	South Dakota	1.75	1.7	1.7
Tennessee	1.79	1.7	Tennessee	1.75	1.7	1.7
Texas	1.79	1.7	Texas	1.75	1.7	1.7
Vermont	1.79	1.7	Vermont	1.75	1.7	1.7
Virginia	1.79	1.7	Virginia	1.75	1.7	1.7
Washington	1.79	1.7	Washington	1.75	1.7	1.7
West Virginia	1.79	1.7	West Virginia	1.75	1.7	1.7
Wisconsin	1.79	1.7	Wisconsin	1.75	1.7	1.7
Wyoming	1.79	1.7	Wyoming	1.75	1.7	1.7

6. COMPARISON OF INDEX OF BUILD IN RECRUITS OF 1917-1918 AND IN VETERANS OF 1919 AND 1864-1865.

Table 59 gives the index of build at demobilization by States. Here, as in Table 58, Alaska and the Dakotas stand at the top. But the other States following them differ a good deal from the mobilization series. Kentucky and Tennessee no longer stand at the bottom, but Florida and Georgia do, though even these States show an increase in robustness.

Table 60 shows the percentage of increase of the index of build of demobilization over mobilization. For the United States as a whole the increase in the index of build amounted to 0.51, or about 1.6 per cent. In the table the State that stands at the top is Colorado, with an increase of 1.38, or 4.3 per cent. Since Colorado men were among the least robust of the recruits, there was the greatest room for improvement. It was suggested that their average lack of robustness on entering the Army was due to the presence of a large

number of persons of tuberculous strains. If so, Army life and exercise in the open air produced a vast improvement in robustness. The increase may have been due to a general improvement or to the selective weeding out of men who were accepted for the Army and subsequently discharged for disability on account of tuberculosis. The second State from the top is Louisiana, in which the recruits also stand relatively low in index of build, 30.55. They had, therefore, a great opportunity for improvement in this respect. Men from Louisiana show the greatest increase in weight of all of the United States proper, while the increase in stature was only medium. This high position of Louisiana in order of increase in index of build is thus due to the increase in average weight of men at demobilization, which is probably due to improved sanitary conditions, whether on the part of white or colored.

The next state in order is Alaska, which showed the greatest increase in weight and also the greatest increase in height. The number of men involved, however, is small. Next follow the States of Oklahoma, Kansas, Idaho, and Nebraska. The Southern States in which the increase in index of build is over 0.75 are Virginia, 1.05; Arkansas, 1; North Carolina, 0.94; Texas, 0.91; Kentucky, 0.87; Tennessee, 0.86; West Virginia, 0.79. In a number of Southern States, however, the increase in index of build of the troops was very slight, as in South Carolina, 0.34; Alabama, 0.24; Georgia, 0.20; Florida, -0.11.

Among the States that lie at the bottom of the list are New Hampshire, with a decrease of 0.60 in the index of build. This agrees with what we have found in respect to the marked decrease in weight and stature in men from this State, a result that probably is due to selection and to the small numbers considered. It is noteworthy that men from Florida on the average showed a decrease in the index of build. The numbers are not large, only 140 men, and these may have been in some way selected, such as being exclusively white or colored troops or from an organization drawn from some particular part of the State.

Next comes Connecticut, which shows practically no change in robustness between mobilization and demobilization, namely -0.08. In this case the numbers are fairly large and the fact suggests that men from this State who are of less than average stature and already above the average in robustness on mobilization had little opportunity to change in this respect. The same remarks may throw light on the low position of New York and the District of Columbia. The lower half of the table includes many of the manufacturing States of the East, such as New Jersey, Maryland, Massachusetts, and Pennsylvania. Rhode Island, which gave a median position in the index of robustness of recruits, retains that position at demobilization.

It will be of interest to compare the index of build by groups of States of veterans of 1865 and 1919. Tables 61 and 62 give the means for such a comparison. By either method of calculating the index of build it appears that the build of veterans is greater in the eastern sections in 1919 than it was in 1865, but less in some western sections.

TABLE 61.—*Comparison of index of build of men at demobilization in 1865 and 1919 (weight divided by first power of height).*

Demobilization, 1919.		Demobilization, 1864-1866 (Gould ? p. 405).	
States.	Index of build.	States.	Index of build.
Rhode Island.....	2.107	New England.....	2.082
Connecticut.....	2.103		
Massachusetts.....	2.093		
Vermont.....	2.123		
New Hampshire.....	2.050		
Maine.....	2.128		
New York.....	2.098	2.107
New Jersey.....	2.096		
Pennsylvania.....	2.126		
Ohio.....	2.111	2.153
Indiana.....	2.138		
Michigan.....	2.155	2.106
Wisconsin.....	2.181		
Illinois.....	2.150		
Kentucky.....	2.121	2.190
Tennessee.....	2.121		

TABLE 62.—*Comparison of index of build of men at demobilization in 1865 and 1919 (weight multiplied by 1,000, divided by square of height).*

Demobilization, 1919.		Demobilization, 1864-65.		
		Gould's later series (pp. 284, 403).	Gould's earlier series (pp. 284, 402).	
States.	Index of build.	Index of build.	States.	Index of build.
Rhode Island.....	31.66	30.87	New England.....	31.54
Connecticut.....	31.34			
Massachusetts.....	31.35			
Vermont.....	31.61			
New Hampshire.....	30.69			
Maine.....	31.69			
New York.....	31.36	31.24	New York.....	32.12
New Jersey.....	31.31		New Jersey.....	31.77
Pennsylvania.....	31.73		Pennsylvania.....	
Ohio.....	31.72	31.68	Ohio and other western States.....	32.12
Indiana.....	31.56			
Michigan.....	32.01	31.34
Wisconsin.....	32.18			
Illinois.....	31.77			
Kentucky.....	31.13	31.91
Tennessee.....	30.92			

7. INDEX OF BUILD BY SECTIONS.

Table 63 gives the index of build of the 156 sections into which the country has been divided, arranged in order of size, the highest index being at the top of the list. This index is obtained by dividing the mean weight $\times 1,000$ of the men in each section by the square of their mean stature. The range is from 32.41 for men from Alaska to 29.88 for men from the hill country of Arkansas, inhabited chiefly by native whites. Considering the table in more detail, we find that of the United States proper, Michigan 1, with a large Scandinavian and Finnish population (only 12 per cent native whites) stands at

the top with an index of 32.15. The position, at the head, of men of Alaska and of a Finnish and Scandinavian section, indicates that people living in the north or derived from northern countries tend to have excessive weight in relation to their height. Thus among the European peoples the Scandinavians are characterized in Table 28, by a weight of 66 kilos, the greatest weight given in the table.

Returning to Table 63, we find next in order California 1. This comprises the agricultural area of central California, whose population is about half whites of native parentage and about 5 per cent Indian, Chinese, and Japanese. The well-known robustness of form of the Orientals may have influenced the result.

Next in order come North Dakota 3, including a large proportion of agricultural Russians and Scandinavians, and Minnesota 3, chiefly Scandinavians and Finns. These are followed by Arizona 1, in which Indians are the prevailing element of the population. It is well known that Indians have an exceptionally robust form; their average body weight being greater than that of any other peoples, according to Martin's table⁵ (p. 238), which gives (from Gould²) the average weight of the Iroquois Indian as 73.8 kilos.

Next in order come South Dakota 3 (Indian) and 2 (characterized again by agricultural Russians, Scandinavians, and Germans). Next are North Dakota 1 with 24 per cent Scandinavians in its population and California 2, a mining area of the middle Sierras, with a population consisting of men selected for their robustness and their ability to withstand rigors of life among the gold diggers. The following sections comprise parts of the States of Minnesota, Wisconsin, North Dakota, South Dakota, Montana, and Oregon, all sections characterized by a high proportion of Scandinavians. This part of the table includes also San Francisco with its 5 per cent of orientals, and Buffalo, N. Y., and vicinity, where have settled many of the lumber and lake men and their descendants. This table brings out vividly the striking robustness of the population of our Northwest.

The sections at the bottom of the table present a great contrast not only in index of build but in geographical and racial elements. At the bottom lies Arkansas 2, a rural hill country with 97 per cent native whites of native parentage. Next comes the mountain region of Tennessee; then, following closely, is the agricultural region of the same State. Next comes a mountainous area of North Carolina. Next comes Illinois 6, including the Negro colony that occupies the territory at the junction of the Ohio and Mississippi Rivers. This population is very tall but decidedly underweight, possibly due to the malaria of the river bottoms. Next come the mountain whites of Kentucky, then the Key West section of Florida, with its mixture of Spanish and West Indian blood, next the mountain whites of Virginia, and next New Mexico 3, a desert region containing many tuberculous whites of native stock and about 14 per cent Mexicans. Next in order come the mountain region of South Carolina, the mountain region of Alabama, and the hill country of Arkansas with 94 per cent native white population. The other sections lying in the lower part of the table are of Missouri, Mississippi, North Carolina, Florida, Georgia, Texas, Kentucky, and Virginia, all of which occupy a low position in the table of the States. Of interest is the low index of robustness of Colorado 6 (30.46). This is the region south of Denver and no doubt contains a considerable tuberculous population.

Other points of interest will be revealed by a comparison of sections from different parts of the table. For example, New York 2, including the most densely populated part of the Western Hemisphere, falls in the upper half of the table with an index of build of 31.36. This high position is in part determined by the small height and stockiness of the population, which comprises a large proportion of south Italians, Greeks, and Polish Jews. Illinois 5, Chicago, with an index of 31.30, lies somewhat below New York, because of the high proportion of men of tall stature, descendants of the pioneers of the West. Pennsylvania 1 (Philadelphia) lies at about the middle of the table, with an index of 31.01. This is due to the lower mean weight of the population of Philadelphia as compared with New York, though the average stature is slightly greater. Again, Massachusetts 4, including Boston, is intermediate between New York and Philadelphia, with a rate of 31.15. Colorado 5, comprising Denver, the section with perhaps the largest number of rejects for tuberculosis, lies near the middle of the list, with an index of 31.01. The cities of Minneapolis and St. Paul (Minnesota 4) have an index of robustness of 31.34, almost exactly equal that of New York City. The average stature is much greater, but the average weight has increased in proportion.

TABLE 63.—*Index of build of recruits, by sections, 1917-1918* $\left(\frac{\text{Weight} \times 1,000}{\text{Stature}^2 (\text{sq.})} \right)$.

State.	Section.	Index.	State.	Section.	Index.	State.	Section.	Index.
Alaska.....		32.41	New York.....	3	31.29	Ohio.....	3	30.78
Michigan.....	1	32.15	Do.....	4	31.38	West Virginia.....	2	30.76
North Dakota.....	3	32.01	Wisconsin.....	1	31.26	Arizona.....	2	30.76
Minnesota.....	3	31.99	Ohio.....	4	31.23	New Mexico.....	1	30.75
South Dakota.....	3	31.94	Nebraska.....	1	31.20	Indiana.....	3	30.73
Do.....	2	31.91	New York.....	8	31.18	Louisiana.....	2	30.72
North Dakota.....	1	31.91	Pennsylvania.....	6	31.15	Florida.....	4	30.69
California.....	2	31.83	New Jersey.....	3	31.15	Texas.....	4	30.68
Minnesota.....	2	31.79	Massachusetts.....	4	31.15	Colorado.....	4	30.67
Wisconsin.....	4	31.78	Vermont.....	All.	31.15	South Carolina.....	6	30.66
California.....	5	31.77	New Jersey.....	1	31.15	South Carolina.....	3	30.64
New York.....	6	31.76	Ohio.....	2	31.14	Tennessee.....	1	30.64
North Dakota.....	2	31.75	Illinois.....	8	31.14	Missouri.....	2	30.64
Pennsylvania.....	5	31.74	New Hampshire.....	2	31.13	Colorado.....	3	30.63
South Dakota.....	1	31.68	Indiana.....	2	31.13	Arkansas.....	1	30.62
Montana.....	1	31.67	Maryland.....	3	31.11	Louisiana.....	1	30.62
Minnesota.....	1	31.65	New Jersey.....	2	31.11	Georgia.....	2	30.59
Oregon.....	1	31.63	New York.....	7	31.11	Florida.....	2	30.58
Michigan.....	4	31.63	Kansas.....	1	31.07	Colorado.....	1	30.58
Pennsylvania.....	3	31.63	Utah.....	2	31.06	Oklahoma.....	2	30.58
Do.....	4	31.59	Maine.....	3	31.06	Missouri.....	1	30.56
Montana.....	2	31.59	Michigan.....	2	31.05	Texas.....	1	30.51
Nevada.....	All.	31.59	Virginia.....	1	31.05	Alabama.....	1	30.48
California.....	1	31.55	Mississippi.....	1	31.04	Oklahoma.....	1	30.48
Washington.....	2	31.53	Alabama.....	4	31.04	Alabama.....	5	30.48
Wisconsin.....	2	31.51	California.....	3	31.03	Colorado.....	6	30.46
Ohio.....	1	31.49	Pennsylvania.....	1	31.01	West Virginia.....	1	30.45
Illinois.....	1	31.49	Utah.....	1	31.01	Louisiana.....	3	30.40
Wyoming.....	All.	31.47	California.....	4	31.01	North Carolina.....	3	30.40
Washington.....	1	31.46	Colorado.....	5	31.01	New Mexico.....	2	30.39
Indiana.....	1	31.44	Massachusetts.....	2	31.01	Virginia.....	3	30.34
Utah.....	3	31.43	Do.....	1	31.00	Georgia.....	1	30.33
Connecticut.....	1	31.42	Michigan.....	5	30.99	Kentucky.....	2	30.33
Iowa.....	1	31.42	Massachusetts.....	3	30.99	Texas.....	2	30.33
Connecticut.....	2	31.42	Iowa.....	2	30.99	Florida.....	1	30.32
Oregon.....	2	31.41	Arizona.....	1	30.97	Georgia.....	1	30.30
Washington.....	3	31.41	South Carolina.....	2	30.97	North Carolina.....	2	30.28
Pennsylvania.....	2	31.40	Rhode Island.....	All.	30.95	Mississippi.....	2	30.28
Do.....	7	31.39	North Carolina.....	4	30.94	Missouri.....	3	30.25
New York.....	1	31.34	Maine.....	2	30.94	Arkansas.....	3	30.25
Maine.....	1	31.37	Delaware.....	All.	30.89	Alabama.....	3	30.24
New York.....	2	31.36	New York.....	5	30.89	South Carolina.....	1	30.20
Minnesota.....	4	31.34	Kansas.....	2	30.88	New Mexico.....	3	30.16
Wisconsin.....	3	31.33	Alabama.....	2	30.88	Virginia.....	4	30.16
Idaho.....	All.	31.33	Colorado.....	2	30.87	Florida.....	3	30.16
New Hampshire.....	1	31.32	Texas.....	5	30.87	Kentucky.....	1	30.07
Maryland.....	1	31.31	Illinois.....	3	30.86	Illinois.....	6	29.98
Nebraska.....	2	31.31	Maryland.....	2	30.85	North Carolina.....	1	29.94
Illinois.....	4	31.31	Illinois.....	7	30.84	Tennessee.....	2	29.91
Michigan.....	3	31.31	North Carolina.....	5	30.83	Do.....	3	29.90
Illinois.....	5	31.30	Missouri.....	4	30.83	Arkansas.....	2	29.88
Do.....	2	31.29	Texas.....	3	30.81			

8. INDEX OF BUILD BY GROUPS OF SECTIONS.

Table 64 gives the index of build for the groups of sections arranged in diminishing order of the index, the largest at the top of the table. From this table it appears that the sections containing 10 per cent or more Finns include the most robust population of the United States. It must be remembered, however, that these sections contain a large proportion of Scandinavians and that they are among the northernmost sections of the United States. The index for this group is only slightly less than that of the Alaskan section. Next come two groups of sections containing a large proportion of Germans and Scandinavians. This is followed by the group containing sections with 10 per cent or more of agricultural Russians (31.59). Then follow two groups characterized by 20 per cent and 15 per cent, respectively, Germans and Austrians. Next comes the sparsely settled group containing a considerable sprinkling of Orientals, who are known to be robust. This is followed by the eastern manufacturing and the commuter groups containing a large proportion of short, stocky people.

At the bottom of the list stand the mountain whites, with an index of 30.07. Just above is the group of native white of Scotch origin. Then come the southern agricultural groups, including a large proportion of native white population. The sections including a large proportion of Negroes stand decidedly above this group. The sparsely settled section containing Indians, that containing Mexicans, and the desert group lie in the lower half of the list, the index of build being depressed, no doubt, by the resort to these regions of the southwest by many tuberculous persons.

TABLE 64.—*Index of build by groups of sections, 1917-18* $\left(\frac{\text{Weight} \times 1,000}{\text{Height (sq.)}}\right)$.

Groups.	Index of build.	Groups.	Index of build.
Finns, 10 per cent.....	32.06	Mountain.....	31.17
Scandinavians, 10 per cent.....	31.64	French Canadians.....	31.02
Germans and Scandinavians, 10 per cent plus.....	31.61	Maritime.....	30.96
Russians, 10 per cent plus.....	31.59	Agricultural, native white, North.....	30.92
Germans and Austrians, 20 per cent plus.....	31.53	Desert.....	30.85
Germans and Austrians, 15 per cent plus.....	31.45	Agricultural, Negro.....	30.78
Sparsely settled.....	31.32	Mexican, sparsely settled.....	30.59
Eastern manufacturing area.....	31.29	Indian, sparsely settled.....	30.58
Commuters.....	31.27	Southern agricultural, native white.....	30.42
Agricultural, mixed foreign and white.....	31.23	Native white, Scotch origin.....	30.33
Mining.....	31.23	Mountain whites.....	30.07

9. THE MEAN INDEX OF BUILD OF EIGHT EUROPEAN RACES OF MEN AT DEMOBILIZATION.

Table 65 gives the index of build of representatives of eight European races as recorded at demobilization. According to this table, the Poles were the most robust people, 32.73. Following them in turn are the Italians, Germans, French, Hebrews, English, Scotch, and Irish. This series indicates that the Mediterranean peoples are more robust than the Nordics. In fact, this difference of build constitutes a striking racial feature.

TABLE 65.—*Index of build of eight European races* $\left(\frac{\text{Weight} \times 1,000}{\text{Height (sq.)}}\right)$.

Race.	Index of build.	Race.	Index of build.
English.....	31.59	French.....	32.28
Scotch.....	31.41	Italian.....	32.63
Irish.....	31.41	Polish.....	32.73
German.....	32.31	Hebrew.....	31.93

10. THE MEAN INDEX OF BUILD OF COLOR RACES.

Finally the index of build has been calculated for white, Negro-mulatto, Chinese, Japanese, and Indian. In order we have:

TABLE 66.—*Index of build of color races* $\left(\frac{\text{Weight} \times 1,000}{\text{Height}^2 (\text{sq.})} \right)$.

Race.	Index of build.	Race.	Index of build.
Indian.....	32.93	Japanese.....	32.00
Chinese.....	32.82	White.....	31.56
Negro-mulatto.....	32.63		

Here, again, a striking likeness appears between the Indian and Chinese. The Japanese resemble, in build, more the whites than the Chinese.

11. EXPLANATION OF PLATES XIV-XIX.

An attempt is made in Plate XIV to show the interrelation of stature, weight, and chest circumference (expiration) in the general population of the first million draft recruits. In the left figure the stature is taken as the controlling factor, the range being from 79 down to 59 inches. The mean stature, 67.49 inches, for the first million draft recruits is shown by the upper heavy horizontal line. Passing downward, the second horizontal line shows the quotient of the average weight in pounds divided by the average stature in inches, which is 2.097 pounds. The corresponding quotient for each class of statures is shown by the vertical divided bars. It is apparent that for the statures from 75 down to 62 inches, the corresponding average weights diminish with the statures closely. However, for the statures 79 to 76 inches, there is a very marked diminution in the proportional weight, for the men with such tall statures are unduly slender. On the other hand, for statures of 61 to 59 inches there is a marked increase in the proportional weight, which is more marked as the stature diminishes. This increase is probably due at least in part to the fact that the local boards sent to the camps only men of such short stature as were unusually robust.

In the third of the horizontal lines, there is shown the quotient of the average chest circumference (expiration) by the average stature in inches. For each stature the corresponding proportional average chest circumference (expiration) is shown by the vertical heavy bars. It is apparent at once that the proportional average chest circumference (expiration) increases as the stature decreases. This increase is due at least in part to the fact that the range of the stature measurements is from 79 to 59 inches, or a total of 21 inches, whereas that of the chest circumference (expiration) is from only 39 to 29 inches, or a total of 11 inches; thus the range of the chest measurements is about 50 per cent of that of the stature measurements, and consequently the quotient of the chest circumference (expiration) divided by the stature increases as the stature decreases. The small chested short men were rejected. The proportional increase of the chest circumference (expiration) to the height is also due in part to the racial increase of robustness of the men of short stature.

In figure 2 which is drawn up in similar manner as figure 1, the weight is taken as the controlling factor with the quotient of the weight divided by the height, and the weight divided by the chest circumference (expiration) shown in the second and third sections below. One sees in a general way the decrease in both parallels the decrease in the weight, but that the quotient of the weight divided by the chest circumference (expiration) follows the downward trend of the weight more closely than does the quotient of the weight divided by the height.

In figure 3 the chest circumference (expiration) is taken as the controlling factor with the quotient of the chest circumference (expiration) divided by the stature, and the weight divided by the chest circumference (expiration) shown in the second and third sections below. It is seen here again that the decrease in both sets of proportional figures parallels fairly closely the downward trend of the chest circumference (expiration), but that the quotient of the weight divided by the chest circumference (expiration) more closely approximates it than that of the chest circumference divided by the stature. In other words, as shown elsewhere, the weight and chest circumference (expiration) are more closely correlated measurements than are the stature and chest circumference (expiration) or stature and weight.

Plate XV is drawn up in a similar manner to Plate XIV, figure 1. There is shown here the interrelation of stature, weight, and chest circumference (expiration) for the men included in the first million draft recruits, distributed by the various States from which they were drafted. It is seen at once that from a number of the States the stature is above the average, but that for many of them the proportional weight and chest circumference (expiration) are below the average. Thus the men from Texas have the greatest average stature, but their proportional weight and chest circumference (expiration) is considerably below the average of the recruits in general. On the other hand, the men from Idaho, South Dakota, Minnesota, and North Dakota not only have great stature, but have also high proportional weight and chest circumference (expiration). The highest proportional weight is found in the men from North Dakota, the lowest proportional chest circumference (expiration) in the men from the District of Columbia, and the highest proportional chest circumference (expiration) in men from Connecticut. The high proportional chest circumference (expiration) in the men from Connecticut, who were much below the average in stature, is due to the fact, as shown in connection with Plate XIV, figure 1, page 177, that the proportion of chest circumference (expiration) to stature increases, as the stature decreases. The lowest average stature is found in men from Rhode Island, next in the men from Connecticut, and then in those from Pennsylvania and New York.

In Plate XVI, as in Plate XIV, figure 2, the weight is taken as the controlling factor. One sees at once that the highest average weights are found in some of the States of the Northwest—South Dakota, North Dakota, Minnesota, Oregon, Montana, and Washington. These States have also high proportional weights to the stature and proportional weights to the chest circumference (expiration). At the extreme left stand Rhode Island and Massachusetts with their large percentage of southern European immigrants. Not

only is the average for these two States below the average, but the proportional weight to the height, and the weight to the chest circumference (expiration) are also below the average.

In Plate XVII, as in Plate XIV, figure 3, the chest circumference (expiration) is taken as the controlling factor. Here, as in Plate XIII, it is some of the States of the Northwest that stand at the extreme right—namely, North Dakota, Nevada, Idaho, Minnesota, Wisconsin, and South Dakota. These States also have higher proportional chest circumference (expiration) to stature, and weight to chest circumference (expiration). The high average of stature, weight, and chest circumference (expiration) of the men from the States of the north central and northwest sections as well as the variations in these measurements found in the men from the other States is, as has been shown elsewhere, the result of racial factors more than of environmental ones.

In Plate XVIII there is shown the interrelation of stature, weight, and chest circumference (expiration) associated with the occupational, physiographic, and population groups of sections. This plate is drawn up in a similar manner to Plate XIV. For figure 1, where the stature is taken as the controlling factor, certain interesting facts are apparent. It is seen at a glance that certain of the "groups" have a stature above the average for the first million draft recruits. However, the proportional weight and chest circumference (expiration) for these "groups" with great statures varies above and below the average. Thus it is seen that the "group" of the mountain whites of the Appalachian Mountains has the greatest stature of all, but that it has a low proportional weight and chest circumference (expiration). The same is also true, though not so markedly so, for the "group" of agricultural native whites of the South. On the contrary, it is apparent that for the German and Scandinavian "groups," while the stature is above the average, their proportional weight and chest circumference (expiration) are likewise so. The "group" composed of the native whites of Scotch origin has a stature greater than the average, with a low proportional weight and a very low proportional chest circumference (expiration). The "group" of Finns, for which people the stature is below the average, has the greatest proportional chest and weight. The lowest average stature is found among the commuters, eastern manufacturing, and French-Canadian "groups." The first two named have average proportional weights, with proportional chest above the average. For the French-Canadians the proportional chest circumference (expiration) is also above the average, but the proportional weight is below it. This high proportional chest circumference (expiration) for these latter three "groups" is due at least in part to the fact that the proportion of the chest circumference (expiration) to the stature increases as the stature decreases (see Plate XIV, fig. 1, p. 177).

In figure 2 the weight is taken as the controlling factor, with the quotient of the weight divided by the stature and the weight divided by the chest circumference (expiration) shown in the second and third sections below. The points that were apparent in figure 1 are further strengthened by the evidence here. Thus the German-Scandinavian, Scandinavian, and Finn "groups" have the greatest mean weight and have also the highest proportional weight

HEIGHT, WEIGHT AND CHEST (EXP.) MEASUREMENTS SHOWING PROPORTIONATE MEASUREMENTS OF TWO OF THEM TO THE TOTAL OF THE THIRD (P.)

HEIGHT WITH RELATIVE WEIGHT AND CHEST (EXP.) WEIGHT WITH RELATIVE HEIGHT AND CHEST (EXP.) CHEST (EXP.) WITH RELATIVE HEIGHT AND WEIGHT

64. (EXP.) - 17. - 12.	57. - 12. - 12.	57. - 12. - 12.
57. - 12. - 12.	57. - 12. - 12.	57. - 12. - 12.
57. - 12. - 12.	57. - 12. - 12.	57. - 12. - 12.

BUILD.

177

PLATE XIV.

TOTAL AND PROPORTIONATE MEASUREMENTS BY STATES (P)

HEIGHT WITH RELATIVE WEIGHT AND CHEST (EXP.)

MEAN HEIGHT

WT. - HT. - LOG.

CA. (112) - WT. - HT.

TOTAL AND PROPORTIONATE MEASUREMENTS BY STATES (P)

CHEST (EXP.) WITH RELATIVE HEIGHT AND WEIGHT

1840 1850 1860 1870 1880 1890 1900 1910 1920 1930 1940 1950 1960 1970 1980 1990 2000

1840 1850 1860 1870 1880 1890 1900 1910 1920 1930 1940 1950 1960 1970 1980 1990 2000

1840 1850 1860 1870 1880 1890 1900 1910 1920 1930 1940 1950 1960 1970 1980 1990 2000

1840 1850 1860 1870 1880 1890 1900 1910 1920 1930 1940 1950 1960 1970 1980 1990 2000

PLATE XVII.

TOTAL AND PROPORTIONATE MEASUREMENTS. GROUPS OF SECTIONS (P)

WEIGHT WITH RELATIVE WEIGHT AND CHEST (XIX) WEIGHT WITH RELATIVE WEIGHT AND CHEST (XIX) CHEST (XIX) WITH RELATIVE HEIGHT AND WEIGHT

WEIGHT WITH RELATIVE WEIGHT AND CHEST (XIX)

WEIGHT WITH RELATIVE WEIGHT AND CHEST (XIX)

WEIGHT WITH RELATIVE WEIGHT AND CHEST (XIX)

BUILD.

181

WEIGHT WITH RELATIVE WEIGHT AND CHEST (XIX)

WEIGHT WITH RELATIVE WEIGHT AND CHEST (XIX)

WEIGHT WITH RELATIVE WEIGHT AND CHEST (XIX)

WEIGHT WITH RELATIVE WEIGHT AND CHEST (XIX)

WEIGHT WITH RELATIVE WEIGHT AND CHEST (XIX)

WEIGHT WITH RELATIVE WEIGHT AND CHEST (XIX)

PLATE XVIII.

TOTAL AND PROPORTIONATE MEASUREMENTS BY EACH SECTION (N)
 (WEIGHT WITH RESPECTIVE WEIGHT AND CREST (N))

PLATE XIX.

TOTAL AND PROPORTIONATE MEASUREMENTS BY EACH SECTION (R). (CONT'D.)
HEIGHT WITH RELATIVE WEIGHT AND CHEST (EXP.)



FIGURE 2

BUILD.

183

PLATE XIX.

DISTRIBUTION, HEIGHT, WEIGHT & CHEST MEAS. STATES OF NATIVITY

¹
MEAN HEIGHT

²
MEAN WT. + MEAN HT. = LBS.

³
MEAN WEIGHT

⁴
MEAN WT. + MEAN CH. (EXP.) = LBS.

⁵
MEAN CHEST (EXP.)

⁶
MEAN CH. (EXP.) + MEAN HT. = IN.





PLATE XX.

divided by the height and the weight divided by the chest circumference (expiration). French-Canadian groups stand at the extreme left of the figure with low absolute and proportional measurements.

In figure 3 the chest circumference (expiration) is taken as the controlling factor. Here again the three "groups" that stood first for mean weight again stand first, but with the order somewhat reversed, it being here Finns, German-Scandinavians, and Scandinavians. In the second and third sections below, which show the quotient of the chest circumference (expiration) divided by the height and the weight divided by the chest circumference (expiration), the superiority of the physique of the Finns is again apparent. At the extreme left stands the "groups" composed of native whites of Scotch origin. They not only have the lowest mean chest circumference (expiration), but also the lowest proportional chest circumference (expiration) to height. The fact that the proportional chest circumference (expiration) to weight reaches the average line is to be accounted for by the exceptionally small divisor, the mean chest circumference (expiration). A further study of Plate XVIII will reveal many interesting facts showing the interrelation of stature, weight, and chest circumference (expiration) associated with the 22 groups of recruits.

Plate XIX, figures 1 and 2, is drawn up in a similar manner to Plate XIV, figure 1. There is shown here the interrelation of stature, weight, and chest circumference (expiration) for the 156 sections into which the United States has been divided for this study, and for that of "Defects Found in Drafted Men."⁹ It is seen at once that the statures for recruits from many of the sections are above the average of the statures obtained for the first million draft recruits. At the extreme left of figure 1, with the highest average stature, there are found certain sections of the South where there is a very high percentage of native-born whites of native origin, many of whom are of Scotch descent. The highest average is found in North Carolina, section 1, and this is followed quite closely by Arkansas 2, Missouri 3, Texas 2, 5, and 3. It is seen from this plate that the relative weight and chest circumference (expiration) varies above and below the average. Thus for the first four of the sections named the relative weight and chest circumference (expiration) are markedly below the average, and the men are tall, slender, and small-chested. The greatest proportional weight is found in Minnesota 1, and North Dakota 3. North Dakota 3, moreover, has the greatest proportional chest circumference (expiration). At the extreme right of the list are found the States whose average stature has been materially reduced by immigration from southern Europe. Reading from the right toward the left, we find Rhode Island (all), New York 2, Pennsylvania 3, Pennsylvania 1, New York 1, Pennsylvania 5, Pennsylvania 7, and Massachusetts 2. The majority of these sections show a proportional weight, either average or slightly above the average, but all of them have a proportional chest circumference (expiration) above the average. Thus again it is made clear, as in Plate XIV, figure 1, that the proportional chest circumference (expiration) to the stature increases as the stature decreases.

VI. PIGNET'S INDEX OF ROBUSTNESS.

This index of the constitution or robustness of individuals depends upon certain relation of stature, weight, and chest circumference (Pignet,²⁰). The index is calculated according to the following formula: Stature in centimeters — (chest circumference in centimeters + weight in kilograms). Pignet offers the following table of standards, by which one can interpret the results obtained by this formula:

Class.

- A.—Under 10: A very powerful constitution.
- B.—11-20: Good constitution.
- C.—21-25: Mediocre constitution.
- D.—26-30: Weak constitution.
- E.—31-35: Very weak constitution.
- F.—Over 36: Bad constitution.

It will be of interest to see how the selection of medical examiners at demobilization boards was influenced by the constitution or index of robustness as determined by the Pignet formula.^a

In an appreciative account of Pignet's formula, Butza²¹ calls it "the criterion of constitution."

It will be observed that Pignet employs the chest "perimeter." It is clear that the chest girth at rest is used: consequently our chest girths of recruits taken at expiration are too small. To use them in Pignet's formula, it is necessary to add certain constants, and those adopted are as follows:

- Chest girth under 32 inches, add 0.50 inch.
- Chest girth 32-34.9 inches (inclusive), add 0.75 inch.
- Chest girth 35-37.9 inches (inclusive), add 1 inch.
- Chest girth 38 and over, add 1.5 inches.

In Table 67 there is considered in classes of stature separated by 2 or 3 inches, the weight in pounds with the number of men measured, circumference of the chest with the number of men measured. In the following columns the stature, chest circumference, and weight are transformed into the metric equivalent. In the last column is given the index of robustness. Under each unit of stature the population is divided into classes containing, respectively, the 5 per cent lightest, the following 10 per cent of greater weight, then the 20 per cent of still greater weight, the 30 per cent of mediocre weight, followed by the 20 per cent of still higher weight, followed by the upper 15 per cent divided into the two classes that include 10 per cent and 5 per cent of the very heaviest men.

Taking first the class of men 59 inches tall, we find that the classes established vary in average weight from 47.4 to 85.6 kilograms, and the corresponding chest circumference increases from 74.9 to 101.3 centimeters. For the 5 per cent shortest men of the smallest weight and chest circumference the index of

^a Pignet's reasoning which led him to suggest the formula given above is as follows:

Chez l'individu normal, le périmètre égale au moins la moitié de la taille, il augmente dans avec elle; de même de poids, dans les organismes normaux, doit s'accroître en même temps que la taille. Ces trois quantités, ayant une marche parallèle, devaient, nous semblait-il, conserver entre elles une différence constante chez les individus normaux, quelle que fut leur taille. Nous eûmes alors l'idée d'additionner le périmètre et les poids et de soustraire de la taille, la somme ainsi obtenue. Soit un homme normal de 1m 54 (154 centimeters) dont le périmètre thoracique est 78 centimeters et le poids 54 kilos. Nous faisons la somme de ce périmètre et de ce poids: 78+54=132. Cette somme est ensuite soustraite de la taille; 154-132=22.

robustness is 27.5, which belongs to the category of weak men of Pignet's classification. In the next higher 10 per cent the index of robustness is 20.31, which belongs to Pignet's median group. The next higher 20 per cent give an index of 13.2, which also belongs to Pignet's good group. The middle 30 per cent, with an index of 3.1, belong to Pignet's class of very good constitution, and the heavier men with larger chest belong to extremely superior members of this category. It appears, then, that camp examiners accepted very few men of the stature of 59 inches who fell into a category below the medium, and indeed all but about 15 per cent belong to the category of good or very good men. This is, of course, to have been expected, as the Army regulations required the elimination of all men under 60 inches. Indeed, we should probably expect no men under 60 inches who did not belong to the category of the exceptionally robust.

Of the men 62 inches (157 centimeters) tall, we find that nearly 5 per cent fall into the category of very weak constitution and an additional 30 per cent into the category of the weak or median. The middle 30 per cent fall into the category of good, whereas the remainder are of strong or very strong constitution.

In the group of men 65 inches in stature (165 centimeters), we find that the average of the lower 15 per cent belong to Pignet's bad category, the next 20 per cent to the weak, and the median 30 per cent to the category of the good. As we pass now to the taller statures, the proportion of men of bad constitution increases until the group of men with a stature of 77 inches, 35 per cent were of bad constitution and only about 35 per cent were better than of median constitution.

Naturally Pignet's index is purely an empirical one and the results have to be interpreted with caution. The formula and the standards established by Pignet do, however, point out the very practical matter, that stature should be considered with weight and chest circumference, and that a satisfactory rating of robustness can be determined only by considering the three together.


In connection with the matter of robustness and military efficiency the statement made by Gould seems important. It is generally held by line officers that men below 60 inches in height are not capable of standing the severe service required in the Army, especially in carrying weight on the back. He says concerning our experiences in the Civil War, "The testimony is overwhelming that very tall men do not bear the fatigues of a campaign so well as persons of ordinary stature; that they are less capable of performing long marches and are more frequently on the sick list at other times." On the whole, the Army ideals of selecting men of medium stature for Army service is justified. In connection with the draft of 1917, efforts were made on more than one occasion to raise the minimum stature to 63 inches. This was due to failure to recognize that there was in this country a great number of short men belonging to the Mediterranean races and to the group of Polish Jews in whom the mean stature is only slightly above 63 inches. Experience in the Italian army indicated that even short men, if they are not too far removed from the standard of their race, are capable of performing excellent military service. In case it ever again becomes necessary to institute a selective draft in this country it should not be forgotten that this country has a great popu-

lation of short men and that it includes many thousands for whom a stature of 60 inches is not a greater departure from the average than a stature of 65 inches is in men of the Nordic races.

TABLE 67.—Comparison of Pignet's index for men of various heights with average chest and weight for certain per cents of the men of each height.

Per-centage of this height.	Height, in inches.	Mean weight, in pounds.	Number of men.	Mean chest (expiration), in inches.	Correc-tion, in inches.	Mean chest (expira-tion), in inches (cor-rected to "rest").	Number of men.	Height, in centi-meters.	Weight, in kilos.	Chest measurement, in centi-meters (cor-rected to "rest").	Pignet's index.
5	59	104.50	194	29.00	0.50	29.50	128	149.86	47.40	74.93	27.53
10	59	114.82	460	30.00	.50	30.50	241	149.86	52.08	77.47	20.31
20	59	124.85	585	31.00	.50	31.50	366	149.86	56.67	80.01	13.18
30	59	137.38	931	32.48	.75	33.23	1,208	149.86	62.32	84.40	3.14
20	59	151.26	605	34.41	.75	35.16	811	149.86	68.63	89.31	- 8.08
10	59	166.30	272	36.31	1.00	37.31	248	149.86	75.44	94.77	-20.35
5	59	188.62	77	38.38	1.50	39.88	84	149.86	85.55	101.30	-36.99
5	62	105.40	1,362	29.00	.50	29.50	850	157.48	47.81	74.93	34.74
10	62	112.00	2,081	30.00	.50	30.50	1,822	157.48	50.80	77.47	29.21
20	62	117.00	2,557	31.00	.50	31.50	2,884	157.48	53.07	80.01	24.40
30	62	124.33	4,774	32.44	.75	33.19	6,313	157.48	56.39	84.30	16.79
20	62	134.01	2,805	34.73	.75	35.48	3,046	157.48	60.79	90.12	6.57
10	62	145.54	1,455	36.00	.75	36.75	541	157.48	66.00	93.35	- 1.87
5	62	166.68	610	37.61	1.00	38.61	392	157.48	75.62	98.07	-16.21
5	65	105.92	1,438	29.00	.50	29.50	2,759	165.10	48.03	74.93	42.14
10	65	115.34	11,770	30.00	.50	30.50	6,757	165.10	52.30	77.47	35.33
20	65	124.36	23,055	31.00	.50	31.50	12,514	165.10	56.43	80.01	28.66
30	65	135.20	22,710	32.97	.75	33.72	33,275	165.10	61.33	85.65	18.12
20	65	145.72	16,741	34.00	.75	34.75	12,347	165.10	66.09	88.27	10.74
10	65	161.92	5,067	35.00	.75	35.75	7,618	165.10	73.43	90.81	8.86
5	65	184.76	1,123	36.75	1.00	37.75	7,156	165.10	83.81	95.89	-14.60
5	67	110.50	2,528	29.00	.50	29.50	2,583	170.18	50.13	74.93	45.12
10	67	120.25	15,679	30.00	.50	30.50	7,589	170.18	54.55	77.47	38.16
20	67	129.76	33,194	31.21	.50	31.71	41,234	170.18	58.83	80.54	30.81
30	67	139.36	35,483	33.00	.75	33.75	26,558	170.18	63.23	85.73	21.22
20	67	149.09	22,983	34.00	.75	34.75	22,018	170.18	67.63	88.27	14.28
10	67	160.64	13,073	35.00	.75	35.75	14,015	170.18	72.86	90.81	6.51
5	67	182.87	4,924	36.66	1.00	37.66	14,294	170.18	82.96	95.66	- 8.44
5	69	115.14	2,032	29.77	.50	30.27	5,585	175.26	52.20	76.89	46.17
10	69	125.40	11,470	31.00	.50	31.50	10,779	175.26	56.88	80.01	38.37
20	69	134.77	26,043	32.00	.50	32.50	18,997	175.26	61.15	82.55	31.56
30	69	144.42	29,999	33.00	.75	33.75	23,133	175.26	65.50	85.73	24.03
20	69	154.09	21,468	34.00	.75	34.75	21,393	175.26	69.89	88.27	17.10
10	69	165.64	14,051	35.37	.75	36.12	23,622	175.26	75.12	91.74	8.40
5	69	185.68	5,445	37.68	1.00	38.68	7,614	175.26	84.24	98.25	- 7.23
5	71	124.06	2,289	29.79	.50	30.29	1,712	180.34	56.30	76.94	47.10
10	71	132.00	3,016	31.00	.50	31.50	3,896	180.34	59.87	80.01	40.46
20	71	139.87	11,368	32.00	.50	32.50	7,731	180.34	63.46	82.55	34.33
30	71	151.86	20,945	33.52	.75	34.27	22,351	180.34	68.00	87.05	24.39
20	71	164.17	8,974	35.00	.75	35.75	8,642	180.34	74.48	90.81	15.05
10	71	175.67	5,470	36.33	1.00	37.33	7,960	180.34	79.66	94.82	5.86
5	71	194.15	2,295	38.42	1.50	39.92	2,317	180.34	88.09	101.40	- 9.15
5	73	128.11	510	29.65	.50	30.15	980	185.42	58.15	78.58	50.69
10	73	140.20	1,852	32.00	.50	32.50	1,652	185.42	63.59	82.55	39.28
20	73	149.73	3,516	33.00	.75	33.75	2,798	185.42	67.87	85.73	31.82
30	73	159.43	3,755	34.00	.75	34.75	3,203	185.42	72.30	88.27	24.85
20	73	170.99	3,711	35.40	.75	36.15	4,758	185.42	77.52	91.82	16.08
10	73	185.57	1,237	37.00	1.00	38.00	995	185.42	84.20	96.52	4.70
5	73	200.30	617	38.48	1.50	39.98	898	185.42	97.79	101.55	-13.92
5	75	128.09	96	30.57	.50	31.07	124	190.50	58.21	78.92	53.37
10	75	145.13	259	32.00	.50	32.50	223	190.50	65.86	82.55	42.09
20	75	154.87	550	33.58	.75	34.33	907	190.50	70.26	87.20	33.04
30	75	166.86	911	35.00	.75	35.75	516	190.50	75.71	90.81	23.98
20	75	179.24	406	36.00	.75	36.75	409	190.50	81.33	93.35	15.82
10	75	190.76	271	37.00	1.00	38.00	271	190.50	86.56	96.52	7.43
5	75	202.00	127	38.42	1.50	39.92	127	190.50	91.63	101.40	- 2.53
5	77	119.06	17	30.60	.50	31.10	20	195.58	54.03	78.99	62.56
10	77	138.28	39	32.00	.50	32.50	29	195.58	62.74	82.55	50.29
20	77	151.80	76	33.00	.75	33.75	61	195.58	68.86	85.73	40.99
30	77	167.88	102	34.48	.75	35.23	132	195.58	76.12	89.48	28.98
20	77	181.09	75	36.00	.75	36.75	61	195.58	82.15	93.35	20.08
10	77	194.69	26	37.00	1.00	38.00	32	195.58	88.32	96.52	10.74
5	77	202.00	25	38.61	1.50	40.11	26	195.58	91.63	101.88	2.07

**PIGNET'S INDEX
MEN OF VARIOUS HEIGHTS (P)
PERCENTAGE DISTRIBUTION OF EACH HEIGHT**



D. SPECIAL ANTHROPOLOGICAL MEASUREMENTS.

1. SITTING HEIGHT.

(a) *General discussion.*—This is the vertical distance from the surface of the bench on which the subject sits to the vertex of his head. It measures the length of trunk, neck, and head, as this length might be measured on a horse. This measurement is readily taken by the same method as standing height, only the zero point is not the floor but the bench level.

This dimension is important because the trunk alone constitutes the most important part of it, so much so that it is sometimes (erroneously) spoken of as the trunk length. From a medical point of view it gives, combined with chest circumference, a better index to trunk robustness than stature and chest. For the purposes of measuring for uniforms it is next in importance to chest circumference in designing blouse pattern of different sizes.

The proportion of sitting height to total stature varies with sex. It is greater in adult females than males, due (in part) to the slightly longer trunk of the former. It diminishes greatly with the changing age from about 66 per cent of stature at birth to 51 per cent at maturity (15 years). It varies with race, being about 51 per cent to 53.1 per cent of total stature in adult Europeans, 49 in Masai of South Africa, 53 in Chinese and North American Indians, up to 55 in Aino. As for Europeans the proportion of sitting height to stature is given for male Ukrainian Jews as 51.4; French, 52; Belgians, 52.2; English, 52.4; and Scandinavians, 53.

(b) *Mean sitting height.*—The mean sitting height of 96,239 white troops is 90.39 centimeters. (See Table LXXXIII). Since the mean stature of white troops is 171.99 centimeters, the relative mean sitting height is 52.55 per cent of stature. This is about the average of adult Europeans. The distribution of frequency of mean sitting height is given in Table LXXXIII, from which it appears that the range in sitting height is between 70 and 107 centimeters, and the mode lies in class 90–91 centimeters.

Thus it appears that sitting height is roughly equal to or slightly in excess of half of the total stature on the average, but this is not true by any means for all individuals. Thus in Table LXXXIII there are five individuals with a sitting height of, say, 76 centimeters and 182 centimeters total stature. For these individuals the relative sitting height is 41.76; that is, in such individuals the sitting height was about two-fifths of the whole stature. In the same table are two individuals of, say, 98 centimeters sitting height and 148 total stature. For such individuals the relative sitting height is 66.51, or two-thirds of the total stature. Such persons have clearly very short legs and might properly be placed in the category of achondroplastic dwarfs, since their legs were only two-thirds of the normal proportional stature. Caution should be observed in making use of such extreme data, for these measurements were possibly inaccurately made or recorded.

(c) *Standard deviation.*—The standard deviation of sitting height as given in Table LXXXIII is 3.51 centimeters. This is over 5 per cent greater than half of the standard deviation of total stature, although the average of sitting height is only $2\frac{1}{2}$ per cent greater than half of the average stature. This indicates that sitting height is a more variable dimension than total stature, and this is partly because the length of the neck and height of the head are both highly variable elements of total stature and they are both included in sitting height. They constitute less important fractions of total height than they do of sitting height.

(d) *Comparison of eight European races.*—The distribution of absolute and proportional frequencies in different classes of sitting height is given for eight European races in Table 69. Table 68 summarizes their constants.

TABLE 68.—*Absolute and relative sitting heights and standard deviations with coefficient of variations in eight European races, demobilization, 1919.*

[Sitting height in centimeters.]

Race.	Number measured.	Sitting height.	Relative sitting height.	Standard deviation.	Coefficient of variation.
		Centimeters.	Per cent.	Centimeters.	Per cent.
Scotch.....	2,074	90.75	52.60	3.47	3.8
English.....	4,199	90.63	52.67	3.45	3.8
Irish.....	6,137	90.46	52.79	3.31	3.7
German.....	7,051	90.36	52.52	3.54	3.9
French.....	1,455	89.47	53.07	3.24	3.6
Polish.....	2,404	89.42	52.78	3.37	3.8
Hebrew.....	1,684	88.06	52.76	3.32	3.8
Italian.....	3,506	87.76	53.13	3.33	3.8

From these comparisons it appears that the Scotch have the tallest sitting height and the Italians the shortest absolute sitting height, but this is because the Scotch and Italians are, respectively, tall and short races. The Germans are the most variable in their sitting height and have the highest coefficient of variation. The French are the least variable although they are by no means the shortest of the races.

The Italians have the greatest relative sitting height, which means that they have the shortest legs, while the Germans have the shortest relative sitting height, which means they have relatively the longest legs. In general, the difference between the relative sitting height and 100 gives the measure of the relative length of legs.

From the foregoing tables it appears that the Nordic races have relatively shorter sitting height, which means relatively longer legs. Since they have as a whole a relatively shorter span than the other peoples, Nordics would seem to have increased length of leg and diminished length of arm; in so far they depart further than any other race from the condition of the anthropoid apes which have short legs and long arms.

(e) *Comparison of color races.*—The mean sitting height of Negro troops is 87.35 centimeters, which is 3.04 centimeters less than the mean sitting height of white troops, and this despite the fact that the mean stature of the corresponding troops, as shown in Tables LXXXIII and LXXXVII, is the same to tenths of a millimeter. This tells us that the Negro troops had shorter trunk, head and neck and longer legs than white troops of the same size. The standard deviation of sitting height is 3.48 for Negro troops, as contrasted with 3.51 for whites, indicating that, just as the average is less, so the variability is smaller. The coefficient of variability of the Negro troops is 39.8 per cent, while that of the white troops is 38.8 per cent. Thus the Negro troops show themselves in respect to sitting height to be slightly more variable than the white troops.

The table below, based on Tables 103, 104, and 107, gives the absolute and relative sitting heights for the five color races.

TABLE 70.—*Absolute and relative sitting height in five color races, demobilization, 1919.*

[Sitting height in centimeters.]			
Race.	Number measured.	Mean sitting height.	Relative sitting height.
White.....	96,239	90.39	52.56
Negro.....	6,433	87.35	50.79
Indian.....	105	90.10	52.53
Chinese.....	22	89.05	52.04
Japanese.....	32	87.88	51.41

Indians and the Oriental races are intermediate in sitting height between the white and the Negro, and the Indian approaches very close to the white in relative sitting height.

2. SPAN.

(a) *General discussion.*—It is a popular assertion that one's span is equal to one's stature. This is seen to be nearly true, *on the average*, for the Irish and Scotch. But it does not hold for the individual. Thus among the white troops (Table LXXXIV) we find a span of 152 centimeters associated with a stature of 177 centimeters, giving a relative span of 0.86. Also, there is a span of 192 centimeters associated with a stature of 168, or 1.14. The most extreme ratios in Table LXXXIV are 79 and 131, respectively; the latter ratio approaches that of the gorilla. There is the possibility that some of these remote extremes are due to errors in measurement; so too much stress must not be laid on them.

(b) *Mean span.*—The mean span for 96,596 white troops at demobilization (1919) is 175.58 centimeters. This is to be compared with the mean stature of the corresponding white troops of 171.99 centimeters (Table LXXXIV). The span is 3.59 centimeters greater than the stature and the relative span is 102.1.

(c) *The comparison of eight European races.*—The absolute mean span and relative spans for the different European races is given in Table 71, based on Table 72.

TABLE 71.—*Absolute and relative span with the standard deviation in eight European races. demobilization, 1919.*

[Span in centimeters.]

Race.	Number meas- ured.	Mean span.	Standard devia- tion.	Relative span.
German.....	7,060	176.66	7.98	102.7
English.....	4,197	175.61	7.95	102.1
Scotch.....	2,066	175.60	7.92	101.8
Polish.....	2,406	174.60	7.53	103.1
Irish.....	6,155	174.10	7.75	101.6
French.....	1,459	172.85	7.72	102.5
Hebrew.....	1,690	170.30	7.42	102.0
Italian.....	3,519	169.19	7.51	102.4

From these comparisons it appears that the Germans of our data have the greatest and the Italians the least absolute span; that the Germans are most variable in this respect and the Hebrews least; that the Polish have the greatest relative span and the Irish the least. Except for the Hebrews, the inhabitants of the British Isles have the shortest relative span. While the central Europeans have the shortest relative span of our recruits, it is in general lower than that given by Martin for the corresponding European races, possibly because the stature of each separate race is greater in the United States than in Europe, due to a selective immigration of tall persons.

(d) *Comparison of the color races.*—The mean span of the Negro troops is 180.76. This is to be compared with the mean stature of the corresponding Negro troops of 171.97 (Table LXXXIX). It exceeds that of the white troops by 5.18 centimeters, or about 2 inches. Thus the span is 8.79 centimeters greater than the stature of the Negro troops; or the relative span is 105.2, a striking increase from the white race of 102.1. This great relative span has been noticed by all observers.

Also the span is decidedly more variable in the colored (8.59) than in the white troops (7.95), and this difference is greater than would be expected, merely from the absolute difference in average span, for the coefficient of variation is 4.75 in Negroes and 4.53 in whites. Since in infants the relative span is about 92 per cent of height, it appears that in the Negro the development of the span has progressed farther beyond the infantile condition than in the whites.

TABLE 73.—Mean absolute and relative spans in the five color races, demobilization, 1919.

[Span in centimeters.]

Race.	Number meas- ured.	Mean span.	Relative span.
White.....	96,596	175.58	102.1
Negro.....	6,441	180.76	105.2
Chinese.....	23	176.41	103.1
Japanese.....	32	177.25	103.7
Indian.....	106	176.86	103.1

The two Oriental races and the Indians are intermediate between the whites and Negro.

3. HEIGHT OF STERNAL NOTCH.

(a) *General discussion.*—The sternal notch, which marks the upper end of the sternum, marks also essentially the upper limit of the trunk. It corresponds closely to the level of cross section No. 22 of trunk in Eycleshymer and Shoemaker's "Cross-section Anatomy."²² The principal viscera that rise above this level are the apices of the lungs and certain large blood vessels. Taken in connection with height of pubis it is useful in measuring the length of the trunk, a measure which is essential for estimating the volume of the trunk, which in turn is a matter of medico-military importance.

The method of measuring the height of the sternal notch is either direct with an anthropometer, or by the use of a plumb bob and cord falling from the end of a pencil or tongue depressor held horizontal at level of the notch. The anthropometer is read direct, the plumb line by reference to the vertical scale on the wall.

The sitting height of the sternal notch is a useful measure because it gives length of trunk direct and is easily made by the anthropometer, of which one end is placed on the bench on which the subject is sitting and the movable arm is brought to the level of the sternal notch. This latter measure ranges in the male and in relation to total height from 30 per cent among the Cochinese through 33 per cent in the French to 35 per cent in certain Negro tribes.

(b) *Mean height of sternal notch.*—The mean height of sternal notch for white troops as shown in Table LXXXV is 141.18 centimeters, which is 82.09 per cent of the mean height (171.99); that is to say, the height of the man from the floor to the sternal notch constitutes over four-fifths of the total stature. Neck and head constitute something less than one-fifth. The relation between sternal notch and total stature, however, is far from constant. Thus in Table LXXXV there are 16 cases of men with a height of, say, 182 centimeters and sternal notch of 138. In these cases the sternal notch height was 75.8 per cent of the total stature. In 27 cases men 166 centimeters tall had a height of sternal notch of 148 centimeters, or 89.2 per cent of the total stature.^a

(c) *Comparison of eight European races.*—The absolute and proportional distributions of sternal notch in eight European races are given in Table 75. A summary table is given in Table 74.

TABLE 74.—*Absolute and relative height of sternal notch in eight European races, demobilization, 1919.*

Race.	Number measured.	Absolute height of sternal notch.	Relative height of sternal notch.
		Centimeters.	Per cent.
Irish.....	6,173	142.28	83.03
Scotch.....	2,066	141.53	82.03
German.....	7,033	141.19	82.07
English.....	4,176	140.87	81.86
Polish.....	2,403	139.15	82.14
French.....	1,456	137.88	81.78
Hebrew.....	1,688	136.93	82.04
Italian.....	3,509	135.37	81.95

From this table it appears that the Irish have relatively the highest sternal notch, whereas, on the other hand, the French have relatively the lowest sternal notch and proportionately the longest head and neck. The English have the greatest variability in respect to the height of the sternal notch, just as they have in many other physical characters. They are greater in degree of variability than the Irish, Scotch, and German. Italians show the least variability, followed by the Polish, Hebrew, and French. Thus the distributions of the relative height of sternal notch and of variability are somewhat irregular in the races of Europe, one outstanding feature being the high sternal notch with the short head and neck among the Irish.

^a There are numbers of obvious errors in recording the height of sternal notch. These are shown by certain irregularities at the extremes of the table. The table as obtained by the tabulators is printed unchanged. It is believed that the few errors will not greatly modify the results.

(d) *Comparison of color races.*—For the Negro troops (Table LXXXIX) the mean height of sternal notch is 142.39 centimeters, which is 82.8 per cent of the total stature. The relative height of the sternal notch is therefore greater in Negroes than in whites, indicating that they have a shorter neck and head, but not as short as the Irish.

A comparison of the height of sternal notch in various color races is given in Table 107. The results of this comparison with the measurements of white and Negro troops are given in the following table:

TABLE 76.—*Absolute and relative height of sternal notch in five color races, demobilization, 1919.*

Race.	Number measured.	Mean measure.	Relative sternal notch.
		Centimeters.	Per cent.
White.....	96,439	141.18	82.09
Negro.....	6,454	142.39	82.80
Chinese.....	22	140.86	82.32
Japanese.....	32	140.44	82.16
Indian.....	107	140.97	82.19

The relative height of sternal notch is seen to be slightly greater in Negro than in white troops. In the Indian and Oriental races the relative height of sternal notch is intermediate between that of white and Negro.

4. HEIGHT OF PUBIC ARCH.

(a) *General discussion.*—This is the vertical distance from the floor to the upper margin of the pelvis at the symphysis of the pubic bones. It is measured by means of an anthropometer of which the movable arm is raised to the required level. The line is sometimes difficult to find, especially in the fat subject, but practically it is readily established, sometimes by following down the front margin of the pelvis from the more lateral position, but also through the practical point that it is the uppermost margin of resistance of the pelvic bone in the middle front line, about 25–30 millimeters above the root of the penis.

The pubic height is important because it is almost exactly (perhaps 35 millimeters below) the level of the center of the acetabulum or the axis of the hinge of the femur. Its height is therefore the length of the physiological leg or the line of rotation of the leg; a matter of prime importance in determining the length of step that requires the least effort. Practically, troops march with less fatigue if soldiers with the same physiological length of leg be grouped in one company or platoon.

Pubic height is also important because it has been nearly universally obtained in the measurement of young men, largely through the influence of Dr. Dudley A. Sargent, director of the Harvard Gymnasium. The height of the pubic arch has been found by Dr. Sargent to range in college men, 16 to 24 years of age, from about 76 centimeters to 99 centimeters and from 43.16 to 56.5 per cent of the stature. The ratio of pubic height to total stature is about 50 per cent. According to the table of Martin⁵ (1914, p. 256, made up from various sources) it is in English males about 49.9 per cent; Laplanders, 50; Poles, 50.7; Belgians, 50.7; Cossacks, 51.4; French, 52.2; of Asiatic peoples

the inhabitants of the Samoyedes Peninsula of Siberia have a relative pubic height of 48.6, the lowest of all races measured. In the Japanese this proportion is 49; Ainos, 49.9; Mongolians, 50.3. In certain African tribes the relative pubic height varies from 49.8 to 52.9, the latter relation being found in the Bushmen and being the highest proportion given. This indicates a relatively extraordinarily long-legged race.

The pubic height was determined by Gould² for 1,013 veterans of the Civil War and found to be 33.26 inches, or 84.48 centimeters, slightly less than the average pubic height found by Dr. Sargent for Harvard University students.

The medical importance of pubic height depends upon the medical significance of long legs and short trunk. As is well known, in certain bone-aplasias and defects of secretions of internal glands the legs are relatively short, as in achondroplastic dwarfs and in cretins. While in different normal families the length of leg (as indicated by pubic height) varies, still this possibly may be a measure of the differences in activities of the internally secreting glands which regulate the growth of the legs.

(b) *Mean pubic height.*—In 91,365 white troops measured at demobilization the height of pubic arch is 86.8 centimeters, which is slightly greater than for Harvard men, owing to the fact that the Harvard men averaged much younger. The relative pubic height is 50.47 per cent of stature.

(c) *Standard deviation of height of pubic arch.*—The standard deviation of pubic height for white troops is, as shown in Table LXXXVI, 5.05 centimeters. The coefficient of variation of height of pubic arch is obtained by dividing this standard deviation of 5.05 by the mean pubic height of 86.82. The result is 5.817 per cent, neither a high nor a low coefficient.

(d) *Comparison of eight European races.*—In the eight European races the mean height of the pubic arch is as indicated in the following table:

TABLE 77.—*Absolute and relative height of pubic arch in eight European races, demobilization, 1919.*

Race.	Number measured.	Absolute pubic height.	Relative pubic height.
		Centimeters.	Per cent.
Scotch.....	1,976	87.30	50.60
English.....	4,061	87.19	50.67
German.....	6,688	86.63	50.35
Irish.....	5,972	86.55	50.51
French.....	1,393	86.80	50.69
Polish.....	2,279	85.27	50.33
Hebrew.....	1,650	83.94	50.29
Italian.....	3,390	82.81	50.13

Our series confirms the results obtained by others, that the French are relatively the longest legged of the European races; the English are second in this respect, followed by the Scotch and Irish. The lowest relative pubic arch is found among the Italians, followed by the Hebrews, Poles, and Germans. We see then, again, that the Nordics and the French have the longest legs, and the peoples of southern and eastern Europe have relatively short legs. Here we have evidence of the relatively greater contrast in this respect between the primates and the Nordics on the one hand, than between primates and the southern European races on the other.

TABLE 78.—Comparative frequency distribution of height of pubic arch in each of eight European races, demobilization, 1919.

SECTION A: ABSOLUTE NUMBERS.

[illegible]

SECTION B: PROPORTIONAL RATIOS PER 1,000.

[illegible]

(e) *Comparison of color races.*—In 6,220 Negro troops, the height of pubic arch is 89.4. In view of the identical average height of white and colored, this shows that the Negro men had, on the average, 2.6 centimeters, or about 3 per cent, higher pubic arch than the white men.

The standard deviation of pubic height for Negro troops is 5.27 centimeters, which is a greater variability than that shown by the whites (5.05); a greater variability which we find in their other dimensions and which is to be explained in part by the greater mean pubic height, but not entirely; and suggests that the mulattoes have had a parentage from diverse races of whites. The coefficient of variability, which is obtained by dividing the standard deviation of the pubic height by the mean pubic height, is in the case of colored troops 5.894 and for the whites 5.817. The relative height of pubic arch is in the case of white troops 50.5 per cent; in the case of colored, 52.01 per cent of the total stature. The Negro group is a long-legged one.

TABLE 79.—Absolute and relative height of pubic arch in five color races, demobilization, 1919.

Race.	Number measured.	Absolute pubic height.	Relative pubic height.
		<i>Centimeters.</i>	<i>Per cent.</i>
White.....	91,365	86.82	50.48
Negro (and mulatto).....	6,220	89.42	52.02
Japanese.....	32	88.31	51.66
Indian.....	105	86.35	50.35
Chinese.....	21	86.12	50.33

The Chinese were found to be the shortest legged of the five races and the Indians to resemble them closely. The Japanese are intermediate between the whites and Negro.

5. NECK CIRCUMFERENCE.

(a) *General discussion.*—Instructions for taking measurements stated that the circumference of the neck was to be taken at the level of the laryngeal prominence. The importance of this measurement is partly medical, since any enlargement of the thyroid gland (as in goiter) would be made manifest by any marked deviation of the neck circumference from the normal. Its military importance is merely in relation to the wearing of the military collar. Physical examination standards repeatedly referred to the necessity of rejecting recruits with enlargement of the neck glands sufficient to interfere with the wearing of the military collar.

(b) *Mean neck circumference.*—Table LXXVIII gives the correlation of neck circumference and chest circumference. According to this table the mean neck circumference for white troops is 35.98 centimeters. Table CV gives the association between the different blouse groups based on chest circumference, sitting height, and neck circumference in the case of white troops. This table shows an extraordinary scattering of large sizes among the small men. The possibility that some of them are due to errors in recording at camps can not be overlooked.

(c) *Standard deviation of neck circumference.*—The standard deviation of neck circumference for white troops is given in Table LXXVIII as 1.8 centimeters. Dividing this by the mean neck circumference we get the coefficient of variation of 5.003 per cent—a low median one.

(d) *Comparison of eight European races.*—The data for the neck circumference of the eight European races was not tabulated.

(e) *Comparison of color races.*—The relation between the neck circumference of white and Negro races is given in the following table:

TABLE 80.—*Absolute and relative neck circumference of white and Negro troops, demobilization, 1919.*

Race.	Number measured.	Mean measurement.	Relative neck circumference.
		Centimeters.	Per cent.
White.....	95,271	35.98	20.9
Negro (and mulatto).....	6,280	36.37	21.2

The neck circumference in Negro troops exceeds that of the white troops by nearly 4 millimeters, or over 1 per cent.

6. BREADTH OF SHOULDER.

(a) *General discussion.*—This is the horizontal transverse distance between the deltoid muscles of the right and left arms at a distance of about four or five centimeters below the acromial processes, or at about the greatest thickening of the deltoid. This measurement was taken in preference to the distance between the acromial processes because of its greater significance in the fitting of uniforms and because it gives a better index of the physiological breadth of shoulder.

This dimension has a certain medical importance inasmuch as the breadth of shoulder is partly dependent upon the breadth of the chest and partly upon the muscular development of the upper part of the arms. Its military importance is probably slight.

The anthropological significance of the breadth of shoulder is considerable, though it must be admitted that anthropologists have more frequently used the distance between the acromial processes than between deltoid muscles as a measure of shoulder breadth. This is partly because this measurement can also be made upon the skeleton. The different measurements of the shoulder breadth as given by Martin,⁵ (p. 141) may be translated as follows:

35. Breadth between the acromia.—To be taken with the anthropometer or "Stangelzirkel" (rod calipers). Care must be taken that the subject stretches the shoulders; that is, does not droop forward, making the measurement too small. One feels the points with the index fingers laid at the apices of the arms of the calipers, direct measurement. Horizontal distance between the two tubercula majora of the humeri; inexact measurement, since the tubercula can rarely be felt through the deltoid muscles. Maximum shoulder breadth (Grosste Schulterbreite) (diamètre bideltoid ou bihumérale), horizontal distance between the two largest projections of the deltoid muscle. Rod caliper, the instrument is not to be firmly pressed in. A very inexact measurement.

In the measurements taken under the direction of Gould² (pp. 239, 260, and 261) on Civil War soldiers at demobilization it was originally provided merely that the breadth of shoulders should be obtained, "whereas it was especially provided in the schedule for the later series that this measure should be taken between the tips of the acromial processes." There were 2,072 measurements of the full breadth of shoulders and 8,796 which gave the distance between the tips of the acromial processes. The mean of the full breadth of shoulder is about 16.35 inches (41.53 centimeters) and ranges between 13 and 19 inches (33 to 48 centimeters). Gould finds that the mean distance between the tips of the acromial processes is 12.73 inches (32.33 centimeters), the individual cases ranging between 9½ and 16½ inches (24.13 to 41.91 centimeters). "Among natives of this country, the mean value is decidedly highest for natives of Kentucky and Tennessee, being 13.51 inches (34.3 centimeters)." Gould notes that "the identification of this apophysis is not easy, and some of our examiners seem to have succeeded here but ill."

As Martin remarks, the breadth between the acromial processes in comparison to trunk length is greater in man than in any other mammal. A great shoulder breadth is also found among the anthropoid apes, in which the shoulders are extraordinarily developed on account of their arboreal or semiarboreal life. Thus in relation to the length of the trunk the shoulder breadth in the orang outang is 59.8; chimpanzee, 54.6; hylobates, 55.5; among Parisians, 77; Germans of Bavaria, 75.3; inhabitants of the Admiral Islands, 71.1; Polish Jews, 66.7, a very low rate among humans. The breadth of shoulder (acromial interval) is sometimes expressed in relation to total stature. Thus expressed, the shoulder breadth is found to be very high among the Eskimos, 24.3; Colorado Indians, 22.5. Among Europeans the relative shoulder breadth is given as follows: Belgians, 23.4; Bavarians, 23; Polish Jews, 22.1; French, about 21; Japanese, about 24; Chinese, 22-24. The absolute breadth of shoulders is stated to increase up to 50 years of life. Thus it is clearly very responsive to activity of the arms and shoulders. The breadth of shoulders as measured between the deltoid muscles also varies much with the general condition and robustness of the body.

(b) *Mean shoulder breadth.*—The mean shoulder breadth of the white troops is, as shown in Table CI, 41.81 centimeters. The relative shoulder width is 24.31 per cent. Thus the mean shoulder width is 0.28 centimeter greater than that of Civil War veterans at demobilization. The ratio is greater than that of the European races given above because the breadth of shoulder is measured between different points.

(c) *Standard deviation of shoulder breadth.*—The standard deviation of shoulder width of white troops, as shown in Table CI, is 2.408 ± 0.0037 centimeter. The coefficient of variation is then 5.7601 per cent, a rather high coefficient of variation. The mean shoulder width of Negro troops is, as shown in Table CIX, 42.89 centimeters. The standard deviation is 2.154 centimeters. We see, therefore, that the mean shoulder width of the colored troops is over 1 centimeter more than that of the white troops and the index of variability is relatively considerably less. The coefficient of variation for the colored troops is 5.013 per cent, or much less than for white troops.

(d) *Comparison of eight European races.*—Table 82 gives the absolute and proportional frequency of occurrence of shoulder breadth in each of the eight races. In Table 81 the third column from the left gives the mean shoulder breadth of the races. It will be recalled that this is the maximum shoulder width and not the space between the acromial processes. Hence the condition of the man plays a considerable rôle in determining the shoulder width. The maximum mean shoulder width, 42.24 centimeters, was found among the Poles; next among the Germans, then follow Scotch, English, and Italians rather close together. The minimum shoulder width, 40.41 centimeters, is found among the French; somewhat greater is the shoulder width of Hebrews and Irish.

Gould found the mean of measurements of "maximum breadth of shoulders" to be about 16.35 inches (41.53 centimeters), which is within 3 millimeters of the mean shoulder width measured in the troops of 1919. In comparison with the figures of 1919, transmuting inches to centimeters, the breadth of shoulders of Civil War veterans from England was 41.17 centimeters, instead of 41.69, showing an increase in the later series. The Scotch gave 42.27 centimeters instead of 41.70, showing a marked decrease half a century later. The Irish of 1866 were 41.83 centimeters, which, contrasted with the 41.43 of 1919, shows something of a decrease in half a century. Veterans of German origin in 1866 gave 41.76 centimeters as compared with the World War data of 42.19, which shows an increase half a century later. How much of this difference is significant of slightly different racial subgroups included in the two sets of measurements, how much to conditions of life, how much to errors of random sampling, can not be stated. It is probable that no important changes in this dimension have occurred in any race during the half century.

The third column from the right of Table 81 gives the standard deviation of shoulder width for the eight races. The greatest deviation is found among the Scotch, 2.11; the lowest among the French, 1.10.

The last column at the right gives the ratio of mean shoulder width to mean stature for each of the races. This column shows that the greatest relative shoulder width occurs among the Italians, 25.21; next among the Poles, and then the Hebrews, followed by the Germans. The smallest relative width is found among the French, 23.97, followed in increasing proportion by the Scotch, Irish, and English. Thus, in general, the Nordics have a smaller shoulder width than the races of southern Europe. If we regard the Nordics as the most aberrant or extremely developed of the human races, then we may say that evolution has been in the direction of diminished shoulder width. This reduction, however, it is to be pointed out, is largely due to the circumstance that the Scotch and English are of taller build than the Italians and Poles and consequently part of their proportional inferiority of shoulder width is due to proportionately larger division. For a comparison we may take the proportions of Gould, which are for the English, 24.6; Scotch, 24.6; Irish, 24.8; French, etc., 25.5; Germans, 25.

(e) *Comparison of color races.*—The following table shows the absolute and relative shoulder breadth in the five color races, demobilization, 1919:

TABLE 83.—*Absolute and relative shoulder breadth in five color races, demobilization, 1919.*

Race.	Number measured.	Mean measurement.	Relative shoulder width.
		<i>Centimeters.</i>	<i>Per cent.</i>
White.....	95,167	41.81	24.3
Negro (and mulatto).....	6,289	42.89	24.9
Chinese.....	21	42.67	24.9
Japanese.....	31	42.00	24.6
Indian.....	104	42.58	24.8

From this table it appears that, as already stated, the Negro troops have a shoulder width that exceeds the whites on the average by about 1 centimeter. The Chinese and Japanese and Indians resemble the Negro troops more than the whites in this respect. The relative shoulder width—that is, shoulder width divided by stature—is also greater in Negro, orientals, and Indians than it is in the whites.

7. TRANSVERSE DIAMETER OF THE CHEST.

(a) *General discussion.*—This measurement was taken at the level of the nipples by means of sliding wooden calipers. The arms of the calipers were held approximately perpendicularly to the axis of the thorax at this level. The measurers were instructed to permit the movable arm of the calipers to remain in contact with the chest during expiration and inspiration and to take the middle distance between the extremes.

This dimension accords very closely with Martin's No. 6 (p. 142): "Transversaler Brustdurchmesser (Frontal-Brustweite; largeur de la poitrine) direct measurement. Horizontal distance between the two most protuberant ribs at the level of the mesosternale."

This measurement has a certain medical importance, especially when used in connection with the measurement of the antero-posterior chest diameter. The ratio of the transverse to the antero-posterior diameter gives the thoracic index (Thorakalindex) of Martin ⁵ (p. 275). This index tends to increase with age; a small one is indicative of an infantile condition of development. Extreme conditions, however, produce the chicken- or pigeon-breasted form, which is a malformation not associated with physical vigor. On the other hand, an extremely low thoracic index (flat chest) should be a warning to observing physicians to look for pulmonary tuberculosis.

The military significance of the chest diameters is largely confined to its medico-military aspect and to its relation to uniforms. In general, however, uniforms are fitted by the chest circumference rather than by the diameter of the chest. The diameters of the chest have a certain anthropological significance. Thus, the transverse diameter for Navajo Indians is given at 27.9 centimeters; for French (with a prevailing shorter stature than Indians), 26.9. The thoracic index for Hova Indians is 143.5; for Bugu Negroes, 124; for African Negroes in general, 138.

(b) *Mean transverse chest diameter.*—The mean transverse chest diameter of 96,583 white troops is, as shown in Table LXXX, 29.02 centimeters. This is a transverse diameter over 1 centimeter greater than that given for the Navajo Indians and over 2 centimeters greater than that of the French. The relative mean transverse chest diameter is 16.87 per cent of the total stature. In the case of Negro troops the mean transverse chest diameter is 29.05, or practically the same as that of white troops. The relative transverse diameter is, therefore, apparently the same in the two races.

(c) *Standard deviation of transverse chest diameter.*—The standard deviation of transverse chest diameter is for the white troops 2.40 centimeters. The coefficient of variation is 8.27 per cent. This is a very high coefficient, and indicates that the diameter of the chest is a very variable dimension. In the case of Negro troops the corresponding coefficient is 7.78 per cent, indicating a slightly smaller variability in the Negro than in the white troops.

As is shown in Tables LXXX and XCVI the standard deviation of the transverse diameter of the chest is markedly greater than that of the antero-posterior diameter. This matter will be discussed when we come to consider the latter dimension.

(d) *Comparison of eight European races.*—Table 86 (summarized in Table 85) gives for each of eight European races the absolute and relative proportional frequency of occurrence of the different transverse chest diameter classes. The third column from the left of Table 85 gives the mean transverse chest diameter for each of the races. The largest diameter, 29.22 centimeters, is found among the Poles, next larger among the Germans, next among the Scotch and English. The smallest transverse chest diameter is found among the Hebrews, followed in ascending order by the French, Italians, and Irish. It is noteworthy that the transverse chest diameter of the Irish stands fifth in the list, whereas the chest circumference of the Irish stands fourth in that list, the fifth place in chest circumference being taken by the Scotch. This indicates either that the Scotch have a relatively broad chest or that the Irish have an exceptionally narrow one. The relative variability of transverse chest diameters is given in the fourth column from the right. We see that the Scotch and French show the highest standard deviation, 2.35, followed by the Germans and Hebrews. The smallest standard deviation, 2.17, is found among the Italians, followed by the English, Polish, and Irish. The third column from the right hand in Table 85 gives the transverse chest diameter in relation to height. From this column we see that the Italians have the greatest relative chest diameter, 17.41; these are followed by the Poles, French, Hebrews, and Germans. The smallest relative transverse chest diameter, 16.78, is found among the English, followed by the Irish and Scotch. Thus, in general, in transverse chest diameter the Nordics are relatively inferior to the Mediterranean races.

8. ANTERO-POSTERIOR DIAMETER OF THE CHEST, AND THORACIC INDEX.

(a) *General discussion.*—The antero-posterior diameter of the chest was taken on the same plane as the transverse diameter, but with the calipers placed antero-posteriorly. The movable arm of the calipers lay over the chest at about the level of the nipples; the fixed arm of the calipers lay on the muscles

of the back, near the top of the scapulæ. The movable arm was kept in contact with the wall of the chest during its rise and fall in respiration, and the middle reading between the extreme was regarded as the antero-posterior diameter of the chest.

The greatest interest of the antero-posterior diameter of the chest lies in relation of the transverse diameter. The index of the thorax is obtained by dividing the transverse diameter by the antero-posterior. In the case of various races, as tabulated by Martin ⁵ (p. 277), the thoracic index is as follows:

TABLE 84.—*Thoracic index of various races.*

	Thoracic index.
Hova Indians.....	143.5
Navajo Indians.....	137.5
French.....	138.6
African Negro.....	138.0
Bugu Negro.....	124.0

The antero-posterior diameter varies in different races partly, of course, in relation to the total stature of the individual. In the case of the Navajo Indians this diameter is given as 216 millimeters on the average; in the case of the French 194 millimeters.

A small antero-posterior diameter in relation to the transverse diameter may indicate pulmonary tuberculosis. Its military significance is probably confined to its medico-military significance.

(b) *Mean antero-posterior chest diameter.*—The mean antero-posterior chest diameter of white troops is, as shown in Table LXXX, 21.58 centimeters. This is markedly less than the transverse chest diameter. The ratio of the larger to the smaller is 134.48, a ratio of the thoracic index which is less than that of the French as given above. For Negro troops the mean antero-posterior chest diameter is 21.21, or slightly less than that of the whites, and the index of 136.96, a ratio still below that of the French given above, though greater than that of the whites. It has been stated by Papillault ²³ that the Negro has a somewhat rounder type of thorax than the European, hence has a large antero-posterior diameter and a relatively small thoracic index. White troops at demobilization had strikingly broad and shallow chests.

(c) *Standard deviation of antero-posterior chest diameter.*—The standard deviation of the antero-posterior diameter of the chest is, as shown in Table LXXX, 1.87 centimeters for white troops, and, as shown in Table XCVI, 1.74 centimeters for colored troops. The corresponding coefficients of variation are 8.665 for white troops and 8.204 for colored. Hence the variability of the antero-posterior chest diameter for colored troops is markedly less than that of the whites.

(d) *Comparison of eight European races.*—Table 87 gives the absolute and proportional frequencies of occurrence in the different classes of antero-posterior chest diameter of the eight European races, summarized in Table 85.

The last column on the right in Table 85 gives the mean antero-posterior chest diameter for these races. From this column it appears that the largest antero-posterior chest diameter, 21.90, is found among the Polish, as was also the case with the transverse chest diameter. The next largest is found among the

Germans, as was also the case with the transverse chest diameter. Third come the Irish, who were fifth in transverse chest diameter, and fourth the Scotch, who were third in transverse chest diameter. Fifth in order are the Italians, who were sixth in transverse chest diameter. Sixth in order of antero-posterior diameter come the English, who were fourth in transverse chest diameter. Seventh are the Hebrews and eighth the French. The French and Hebrews have exchanged places in antero-posterior chest diameter as compared with transverse. It will be interesting to compare the thoracic index to be obtained by dividing the transverse diameter $\times 100$ by the antero-posterior. The results are expressed in per cents in next to the right-hand column of Table 85. The column of transverse diameters divided by antero-posterior shows that the Irish have the smallest thoracic index (index 133.22) and the English the greatest (134.59). The order from the smallest to the greatest is as follows: Irish, Hebrew, Polish, French, German, Italian, Scotch, and English. Thus in general, excluding the Irish (who are only in part Nordic), and including the Polish (who are Nordic to a considerable degree), it appears that the Nordic races are characterized by greatest chest index. Comparing the variability of the antero-posterior diameters, it appears that the standard deviation is greatest, 1.76, among the English, followed by the Scotch, German, and Hebrew. The standard deviation is least, 1.66, among the Italians and Polish, followed by the French and Irish.

TABLE 85.—*Absolute and relative transverse diameter of chest with the standard deviation; also antero-posterior diameter of the chest with the thoracic index multiplied by 1,000 (transverse diameter divided by the antero-posterior diameter), in eight races, demobilization, 1919.*

Race.	Number measured.	Transverse diameter of chest.	Standard deviation.	Relative transverse diameter.	Thoracic index $\times 100$.	Antero posterior diameter of chest.
		<i>Centimeters.</i>	<i>Centimeters.</i>	<i>Per cent.</i>		<i>Centimeters.</i>
English.....	4,192	28.87	2.22	16.78	134.59	21.45
Scotch.....	2,057	29.01	2.35	16.81	134.43	21.58
Irish.....	6,135	28.77	2.30	16.79	133.19	21.60
German.....	7,074	29.12	2.32	16.93	133.64	21.79
French.....	1,433	28.58	2.35	16.95	133.61	21.39
Italian.....	3,514	28.76	2.17	17.41	133.89	21.48
Polish.....	2,406	29.22	2.26	17.25	133.49	21.60
Hebrew.....	1,690	28.25	2.31	16.92	133.22	21.42

9. WAIST CIRCUMFERENCE.

(a) *General discussion.*—The waist circumference was taken at the level of the umbilicus. The waist circumference in relation to stature is somewhat variable in different races. As given in Martin's table (p. 288), in different races of Africa it varies in the males from 43 to 49 per cent. In young men of the French race it is about 42 per cent.

(b) *Mean waist circumference.*—The mean waist circumference of 96,157 white troops, as shown in Table 103, is 77.87 centimeters. The relative waist circumference is 45.28 per cent. This is slightly larger than the relative waist circumference of young French males. The mean waist circumference of 6,445 colored troops is, as shown in Table 104, 77.83 centimeters, or practically the same as for whites. The relative circumference is, therefore, practically the same, since the stature of white and colored troops is practically equal.

(c) *Standard deviation of waist circumference.*—The standard deviation of waist circumference for white troops, as shown in Table LXXXI, is 6.00 centimeters while that for colored troops is 5.76. In view of the practical equality of the means, this indicates a greater variability of the waist circumference in white troops as compared with colored troops. This relation is brought out more clearly by the coefficients of variation which are, in the case of white troops, 7.705, and in the case of colored troops 7.40.

(d) *Comparison of eight European races.*—Table 89 gives the frequencies and proportional distributions in the different classes of waist circumference for each of the eight races. Table 88 gives in the fourth column from the right the average waist circumference of the different races. It appears from this column that the Germans have the largest waist circumference, 78.46 centimeters, the Polish second, Irish third, and English fourth. On the other hand, the Hebrews have the smallest average waist circumference, followed in order by the Italians, French, and Scotch. The Germans stand second in chest circumference and the Poles first, whereas the Germans stand first in waist circumference and the Poles second. It is clear that there is a relatively greater abdominal development in the Germans than in the Poles. The second column from the right gives the standard deviation as a measure of variability of the different races in respect to waist circumference. The standard deviation is highest, 6.26, among the Irish, next among the Hebrews, then the Germans and English. It is least among the Polish, 5.48, next higher among the French, Italians, and Scotch. The relation of waist circumference to stature is given in the right-hand column in the table. From this column it appears that in relation to stature the Italians have the largest waist circumference, 46.71; they are followed by the Poles, Hebrews, and French. On the other hand, the English have the smallest waist circumference in relation to stature, followed in ascending series by the Scotch, Irish, and Germans. Thus the Nordic race is characterized by small waist circumference as compared with the Mediterranean, Polish, and Hebrew.

TABLE 88.—*Absolute and relative waist circumference with the standard deviation and the coefficient of variation in eight European races, demobilization, 1919.*

Race.	Number measured.	Absolute waist circumference.	Standard deviation.	Coefficient of variation.	Relative waist circumference.
		<i>Centi-meters.</i>	<i>Centi-meters.</i>	<i>Per cent.</i>	<i>Per cent.</i>
English.....	4,195	76.69	6.09	7.941	44.57
Scotch.....	2,061	77.53	6.00	7.739	44.93
Irish.....	6,152	77.70	6.26	8.057	45.34
German.....	7,073	78.46	6.10	7.775	45.61
French.....	1,455	77.32	5.84	7.553	45.86
Italian.....	3,520	77.16	5.87	7.608	46.71
Polish.....	2,405	78.38	5.48	6.992	46.27
Hebrew.....	1,687	76.71	6.11	7.965	45.96

TABLE 89.—*Comparative frequency distribution of waist circumference in each of eight European races, demobilization, 1919.*

SECTION A: ABSOLUTE NUMBERS.

Race.	Total.	Waist circumference, in centimeters.											
		50-63.	64-67.	68-71.	72-75.	76-79.	80-83.	84-87.	88-91.	92-95.	96-99.	100-103.	104 and over.
English.....	4,195	32	67	696	1,121	1,196	531	314	158	48	24	7	1
Scotch.....	2,061	10	34	233	533	589	359	178	81	26	16	1	1
Irish.....	6,152	31	119	650	1,570	1,756	1,110	513	225	94	51	20	13
German.....	7,073	42	98	533	1,609	2,031	1,517	727	320	113	56	19	8
French.....	1,455	9	27	150	388	452	239	111	52	17	7	1	2
Italian.....	3,520	17	73	400	1,004	952	626	275	108	40	15	9	1
Polish.....	2,405	10	19	169	535	720	587	233	94	27	9	2
Hebrew.....	1,687	7	50	240	489	439	251	125	52	21	8	1	4
Number measured..	28,548	158	487	3,071	7,249	8,135	5,220	2,476	1,090	386	186	60	30
Not measured.....	122
Total.....	28,670

SECTION B: PROPORTIONAL RATIOS PER 1,000.

Waist circumference, in centimeters.														
Race.	Total.	50-53.	64-67.	68-71.	72-75.	76-79.	80-83.	84-87.	88-91.	92-95.	96-99.	100-103.	104 and over.	Total.
English.....	4,195	7.63	15.97	165.91	267.21	285.10	126.58	74.85	37.66	11.44	5.72	1.67	0.24	1,000
Scotch.....	2,061	4.85	16.50	113.05	258.62	285.80	174.19	86.37	39.30	12.62	7.76	.49	.49	1,000
Irish.....	6,152	5.04	19.34	105.66	255.21	285.44	180.42	83.39	36.57	15.28	8.29	3.25	2.11	1,000
German.....	7,073	5.94	13.86	75.36	227.49	287.14	214.48	102.78	45.24	15.97	7.92	2.69	1.13	1,000
French.....	1,455	6.19	18.56	103.10	266.68	310.65	164.26	76.29	35.74	11.68	4.81	.69	1.37	1,000
Italian.....	3,520	4.83	20.74	113.63	285.23	270.48	177.85	78.12	30.68	11.36	4.26	2.56	.28	1,000
Polish.....	2,405	4.16	7.90	70.27	222.45	299.36	244.08	96.88	39.09	11.23	3.74	.83	1,000
Hebrew.....	1,687	4.15	29.64	142.27	289.88	260.22	148.79	74.10	30.82	12.45	4.74	.59	2.37	1,000
Number measured.....	28,548	5.53	17.06	107.58	253.93	284.96	182.85	86.74	38.18	13.52	6.52	2.10	1.05	1,000
Not measured.....	122													
Total.....	28,670													

(e) *Comparison of white and colored races.*—A comparison of white and Negro troops with reference to waist circumference has been made in earlier paragraphs and shows no important differences between the races in this respect. Despite the greater circumference of the chest in the white troops, the waist circumference is practically the same in white and colored. This shows that

the Negro troops have the more nearly cylindrical body, the white troops more conical, the increase of the chest over the waist being 102 millimeters in Negro troops and 109 millimeters in white troops.

10. TRANSVERSE DIAMETER OF THE PELVIS.

(a) *General discussion.*—This was measured as the maximum distance between the crests of the ilium. It is measurement No. 40 of Martin⁵ (p. 143):

Grösste Breite zwischen den Darmbeinkämmen (Beckenbreite, Cristalbreite, Distantia inter-cristalis; largeur maximum des hanches, diametre bi-iliaque externe; distance between iliac tubercles). Direct measurement, horizontal distance between the two ilio-cristalia, rod calipers.

The measurement is thus taken against the labium externum of the crista iliaca and the arms of the instrument slightly pressed upon the flesh.

The measurements of 100,000 soldiers were taken practically in accordance with these directions.

The medical importance of this measurement is comparatively unimportant in the case of the male. It may have some relation to hernia, however, not yet determined. The military importance of this measurement is probably confined to its relation to uniforms. The breeches, constricted by the belt, are largely supported by the crest of the pelvis. In the case of slender soldiers the diameter of the body at the waist is less than at the pelvis; in the case of fat men it is greater. It is possible that the relation between circumference of waist and tranverse diameter of pelvis may come to have a medico-military significance, not only as an index of the nutrition of the soldier, but also because of its importance in relation to glandular disturbances that cause the deposition of fat on the omentum and in the body wall of the waist region.

The anthropological significance of pelvic diameter is very great. As Martin points out, this diameter is considerable in man and anthropoids. In Bavarians the breadth of pelvis is about 56 per cent of the length of the trunk (in women nearly 60 per cent). In the gorilla it is even larger, 66.5 per cent, in the chimpanzee 42 per cent, among the lower monkeys 37–25 per cent.

In general the species with broad pelvis have also broad shoulders, producing a rectangular form of the trunk.

The breadth of pelvis may also be expressed in relation to the total height. Here again the difference between the sexes is marked and the figures given here refer only to males. Thus, following Martin's (1914, p. 269) table, among European races the ratio of pelvic diameter to stature is: Jews, 16 per cent; Russians, 16.3 per cent; Poles, 16.4; French in general, 16.5; Parisians, 16.9; Germans in general, 17.0; Roumanians, 17.2. Among Asiatics, the south Chinese have the smallest pelvis, 14.7; Japanese, 15.3 to 16.6; northern Chinese, 17–18.3. Many African tribes have relatively small pelves; Fiot, 14.2; Batua, 14.4; Bushmen, 16.4. Thus Negroes, South Chinese, and Jews have the smallest pelvic diameter of their respective continents. The maximum pelvic diameter is found among the Iroquois Indians, viz., 18.9.

Gould² secured the measurement of the breadth of pelvis of several thousand soldiers and sailors. He gives as a mean dimension 11.92 inches, or 30.28 centimeters, the mean result for men in usual vigor being greater by 0.14 inch (or 0.36 centimeter) than for men in poor health.

Gould² found the following mean values for the breadth of pelvis for men in different parts of the country:

TABLE 90.—*Absolute transverse diameter of the pelvis, by sections, demobilization, 1865.*

Nativity.	Number of men.	Mean value.	Probable variation per individual.	Probable error.	Mean value in centimeters.
		<i>Inches.</i>	<i>Inches.</i>	<i>Inches.</i>	
New England.....	976	11.890	0.675	0.022	30.20
New York alone.....	2,085	12.046	.628	.012	30.60
New York, New Jersey, and Pennsylvania.....	3,119	12.014	.523	.009	30.51
Ohio and Indiana.....	1,417	11.890	.474	.013	30.20

(b) *Mean transverse diameter of the pelvis.*—The mean transverse pelvic diameter of the 95,658 white troops is 29.43 centimeters. The relative pelvic diameter is 17.11. Thus the transverse diameter of the body at the pelvis is 0.23 centimeter greater than the transverse chest diameter in white troops. This increase amounts to 1.36 per cent.

The mean transverse diameter of the pelvis of colored (Negro) troops is 28.42 centimeters, which is 1.01 centimeters less than that of white troops, despite the fact that the stature of the two races is practically the same. The transverse diameter of the pelvis is thus 0.63 centimeter less than the mean transverse diameter of the chest, or 2.169 per cent. The difference between the diameter of the chest and the pelvis is thus greater in colored than in white troops, despite the fact that the body form is more nearly cylindrical in the colored troops. This indicates then that the Negro troops have relatively narrower hips than the white troops and equal waists, but slightly smaller chest circumference. It may be remarked that casual observation of large numbers of Negro troops indicated the frequent presence of individuals with remarkably small pelvic diameter.

(c) *Standard deviation of transverse diameter of the pelvis.*—The standard deviation of transverse pelvic diameter for white troops is 2.85 centimeters and for colored 2.35, indicating a much greater absolute variability in white than in colored troops in this dimension. The coefficient of variation in this dimension is for white troops 9.684 per cent and for colored troops 8.269. Thus the pelvic diameter of colored troops is relatively as well as absolutely much less variable than that of white troops.

(d) *Comparison of eight European races.*—Table 92 gives the absolute and proportional frequencies of the different classes of transverse diameter of the pelvis for each of the eight races. From Table 91, fourth column from the right, it appears that the largest mean transverse diameter of the pelvis is found in the Germans, 29.80; next in the Poles, 29.55, followed by the Scotch and English.

TABLE 91.—*Absolute and relative transverse pelvic diameter, with the standard deviation and the coefficient of variation, in eight European races, demobilization, 1919.*

Race.	Number measured.	Absolute transverse pelvic diameter.	Standard deviation.	Coefficient of variation.	Relative transverse pelvic diameter.
		<i>Centimeters.</i>	<i>Centimeters.</i>	<i>Per cent.</i>	<i>Per cent.</i>
English.....	4,169	28.28	2.73	9.324	17.02
Scotch.....	2,053	28.38	2.84	9.666	17.03
Irish.....	6,108	28.92	2.99	9.302	16.88
German.....	7,051	28.80	2.87	9.631	17.32
French.....	1,429	28.70	2.65	9.233	17.02
Italian.....	3,501	28.62	2.61	9.120	17.33
Polish.....	2,396	28.55	2.64	9.934	17.44
Hebrew.....	1,688	28.34	2.60	9.174	16.98

The average transverse diameter of the pelvis is smallest in the Hebrew, 28.34; next in the Italians, then the French and Irish. The standard deviation as an index of variation is given in the third column from the right. This shows that in respect to transverse diameter of the pelvis, the Germans are the most variable, 2.87; Scotch next, followed by the English and Irish. The Hebrews are the least variable, 2.60, and then in ascending order come the Italians, Polish, and French. The last column at the right gives the relation of the average transverse diameter of the pelvis to average stature of each of the races. From this column it appears that the Poles have the relatively largest pelvic diameter, 17.44; followed by the Italians and Germans. The Irish have the relatively smallest pelvic diameter, 16.88, followed in ascending order by the Hebrew, French, English, and Scotch.

(e) *Comparison of color races.*—The following summary table gives the means of comparing the diameter of the pelvis of five color races:

TABLE 93.—*Absolute and relative transverse diameter of the pelvis in the five color races, demobilization, 1919.*

Race.	Number measured.	Mean diameter, in centimeters.	Relative transverse diameter of pelvis.
White.....	95,658	29.43	17.1
Negro (and mulatto).....	6,354	28.42	16.5
Chinese.....	22	30.00	17.5
Japanese.....	32	28.88	16.9
Indian.....	107	29.71	17.3

The above table shows the comparative transverse pelvic diameter in the different color races. The mean diameter is seen to be 29.4 in the white troops and in Negro troops 28.4. There is, therefore, a difference of over 3 per cent—a deficiency in the Negro troops. The pelvic diameter of the Indians and Chinese is still greater than that of the whites, attaining 30 centimeters in the latter. The pelvic diameter of the Japanese, on the other hand, is only slightly greater than that of the Negro.

11. ARM LENGTH.

(a) *General discussion.*—The length of the arm was measured as the tailor measures it—from the second dorsal vertebra to the styloid process of the ulna of the right arm, the forearm being flexed. The arm length is, therefore, properly not such, but the half-diameter of the chest at the level of the axilla plus the length of the arm as far as the styloid process. This measure is perhaps useful only for tailors, as anthropologists usually measure the length of the arm from the acromion. The length of the arm in the strict sense may be approximately secured by subtracting one-half the transverse diameter of the chest. The relative arm length as measured from the acromion varies widely in different races, as is indicated by the table of Martin ⁵ (1914, p. 294). This is in the case of Bavarians, 35.4 per cent; French, 35 per cent; African Negroes, 35.5 per cent; Mawambu pygmies, 33.3 per cent; Lolo in Hunan, 32.4 per cent.

(b) *Mean arm length.*—The mean “arm” length in the Army measurements was for white troops 78.42 centimeters (Table LXXXII), and for Negro troops, 80.56 centimeters (Table CXIV). Thus it will be seen that with the same mean stature the Negro troops have “arms” which averaged 2.14 centimeters longer than white troops. The difference in relative arm length will be the same as the absolute arm length because of the similarity of height of the two races. If now we subtract the half of the transverse chest diameter from the “arm” length of white troops, we find it to be 78.42 minus 14.51, or 63.91. In the case of Negro troops, it is 80.56 minus 14.53, or 66.03. Thus a comparison of the arm length in the strict sense shows that of the Negro troops to be over 2 centimeters greater than that of white troops. The relative arm length will be secured by dividing these strict arm lengths by the stature. It is 37.16

in the case of whites and 38.40 in the case of Negro troops. From this point of view the relative arm length of Negro troops exceeded that of the white troops by about 3 per cent. This is in accordance with other observations, since, as shown in Martin's tables⁵ (p. 293), there are three African races (Ba-Binga, Lobi, and Bugu) which have a relative entire arm length (including the finger) that is greater than that of any European races.

(c) *Standard deviation of arm length.*—The standard deviation of the "arm" length of the white troops is 4.69; of Negro troops, 4.76. Thus, absolutely, the latter are the more variable. A comparison of the coefficients of variation, however, gives 5.981 per cent for the whites and 5.909 per cent for the Negro troops. Thus the Negro troops are relatively less variable than the whites in this respect.

Double the arm length plus length of wrist and fingers is approximately equal to span. We have seen that span is greater in colored than in whites, just as "arm" length is. Also, both measures are absolutely more variable in the colored troops. Thus by both tests the arms of the colored are longer and absolutely more variable than those of white troops.

12. FOREARM LENGTH.

Table LXXXII shows the correlation between "total arm length" and that of forearm in white troops. The mean length of the forearm (that is, from the olecranon process to the styloid process) is, in the case of white troops, 26.91 centimeters, and in the case of colored troops, 28.20 centimeters. Dividing the mean forearm by the total "arm length," minus half the transverse chest diameter, we find that for white troops the forearm constitutes 42.01 per cent of the whole arm length and for colored troops, 42.71 per cent. Thus the forearm length is not only absolutely greater in colored than in whites but also constitutes a relatively larger proportion of the arm length.

The *relative* length of forearm (i. e., in proportion to stature) is in the case of white troops 15.65 per cent and in the case of Negro troops 16.40. In Martin's table (1914, p. 297) it appears that some of the African Negroes have a relatively greater arm length than any of the European races listed, even as great as 17.7 per cent. The relative arm length of these European races varies from 14.3 per cent (Parisians) to 15.5 per cent (Bavarians) and 15.9 per cent (Germans and Jews). Martin also notes that in exceptionally long arms excess length is especially due to the great length of forearm.

13. LEG LENGTH.

(a) *General discussion.*—The measurement here called leg length is actually the distance from the gluteal fold to the tip of the internal malleolus of the tibia, as measured by a tape. It is to be noted that this dimension added to the sitting height falls about 10 centimeters short of the total stature. The difference is due, on the one hand, to the height of the internal malleolus above the floor, which is about 8 centimeters. The remaining 2 centimeters are accounted for by the sag of the gluteal muscles in the standing subject, so that the gluteal fold lies about 2 centimeters farther from the vertex in the standing subject than in the sitting subject.

The leg length as thus measured is not the physiological leg length, but primarily of interest to the manufacturer of uniforms and other clothing. It is much less valuable from a military point of view than the total leg length as indicated by the height of the pubic arch. The leg length may also be secured by subtracting the sitting height from the total stature.

(b) *Mean leg length.*—The mean "leg length" as defined is for white troops 71.69 centimeters, as indicated by Table LXXVI. For Negro troops it is 74.38, as shown in Table XCII. Thus there is a difference of 2.69 centimeters between white and Negro troops, or 3.75 per cent of the "leg length" of the whites. The relative "leg length" is 41.68 per cent of height for white troops.

We may compare the leg length found by subtracting the sitting height from the total stature. In white troops this is 171.99 minus 90.39, or 81.60 centimeters. In the case of Negro troops it is 171.97 minus 87.35, or 84.62. Thus, by these means also we find an excess of 3.57 per cent in the leg length of Negro troops as compared with whites. Since the anthropoid apes are characterized by relatively short legs, the Negro in this respect represents a greater departure from the anthropoid types than do the whites. The *relative* leg length, determined by the method of subtracting sitting height from body height, is in the case of white troops 47.45 per cent and in the case of Negro troops 49.21 per cent. These figures are in good agreement with those given in Martin's table (p. 312), where the relative leg length obtained in this way is for Europeans mostly between 47.0 and 48.5 per cent, while for different African tribes it varies from 47.2 to 49.7 per cent. Armenians and Tartars have a relative leg length below the average; the American Indians show a great range in this respect.

(c) *Standard deviation of leg length.*—The standard deviation of "leg length" is for white troops, as shown in Table 103, 4.71, and for colored troops (Table 104) 4.59. The corresponding coefficients of variation are for white troops 6.57 and for colored 6.17. This shows again a lower relative variability in colored than in white troops in respect to this dimension.

(d) *Comparison of eight European races.*—Table 95 gives the distribution of absolute and proportional frequencies of "leg length" in the eight races. It will be recalled that the leg length is the distance from the gluteal fold to the internal malleolus and includes, therefore, the sum of the thigh and lower leg, excluding the foot. Table 94 shows in the third column from the right the average leg length of the different races. This is greatest in the Scotch, 71.68, and next in the Germans, followed by the English and Irish. It is lowest in the Italians, followed by the Hebrews, French, and Poles. The third column from the right gives the variability of leg lengths for the different races. This is seen to be greatest among the English, next among the Scotch, and then in order among the Germans and Irish. It is least, 4.19, among the Italians, and then slightly greater in order among the Hebrews, Poles, and French.

TABLE 94.—*Absolute and relative leg length with the standard deviation and the coefficient of variation in eight European races, demobilization, 1919.*

Race.	Number measured.	Absolute leg length.	Standard deviation.	Coefficient of variation.	Relative leg length.
		<i>Centimeters.</i>	<i>Centimeters.</i>	<i>Per cent.</i>	<i>Per cent.</i>
English.....	4,152	71.34	4.59	6.434	41.46
Scotch.....	2,038	71.68	4.56	6.362	41.54
Irish.....	6,110	70.91	4.39	6.191	41.38
German.....	7,012	71.47	4.51	6.310	41.54
French.....	1,438	69.22	4.34	6.270	41.06
Italian.....	3,446	67.84	4.19	6.176	41.07
Polish.....	2,377	70.16	4.30	6.129	41.41
Hebrew.....	1,664	68.93	4.29	6.224	41.30

Since leg length is partly dependent upon stature, the right-hand column shows that the Scotch and Germans have relatively the greatest leg length, 41.54, as above defined; they are followed by the English and Poles. The French have the least relative leg length, 41.06, as above defined, being in this respect close to the Italians. Considerably above them stand the Hebrews and Irish. Since the French have relatively the greatest height of pubic arch and the shortest relative leg length, it would follow either that the ankle is relatively high in the French or that the symphysis pubis is placed relatively high.

TABLE 95.—Comparative frequency distribution of leg length in each of eight European races, demobilization, 1919.

SECTION A. ABSOLUTE NUMBERS.

Race.	Total.	Leg length, in centimeters													
		60-61	62-63	64-65	66-67	68-69	70-71	72-73	74-75	76-77	78-79	80-81	82-83	84-85	86-87
English.....	4,162	31	124	245	467	598	721	662	557						14
Scotch.....	2,038	20	43	105	192	313	375	354	245						4
Irish.....	6,110	61	156	379	723	1,045	1,184	947	700						7
German.....	7,012	57	166	368	655	1,046	1,343	1,100	691						15
French.....	1,438	41	88	152	225	286	246	175	109						1
Italian.....	3,446	174	351	543	638	619	477	328	187						2
Polish.....	2,377	37	88	164	341	459	447	359	244						3
Hebrew.....	1,664	52	106	200	270	333	274	194	104						2
Number measured.....	28,237	473	1,122	2,204	3,527	4,659	5,077	4,209	3,007						48
Not measured.....	433														
Total.....	28,670														

SECTION B. PROPORTIONAL RATIOS PER 1,000.

Race.	Total.	Leg length, in centimeters.														Total.
		60-61	62-63	64-65	66-67	68-69	70-71	72-73	74-75	76-77	78-79	80-81	82-83	84-85	86-87	
English.....	4,162	7.47	29.87	59.01	112.46	144.08	176.06	169.44	134.15	86.23	46.85	22.88	10.90	4.66	3.37	1,000
Scotch.....	2,038	9.81	21.10	51.52	99.31	148.66	194.00	173.70	120.22	92.25	56.37	31.40	13.25	3.43	1.96	1,000
Irish.....	6,110	9.96	26.53	62.04	119.97	171.03	193.79	166.00	114.57	78.44	50.45	17.84	9.49	3.76	1.15	1,000
German.....	7,012	8.13	23.67	55.47	94.94	146.16	191.83	166.71	127.07	83.81	44.78	23.24	12.55	6.13	2.14	1,000
French.....	1,438	28.51	61.20	105.70	156.47	193.59	171.07	121.70	75.80	43.81	21.56	9.04	4.87	2.70	1.00	1,000
Italian.....	3,446	50.50	101.86	157.59	185.14	179.63	128.42	95.19	45.56	26.70	13.05	4.06	1.45	0.29	0.53	1,000
Polish.....	2,377	15.57	37.02	77.41	143.46	190.48	168.08	161.03	102.65	54.80	26.24	12.62	4.21	4.21	1.26	1,000
Hebrew.....	1,664	31.25	63.70	120.19	165.87	200.12	164.67	116.59	62.50	40.67	22.84	7.21	2.40	0.60	1.20	1,000
Number measured.....	28,237	16.75	36.73	78.05	124.91	163.00	179.80	149.06	105.49	70.55	37.43	18.20	8.61	3.72	1.70	1,000
Not measured.....	433															
Total.....	28,670															

(e) *Comparison of color races.*—The following table gives a summary of the absolute and relative leg lengths of the five colored races measured:

TABLE 96.—*Absolute and relative leg length in five color races, demobilization, 1919.*

Race.	Number measured.	Mean length.	Relative leg length.
		<i>Centimeters.</i>	<i>Per cent.</i>
White.....	76,141	71.69	41.7
Negro (and mulatto).....	5,595	74.38	43.3
Chinese.....	22	70.86	41.4
Japanese.....	29	74.22	43.4
Indian.....	106	71.63	41.8

The distance from the gluteal fold to the internal malleolus in the different races is shown in the table above. We see from this table again that the leg length is over 2.69 centimeters greater in Negro troops than in white, despite the practical equality in total stature. The relative leg length is 43.3 per cent among the Negro troops, 41.7 per cent among the whites, and 41.4 per cent among the Chinese; the Japanese, 43.4, and Indian, 41.8. The Japanese in this respect are more like the Negro troops.

14. KNEE HEIGHT.

(a) *General discussion.*—Knee height was taken as the distance from the floor to the top of the patella. It has relatively small military importance, excepting in so far as by subtracting it from the "leg length" plus 8 centimeters the length of the thigh will be given, from which may be estimated the corresponding dimensions of the breeches.

(b) *Mean knee height.*—The mean knee height for white troops is 47.08 (Table 103); for colored troops, 47.26 (Table 104). That of the colored troops is sensibly greater than that of the white troops. In the case of white troops the knee height constitutes 65.67 per cent of leg length, and in the case of the colored troops 63.54 per cent. Thus in the colored troops the lower leg is relatively a smaller proportion of the whole leg length than in the case of the white troops; consequently the thigh is relatively long. This is in striking contrast to the conditions found in the upper appendage, where the forearm (exclusive of the hand) proves to form a relatively larger proportion of the whole arm in colored than in white troops. Since the proportion of knee height to total stature is, in the case of white troops, 27.38 per cent and 27.48 in the case of colored, in relation to total stature the lower leg of the colored troops is greater than that of white troops, and this despite the fact that it constitutes a smaller fraction of the total "leg length."

The index giving the relation of upper leg to lower leg (excluding the foot) may be calculated as follows:

	White.	Colored.
Pubic height.....	86.82	89.42
Knee height.....	47.08	47.26
Thigh.....	39.74	42.16

Also the lower leg length in the strict sense, excluding the foot, may be approximately determined by subtracting 8 centimeters from the knee height. This gives us, then, in the case of whites, a net lower leg length of 39.08 centimeters; in the case of colored, 39.26 centimeters.

(c) *Standard deviation of knee height.*—This is 3.62 centimeters in white troops and 3.64 centimeters in colored; the length of lower leg and foot is absolutely more variable in white than in Negro troops, despite their shorter length in whites. The coefficient of variability of this dimension is in white troops 7.689 per cent and in colored 7.702 per cent. This is a relatively high coefficient.

TABLE 97.—*Absolute and relative knee height with the standard deviation and the coefficient of variation in eight European races, demobilization, 1919.*

Race.	Number measured.	Absolute knee height.	Standard deviation.	Coefficient of variation.	Relative knee height.
		<i>Centimeters.</i>	<i>Centimeters.</i>	<i>Per cent.</i>	<i>Per cent.</i>
English.....	3,171	47.74	4.14	8.672	27.74
Scotch.....	1,651	47.83	3.91	8.175	27.72
Irish.....	4,708	46.59	3.72	7.985	27.19
German.....	5,646	47.22	3.74	7.920	27.45
French.....	701	46.88	3.84	8.200	27.78
Italian.....	2,880	45.13	3.51	7.778	27.32
Polish.....	1,917	46.09	3.66	7.939	27.56
Hebrew.....	1,468	45.57	3.50	7.878	27.30

15. THIGH CIRCUMFERENCE.

(a) *General discussion.*—Measurers were instructed to secure the maximum circumference of the thigh by means of a tape passed around the upper portion of the thigh and moved slightly upward until it reached the level of the gluteal fold.

The military importance of this measurement is probably not great, though there is possibly a correlation (never determined, however) between the thigh girth and the capacity of the soldier to make prolonged marches and carry heavy burdens. The circumference of the thigh was used in the table of manikin dimensions (Table 122) to secure the greatest breadth of the "hips" or greatest transverse diameter at the level of the gluteal fold. This was obtained by taking twice the quotient of the circumference of the thigh divided by π , or 3.1416.

The thigh girth in relation to stature varies in different races. It attains its smallest dimensions in certain African tribes. Thus in the Ba-Tua the relative thigh girth is given as 28.2 (Martin,⁵ p. 322). The length of the thigh divided by its circumference gives an index which varies markedly during developmental years. In the case of children 14–15 years this ratio is about 52 per cent.

(b) *Mean thigh circumference.*—The mean thigh circumference for white troops is 52.709 centimeters, as shown in Table 103. The corresponding measurement for colored troops is 54.077 (Table 104). Thus in the colored troops it is 1.3 centimeters greater than in white troops. The relative thigh circumference is 30.65 per cent of stature in the case of white troops, about the same as for the average European (Martin,⁵ p. 322). In the case of colored troops it is 31.45, about the same as for the Ba-Binga, as shown in Martin (1914, p. 322). The length divided by the circumference is 75.60 per cent in the white troops and 77.96 in the colored.

(c) *Standard deviation of thigh circumference.*—The standard deviation of thigh circumference is for white troops 3.73 centimeters, as shown in Table 103, and for colored troops 3.72, or practically the same. Since the mean circumference is greater for the colored troops than for the whites, the coefficient of variability of the colored troops (6.88) is less than that for the whites (7.08).

(d) *Comparison of eight European races.*—Tables 100 give the absolute and proportional frequencies of each of the different classes of thigh girth for each of the eight races. In Table 99 the third column from the left gives the mean thigh girth for each of these races. From this column it appears that at demobilization the men of German origin showed the greatest thigh girth, 53.19. These were followed by the Poles, English, and Scotch. On the other hand, the French have the smallest thigh girth, 51.98, followed in ascending order by the Italians, Hebrews, and Irish. The third column from the right gives the standard deviation as an index of variability in these races. From this column it appears that the thigh girth is most variable in the Irish, 3.68; next in the English, 3.66; then in the Germans, Italians, and Hebrews. It is least variable in the French, 3.44; followed in ascending series by Polish and Scotch. Thigh circumference in relation to stature is given in the right-hand column of

Table 99. From this column it appears that the Italians have the relatively largest thigh girth, 31.50, followed by the Hebrews, Polish, and Germans. The Scotch have the relatively smallest thigh girth, 30.35, followed in ascending order by the English, Irish, and French. Thus in general the Mediterranean peoples and Hebrews have the largest relative thigh girth; the Nordic races and the French the relatively smallest thigh girth. This is another index of the slenderness of the Nordics.

TABLE 99.—*Absolute and relative thigh circumference with the standard deviation and the coefficient of variation in eight European races, demobilization, 1919.*

Race.	Number measured.	Absolute thigh circumference.	Standard deviation.	Coefficient of variation.	Relative thigh circumference.
		<i>Centimeters.</i>	<i>Centimeters.</i>	<i>Per cent.</i>	<i>Per cent.</i>
English.....	4, 146	52.38	3.66	6.987	30.44
Scotch.....	2, 037	52.36	3.56	6.799	30.35
Irish.....	6, 070	52.27	3.68	7.040	30.50
German.....	6, 960	53.19	3.62	6.806	30.92
French.....	1, 451	51.98	3.44	6.618	30.83
Italian.....	3, 489	52.03	3.59	6.900	31.50
Polish.....	2, 385	52.46	3.45	6.576	30.97
Hebrew.....	1, 664	52.18	3.58	6.861	31.26

16. CALF CIRCUMFERENCE.

(a) *General discussion.*—The instructions to anthropologists at camps called for the measurement of the maximum circumference of the calf.

Martin⁵ states (p. 322) that the stronger or weaker development of the calf rests either upon the development of the *musculus triceps suræ* or on the degree of enlargement of the *panniculus adiposus*. The latter factor contributes more to the circumference of the calf in the female, the former in the male sex. Strongly muscular calves indicate a highly placed belly of the *gastrocnemius* muscle, while calves of smaller circumference are characterized by a *gastrocnemius* with longer muscle fibers but smaller cross section. Also there is a correlation with the length of the tibia, since with a shorter tibia there is found prevalently a *gastrocnemius* with short-muscle belly and longer tendons; with the longer bone, on the contrary, the muscle with a long belly and short tendons. The calf of small circumference (i. e., a slight development in breadth and thickness of the *musculus triceps suræ*, and with a low lying transition of the muscle into the terminal tendons) is found especially in the Negro groups, among the Egyptians, Australians, Dravida, and Weddas; while thicker and shorter calves are characteristics of most European groups, and of the Mongoloid and Malay varieties.

The military importance of the circumference of the calf is slight. It measures something of the degree of development of the *gastrocnemius* muscle, which is of great importance in marching.

(b) *Mean calf circumference.*—The mean calf circumference of white troops is 34.09 centimeters and that of the colored troops 34.71 centimeters, which is 0.62 centimeter greater than that of the white troops. This is the more remarkable in view of the general slenderness of calf in African tribes. The circumference of the calf in relation to total stature is found from the data given in Tables 103 and 104 for white and colored troops, respectively. In the case of the former it is 19.82 per cent, which is somewhat less than the average European, placed by Martin⁵ (p. 322) at 20.5. In colored troops it is 20.18.

The relation between the maximum calf circumference and thigh circumference is, in the case of white troops, 64.7 per cent, and in the case of colored troops, 64.2 per cent. These are rather low ratios compared with those given by Martin⁵ (p. 322), which lie between 66.3 and 70 per cent in the male.

(c) *Standard deviation of calf circumference.*—The standard deviation of calf circumference in white troops is 2.019 centimeters and in the case of colored 2.01. The coefficient of variation is, in the case of white troops, 5.93 per cent, and in the case of colored, 5.79 per cent. Thus the calf circumference is much more variable in colored than in white troops.

(d) *Comparison of eight European races.*—Table 102 gives the absolute and proportional frequencies of occurrence of the different classes of calf circumference for each of the eight races.

The third column from the left of Table 101 gives the mean calf circumference. This varies in the different races from a maximum in the Polish of 34.44, followed in descending order by the Germans and Scotch. The minimum average calf circumference, 33.68, is found among the Hebrews, followed in ascending order by the French, Italians, Irish, and English. The relative variability in this dimen-

sion in the various races is indicated by the standard deviation, third column from the right. According to this the English and Irish have the greatest variability in calf circumference, 2.07, followed by the Scotch. Relatively slight variability is found in the Polish, 1.93, followed in increasing order by the French, Germans, Italians, and Hebrews. Thus, the more northern races show greater variability in respect to this dimension. In the right-hand column of Table 101 is given the calf circumference in relation to stature. The relatively greatest calf circumference is found among the Italians, 20.41; followed by the Polish, Hebrews, Germans, and French. The relatively smallest calf circumference is found among the English, 19.70; followed in ascending order by the Scotch and Irish. Thus the northern races show the smallest relative calf circumference, which is in accordance with the generally slender build of these people.

TABLE 101.—*Absolute and relative calf circumference, with the standard deviation and the coefficient of variation in eight European races, demobilization, 1919.*

Race.	Number measured.	Mean absolute calf circumference.	Standard deviation.	Coefficient of variation.	Relative calf circumference.
		Cm.	Cm.	Per cent.	Per cent.
English.....	4,214	33.90	2.07	6.106	19.70
Scotch.....	2,079	34.04	2.06	6.052	19.73
Irish.....	6,174	33.83	2.07	6.119	19.74
German.....	7,094	34.40	2.02	5.872	20.00
French.....	1,463	33.68	1.96	5.820	19.98
Italian.....	3,532	33.71	2.04	6.052	20.41
Polish.....	2,417	34.44	1.93	5.604	20.33
Hebrew.....	1,697	33.66	2.04	6.061	20.17

TABLE 102.—*Comparative frequency distribution of calf circumference in each of eight European races, demobilization.*

SECTION A: ABSOLUTE NUMBERS.

Race.	Total.	Calf circumference, in centimeters.											
		29	30	31	32	33	34	35	36	37	38	39	40
English.....	4,214	29	140	320	609	766	802	662	431	232	119	92	12
Scotch.....	2,079	15	57	136	285	369	382	350	239	139	51	50	6
Irish.....	6,174	70	218	462	865	1,177	1,211	950	598	327	149	126	21
German.....	7,094	32	131	314	731	1,172	1,415	1,245	992	572	290	184	16
French.....	1,463	15	47	116	243	281	284	228	129	75	23	21	1
Italian.....	3,532	39	141	295	533	672	524	352	159	82	61	2	
Polish.....	2,417	6	34	102	230	403	484	468	340	204	92	51	3
Hebrew.....	1,697	24	58	149	253	350	322	263	137	63	36	39	3
Total.....	28,670	230	826	1,894	3,749	5,190	5,572	4,690	3,218	1,771	842	624	64

SECTION B: PROPORTIONAL RATIOS PER 1,000.

Race.	Total.	Calf circumference, in centimeters.												Total.
		29	30	31	32	33	34	35	36	37	38	39	40	
English.....	4,214	6.88	33.22	75.94	144.52	181.78	190.32	157.10	102.28	55.06	28.24	21.83	2.85	1,000
Scotch.....	2,079	7.22	27.42	65.42	137.10	177.49	183.74	168.35	114.96	66.86	24.53	24.05	2.89	1,000
Irish.....	6,174	11.34	35.31	74.83	140.10	190.64	196.15	153.87	96.87	52.97	24.13	20.41	3.40	1,000
German.....	7,094	4.51	18.47	44.26	103.04	165.21	199.46	175.50	139.84	80.63	40.88	25.94	2.26	1,000
French.....	1,463	10.25	32.13	79.29	166.10	192.07	194.12	155.84	88.18	51.27	15.72	14.35	.68	1,000
Italian.....	3,532	11.04	39.92	83.52	150.90	190.26	190.26	148.36	99.66	45.02	23.22	17.27	.57	1,000
Polish.....	2,417	2.48	14.07	42.20	95.16	166.73	200.25	193.63	140.67	84.40	38.06	21.10	1.24	1,000
Hebrew.....	1,697	14.14	34.18	87.80	149.09	206.25	189.75	154.98	80.73	37.12	21.21	22.98	1.77	1,000
Total.....	28,670	8.02	28.81	66.06	130.76	181.02	194.35	163.59	112.24	61.77	29.37	21.76	2.23	1,000

17. SUPRAPATELLA CIRCUMFERENCE.

(a) *General discussion.*—The directions required the anthropologists to take the circumference of the leg above the patella. The importance of this measurement seemed to be primarily for uniforms, as these fit closely at this part of the leg.

(b) *Mean suprapatella circumference.*—The mean suprapatella circumference was 37.336 centimeters for white troops, and 37.611 centimeters for Negro, thus about 0.3 centimeter greater in Negro troops, corresponding with their generally greater girth of leg. Since the stature of the white and Negro troops is the same, the relative circumference of the suprapatella region is in the same proportion as the mean.

(c) *Standard deviation of suprapatella circumference.*—The standard deviation of suprapatella circumference is 2.45 centimeters for white troops and 2.43 centimeters for Negro troops, or nearly the same. The coefficient of variation of suprapatella circumference in white troops is 6.56, a relatively high variability, and in the case of Negro troops it is 6.46, a strikingly lower variability.

18. KNEE-PATELLA CIRCUMFERENCE.

(a) *General discussion.*—The instructions for anthropologists called for the measurement of the knee at the level of the patella. This measurement was taken primarily for the fitting of uniforms.

(b) *Mean knee-patella circumference.*—The mean patella circumference among white troops is 36.21 centimeters, and in colored troops 36.52 centimeters. Thus the patella circumference of the colored troops exceeds markedly that of the white troops which is in accordance with the greater girths of other parts of the leg.

(c) *Standard deviation of knee-patella circumference.*—The standard deviation of patella circumference is for white troops 1.979, and for Negro troops 1.987. The relative variability in the whites is 5.47 per cent and in the Negro troops 5.45 per cent. Here again this dimension shows itself relatively less variable in the Negro than in the white troops.

19. COMPARISON OF DIMENSIONS OF WHITE AND NEGRO TROOPS.

(a) *Comparison of means of whites and Negroes.*—In the preceding sections there have been given for many of the dimensions the averages found in the color races. The numbers involved are small in the case of Japanese, Chinese, and Indians, but are so considerable in the case of white and Negro troops as to make a comparison significant.

Tables 103 and 104 give the differences in means and standard deviations of 20 dimensions of white and Negro troops. The results of these tables are shown graphically in Plate I. From the tables and the plate it appears that whereas the average height of white and Negro soldiers is practically the same the Negro men exceeded, on the average, the white men in the following dimensions:

(b) *Span.*—The total span of the Negroes is about 3 per cent greater than that of white men.

(c) *Leg length*.—Since the lengths of arm and leg are correlated in animals generally, it is in accordance with expectation to find that the leg is longer in the Negro than in the white troops, showing an excess of about 3 per cent.

(d) *Arm length*.—As this constitutes an important part of the span, we may expect, as we find, that arm length will be greater in the Negro than in the white troops.

(e) *Pubic height*.—This measures the physiological length of leg and shows about the same excess as leg length.

(f) *Knee height*.—As a component of leg length, knee height shows a slight excess in Negro over white troops.

(g) *Forearm*.—This, as in the segments of the arm length, shows an excess in the Negro troops.

(h) *Sternal notch*.—This is slightly greater in Negro than in white troops. Consequently the height of neck and head together must be less in Negro than in white troops.

(i) *Sitting height*.—Since the total height is the same and the leg length greater in Negro than in white troops (Gould, 1865, pp. 253, 255, 299; also our Table 108), it is clear that sitting height must be less in Negro than in white troops, and such proves to be the case. This smaller sitting height is due in part to the smaller length of head and neck in Negro troops as compared with white troops, but also the length of the trunk from the gluteal fold to sternal notch is relatively less in Negro than in white troops.

In contrast with the vertical dimensions the circumferences and diameters show for the most part relatively slight differences between white and Negro troops; largely because they are smaller dimensions. However, certain differences are clearly shown. The circumferences of the trunk, whether taken at chest or at waist, are slightly less in Negro than in white troops. The transverse diameter of the pelvis is strikingly less in Negro troops. The breadth of the shoulder is, however, somewhat greater in Negro than in white troops and the same is true of the circumference of the neck, thigh, and calf.

Despite approximately the same height, Negro troops weighed nearly 5 pounds more than white troops. The index of build of the Negro troops was about 32.7 as compared with 31.6 for white troops.

The general comparative picture we get of the white troops (including a great variety of races) and the Negro troops is this: The Negro troops have relatively longer legs and arms; shorter trunks; smaller circumference of the waist; more nearly parallel outlines of the trunk; the waist is less marked because of the relatively small transverse diameter of the pelvis and chest; less nearly circular ellipse on cross section of the chest; larger, shorter necks; larger leg girth; and greater weight than the whites. The Negro seems more powerfully developed from the pelvis down and the white more powerfully developed in the chest.

In summary, then, the main differences of shape between Negro and white troops are that the former have relatively longer appendages, shorter trunk, head, and neck, broader shoulders, narrower pelvis, and greater girth of neck, thigh, and calf than the latter.

TABLE 103.—Summary of dimensions of approximately 100,000 white troops, demobilization.

	Number of men measured.	Mean.		Relation to height (centimeters).	Standard deviation (centimeters).	Probable error (centimeters).	Coefficient of variation (centimeter) (per cent).	Standard deviation (inches).	Table from which figures were taken.
		Centimeters.	Inches.						
Stature.....	96,596	171.99	67.72	6.66	± 0.0102	3.872	2.62	LXXXXIII
Span.....	96,596	175.58	69.13	102.10	7.95	± .0122	4.528	3.13	LXXXXIV
Sitting height.....	96,239	90.39	35.59	52.55	3.51	± .0054	3.883	1.38	LXXXXIII
Pubic arch, height.....	91,365	86.82	34.18	50.47	5.05	± .0080	5.817	1.99	LXXXXVI
Sternal notch, height.....	96,439	141.18	55.58	82.09	5.91	± .0091	4.186	2.33	LXXXXV
Leg length.....	76,141	71.69	28.22	41.68	4.71	± .0081	6.570	1.85	LXXXXVI
Knee height.....	76,141	47.08	18.54	27.38	3.62	± .0063	7.689	1.43	LXXXXVI
Arm length.....	94,940	78.57	30.87	45.00	4.69	± .0053	5.981	.90	LXXXXVI
Forearm.....	82,492	26.91	10.59	15.65	1.73	± .0029	6.429	.68	LXXXXII
Chest circumference.....	96,867	88.79	34.96	51.62	5.09	± .0078	5.733	2.00	XCIX
Chest transverse.....	96,583	29.02	11.42	16.87	2.40	± .0037	8.270	.94	LXXX
Chest, antero-posterior.....	96,583	21.58	8.50	12.55	1.87	± .0029	8.665	.74	LXXX
Shoulder width.....	95,167	41.81	16.46	24.31	2.41	± .0037	5.764	.95	CJ
Neck circumference.....	95,271	35.98	14.16	20.92	1.80	± .0028	5.003	.71	LXXXVIII
Waist circumference.....	96,157	77.87	30.66	45.28	6.00	± .0093	7.705	2.36	LXXXXI
Transverse pelvic diameter.....	95,658	29.43	11.59	17.11	2.85	± .0044	9.684	1.12	LXXXXI
Thigh circumference.....	95,188	52.71	20.75	30.65	3.73	± .0058	7.076	1.47	CXVIII
Suprapatella.....	96,157	37.34	14.70	21.71	2.45	± .0056	6.561	.96	CXIX
Knee patella.....	96,157	36.21	14.26	21.05	1.98	± .0030	5.468	.78	CXX
Calf circumference.....	96,087	34.09	13.42	19.82	2.02	± .0045	5.925	.80	CXXI
Weight.....	79,706	a 65.62	b 144.67	c 31.56	a 7.67	± .0424	3.587	b 16.92	

a Kilograms.

b Pounds.

c Weight in pounds×1,000.
(Height in inches.)³

TABLE 104.—Summary of dimensions of approximately 6,000 colored troops, demobilization.

	Number of men measured.	Mean.		Relation to height (centimeters).	Standard deviation (centimeters).	Probable error (centimeters).	Coefficient of variation (centimeter) (per cent).	Standard deviation (inches).	Table from which figures were taken.
		Centimeters.	Inches.						
Stature.....	6,441	171.97	67.70	6.91	± 0.0410	4.018	2.72	LXXXXIX
Span.....	6,441	180.76	71.17	105.10	8.59	± .0510	4.752	3.38	LXXXXVIII
Sitting height.....	6,443	87.35	34.39	50.79	3.48	± .0207	3.964	1.37	LXXXXVII
Pubic arch, height.....	6,220	89.42	35.21	52.00	5.27	± .0319	5.894	2.07	XC
Sternal notch, height.....	6,454	142.39	56.06	82.80	6.05	± .0359	4.249	2.38	LXXXXIX
Leg length.....	5,596	74.38	29.28	43.25	4.59	± .0292	6.171	1.81	XCII
Knee height.....	5,725	47.26	18.61	27.48	3.64	± .0229	7.702	1.43	XCII
Arm length.....	6,135	80.56	31.72	46.85	4.76	± .0213	5.909	.93	CXIV
Forearm.....	5,514	28.20	11.10	16.40	2.03	± .0131	7.199	.80	CXVIII
Chest circumference.....	6,355	87.99	34.64	51.17	4.76	± .0285	5.410	1.48	CVII
Chest transverse.....	6,450	29.05	11.44	16.99	2.26	± .0134	7.780	.89	XCVI
Chest, antero-posterior.....	6,450	21.21	8.35	12.33	1.74	± .0103	8.204	.69	CIX
Shoulder width.....	6,289	42.89	16.88	24.94	2.15	± .0130	5.013	.85	XCIV
Neck circumference.....	6,280	36.37	14.32	21.15	1.72	± .0103	4.729	.68	CXXII
Waist circumference.....	6,445	77.83	30.64	45.25	5.76	± .0342	7.401	2.27	XXV
Transverse pelvic diameter.....	6,354	28.42	11.19	16.53	2.35	± .0140	8.269	.93	CXXVII
Thigh circumference.....	6,367	54.08	21.29	31.45	3.72	± .0330	6.879	1.46	XXV
Suprapatella.....	6,443	37.61	14.81	21.87	2.43	± .0214	6.461	.96	CXXVI
Knee patella.....	6,444	36.52	14.38	21.24	1.99	± .0175	5.449	.78	CXXVII
Calf circumference.....	6,444	34.71	13.67	20.18	2.01	± .0125	5.791	.79	CXXVIII
Weight.....	3,319	a 67.83	b 149.53	c 32.65	a 8.00	± .1452	3.203	b 17.53	

a Kilograms.

b Pounds.

c Weight in pounds×1,000.
(Height in inches.)³

TABLE 105.—Average dimensions in color races, demobilization, 1919.

Dimension.	96,500 white. ^a	6,400 colored. ^b	22 Chinese. ^c	29 Japa- nese. ^c	106 Indians. ^c
Stature.....centimeters	171.99	171.97	171.11	170.94	171.51
Weight.....pounds	144.67	148.53	148.94	144.92	150.13
Index of build.....	31.55	32.55	32.82	32.00	32.93
Sitting height.....centimeters	90.39	87.35	89.05	87.88	90.10
Span.....do	175.58	180.76	176.41	177.25	176.86
Sternal notch.....do	141.18	142.39	140.86	140.44	140.97
Pubic height.....do	86.82	89.42	86.12	88.31	86.35
Knee height.....do	47.08	47.26	46.20	46.71	46.97
Leg length.....do	71.69	74.38	70.86	74.22	71.63
Arm length.....do	78.42	80.56			
Forearm.....do	26.91	28.20			
Chest circumference.....do	88.79	87.99			
Chest transverse.....do	29.02	29.05			
Chest antero-posterior.....do	21.58	21.21			
Shoulder width.....do	41.81	42.99	42.67	42.00	42.58
Pelvic width.....do	29.43	28.42	30.00	28.88	29.71
Neck circumference.....do	35.98	36.37			
Waist circumference.....do	77.87	77.83			
Thigh circumference.....do	52.71	54.08			
Suprapatella.....do	37.34	37.61			
Patella.....do	36.21	36.52			
Calf circumference.....do	34.09	34.71			

^a See Table 103.

^b See Table 104.

^c See Table 107.

TABLE 106.—Relative dimensions in color races, demobilization, 1919.

[Percentage rates.]

Dimension.	White.	Colored.	Indians.	Chinese.	Japanese.
Weight (in lbs.) [Stature (in.)] ^a	31.56	32.65	32.93	32.82	32.00
Sitting height (cm.) Stature (cm.).....	52.55	50.79	52.53	52.04	51.41
Sternal notch (cm.) Stature (cm.).....	82.09	82.80	82.19	82.32	82.15
Pubic height (cm.) Stature (cm.).....	50.47	52.00	50.35	50.33	51.66
Leg length (cm.) Stature (cm.).....	41.68	43.25	41.76	41.41	43.42
Knee height (cm.) Stature (cm.).....	27.38	27.48	27.39	27.00	27.33
Span (cm.) Stature (cm.).....	102.10	106.10	103.10	103.10	103.70
Shoulder width (cm.) Stature (cm.).....	24.31	24.94	24.83	24.94	24.57
Chest circumference (cm.) Stature (cm.).....	51.62	51.17			
Chest circumference (cm.) Weight in lbs.....	61.37	58.84			
Transverse chest (cm.) Antero-post.chest (cm.).....	134.48	136.96			
Antero-post.chest (cm.) Stature (cm.).....	12.55	12.33			
Waist circumference (cm.) Stature (cm.).....	45.28	45.25			
Pelvic diameter (cm.) Stature (cm.).....	17.11	16.53	17.32	17.53	16.90
Thigh circumference (cm.) Stature (cm.).....	30.65	31.45			
Calf circumference (cm.) Stature (cm.).....	19.82	20.18			

SECTION C: SPAN.

[illegible]

SECTION D: HEIGHT OF STERNAL NOTCH.

[illegible]

SECTION G: LEG LENGTH.

Race.	Total.	Leg length, in centimeters.																	Mean leg meas-ure.
		60-61	62-63	64-65	66-67	68-69	70-71	72-73	74-75	76-77	78-79	80-81	82-83	84-85	86-87	88-89	90-91	92-93	
Chinese.....	22			2	5	2	3	3	3	3	1								70.86
Japanese.....	29			2	1	2	3	4	6	2	3	6							74.28
Indian.....	106	2	1	5	12	14	17	11	24	16	2		2						71.63
‡ Black.....	301	2	5	3	19	27	38	47	48	38	33	20	7	8	5			1	74.05
Number measured.....	458	4	6	12	37	45	61	65	81	59	39	26	9	8	5			1	73.35
Not measured.....	9																		
Total.....	467																		

Race.	Total.	Knee height, in centimeters.												Mean knee height.
		38-39	40-41	42-43	44-45	46-47	48-49	50-51	52-53	54-55	56-57	58-59	60-61	
Chinese.....	20		1	5	2	5	3	3	1					
Japanese.....	20			4	4	12	4	4						
Indian.....	102		4	11	26	11	20	11	7					
‡ Black.....	289	2	7	16	48	61	72	42	13	4	2	1		
Number measured.....	420	4	12	36	83	89	99	60	23	8	5	2		
Not measured.....	47													
Total.....	467													

20. GENERAL COMPARISON OF OTHER COLOR RACES.

Unfortunately the numbers of Indians, Chinese, and Japanese measured were so small that the value of the comparison of the measurement for them with whites and Negroes is much reduced (Table 105). Nevertheless, some results are fairly clear. Of all three races the Indians are the tallest and the Japanese the shortest, but the height of Indians averaged less than that of the white or Negro troops. In average weight and build the Indians exceeded any other race. Next to the whites the Japanese have the lowest index of build. The sitting height of the Indians exceeded that of any of the color races except white, despite the fact that their stature is inferior to that of the Negroes. Their leg length is less than that of whites, Negroes, and Japanese, but greater than that of the Chinese. The shoulder width is greater than that of whites and less than that of Negro troops. The pelvic width of the Indians is greater than that of any of the other races, except the Chinese. On the whole, the 106 Indians measured resembled, in their proportions, more the 22 Chinese measured than any other race.

A comparison of the *relative* dimensions of the color races (Table 106) offers points of interest. The Negro troops have the stockiest build, the Indians come next, and the whites are last. The white troops have the relative largest sitting height (trunk, head, and neck), the Indians about the same, and the Negro troops least. The relative height of the sternal notch is greatest in the Negro troops and least in white troops. Pelvic height also is greatest in Negro troops and less in Indians and Chinese than in the whites; the whites are intermediate in leg length. The white troops have the relatively shortest span and the Negro troops have the longest. The relative shoulder width is greatest in Negro and Chinese troops and least in white. The relative transverse pelvic diameter is least in the Negro troops and greatest in the Chinese. The chest of the Negro troops is more elliptical on cross section than that of the whites.

TABLE 108.—*Comparative measurements at demobilization, Civil and World Wars.*

WHITE TROOPS.			
Measurements.	Number of men measured.	Mean.	Remarks. ^b
Height (demobilization, 1919).....	98,598	67.72	
Sitting height:			
Gould ^a	10,876	36.08	Sitting height is made up of head and neck 9.94 + body length 26.16; pages 253 and 255.
Demobilization, 1919.....	98,239	35.59	
Pubic height:			
Gould ^a	1,061	33.27	Sailors; pages 290 and 291.
Demobilization, 1919.....	91,365	34.18	
Leg length:			
Gould ^a	10,876	28.49	Leg length is the difference between the total leg length 31.06, and the thickness of the foot, 2.57; pages 257 and 274.
Gluteal fold to apex internal malleolus (demobilization, 1919).....	76,141	28.22	
Knee height:			
Gould ^a	10,848	18.61	Knee height; page 258.
Demobilization, 1919.....	76,141	18.54	
Chest circumference:			
Gould ^a	10,874	34.49	Chest circumference; page 263.
Demobilization, 1919.....	95,867	34.96	
Neck circumference:			
Gould ^a	9,300	13.63	Neck circumference; page 260.
Demobilization, 1919.....	95,271	14.16	
Waist circumference:			
Gould ^a	10,876	31.47	Waist circumference; page 266.
Demobilization, 1919.....	95,157	30.66	
Weight:			
Gould ^a	10,757	141.38	Weight; Table III, page 403.
Draft, 1917-1918.....	873,159	141.54	
Demobilization, 1919.....	79,706	144.67	
Height:			
Volunteer recruits (Gould) ^a	1,104,841	67.64	Height, white and colored; Table VI, page 105.
Draft, 1917-1918.....	873,159	67.49	
Baxter.....	501,068	67.30	Height, white and colored draft recruits; Baxter, Volume I, page 23.
NEGRO TROOPS.			
Height (demobilization, 1919).....	6,441	67.70	
Sitting height:			
Gould ^a	6,441	34.11	Sitting height is made up of head and neck 9.92 + body length 24.49; page 299.
Demobilization, 1919.....	6,443	34.30	
Pubic height:			
Gould ^a	2,020	34.30	Pubic height; pages 299 and 300.
Demobilization, 1919.....	6,220	35.21	
Leg length:			
Gould ^a	2,020	29.43	Leg length is the difference between the total leg length, 32.10, and the thickness of the foot, 2.57; Table V, pages 303 and 305.
Gluteal fold to apex internal malleolus (demobilization, 1919).....	5,595	29.28	
Knee height:			
Gould ^a	2,020	19.14	Knee height; Table V, page 314.
Demobilization, 1919.....	5,725	18.61	
Chest circumference:			
Gould ^a	2,020	34.28	Chest circumference; Table V, page 304.
Demobilization, 1919.....	6,355	34.64	
Neck circumference:			
Gould ^a	2,020	13.92	Neck circumference; Table V, page 304.
Demobilization, 1919.....	6,280	14.32	
Waist circumference:			
Gould ^a	2,020	30.30	Waist circumference; Table V, page 304.
Demobilization, 1919.....	6,445	30.64	
Weight:			
Gould ^a	2,001	144.58	Weight; Table I, page 402.
Demobilization, 1919.....	3,319	149.53	

^a Demobilization, 1865 (Gould, 1869).^b Except where specified the references are to Gould.

21. COMPARISON OF THE SOMATIC PROPORTIONS IN THE EIGHT EUROPEAN RACES.

(a) *General discussion.*—The number of races in the United States of which representatives were measured at demobilization is very great. Provision was made in coding for some 78 countries and subdivisions of the populations of countries. But when the final results were tabulated it was found that there

were only eight of the European nations native-born representatives of which were included in our statistics in sufficient frequency to make the analysis worth while. These races are:

TABLE 109.—*Approximate number of men measured in 8 European races, demobilization, 1919.*

Race.	Approximate num- of men measured.	Race.	Approximate num- of men measured.
English.....	4,204	French.....	1,457
Scotch.....	2,074	Italian.....	3,519
Irish.....	6,164	Polish.....	2,408
German.....	7,077	Hebrew.....	1,692

For the above races the principal dimensions as given in Table 110 were drawn up.

TABLE 110.—*Absolute dimensions in 8 European races, demobilization, 1919.*

Dimension.	English.	Scotch.	Irish.	German.	French.	Italian.	Polish.	Hebrew.
Number men measured.....	4,204	2,074	6,164	7,077	1,457	3,519	2,408	1,692
Height.....cm.	172.08	172.54	171.36	172.04	168.59	165.18	169.41	166.91
Weight.....lbs.	144.98	144.93	142.98	148.20	142.16	127.99	145.62	137.85
Index of build.....	31.59	31.41	31.41	32.31	32.37	32.63	32.73	31.93
Sitting height.....cm.	90.63	90.75	90.46	90.36	89.47	87.76	89.42	88.06
Span.....cm.	175.61	175.60	174.10	176.66	172.85	169.19	174.60	170.30
Sternal notch.....cm.	140.87	141.53	142.28	141.19	137.88	135.37	139.15	136.93
Pubic height.....cm.	87.19	87.30	86.55	86.63	85.90	82.81	85.27	83.94
Knee height.....cm.	47.74	47.83	46.59	47.22	46.83	45.13	46.69	45.57
Leg length.....cm.	71.34	71.68	70.91	71.47	69.22	67.84	70.16	68.93
Chest circumference.....cm.	88.18	88.57	88.67	89.52	88.49	88.87	90.42	87.53
Transverse chest.....cm.	28.87	29.01	28.77	29.12	28.58	28.76	29.22	28.26
Antero-posterior chest.....cm.	21.45	21.58	21.60	21.79	21.39	21.48	21.90	21.42
Shoulder width.....cm.	41.69	41.70	41.43	42.19	40.41	41.64	42.24	41.42
Pelvic width.....cm.	29.28	29.38	28.92	29.80	28.70	28.62	29.55	28.34
Waist circumference.....cm.	76.69	77.53	77.70	78.46	77.32	77.16	78.38	76.71
Thigh circumference.....cm.	52.38	52.36	52.27	53.19	51.98	52.03	52.46	52.18
Calf circumference.....cm.	33.90	34.04	33.83	34.40	33.68	33.71	34.44	33.66
Flaxen hair.....per 1,000.	55.05	52.81	37.80	68.49	27.19	6.02	75.77	16.01
Light brown hair.....per 1,000.	235.70	228.85	188.54	306.08	138.77	59.06	333.47	110.31
Clear red hair.....per 1,000.	13.82	20.05	25.42	6.79	7.67	1.72	7.08	8.90
Clear blue eyes.....per 1,000.	441.57	477.29	533.70	426.14	342.90	111.59	468.50	230.80

(b) *Stature*.—Table 25 gives the proportional distribution of different classes of stature. In order of mean stature the Scotch stand first, 172.54 centimeters. They are followed by the English, 172.08; German, 172.04; Irish, 171.36; Polish, 169.41; French, 168.59; Hebrews, 166.91; and Italians, 165.18. The standard deviation in stature is least in the Italians (probably because they are shortest) and greatest in the English, indicating a great admixture of race statures in that people. Other high standard deviations are: German, 6.61; French, 6.50. Next to the Italians in limited stature variability stand the Polish with a standard deviation of 6.12, and the Hebrews with 6.20. The Irish have a standard deviation of 6.31, and the Scotch of 6.39.

Corresponding to their tall stature, we find among the Scotch a larger proportion of men of stature class 172–173 centimeters than among any other people. Indeed, this constitutes the modal class for the Scotch. For the English 170–171 is the modal class and the same holds for the German, Irish, Polish, and French. For Hebrews and Italians, however, the modal class is 164–165 centimeters. Using the English system of measures, the average

stature of the Scotch is about 68 inches (172.54 centimeters), of the Italians 65 inches (165.18 centimeters).

(c) *Weight*.—Table 37 gives the distribution of absolute weights and the relative proportion of the different weight classes for the eight European races.

From Table 110 we learn that though the Scotch have the tallest average stature they have not the greatest average weight. This greatest average weight is found in the Germans, 148.20 pounds; second in order come the Polish, 145.62 pounds; then the English and Scotch, respectively, 144.98 and 144.93 pounds; the Irish, 142.96 pounds; French, 142.16 pounds; and at the bottom of the list the Hebrews and Italians, respectively, 137.85 and 137.99 pounds.

In variability of weight the Scotch (standard deviation 17.41) stand at the top, followed by the English, Irish, and Germans. The Polish stand at the bottom of the list (standard deviation 15.29), with Italians, French, and Hebrews above.

(d) *Index of build*.—The index of build of the eight races is obtained by dividing the mean weight by the square of the mean stature.

TABLE 111.—*Index of build in eight European races, obtained by dividing weight by stature and by the square of stature, demobilization, 1919.*

Race.	Weight× 1,000.	Weight.
	Stature ² .	Stature.
Polish.....	32.73	2.183
Italian.....	32.63	2.122
German.....	32.31	2.188
French.....	32.37	2.142
Hebrew.....	31.93	2.096
English.....	31.59	2.140
Scotch.....	31.41	2.134
Irish.....	31.41	2.119

Table 111 gives the index of build, using both the first and second powers of the statures as divisors. The races are arranged in order of the quotient of weight divided by stature squared. It appears from this calculation that the Polish have the largest index of build, followed in order by the Italians, Germans, French, Hebrew, English, Scotch, and Irish. If it is contended that the larger races are given an unduly small index of build, due to the squaring of the stature, a comparison may be made of the weight divided by the first power of the stature, in which the order of robustness is German, Polish, French, English, Scotch, Italian, Irish, and Hebrew. From other evidence it seems probable that the first series is the more significant.

(e) *Summary*.—Without calling attention in detail to all the striking results shown in Tables 110 and 111, it may be said in summary that in absolute dimensions of the five groups—Scotch, English, Irish, German, and Polish—the Scotch lead in stature, sitting height, pubic height, knee height, and leg length—thus in vertical dimensions. The English are usually second in these respects. The Poles are first in horizontal dimensions—in index of build, chest circumference, shoulder width, and calf circumference. The Germans are first in absolute weight and second in the horizontal dimensions, but not in index of build.

Of the four groups—Polish, French, Hebrew, and Italian—the Poles are in a class by themselves in absolute dimensions. The French are next, being first in sitting height, pubic and knee heights, and second in the other dimensions. The Italian group stands first in nothing; they are mostly inferior in absolute vertical dimensions to the Hebrews but exceed them in horizontal ones, such as index of build and dimensions of chest, shoulder width, and calf circumference.

TABLE 112.—*Relative dimensions in eight European races, demobilization, 1919.*^a
[Per cent rates.]





Dimensions.	English.	Scotch.	Irish.	German.	French.	Italian.	Polish.	Hebrew.
Weight, in pounds } Stature (inches) }	31.59	31.41	31.41	32.31	32.27	32.63	32.73	31.92
Sitting height } Stature }	52.67	52.60	52.79	52.52	53.07	53.13	52.78	52.76
Sternal notch } Stature }	81.86	82.08	83.03	82.07	81.78	81.95	82.14	82.04
Pubic height } Stature }	50.67	50.60	50.51	50.35	50.89	50.14	50.33	50.29
Leg length } Stature }	41.46	41.54	41.38	41.54	41.06	41.07	41.41	41.30
Knee height } Stature }	27.74	27.72	27.19	27.45	27.78	27.32	27.56	27.30
Span } Stature }	102.10	101.80	101.60	102.70	102.60	102.40	103.10	102.00
Shoulder width } Stature }	24.23	24.17	24.18	24.52	23.97	25.21	24.93	24.82
Chest circumference } Weight (in pounds) }	60.82	61.11	62.02	60.40	62.25	64.40	62.09	63.50
Transverse chest } Antero-posterior chest }	134.59	134.43	133.19	133.64	133.61	133.99	133.49	131.88
Antero-posterior chest } Stature (pounds) }	12.47	12.51	12.61	12.67	12.69	13.00	12.93	12.83
Waist circumference } Stature (pounds) }	44.57	44.93	45.34	45.61	45.86	46.71	46.27	45.96
Pelvic diameter } Stature }	17.02	17.03	16.88	17.32	17.02	17.33	17.44	16.98
Thigh circumference } Stature }	30.44	30.35	30.50	30.92	30.83	31.50	30.97	31.26
Calf circumference } Stature }	19.70	19.73	19.74	20.00	19.98	20.41	20.33	20.17

^a Unless specified all measurements are in centimeters.





In relative dimensions (Table 112) in the five groups—Scotch, English, Irish, Germans, and Polish—it appears that the Polish stand first in all relative horizontal dimensions and the Germans second. In many of such dimensions the English stand last, as the most slender, although the Scotch and Irish are close competitors for this place. Of relative vertical dimensions the English stand first in relative pubic and knee height, the Irish in relative sitting height and sternal notch, and the Scotch and German in leg length. On the other hand, the Irish stand at the bottom in leg dimensions, and the Germans at or near the bottom in relative sitting and pubic heights. In build the Poles are first and the Scotch and Irish last. The Irish chest is most nearly cylindrical (infantile) and the English flattest.

In the four groups—Polish, French, Hebrew, and Italian—the Italians are first in relative horizontal dimensions, while the Poles are frequently last. In relative vertical dimension the Poles are first in sternal height, while the French exceed in relative pubic height and knee height. The Hebrews are last in relative sitting height and knee height. The chest rotundity decreases from the Hebrews, through Polish, and French to Italians, who are in this group the most like the English in this respect.





AVERAGE DIMENSIONS, EIGHT RACES, DEMOBILIZATION-1919 **MEAS. IN CMS.; WT. IN LBS.**



AVERAGE DIMENSIONS, EIGHT RACES, DEMOBILIZATION-1919 MEAS. IN CENTIMETERS




AVERAGE DIMENSIONS, EIGHT RACES, DEMOBILIZATION-1919 MEAS. IN CENTIMETERS




AVERAGE DIMENSIONS, EIGHT RACES, DEMOBLIZATION-1919 MEAS. IN CENTIMETERS


1-SHOULDER WIDTH




2-PELVIC DIAMETER, TRANS.







3-CHEST DIAMETER, TRANS.







4-CHEST, ANTERO-POSTERIOR







RELATIVE DIMENSIONS, EIGHT RACES, DEMOBILIZATION - 1919 PERCENTAGE RATES BASED ON MEAS. IN CMS.



RELATIVE DIMENSIONS, EIGHT RACES, DEMOBILIZATION-1919
PERCENTAGE RATES BASED ON MEAS. IN CMS.



RELATIVE DIMENSIONS, EIGHT RACES, DEMOBILIZATION-1919 **PERCENTAGE RATES BASED ON MEAS. IN CMS.**



E. CORRELATIONS BETWEEN MEASUREMENTS.

1. CORRELATIONS BETWEEN MEASUREMENTS FOR WHITE AND NEGRO TROOPS.

(a) *General description of tables.*—From the foregoing sections it is clear that height, weight, and chest circumference are not independent, but, on the contrary, closely interdependent measurements. In order to understand the law of their associations, it is necessary to apply correlation tables. Such correlation tables are given in Tables I, II, and III. Table I shows the correlation between stature and weight. It answers the question, How were the weights of men of the stature of 59 inches distributed; how those of the stature of 60, 61, and 62 inches, etc.? The distribution of weights for men of different stature is given by reading in horizontal lines across the table. The table also gives the relation between the different statures of men for a given weight. It answers the question, What proportion of men weighing 105–109 pounds are 60 inches, 61, 62 inches, etc., tall? The distribution of statures among men of a given weight is given by reading down in the vertical columns. It will be observed that the entries become larger in the middle of the table; this is because men of medium stature and medium weight are much commoner than those of extreme stature or those of extreme weight. Thus, in Table I the largest entry is 18,930, which means that that number of men out of 868,445 had a stature about 67 inches and 135–139, inclusive, pounds. This combination was then the commonest one among the early recruits into the United States Army.

Table I answers the question, What proportions of men of a given stature, such as 69 inches, fall into each of the different classes of weight? These proportions per 1,000 are obtained by reading along the horizontal lines. Table I answers the question, What proportions of men of different classes of weight, as for instance 145–149 pounds, fall into each of the different classes of stature? The answer is given again by reading along the horizontal lines. It must be remembered in applying these tables that the frequencies in the extreme classes fail to give a good picture of the distribution of weights and statures in that part of the population. This is because there was a selection against men of under 63 inches, and this selection was especially marked in the case of men under 60 inches tall. A few short men were taken, provided they had an exceptionally fine physique, were especially robust, and had a relatively high weight. Consequently we actually find a larger proportion of men of 59 inches with a weight of 125–129 pounds than we do of men of 60 inches. Similarly for selected men of 60 inches, the most frequent weight was between 115–119 pounds, which is the same as the most frequent weight for men of 61 or even 62 inches. This shows that even for men of 60 inches a disproportionately large number of lightweighters were rejected. On the other hand, few men were accepted who were 76 inches tall. Some such were indeed accepted if they were not obese, so it occurs that the proportion of men weighing 115–119 pounds actually increases as the stature increases from 74 to 79 inches. This

shows that as the stature increased there was a tendency to reject a disproportionately larger number of heavy men. The same thing is shown in the men of the weight of 120–124 pounds. Between the limits of 62 inches and 75 inches and 100 and 199 pounds, inclusive, the table represents, however, nearly the conditions found in the general population.

(b) *Correlations between stature and weight.*—By means of a mathematical treatment proposed by Francis Galton and elaborated by Karl Pearson, it is possible to find a single numerical expression for the correlation between pairs of dimensions related like stature and weight. By applying the proper mathematical formula it is determined that the correlation of stature to weight (using the entire Table I) is 0.4810. This may be interpreted as indicating that as there is an increase of one stature class, there tends to be an increase of about 0.48, on the average, of the weight class. If the correlation were perfect, any one height class would be accompanied by only one correlated weight class, but it is clear that this is not the case, that the weights of men of successive classes are very variable, and, as weight increases with the increase of stature, that there is a tendency for the individuals to mass themselves around a central point in the table. For English undergraduates a corresponding correlation has been found of the value 0.49 (Pearson, 1899).

The coefficient of correlation, 0.48, is a fairly high one, as correlations go. The relation between breadth and length in a collection of German skulls has been found to be 0.49. The relation between capacity and breadth of German skulls has been placed at 0.67. The relation of length of radius and stature is about 0.70, whereas that for arm and stature is only 0.37, and clavicle and scapula, 0.12 to 0.16.

Table I gives for each class of stature the mean weight. This table is of interest in comparison with the statistics obtained by Gould² at the end of the Civil War and given in his Table IX (p. 408). This Table IX is indeed for white men only, whereas our Table I includes about 6 per cent colored. In Gould's table half inches are tabulated as well as whole inches, and in order to make comparison with our table it is necessary to combine the half inch with the following full inch in his table. It has been done in our Table 113, which shows in parallel columns the average weight of men of a given height, 1865 and 1917–1918.

TABLE 113.—*Comparative weight of men of different statures among white soldiers of 1865 at demobilization and white and colored soldiers at demobilization, 1919.*

Height, in inches.	Mean weight, in pounds.		Height, in inches.	Mean weight, in pounds.	
	White soldiers at demobilization, 1865. ^a	White and colored at demobilization, 1919. ^b		White soldiers at demobilization, 1865. ^a	White and colored at demobilization, 1919. ^b
60.....	111.79	123.00	68.....	144.93	145.52
61.....	117.59	125.66	69.....	149.04	149.39
62.....	120.77	127.10	70.....	153.19	153.30
63.....	122.95	129.78	71.....	158.21	155.91
64.....	128.43	131.84	72.....	162.47	159.84
65.....	132.12	135.20	73.....	165.40	164.03
66.....	136.06	139.26	74.....	168.98	165.54
67.....	140.77	142.71	75.....	170.39	168.00

^a Calculated from Gould,² p. 408.

^b Calculated from Table I.

Comparing the two series, with due allowance for the latter including about 6 per cent negro troops, it appears that, on the average, men of 70 inches and under in stature are heavier among the demobilized soldiers of 1919, but men over 70 inches are lighter than they were in 1866. This indicates that there has been a change in our population through the addition of short stout men and tall lank men. There has indeed clearly been an addition to our population of short thick-set persons from southeastern Europe and from French Canada, and our tall population (including the mountain whites and many of the tall men of the Southern States) has become lanker, through the inclusion of a larger percentage of this lank type in the 1917 data than in the 1866 data.

We may seek a check on this conclusion by comparing our measurements of the draft boards on recruits of 1917-1918 with figures for 6,359 American-born white drafted men accepted for military service by recruiting officers during the draft period of the Civil War, as given in Baxter¹ (Vol. II, pages 300-315). (See Tables 11 and 12, p. 74.)

Table 12, based on Baxter's, indicates that for Civil War drafted recruits, white native Americans, the commonest combination was 120-139 pounds weight and 65-67 inches of stature, and this combination was found in 171 per 1,000 men. Of World War recruits the commonest combination is 120-139 pounds and 67-69 inches stature, and this combination was found in 134 per 1,000 men.

A comparison of Table 12 with Table 11, giving the statistics for the recruits of 1917-1918, all colors and nativities and in the same form as for 6,359 American-born whites, Civil War tables, shows that there was a considerably larger proportion of men 69-73 inches and over in the recruits of 1917-1918. Assuming the basis of selection to have remained the same, then it would appear that there is in the population now a smaller proportion of very short men and a larger proportion of very tall men as compared with 50 years ago. However, the comparability of these tables is very limited, since the one for the Civil War includes only native-born white Americans and the other includes all colors and nativities drafted in 1917-1918. Also the army of the Civil War contained many boys of 18 years or under.

There is shown in the tables a considerable decrease of men of small weight, under 140 pounds, and an increase of men of large weight, over 140 pounds. The tables as they stand indicate an increase of short thick-set men and tall slender men. The former group is doubtless made up largely of recent immigrants from southeastern Europe, who are excluded in the Civil War table. The latter is largely due to the inclusion in our statistics, of the tall lank type from the Southern States who were to a large extent also excluded in the Civil War table. Whether this type is racial or due to other causes does not concern us now. It is largely through the inclusion of these men from the Southern States that there is a larger percentage of high statures among the recruits of 1917-1918 than among those of the Civil War. At least this influence has been added to that of the immigration of Scandinavians. As already stated, the value of this comparison is very limited, since Baxter's figures are for draft recruits, American-born recruits, and those for 1917-1918 are for the mixed population.

It is very difficult to answer the question whether the physique of our young men has changed in the last 50 years. Indeed, the question thus unqualified has little meaning. Had the racial constitution of the population remained constant, that is, had there been no heavy immigration, then the question would have more meaning; but in view of the tremendous immigration, amounting in some years to nearly a million persons, the physical changes of the racial constitution of our stock have been so great as to mask entirely any slight alteration that may have occurred in the physique of the stock of 50 years ago, either through improvement or deterioration of environmental or economic conditions.

TABLE 114.—*Correlation between stature and chest circumference, Civil War recruits (Baxter,¹ Vol. II, p. 166).*

Stature, in inches.	Chest circumference (expiration), in centimeters.						Total.
	Under 29.	29-30.9	31-32.9	33-34.9	35-36.9	37 and over.	
Under 61.....	1.391	2.257	2.060	1.203	0.443	0.124	7.478
61-62.....	4.365	12.657	13.739	7.797	2.475	.555	41.588
63-64.....	6.512	30.083	50.107	38.755	13.425	2.892	141.774
65-66.....	4.530	32.409	85.775	91.375	40.017	9.506	263.612
67-68.....	1.852	19.105	77.520	111.183	61.263	17.760	288.683
69-70.....	.559	7.033	36.219	68.695	47.688	17.012	177.206
71-72.....	.150	1.569	10.083	23.119	20.562	9.027	64.490
73 and over.....	.018	.261	1.708	4.834	5.225	3.127	15.173
Total.....	19.377	105.404	277.161	346.961	191.098	60.003	1,000.004

Total strength, 501,068 drafted men, includes substitutes and late volunteers of all nationalities.

TABLE 115.—*Correlation between stature and chest circumference, recruits, 1917-1918 (per 1,000).*

Stature, in inches.	Chest circumference, in centimeters.					Total.
	29-30	31-32	33-34	35-36	Over 37.	
Under 61.....	0.990	2.339	2.288	0.971	0.292	6.890
61-62.....	4.611	10.772	8.016	2.768	.655	26.822
63-64.....	13.532	36.977	31.306	11.358	3.177	96.350
65-66.....	23.029	77.475	79.057	32.424	9.329	221.314
67-68.....	21.203	91.602	113.965	53.286	16.468	296.524
69-70.....	10.224	57.215	89.970	49.593	16.328	223.330
71-72.....	2.838	20.099	40.106	26.010	9.591	98.644
73 and over.....	.515	4.351	11.536	9.612	4.117	30.131
Total.....	76.942	300.830	375.244	186.022	59.957	999.995

Total strength, 873,159 drafted men of all nationalities.

(c) *Correlation between stature and chest circumference.*—Table II gives the correlation of stature to chest circumference; that is, it shows the absolute number of men who were 59 inches tall and who belonged to each of the respective classes of chest circumference from 29 to 39 inches, and the same for each class of stature from 59 to 79 inches. The ratios per 1,000 for each of the separate statures is given in Table II, and similarly the distribution per 1,000 of each of the separate chest measurements of the different statures is also given in Table II. Table II shows that in 873,159 men measured the commonest combination among drafted men was 68 inches stature and 33 inches chest circumference.

From Table II it appears that there was actually a larger proportion of men that were 59 inches tall who had a chest circumference of 33 inches or over than of men that were 60 inches tall, and correspondingly of the men 60 inches tall there was a larger proportion with a chest circumference 33 inches and over than there was of men 61 and 62 inches tall. The reason for this is that there was a selection for Army purposes of the stoutest men of small stature. The men of short stature who had a chest circumference of only 30 inches were largely eliminated. For men of 62 inches stature and above, the effects of this selection is no longer obvious. Similarly in the case of men over 75 inches tall we find the chest circumference not increasing with the stature. Indeed, the chest circumference in Table II tends slightly to decline in the case of the very tall men. This is apparently due to the selective elimination from military service of the very heavy men among the tall men who were examined. The question arises whether there has been a change in physique of men of military age since the Civil War. Some light is thrown on the subject by a comparison of Table 114 and Table 115. Table 114 gives the per mille distribution of the different combinations of stature and chest circumference classes from 501,068 men of all nationalities, draft recruits for the Civil War, taken from a population already depleted by volunteer enlistments. Table 115 gives similarly the per mille distribution of the combination of stature and chest circumference classes for 873,159 recruits for the World War. Assuming that no men under 29 inches chest circumference were accepted for the World War, we may compare the remaining classes of chest circumference with each stature class in the two tables. The most frequent combination in both the Civil War and the World War is 67-68 inches stature and 33-34 inches chest circumference. This group contains 111 per 1,000 in the Civil War recruits and 114 per 1,000 in the World War recruits. Taking the men with the commonest chest circumference, 33-34 inches, it appears that in the World War there was a larger proportion of the tall statured men of this chest circumference, indicating the larger proportion of slender men. For the 31-32 inches we find similarly a larger proportion of tall slender men. Taking the group 35-36 inches, we find again an excess of the taller men. It is indeed only in the group of short slender men^a that we find a deficiency in the World War recruits. There were proportionately more of the tall slender men in the World War than in the Civil War. This result accords with what has been found already and doubtless is due to the fact that the draft for the World War covered the Southern States, the home of tall slender men, whereas these were naturally not included in the recruits for the Civil War, Federal Army.

Another matter of importance that comes from a consideration of Table II is the coefficient of correlation. This is found to be 0.2304, about half the correlation that exists between stature and weight. This shows that the relation between stature and weight is twice as close as that between stature and chest circumference. In other words, men of a given stature are less variable in respect to their weight measurements than in respect to their chest circumference.

^a The common type of very young men found in the Civil War statistics.

(d) *Correlation between weight and chest circumference.*—Table III gives the correlation between weight and chest circumference. The coefficient of correlation is found to be 0.6907, which is a much higher correlation than between stature and weight and stature and chest circumference. This is in accordance with common experience, namely, that chest measurement varies closely with weight in a given stature;—the heavier the man the greater his chest circumference. This table shows that the commonest chest circumference is 33 inches and the commonest weight 135–139 pounds. The most frequent combination of chest circumference and weight is 33 chest and 135–139 weight. This then corresponds to the condition of the typical man of military age. Since the most frequent height is 68 inches, the most frequent combination of these three dimensions found in recruits of military age is the following: Stature, 68 inches; weight, 135–139 pounds; chest, 33 inches. The corresponding average measurements are: Stature, 67.49 inches; chest circumference (expiration), 33.22, and weight 141.54 pounds.

Table III (B) gives the ratio per 1,000 of the different weights to chest measurement, and Table III (C) gives the ratio per 1,000 of the separate chest measurements to each weight class.

(e) *Correlation between stature and waist circumference.*—Table LXXV gives the correlation between stature and waist circumference for 103,410 white and colored troops consolidated. The stature groups range from 148 to 205 centimeters, the mode being in the class 170–171 centimeters, a class which contains 12 per cent of all cases. The average stature is 171.99 centimeters, with a standard deviation of 6.68 ± 0.01 .

The waist circumference ranges from 50 to 104 centimeters and over, the modal class being 76–77 centimeters, and the mean waist circumference 77.84 centimeters, with a standard deviation of 5.91 ± 0.01 . The correlation between stature and waist circumference is 0.1923 ± 0.0019 . This is not a high correlation, such as is found, in a symmetrical figure, on the two sides of the sagittal plane. It is well known that persons who are very tall are large in all dimensions; still, there are so many short persons that are stout and so many tall persons that are thin, as measured by the waist circumference, that the first obvious relation is obscured by the second one.

From the table we see that the commonest relation of stature and waist circumference is that of stature of 170–171 centimeters, and a waist circumference of 76–77 centimeters. This condition is found in about 19 per 1,000 of the men measured.

2. CORRELATION BETWEEN MEASUREMENTS FOR WHITE TROOPS (DEMOBILIZATION).

(a) *Correlation between chest circumference and transverse diameter of pelvis between cristæ.*—Table LXXIX shows the correlation between chest circumference and transverse diameter of pelvis between cristæ. The table shows that the modal diameter of the pelvis is 29 centimeters, a class that includes about 16 per cent of all. The commonest combination of chest circumference and transverse diameter of the pelvis is: 86–89 centimeters chest circumference

and 29 centimeters pelvic diameter, giving a combination found in nearly 6 per cent of recruits. The mean diameter of the pelvis for the white troops is 29.45; the standard deviation of this dimension is 2.90. The correlation is $.3073 \pm 0.0021$.

(b) *Correlation between stature and sitting height.*—Table LXXXIII shows the correlation between total stature and sitting height for white troops. As has been pointed out earlier, sitting height is usually about 53 per cent of the total stature. In Table LXXXIII it appears that the commonest sitting height is 90–91 centimeters, while the commonest stature is 170–171 centimeters; the sitting height here also is about 53 per cent of the stature. The average sitting height is 90.39, with a standard deviation of 3.51; the mean stature in this table is 171.99 centimeters, with a standard deviation of 6.66. The range in the relation of sitting height to stature is, however, great, as indicated in the table. Thus there were 4 per 1,000 of the recruits with a stature of 162.5 centimeters, and sitting height of 90.5 centimeters. For these the relative sitting height is 55.6 per cent of the total stature. One per 1,000 of the men had a stature of 184.5 and a sitting height of 90.5; here the relative sitting height is 49.1 per cent of the total stature. Similarly, of men 170.5 centimeters in stature, there were 0.6 per 1,000 who had a sitting height of 80.5; thus their relative sitting height was 47.2 per cent of the stature. Again, 0.5 per 1,000, with a sitting height of 98.5 and a total stature of 170.5, had a relative sitting height of 57.8 per cent of the total stature. The correlation between stature and sitting height is found to be 0.6626 ± 0.0012 , a high correlation, as was to be expected, since sitting height is a segment of total stature.

(c) *Correlation between stature and height of sternal notch.*—Table LXXXV shows the correlation between stature and height of sternal notch from the floor (in centimeters). The commonest height of the sternal notch is 140–141 centimeters, and the mean height of sternal notch is 141.18; standard deviation, 5.91 centimeters. The table shows for each of the different statures the absolute distribution and the frequency of different heights of sternal notch.

Since the height of the sternal notch is an important element of the total stature, it is to be expected that there is a close relation between the two dimensions. The coefficient of correlation is calculated from Table LXXXV as 0.8567 ± 0.0006 , a very high correlation. The ratio of height of sternal notch to total stature is as 141.18 : 171.99, or 82.09, or about five-sixths of the total stature.

(d) *Correlation between stature and height of pubic arch.*—Table LXXXVI gives the correlation between stature and height of pubic arch for white troops. The modal height of pubic arch is 86–87 centimeters, a group which contains about 15 per cent of all. The average height of pubic arch is 86.82 centimeters; standard deviation, 5.05 centimeters. It will be observed from this table that the relation of mean pubic height to mean stature is as 86.82 : 172.02, or 50.47 per cent. Thus we see that in this series height of pubic arch is almost precisely one-half total stature.

Since height of pubic arch constitutes about one-half of the total stature, it is to be expected that the correlation between the two would be fairly high. It proves to be 0.6960, or over two-thirds, naturally less than the correlation

between sternal height and stature, because sternal height is a larger component of total stature.

(e) *Correlation between stature and span.*—Table LXXXIV gives the correlation between stature and span for white troops. The modal span is seen to fall in the class 174–175 centimeters, which contains about 10 per cent of all men measured. The ratio of mean span to mean stature is as 175.58 : 171.99, or 102.09 per cent. Thus mean span is seen to be slightly greater than mean stature. There is, however, a good deal of variation in this respect. Thus in the case of men with a span of 168.5 centimeters the most frequent stature is 166.5 centimeters, giving 101.2 per cent. However, there is at one extreme a number of men of this same span who have a height of only 154.5, giving a ratio of 109.06. In this group the span is 9 per cent greater than stature. On the other hand, in six cases the stature of men with a span of 168.5 was 186.5, giving a ratio of 90.35 per cent. In this case the span is about 10 per cent less than the stature.

The correlation between stature and span is 0.7944, a high correlation, as a glance at the correlation surface shows must be the case. For English fathers the correlation between these two dimensions was found by Pearson (1903) to be 0.783; for the sons of such fathers, 0.802. Our result is almost intermediate between Pearson's two figures.

(f) *Correlation between chest circumference and weight.*—Table LXXVII gives the correlation between chest circumference in centimeters and weight in pounds for white troops. The commonest combination is seen to be a chest circumference of 86–89 centimeters, and a weight of 140–149 pounds. This class contains about 10 per cent of all men measured. As chest circumference and weight are more or less independent measures, it is not to be expected that the correlation between them will be very high, but it proves to be 0.6598 ± 0.0013 . This is a fairly high correlation and indicates that the development of muscles and the deposition of fat upon the chest go hand in hand with increasing weight, so that the two are closely interdependent. It will be noted that this correlation is distinctly less than that found (p. 426) in the case of recruits

(g) *Correlation between chest circumference and neck circumference.*—Table LXXVIII gives the correlation between chest circumference and neck circumference in white troops. The modal class for neck circumference is seen to be 36 centimeters, for chest circumference 86–89 centimeters. The mean neck circumference is 35.98; standard deviation 1.80 centimeters. The mean chest circumference is 88.79; standard deviation 5.18. Thus, in this group the neck circumference is to chest circumference as 35.98:88.79, or 40.52 per cent. That is, the neck circumference is about 40 per cent or two-fifths of the chest circumference. The correlation between these two dimensions is 0.5061 ± 0.0016 . This fairly high correlation indicates that the same developmental factors that determine a robust trunk also determine to a considerable extent a large neck. Since chest circumference is so closely correlated with weight, it is probable that the neck circumference is also somewhat correlated with weight, though the actual correlation was not calculated.

(h) *Correlation between transverse and antero-posterior diameters of the chest.*—Table LXXX gives the correlation between transverse and antero-posterior chest diameters in white troops. The modal class for transverse chest diameter is 28–29 centimeters, and for antero-posterior 20–21 centimeters. The average transverse diameter of the chest is 29.02; standard deviation 2.40. The mean antero-posterior chest diameter is 21.58; standard deviation 1.87. Thus, the antero-posterior diameter is to the transverse diameter as 21.58:29.02, or 74.36 per cent. In other words, the antero-posterior diameter is, on the average, about three-fourths the transverse diameter of the chest. There is, however, a good deal of variation in this regard. Thus the transverse diameter of the chest is seen to range from 18 to 49 centimeters, the larger diameter being 2.5 times the smaller diameter. Since the larger chest circumference is more than twice the smaller chest circumference, this great variation in transverse chest diameter indicates that the length of the axes of the chest is very much more variable than the total circumference. The capacity of the chest is much more constant than its form.

The table gives the range of antero-posterior diameter as extending from 14 to about 40 centimeters. Here we see that the largest class of antero-posterior diameter is 2.7 times the smaller antero-posterior diameter. Thus the range is somewhat greater in per cent than the variability in the transverse chest diameter. The variability of the transverse diameter is, however, seen to be somewhat greater than that of the antero-posterior diameter, as 2.40:1.87. This is, however, largely because the transverse diameter is a greater dimension than the antero-posterior diameter. The coefficient of variation, which is obtained by dividing the standard deviation by the mean, is for the transverse diameter of the chest 8.27 per cent and for the antero-posterior diameter 8.67 per cent. Thus taking into account the differences in mean dimension, the antero-posterior diameter is more variable than the transverse. This will be easily understood by those who have measured a large number of men. Even among those accepted, there are many cases of chicken-breasted individuals with prominent sternum, greatly increasing the antero-posterior diameter. The correlation between the two diameters is relatively small, 0.2714. This small correlation is no doubt the resultant of two factors, one which tends to keep the shape of the thorax constant and the other which tends to preserve a fairly constant volume, at least for men of a given size. The correlation of the first set of factors is positive, of the latter negative; that is to say, a long transverse diameter would be correlated with a relatively shorter antero-posterior diameter.

(i) *Correlation between chest circumference and transverse pelvic diameter.*—Table LXXIX shows the correlation between chest circumference and breadth of the pelvis (between cristæ) for white troops. Of chest circumference, the modal class is seen to be 86–89 centimeters, the mean 88.78, and standard deviation 5.17. Of pelvic diameter the modal class is 29 centimeters, and the mean 29.45, and standard deviation 2.90. The relation of mean chest circumference to mean pelvic diameter is thus 88.79:29.43, or 33.14 per cent. Thus for white troops the pelvic diameter is almost exactly one-third of the chest girth, while it is 38 per cent of waist girth, indicating again the fact that chest

girth exceeds waist girth in these veterans. The correlation between these dimensions is 0.3073 ± 0.0021 as compared with 0.3510 ± 0.0019 for waist and pelvis. This suggests that pelvic diameter has a slightly closer relation with waist girth on the one hand than with chest girth on the other; doubtless due to the closer proximity of the two dimensions.

(j) *Correlation between waist circumference and transverse pelvic diameter.*—Table LXXXI shows the correlation between waist circumference and transverse diameter of the pelvis (between cristæ) for white troops. The modal class of waist circumference is seen to be 76–79 centimeters; the mean is 77.87; standard deviation 6.08. The modal class for transverse pelvic diameter is 29 centimeters, mean transverse pelvic diameter 29.43; standard deviation 2.85. The relation of mean pelvic diameter to mean waist circumference is thus seen to be 37.8 per cent. This relation, however, is less significant than the relation between the transverse pelvic diameter and the transverse diameter of the chest. This is as 29.43: 29.02, or 101.41. That is to say, on the average, the transverse pelvic diameter is about 1.4 per cent greater than the transverse chest diameter. The correlation between the above two dimensions is 0.3510 ± 0.0019 . This correlation is to be expected, since both dimensions depend upon the form of the trunk which constitutes roughly a cylinder of which the diameter as well as the length varies. However, the fact that the coefficient of correlation deviates so far from unity proves that the capacity of the chest and the transverse diameter of the pelvis are to a considerable extent independently variable, and this is understandable in view of the comparative rigidity of the pelvis and the great elasticity of the chest. For the chest is capable of very great extension and development in such training as was given to military men.

(k) *Correlation between arm length and forearm.*—Table LXXXII gives the correlation between total arm length (a measurement which extends from the spines of the vertebral column along the outside of the flexed arm to the styloid process at the wrist) and the forearm (or the distance from the olecranon process at the elbow to the styloid process). Thus the forearm is a part of the total "arm length" measurement.

The modal class of arm length is 78–79 centimeters; the average arm length is 78.42; standard deviation 4.69 centimeters. The modal class of forearm length is 27 centimeters; the mean forearm is 26.91; standard deviation 1.73. Thus the forearm measurement constitutes 34.32 per cent of the total "arm length," or slightly more than one-third. Of the total arm length measurement, then, about two-thirds is the distance from the elbow to the vertebral column. The average transverse diameter of the chest is 29.02, half the chest diameter is 14.51. Subtracting the sum of half the mean transverse chest diameter and mean length of the forearm ($14.51 + 26.91 = 41.42$) from the total arm length, we get 37.16 centimeters as the length of the upper arm. This makes the relation of the length of the forearm to the length of the upper arm as 26.91: 37.16, or 72.42 per cent. Calling the total "arm length" 100, then the relative length of the segments to be assigned to the half chest diameter, upper arm and forearm as far as the styloid process, are 18.46, 47.29, and 34.25, or very roughly 1, 3, and 2, respectively.

Since the forearm is part of the measurement of arm length, a high correlation between the two parts is to be expected. This is found to be 0.5837, which is a fairly high correlation. That it is not higher is no doubt due to the fact that it is the resultant of two independently working factors, one which influences the arm as a whole and all its parts and tends to create a positive correlation, and the other which, with constant arm length, tends to alter the relative position at which the division between fore and upper arm shall occur. This tends toward a negative correlation.

(l) *Correlation between leg length and knee height.*—Table LXXVI gives the correlation between the length of the leg and the height of the knee for white troops. As indicated elsewhere, the length of the leg is measured from the gluteal fold (which is the posterior continuation of the perineum and marks approximately the lower end of the sitting height dimension) to the apex of the internal malleolus. The knee height, on the contrary, is measured from the floor to the top of the patella. Thus the knee height is included in part in the leg length, but is not completely included in it.

The modal class of leg length is 70–71 centimeters, the mean leg length is 71.69; standard deviation 4.71. The modal class of knee height is 46–47 centimeters. The mean knee height is 47.08; standard deviation 3.62 centimeters. The mean leg length is 71.69 centimeters; standard deviation, 4.71 centimeters. Thus the leg length is seen to be more variable than the knee height, which, however, is to be expected, owing to its greater length. If we divide the two standard deviations by the mean length of the corresponding parts, we get a coefficient of variation for leg length of 6.57 per cent and a coefficient of variation for knee height of 7.69. That is to say, knee height is a relatively more variable dimension than the leg length. This suggests that in addition to the variation in the knee height, correlated with variations in the leg length and the size of the body as a whole, there is also a variation in the knee height (assuming the leg length constant) due to the fact of variation in the relative position of the knee, which is sometimes at a relatively higher sometimes at a relatively lower point on the leg.

The correlation between knee height and leg length is 0.4178, a fairly high correlation, because the knee height is a part of leg length. That it is not larger is due to the fact, as pointed out above, that the knee height is not entirely included in the leg length. Variation in the relation of knee height to leg length is considerable. Thus with a constant leg length of 70.5 centimeters, we have on the one hand a knee height of 38.5 centimeters, and on the other of 58.5 centimeters. In the first case the ratio of knee height to leg length is 54.61 per cent, in the second 82.98 per cent. Adding 8.5 centimeters to the mean leg length to give the height of the internal malleolus from the floor, we have a mean leg length of 80.19. Using this as a divisor, we have a ratio for the short knee height of 48.01 per cent and for the longer height of 72.95 per cent. That is, in the shorter knee height the lower leg is less than half of the total leg length; in the greater knee height it approaches three-fourths of the total leg length. In such cases, then, the thigh would constitute only about one-fourth of the total leg length.

If one subtracts from the average knee height 8.5 centimeters, being the average distance from the internal malleolus to the sole of the foot, then the average height of the lower part, of the leg is 38.6 centimeters, which, divided by the leg length (71.69), gives 53.84 per cent as the average relation of the lower leg to total leg length. This is a relatively high proportion as compared with the dimensions given in Martin⁵ (pp. 314-315), where at the age of 13 years in the male the "Unterschenkel" is about 42 per cent; in the case of adult Chinese 42.7 per cent. The high per cent of leg length found in our table is no doubt partly due to the circumstance that the measurement was made to the top of the patella, whereas in Martin's measurement, it was made only to the head of the tibia, which is located about 5 centimeters below the top of the patella. Subtracting these 5 centimeters + 8.5 (the height of the internal malleolus), or 13.5 altogether, from the mean knee height, we have 33.6 remaining, which, divided by 71.69, gives 46.87 per cent. Even this gives a relatively long lower leg, due, again, to the fact that our divisor "leg length and foot" is still too short, being height of gluteal fold instead of height of trochanter or iliospinale. For trochanter leg length about 5 centimeters has to be added to our "leg length and foot," which gives a relative knee height of 43.8 per cent.

(m) *Correlation between leg length and waist circumference.*—Table CXV shows the correlation between waist circumference and leg length for white troops. This is the basal table used in forming the breeches groups for uniforms. The modal class of waist circumference is 76-79 centimeters. The mean is 77.87; standard deviation 6.08 centimeters. The modal class of leg length is 70-71 centimeters; mean leg length 71.44 centimeters. This mean leg length is clearly to be preferred to that obtained from Table LXXXVI, which is based on 20,000 fewer measurements. The coefficient of correlation between waist circumference and leg length is 0.1591 ± 0.0021 , a low correlation but positive, indicating that, through the operation of factors that influence the size of the body as a whole, on the average, men with larger waist circumference have longer legs. That the correlation is so low is due in large part to the fact that shorter men are, on the average, more robust (have relatively larger waist and chests) than taller (longer-legged) men.

3. CORRELATION BETWEEN MEASUREMENTS.—NEGRO TROOPS.

In the following paragraphs the correlations are given between various pairs of dimensions for Negro troops. The numbers are unfortunately small, under 6,500, but the means and correlations obtained from them are doubtless significant for comparison with white troops.

(a) *Correlation between stature and sitting height.*—Table LXXXVII gives the correlation between stature and sitting height for 6,433 colored troops. The modal class of sitting height is 86-87 centimeters, the mean sitting height is 87.35; standard deviation 3.48. The mean stature is 171.99; standard deviation 6.90 centimeters. The relation of mean sitting height to stature is 50.79 per cent. Considering only the classes which contain more than 10 individuals, the range of relative sitting height for men of stature 170-173 is from 46.7

per cent to 53.6 per cent. The coefficient of correlation between stature and sitting height is 0.6088.

(b) *Correlation between stature and height of sternal notch.*—Table LXXXIX gives the correlation between stature and height of sternal notch in 6,454 colored troops. The modal class for sternal notch is 142–143 centimeters; the average is 142.39; standard deviation 6.05. The relation of height of sternal notch to mean stature is 82.8 per cent. If the standard deviation of the mean stature (in this table, 6.91 centimeters) is somewhat more variable than the height of sternal notch, it may be because of the greater number of units involved in mean stature. Dividing each standard deviation by the mean in order to secure the coefficient of variation, we find that this is for the mean stature 4.25 per cent, and for sternal notch 4.01 per cent. Thus, the height of the sternal notch proves to be also a relatively less variable dimension. The coefficient of correlation between these two dimensions is 0.8582.

(c) *Correlation between stature and height of pubic arch.*—Table XC gives the correlation between stature and height of pubic arch in the case of 6,220 colored troops. The modal class of pubic height is 90–91 centimeters, the mean pubic height is 89.42; standard deviation 5.27. The relation of mean height of pubic arch to mean stature is 52.02 per cent. The variability in this respect is considerable. Thus the men with a stature of 172–173 centimeters have a relative pubic height ranging (if we include only the more frequent classes) from 46.67 to 55.94. The coefficient of correlation is 0.6948.

(d) *Correlation between stature and knee height.*—Table XCI shows the correlation between stature and knee height for 5,725 colored troops. The modal class of knee height is 46–47 centimeters. The average is 47.26; standard deviation 3.64. Mean height is 172.05; standard deviation 6.90. The average knee height constitutes 27.47 per cent. The coefficient of correlation between the two dimensions is 0.4763.

(e) *Correlation between stature and span.*—Table LXXXVIII gives the correlation between stature and span in the case of 6,441 colored troops. The modal class of span is 182–183 centimeters; the average span is 180.76; standard deviation 8.59. The relation of span to height is 105.16 per cent. The range in this respect is seen to be considerable. Thus of men with an average stature of 170.5 we have some with a span of 168.5, or 98.83 per cent. At the other extreme we have men with a span of 190.5, or 1.118 times the stature. The coefficient of correlation between the two dimensions is 0.7292; less than in whites.

(f) *Correlation between chest circumference and weight.*—Table XCIII gives the correlation between chest circumference and weight for 3,319 colored troops. The number is small because in one of the camps, for a period, the colored men were not weighed. The modal class of weight is 140–149 pounds and modal chest circumference 86–89 centimeters. The mean weight is 149.53; standard deviation 17.53 pounds. The mean chest circumference is 88.14; standard deviation 4.79 centimeters. The range of weight is from 100–200 pounds and over. Of the 3,319 men, 23 weigh 200 pounds or over, or 6.93 per 1,000. The chest circumference ranges from around 70 to over 105 centimeters, the largest number being 50 per cent greater than the smallest. The correlation between chest circumference and weight is 0.6559 ± 0.0067 , a high correlation

because, as pointed out in another connection, the chest circumference varies directly with weight since extra weight is apt to be laid down on muscles and fatty tissues of the chest. The correlation is the same as in whites.

(g) *Correlation between chest circumference and sitting height.*—Table CVII gives the correlation between chest circumference and sitting height in the case of 6,355 colored troops. The modal class for sitting height is 86–87 centimeters. Mean sitting height is 87.35; standard deviation 3.43. The modal class for chest circumference is 86–87 centimeters; mean chest circumference 87.99; standard deviation 4.76. We see here a very close relation between chest circumference and sitting height, the ratio of the one to the other being as 1.007:1. The range in chest circumference, even excluding the extreme classes with fewer than 5, is very great, from 70 centimeters to 105, or an increase of 150 per cent. For men with a sitting height of 86–87 centimeters there is a range of classes containing 10 or more from 76–77 to 98–99 centimeters. In the slenderest group this gives a ratio of chest circumference to sitting height of 88.44 per cent; for the stoutest men the ratio is 113.87 per cent. The correlation between chest circumference and sitting height is 0.3012.

(h) *Correlation between chest circumference and neck circumference.*—Table XCIV gives the correlation between chest circumference and neck circumference for 6,280 colored troops. The neck circumference ranges from 29 to 44 centimeters, the modal class being 36 centimeters. The average neck circumference is 36.37; standard deviation 1.72. The mean chest circumference is 87.97; standard deviation 4.84. The relation of neck circumference to chest circumference is obtained by dividing the mean of the former by the mean of the latter, or 41.34 per cent. Taking the class of 83.5 chest circumference, we find the extremes of neck circumference having more than 5 in the class as follows: For the smallest neck circumference, 31 centimeters, or 37.15 per cent of chest circumference; for the largest neck circumference, 40 centimeters, or 47.90 per cent. The correlation between neck circumference and chest circumference is 0.5172 ± 0.0062 ; practically as in whites.

(i) *Correlation between transverse and antero-posterior diameters of the chest.*—Table XCVI gives the correlation between transverse and antero-posterior chest diameters in the case of 6,450 colored men. The antero-posterior diameter ranges from 14 to 35 centimeters, with a modal class at 20–21 centimeters. The mean antero-posterior diameter is 21.21; standard deviation 1.74. The transverse chest diameter ranges from 18 to 45 centimeters, with a modal class at 28–29 centimeters, and an average of 29.05; standard deviation 2.26. The antero-posterior diameter is, therefore, to the transverse as 21.21:29.05, or 73.01 per cent. For men of antero-posterior diameter of 20.5 centimeters, there is a considerable range of transverse diameter from 20.5 to 38.5 centimeters. In the narrowest chest, the relation of antero-posterior to transverse diameter is 100 per cent. In the broadest chest it is 53.25 per cent. The corresponding thoracic indices are 100 and 188.

The standard deviation of transverse diameter is greater than that of the antero-posterior, but this may be due to the greater average size of the transverse dimension. The coefficient of variability of the transverse diameter is 7.78; of antero-posterior diameter it is 8.20. This indicates that the antero-

posterior diameter is relatively the more variable. The coefficient of correlation between transverse and antero-posterior chest diameters is 0.2267.

(j) *Correlation between chest circumference and transverse diameter of pelvis.*—Table XCV gives the correlation between chest circumference and transverse diameter of pelvis in the 6,345 colored troops. The range of diameters of pelvis, including classes containing more than 10, is from 21 to 39 centimeters. The modal class is 28 centimeters, and the average diameter is 28.54; standard deviation 2.64. Taking the class of men averaging 87.5 centimeters chest circumference, including only the groups containing 10 or more, we find a range from 23 to 34 centimeters. The relation of mean transverse diameter of pelvis to mean chest circumference is 32.44 per cent. For the men of smallest pelvic diameter referred to above (23 centimeters) it is 26.29 per cent; for the men with greatest pelvic diameter (34 centimeters) it is 38.86 per cent.

More significant, perhaps, is the ratio of transverse diameter of pelvis to transverse chest diameter, 98.24 per cent. Thus the transverse diameter of the pelvis is slightly less than the transverse diameter of the chest. The correlation between chest circumference and transverse diameter of pelvis is 0.3297 ± 0.0075 .

(k) *Correlation between waist circumference and transverse diameter of pelvis.*—Table XCVII gives the correlation between waist circumference and transverse diameter of pelvis in 6,354 colored troops. The most frequent combination of measures is 76–79 waist circumference with 28 centimeters diameter of pelvis. The mean diameter of pelvis is for this group 28.42; standard deviation 2.35. The mean waist circumference is 77.82; standard deviation 5.71. The ratio of diameter of pelvis to waist circumference is thus 36.52 per cent—that is, the waist is relatively smaller with relation to the hips than the chest is. The standard deviation of the waist circumference is greater than that of the transverse diameter of the pelvis as 5.71:2.35. The coefficient of variation, however, is in the one dimension 7.40 per cent and the other 8.27 per cent. Thus, rather remarkably, the diameter of the pelvis seems to show a relatively greater variability than the circumference of the waist. (Note the greater variability of pelvic diameter and waist circumference in whites than in colored). The correlation between waist circumference and transverse diameter of pelvis is 0.4456 ± 0.0068 .

(l) *Correlation between arm length and forearm.*—Table XCVIII gives the correlation between arm length and forearm for 5,514 colored troops. The arm length, as will be remembered, is defined as the distance from the spines of the vertebral column to the styloid process. The forearm is from the elbow to the same process. The modal class for arm length is 80–81 centimeters; for forearm 28 centimeters. The average arm length is 80.79; standard deviation 4.76. The average length of the forearm is 28.20; standard deviation 2.03. The mean forearm is to the mean arm length as 28.20:80.79, or 34.91 per cent. Taking arm-length class 78.5, and considering only those classes which contain 5 or more individuals, the relatively shortest forearm is 24 centimeters, or 30.57 per cent; the longest forearm is 31 centimeters, or 39.48 per cent of "arm length."

The total arm length may be divided into three sections, including half the transverse diameter of the chest, upper arm and forearm. The average half

transverse diameter of chest is 14.53. If we add to this the mean forearm 28.20, there remains 38.06 for the approximate length of the upper arm. In relation to the total mean arm length of 80.79, these dimensions are, respectively, 17.98, 47.11, and 34.91 per cent. The correlation between arm length and forearm is 0.5782, a relatively high correlation, because one measurement is included in the other.

(*m*) *Correlation of leg length and knee height.*—Table XCII gives the correlation of leg length and knee height for 5,595 colored troops. Leg length has been defined as the distance from the gluteal fold to the internal malleolus, and knee height as the distance from the sole of the foot to the top of the patella. The two measurements therefore overlap and one is not wholly included in the other. The modal class of leg length is 74–75 centimeters, and that of knee height is 46–47. The average leg length is 74.38; standard deviation 4.59. The average mean knee height is 47.32; standard deviation 3.37. It is probable from the table that there are some adult males who have a smaller knee height than 38 centimeters and a greater knee height than 57 centimeters.

To compare the leg length and knee height, we may subtract from the knee height 8 centimeters, in order to get the length of the lower leg from the top of the patella down. As thus defined, the knee height from the top of the patella to the malleolus is 39.3. If we subtract further 6 centimeters for the distance from the top of the patella to the head of the tibia, we get 33.3 centimeters as the length of the lower part of the leg. This distance divided by the leg length gives the proportion of the lower leg to total length of leg as 44.77 per cent. The knee height as measured constitutes 63.62 per cent of the total leg length. The correlation between knee height and leg length is 0.4305.

4. COMPARISON OF CORRELATION BETWEEN WHITES AND NEGROES.


Tables 103, 104, and 116 give the comparative measurements and correlation of parts in the white and Negro troops. These tables show at a glance the means of the various dimensions, their standard deviations, and the correlation of certain pairs. We see, for example, that the stature of the Negro troops is more variable than of the white troops, but that the sitting height is 1 per cent less variable in the Negro than in the white. Similarly, the span is more variable in the Negro than in white troops, but the correlation between stature and span is less. This relation between size of standard deviation and correlation is to be expected, since the smaller the variability of each of two dimensions the greater the correlation is apt to be between them. Table 116 shows that the correlation between stature and height of sternal notch is about the same in the two races, slightly greater in the Negro than in the white. Between stature and height of pubic arch it is about the same in the two races. Between leg length and knee height the correlation is much greater in the Negro than in the white; between chest circumference and sitting height the correlation is markedly greater in Negro than in white troops; between transverse and antero-posterior chest diameters the correlation is much greater in the white than in the Negro. This is perhaps associated with the greater similarity in white troops than in Negro troops of the axes of the ellipse made by the cross section of the chest. The correlation between pelvic diameter and waist girth is greater in Negro than in white troops,

perhaps associated with the smaller pelvic diameter. The correlation between chest circumference and pelvic diameter is also greater in the Negro than in the white troops, perhaps associated with the smaller size of the latter dimensions in the Negro race. (See Plate XXIX, page 253.)

TABLE 116.—*Correlations, summary of white and colored troops, demobilization, 1919.*

Dimension.	Demobilization.			Mobilization, white and colored.	Taken from tables—		
	White.	Colored.	White and colored.		White.	Colored.	White and colored.
Stature and waist.....			0.1923±0.0019				LXXV
Stature and weight.....			.5198±.0017	0.4810±0.0006			LXXIV
Stature and chest.....				.2304±.0007			I
Weight and chest.....				.6907±.001			II
Stature and sitting height.....	0.6626±0.0012	0.6088±0.0053			LXXXIII	LXXXVII	III
Stature and sternal notch.....	.8567±.0006	.8582±.0022			LXXXV	LXXXIX	
Stature and pubic arch.....	.6960±.0012	.6948±.0044			LXXXVI	XC	
Stature and knee height.....		.4763±.0069				XCI	
Stature and span.....	.7944±.0006	.7292±.0034			LXXXIV	LXXXVIII	
Chest circumference and weight.....	.6598±.0013	.6559±.0067			LXXVII	XCIH	
Chest circumference and sitting height.....	.2415±.0021	.3012±.0077			XCLX	CVII	
Chest circumference and neck circumfer- ence.....	.5061±.0016	.5172±.0062			LXXVIII	XCIV	
Chest transverse and antero-posterior.....	.2714±.0020	.2267±.0080			LXXX	XCVI	
Chest circumference and pelvis, transverse.....	.3073±.0021	.3297±.0075			LXXIX	XCV	
Waist circumference and pelvis, transverse.....	.3510±.0019	.4456±.0068			LXXXI	XCVII	
Arm length and fore- arm.....	.5837±.0015	.5782±.0060			LXXXII	XCVIII	
Leg length and knee height.....	.4178±.0020	.4305±.0073			LXXVI	XCH	
Leg length and waist circumference.....	.1591±.0021				CXV		

CORRELATIONS, WHITE AND NEGRO TROOPS DEMobilIZATION - 1919



F. PATTERNS FOR UNIFORMS.

The measurements ordered by the War Department were for the purpose of securing patterns for uniforms. This purpose guided the set of measurements taken and has influenced the statistical treatment of the data secured. It is believed, however, that this fact will not diminish their importance for general anthropological purposes.

The uniform of the soldier consists of two more or less independent pieces, the blouse for the upper part of the body and the breeches for the lower part. The problem, therefore, is different from that of fitting a single suit—like a union suit—to the soldier, and the matter of precise length of trunk is of relatively less importance in uniforms than it would be for single-piece suits.

1. MEASUREMENTS FOR BLOUSES.

(a) *General discussion.*—Our first purpose, then, was to secure measurements which would serve first for making patterns for the blouse and secondly for making patterns for breeches. One limitation was prescribed by the office of the quartermaster, namely, that uniforms would not be made for any group which contained fewer than 5 per 1,000 men. Consequently, it became necessary to combine, for the purpose of this study, many of the smaller classes to fit the needs of the series. The construction of the blouse groups is shown in Tables XCIX and CVII, which give the correlation between chest circumference and sitting height. These two measures were taken as of primary importance in considering the blouse. The chest circumference is the primary basis of classification, and the length of the trunk, as measured by sitting height, is of secondary importance.

The correlation Tables XCIX and CVII were divided, as indicated in the tables, into 22 groups. The first included all chest circumferences under 78 centimeters. The last three groups included all chest circumferences of 102–105, 106–109, and 110–117, respectively. The last two groups, indeed, do not contain the prescribed 5 per 1,000. The division was made rather to meet anthropological interests. All of the other chest circumference groups were classes with a range of 4 centimeters. These groups are 78–81, 82–85, 86–89, 90–93, 94–97, 98–101. The division of each of these chest circumference classes was made so as to provide approximately 20 per cent in each of the extreme groups and 60 per cent in each of the median groups. The group with the shortest sitting height was designated by the initial "S," for short; that with median sitting height by "M," for median; and that with longest sitting height by "L," for long. The 22 groups thus constructed were called blouse groups, and their association with other dimensions was determined.

(b) *Chest circumference.*—Table XCIX gives for white troops approximately the frequency per 1,000 men of each of the different chest circumferences for

each sitting height. Thus of men of the sitting height of 86–87 centimeters there were 2 with a chest circumference of 68–69 centimeters; there were 5 with chest circumference 70–71; 21 with chest circumference 72–73 centimeters; and 34 with chest circumference 74–75 centimeters, etc. The modal chest circumference for men of this sitting height was 76–77 centimeters. Taking the distribution as a whole, we find that the commonest sitting height is 90–91 centimeters, the commonest chest circumference is 88–89 centimeters, and the commonest combination is that of 88–89 chest circumference and 90–91 sitting height. This group includes about 3.33 per cent of the individuals of the table. The central blouse group is that with chest circumference 86–89 centimeters and sitting height of 88–93 centimeters, and includes about 200 per 1,000, or 20 per cent. Since Table XCIX gives absolute numbers for 95,867 persons, the numbers have to be increased about 4.3 per cent to give exact ratios per 100,000.

(c) *Weight*.—Table C gives the association between the different blouse groups and the weight of the individual for white troops. Thus for 79,706 of such troops the total distribution is shown in the second column from the left of the table. The next column gives the findings for blouse group 1 with chest circumference of 68–77 centimeters, inclusive. The remaining columns give the absolute frequencies of the different weight classes for each blouse group; also the mean weight of men for each blouse group.

As the table shows, there are naturally more light-weight men associated with the small-size blouses and an excess of heavy-weight men associated with the large-size blouses.

Table 117 (p. 273) gives the average measurements of white men belonging to each of the 22 blouse groups. It is upon this table that the table of dimensions of manikins (Table 122, p. 276) is, in part, made up.

TABLE 117.—*Dimensions associated with the "blouse" groups, white troops, demobilization.*

(From Tables XCIX-CVI.)

Blouse group designation.	Average chest circumference.	Average sitting height.	Average stature.	Average sternal notch.	Average length of head and neck.	Average trunk height.	Average arm length.	Average neck circumference.	Average shoulder width.	Average chest transverse diameter.	Average chest anteroposterior diameter.	Average diameter pelvis (transverse diameter).	Average weight.	Rate per 1,000 of each group.
	Cm.	Cm.	Cm.	Cm.	Cm.	Cm.	Cm.	Cm.	Cm.	Cm.	Cm.	Cm.	Lbs.	
1.....	75.0	88.8	169.5	139.3	30.2	58.6	76.8	34.4	39.6	27.0	20.0	28.1	129	8.8
2.....	80.0	83.6	163.7	134.6	26.1	54.5	74.2	34.2	39.2	26.8	19.9	27.4	120	7.9
2 m.....	80.0	88.6	169.2	139.0	30.2	58.4	76.1	34.5	39.7	27.1	20.1	27.9	126	34.9
21.....	80.0	93.5	176.4	144.5	31.9	61.6	78.1	34.8	40.6	27.6	20.5	28.7	135	10.4
3.....	83.6	83.6	163.7	134.6	29.1	54.5	74.9	34.9	40.0	27.5	20.5	27.9	125	21.7
3 m.....	83.7	88.8	169.5	139.3	30.2	58.6	76.7	35.0	40.5	27.8	20.7	28.4	132	125.3
31.....	83.8	93.7	176.7	144.8	31.9	61.8	78.7	35.2	41.2	28.2	20.8	29.1	140	50.6
4.....	87.5	85.4	165.2	135.8	29.4	56.0	76.2	35.6	41.0	28.5	20.8	28.6	134	62.2
4 m.....	87.6	90.5	172.0	141.3	30.7	59.8	78.2	35.7	41.5	28.8	21.3	29.2	141	206.2
41.....	87.6	95.4	178.9	146.5	32.4	63.0	80.1	35.9	42.1	29.1	21.5	29.9	150	50.3
5.....	91.3	85.4	165.2	135.8	29.4	56.0	77.1	36.3	41.9	29.4	21.9	29.2	142	36.6
5 m.....	91.4	90.6	172.1	141.4	30.7	59.9	79.0	36.4	42.3	29.5	22.0	29.8	150	162.4
51.....	91.4	95.5	179.0	146.6	32.4	63.1	81.1	36.6	42.9	29.8	22.1	30.5	158	54.0
6.....	95.2	85.4	165.2	135.8	29.4	56.0	77.9	37.1	42.8	30.0	22.6	30.1	150	12.7
6 m.....	95.2	91.5	173.5	142.4	31.1	60.4	76.9	37.2	43.3	30.4	22.7	30.6	160	91.8
61.....	95.3	97.3	180.9	148.3	32.6	64.7	82.7	37.4	44.0	30.8	22.9	31.3	170	12.7
7.....	99.1	87.3	167.6	137.7	29.9	57.4	78.9	37.8	43.5	30.9	23.5	31.0	162	8.1
7 m.....	99.1	92.4	174.8	143.2	31.6	60.8	81.1	37.9	44.1	31.2	23.5	31.5	170	24.3
71.....	99.1	97.4	181.1	148.5	32.6	64.8	82.8	38.0	44.5	31.3	23.7	31.7	179	5.4
8.....	103.1	92.4	174.8	143.2	31.6	60.8	81.3	38.8	45.0	32.2	24.5	32.4	181	8.4
9.....	107.2	92.6	175.1	143.4	31.7	60.9	81.6	39.7	45.7	33.2	25.5	33.4	189	2.3
10.....	112.1	92.6	175.1	143.4	31.7	60.9	81.7	40.8	46.2	34.3	26.4	34.7	191	1.0

TABLE 118.—*Dimensions associated with the "blouse" groups, colored troops, demobilization.*

(From Tables CVII-CXIV.)

Blouse group designation.	Average chest circumference.	Average sitting height.	Average stature.	Average sternal notch.	Average length of head and neck.	Average trunk height.	Average arm length.	Average neck circumference.	Average shoulder width.	Average chest transverse diameter.	Average chest anteroposterior diameter.	Average diameter pelvis (transverse diameter).	Average weight.	Rate per 1,000 of each group.
	Cm.	Cm.	Cm.	Cm.	Cm.	Cm.	Cm.	Cm.	Cm.	Cm.	Cm.	Cm.	Lbs.	
1.....	71.4	86.5	170.9	141.7	29.2	57.3	79.0	35.4	41.7	27.7	20.6	27.1	135	3.2
2.....	75.9	84.9	168.6	139.7	28.9	56.0	76.3	34.4	40.4	26.4	19.4	26.7	127	9.0
3.....	80.0	81.6	165.1	136.9	28.2	53.4	76.1	34.5	40.4	26.9	19.8	26.8	126	14.3
3 m.....	79.9	86.1	170.3	141.1	29.2	56.9	77.9	34.9	41.0	27.3	20.0	27.1	131	41.0
31.....	80.1	91.6	177.5	146.7	30.8	60.8	79.2	35.2	41.3	27.5	20.4	28.0	141	7.1
4.....	83.7	81.7	165.2	137.0	28.2	53.5	77.5	35.3	41.3	27.8	20.4	27.1	130	42.2
4 m.....	83.8	86.4	170.8	141.6	29.2	57.2	79.1	35.5	41.8	28.1	20.4	27.7	136	147.1
41.....	83.9	91.6	177.5	146.7	30.8	60.8	80.3	35.7	42.3	28.5	20.7	28.4	146	35.9
5.....	87.3	81.6	165.1	136.9	28.2	53.4	78.6	36.1	42.2	28.5	21.0	28.0	140	40.8
5 m.....	87.5	87.3	171.9	142.6	29.3	58.0	80.4	36.3	42.7	29.0	21.1	28.4	147	276.6
51.....	87.7	93.3	179.3	148.4	30.9	62.4	82.2	36.4	43.3	29.1	21.3	29.2	154	30.5
6.....	91.8	83.0	168.4	138.1	28.3	54.7	80.2	36.9	43.5	29.7	21.7	28.7	151	27.5
6 m.....	91.3	87.9	172.6	143.2	29.4	58.5	82.0	37.0	43.7	29.6	21.7	29.2	157	172.9
61.....	91.4	92.5	179.5	148.6	30.9	62.6	83.4	37.1	44.0	29.9	21.8	29.6	163	32.3
7.....	95.6	84.5	168.0	139.2	28.8	55.7	81.8	37.6	44.3	30.4	22.2	29.6	164	17.3
7 m.....	95.1	88.9	174.0	144.0	30.0	58.9	83.2	37.9	44.7	30.4	22.4	29.8	168	53.5
71.....	95.2	93.6	179.6	148.7	30.9	62.7	84.1	38.0	45.4	30.7	22.6	30.1	176	22.5
8.....	99.6	84.7	168.2	139.4	28.9	55.8	81.2	37.8	44.3	31.6	23.3	30.4	165	3.3
8 m.....	99.0	89.6	178.0	144.8	30.2	59.4	84.5	38.9	45.7	31.5	23.2	30.8	182	14.8
81.....	99.1	95.4	181.6	150.3	31.3	64.1	86.2	38.7	46.3	31.9	23.1	31.1	180	3.5
9.....	104.2	90.4	176.4	145.8	30.6	59.8	83.5	39.4	47.6	32.6	24.9	32.9	193	4.9

2. MEASUREMENTS FOR BREECHES.

The primary classification of breeches is made on the circumference of the waist; the secondary division is length of leg. The method of taking these measurements has been already described (p. 57). In order to determine the number and limits of groups to which the breeches patterns should be cut, Table CXV was drawn up. This gives the different classes of waist circumference from 63 and under to 110 centimeters for white troops. Groups 1 and 2 were not subdivided, on account of small size. Group 9 remained undivided for the same reason, and the following three larger classes of waist circumference, containing few individuals, were grouped into one breeches group. On the other hand, waist circumference 68-71, 72-75, 76-79, 80-83, 84-87, 88-91, were each divided into three groups, short, median, and long, because of the number of men falling into these classes of waist circumference. This makes 22 classes of breeches groups. Table 121, derived from Tables CXV and CXXII, gives the relative frequency per 1,000 of each of the breeches groups for white and colored troops.

Tables 119 and 120 were prepared to give the association between the various breeches groups and dimensions of various parts of the body, for both white and colored troops. It is believed that these should be used in the making of uniforms. The more important anthropometric conclusions have been drawn from them in the earlier part of this book, under the respective parts.

TABLE 119.—*Dimensions (in centimeters) associated with the "breeches" group, white troops, demobilization.*

From Tables CXV-CXXI.]

Breeches group designation.	Average circumference of waist.	Average length of leg.	Average thigh circumference.	Average supra-patella circumference.	Average patella circumference.	Average calf circumference.	Average knee height.	Average transverse pelvic diameter.	Ratio per 1,000 of groups.
1.....	61	70.9	49.2	36.3	34.9	33.6	46.6	28.6	4.7
2.....	66	69.3	48.0	34.9	34.2	32.1	45.8	27.2	15.6
3 s.....	70	63.2	49.5	35.4	34.3	32.5	44.4	27.0	14.5
3 m.....	70	70.2	49.6	35.7	35.9	32.6	46.1	28.1	73.6
3 l.....	70	78.1	49.7	35.7	35.4	32.8	48.3	29.4	11.0
4 s.....	74	63.1	50.9	36.3	34.9	33.0	44.3	27.5	27.9
4 m.....	74	70.5	51.0	36.4	35.5	33.3	46.4	28.6	179.4
4 l.....	74	78.2	51.1	36.5	36.0	33.5	48.7	29.8	34.6
5 s.....	77	65.1	52.4	37.1	35.6	33.8	44.9	28.4	52.6
5 m.....	77	71.4	52.5	37.3	36.2	34.0	46.9	29.3	183.6
5 l.....	77	78.3	52.5	37.3	36.6	34.3	49.2	30.5	53.5
6 s.....	81	65.0	54.2	38.1	36.4	34.6	45.1	29.1	29.5
6 m.....	81	72.3	54.2	38.2	36.9	34.8	47.4	30.1	143.5
6 l.....	81	80.4	54.1	38.3	37.4	35.0	50.1	31.1	22.3
7 s.....	85	65.0	55.6	38.9	36.9	35.2	44.9	29.8	11.9
7 m.....	85	72.4	55.7	39.0	37.5	35.5	47.7	30.8	67.9
7 l.....	85	80.3	55.4	38.9	37.7	35.7	50.8	32.1	13.6
8 s.....	89	64.9	56.9	39.5	37.4	35.6	45.0	30.5	4.9
8 m.....	89	72.5	57.0	39.6	38.0	36.0	47.8	31.4	27.6
8 l.....	89	80.4	56.7	39.4	38.1	36.1	50.9	32.4	6.1
9.....	93	72.3	58.7	40.2	38.3	36.5	47.7	32.1	12.7
10.....	97	72.3	59.2	40.3	38.3	36.5	47.8	32.4	5.8
11.....	101	72.5	61.1	40.7	39.1	37.4	48.3	33.4	2.0
12.....	104	72.0	61.8	41.1	39.7	36.9	47.1	34.1	1.4

TABLE 120.—*Dimensions (in centimeters) associated with the "breeches" group, colored troops, demobilization.*

[From Tables CXXII-CXXVIII.]

Breeches group designation.	Average circumference of waist.	Average length of leg.	Average thigh circumference.	Average supra-patella circumference.	Average patella circumference.	Average calf circumference.	Average knee height.	Average transverse pelvic diameter.	Ratio per 1,000 of groups.
1.....	61	74.2	50.6	36.7	26.2	34.2	47.3	28.0	5.28
2.....	66	71.6	48.9	34.8	34.2	32.3	45.7	26.4	11.02
3s.....	70	65.2	49.9	35.5	34.3	32.5	43.3	25.9	11.48
3m.....	70	72.4	51.0	35.6	35.0	33.0	45.8	26.9	65.79
3l.....	70	80.3	51.0	35.8	35.4	33.0	48.4	27.4	13.65
4s.....	74	66.9	52.5	36.7	35.3	33.5	44.4	27.0	38.95
4m.....	74	73.5	52.4	36.5	35.7	33.8	46.4	27.5	158.26
4l.....	74	80.1	52.3	36.4	36.2	33.8	48.7	28.1	38.48
5s.....	77	66.8	53.9	37.3	35.8	34.3	45.3	27.4	40.96
5m.....	77	73.7	53.9	37.7	36.5	34.8	47.2	28.3	206.36
5l.....	77	80.5	53.8	37.5	36.9	34.8	49.7	28.8	76.96
6s.....	81	67.1	55.4	38.7	36.7	35.4	45.4	28.4	17.84
6m.....	81	74.6	55.5	38.6	37.3	35.5	47.8	29.3	147.25
6l.....	81	82.1	55.7	38.5	37.8	35.9	50.5	29.8	32.89
7s.....	85	67.5	57.1	39.1	37.4	36.2	45.4	29.4	8.07
7m.....	85	75.0	57.1	39.5	37.8	36.1	47.8	30.0	61.44
7l.....	85	82.3	57.4	39.5	38.0	36.3	50.6	30.3	13.65
8s.....	89	64.0	57.7	39.4	37.1	36.1	44.7	29.8	2.48
8m.....	89	74.5	58.7	40.5	38.7	36.7	47.8	30.4	22.81
8l.....	89	82.3	59.5	40.2	38.6	37.3	51.2	31.2	7.14
9.....	93	76.6	60.1	40.5	38.6	37.0	48.2	31.2	9.31
10.....	97	75.0	61.8	40.4	38.5	37.1	48.0	31.2	5.90
11.....	101	73.6	62.9	41.1	39.4	38.1	47.8	32.7	4.03

TABLE 121.—*"Blouse" and "breeches" groups, white and Negro troops.—Designation of each group, basic measurements adopted, and proportional number of each group of the total number of men measured at demobilization.*

"Blouse" groups.						"Breeches" groups.					
White. ^a			Colored. ^b			White. ^c			Colored. ^d		
Designation.	Chest circumference (rest).	Sitting height.	Proportional number of total men.	Designation.	Chest circumference (rest).	Sitting height.	Proportional number of total men.	Designation.	Waist circumference.	Leg length.	Proportional number of total men.
1.....	68-77	76-101	8.80	1.....	68-73	78-93	3.15	1.....	50-63	58-97	4.72
2s.....	78-81	78-85	7.91	2.....	74-77	80-91	8.97	2.....	64-67	50-99	15.60
2m.....	78-81	86-91	34.86	3s.....	78-81	78-83	14.32	3s.....	68-71	50-65	14.47
2l.....	78-81	92-101	10.43	3m.....	78-81	84-89	40.91	3m.....	68-71	66-75	73.57
3s.....	82-85	76-85	21.67	3l.....	78-81	90-99	7.08	3l.....	68-71	76-99	11.00
3m.....	82-85	86-91	125.34	4s.....	82-85	78-83	42.17	4s.....	72-75	50-65	27.91
3l.....	82-85	92-107	50.62	4m.....	82-85	84-99	147.13	4m.....	72-75	66-75	179.38
4s.....	86-89	76-87	62.24	4l.....	82-85	90-99	35.88	4l.....	72-75	76-101	34.56
4m.....	86-89	88-93	208.24	5s.....	86-89	78-83	40.76	5s.....	76-79	50-67	52.50
4l.....	86-89	94-107	50.23	5m.....	86-89	84-91	276.63	5m.....	76-79	68-75	183.55
5s.....	90-93	76-87	36.60	5l.....	86-89	92-99	30.53	5l.....	76-79	76-101	53.48
5m.....	90-93	88-93	182.42	6s.....	90-93	78-85	27.54	6s.....	80-83	50-67	29.52
5l.....	90-93	94-107	53.98	6m.....	90-93	84-91	172.94	6m.....	80-83	68-77	143.54
6s.....	94-97	76-87	12.72	6l.....	90-93	92-99	32.26	6l.....	80-83	78-101	22.26
6m.....	94-97	88-95	91.81	7s.....	94-97	78-87	17.31	7s.....	84-87	50-67	11.93
6l.....	94-97	96-107	12.74	7m.....	94-97	86-91	53.50	7m.....	84-87	68-77	67.94
7s.....	98-101	76-89	8.06	7l.....	94-97	92-99	22.50	7l.....	84-87	78-101	13.59
7m.....	98-101	90-95	24.33	8s.....	98-101	78-87	3.30	8s.....	88-91	50-67	4.89
7l.....	98-101	96-106	5.37	8m.....	98-101	86-93	14.79	8m.....	88-91	68-77	27.59
8.....	102-105	76-106	8.40	8l.....	98-101	94-99	3.46	8l.....	88-91	78-99	6.05
9.....	106-109	78-101	2.33	9.....	102-109	82-97	4.88	9.....	92-95	50-99	12.72
10.....	110-117	84-101	.86					10.....	96-99	64-85	5.07
								11.....	100-103	58-89	2.02
								12.....	104-109	58-89	1.38
			1,000.00				1,000.00				1,000.00
											1,000.00

^a Table XCIX.^b Table CVII.^c Table CXV.^d Table CXXII.^e 63 and under.

/ 100 and over.

3. DIMENSIONS OF MANIKINS.

The original orders authorizing the measurement of 100,000 soldiers provided for the construction of manikins from the measurements. Consequently Tables 117 to 120, inclusive, have been drawn up giving data for making such manikins. Tables 117 and 118 give the measurements for the upper part of the body, required for fitting blouses. Tables 119 and 120 give the measurements for the body, from the waist down, for fitting breeches.

Later it was desired to construct entire human figures, and these could not be obtained by piecing together the half figures of which the dimensions are given in the above tables. To construct these entire manikins a slight proportional adjustment had to be made in the "long" and "short" groups. The results are shown in Table 122, which is that of the dimensions of 21 complete manikins for white troops. Whether Tables 117 and 119 or Table 122 shall be used in the manufacture of uniforms and other clothing depends, curiously enough, on the esthetic choice between having the lower edges of the blouses (in case of men of the same stature but different trunk lengths) reach a common level from the floor or reach a common anatomical level (e. g., the trochanters) on the body. Those who regard the former as desirable will use the table of total manikins; those who prefer the latter will use the two tables for blouse groups and breeches groups, respectively.

TABLE 122.—*Dimensions of the 21 manikins (in centimeters), white troops.*

4. SIZES AND PROPORTIONS OF MEN IN THE DISTRIBUTION ZONES, Q. M. C.

One aim of the measurements of the 100,000 men was to secure manikins for the construction of patterns for uniforms. The second aim was to secure the proper distribution of sizes of uniforms to the different areas covered by the distribution zones of the Quartermaster Corps. Certain of these zones are

designated largely because of the storage capacity of certain large cities or other special relation to the quartermaster's activities. Such are the cities of Philadelphia (D. Z. 3), Baltimore (D. Z. 4), Jeffersonville (D. Z. 6), and the District of Columbia (D. Z. 15). In addition there are 10 distribution zones covering certain large sections of the country or groups of States. These zones may be defined by their included States as follows:

ZONE 1.	ZONE 5.	ZONE 8.	ZONE 12.
Maine. New Hampshire. Vermont. Massachusetts. Rhode Island.	North Carolina. South Carolina. Georgia. Florida. Alabama. Tennessee.	Kansas. Missouri. Oklahoma. Arkansas. Illinois, southern half.	New Mexico. Arizona.
ZONE 2.	ZONE 7.	ZONE 9.	ZONE 13.
Connecticut. New York. New Jersey. Pennsylvania.	West Virginia. Kentucky. Ohio. Indiana. Michigan. Wisconsin. Minnesota. Iowa. Illinois, northern half.	Mississippi. Louisiana.	Montana. Idaho. Nevada. Washington. Oregon. California.
ZONE 3.		ZONE 10.	ZONE 15.
Philadelphia.		Texas.	District of Columbia.
ZONE 4.		ZONE 11.	
Delaware. Maryland. Virginia.		North Dakota. South Dakota. Nebraska. Wyoming. Colorado. Utah.	

Table CXXXIV shows the distributions of frequencies of different statures for the different distribution zones, for a total of 102,061 men. This table also gives the proportional frequency of the different statures in each zone. Arranging the zones in order of average stature of the men, we have the following: Zone 10 (Texas), 174.23; zone 5 (Southern States from North Carolina to Alabama, including Tennessee), 173.90; zone 13 (Pacific Coast States, Nevada, Idaho, and Montana), 173.51; zone 8 (Missouri, Arkansas, Kansas, and Oklahoma), 173.48; zone 11 (North and South Dakota, Colorado, Wyoming, and Utah), 173.44; zone 9 (Mississippi and Louisiana), 173.33; zone 12 (New Mexico and Arizona), 172.73; zone 7 (Central States, including also West Virginia, Kentucky, Wisconsin, Minnesota, and Iowa), 172.06; zone 4 (Delaware, Maryland, Virginia), 171.88; zone 2 (Connecticut, New York, New Jersey, and Pennsylvania), 170.10; zone 1 (New England except Connecticut), 169.78.

Arranging the different zones in order of variability as measured by the standard deviation, we have the following: Zone 12 (Desert States), 6.686; zone 2 (Middle States), 6.622; zone 11 (the Dakotas and Mountain States), 6.612; zone 9 (Mississippi, Louisiana), 6.572; zone 4 (Delaware, Maryland, Virginia), 6.566; zone 7 (Central States), 6.500; zone 5 (Southeastern States), 6.484; zone 1 (New England, except Connecticut), 6.460; zone 13 (Pacific and Northwestern States), 6.412; zone 8 (Missouri, Arkansas, Kansas, Oklahoma), 6.356; zone 10 (Texas), 6.304. Thus it appears that, as other parts of the study have shown, Texas contains among the tallest men of the country and they prove to be the most homogeneous in stature. New England contains the

shortest men and they are fairly uniform in this respect. The greatest variability occurs in the Desert States of New Mexico and Arizona, where there is an admixture of Indians, Mexicans, and white Americans of European origin.

Table CXXXIV-B gives the proportional distribution of the different statures for each of the different zones. Thus for zones 1, 2, and 7 the modal stature is 170-171 centimeters; for zones 4, 5, 8, 9, 10, and 13 it is 172-173 centimeters; for zones 11 and 12 it is 174-175 centimeters. Thus Table CXXXIV-B tells the quartermaster what proportion out of every 1,000 suits of uniforms sent to the different zones should fit men of the respective statures.

Since, however, the blouses and breeches are separate garments, it is more important to know the proportion of men of different chest dimensions and waist dimensions, respectively, that occur in the different zones. The required information is given in Tables CXXXVI and CXXXVII. Table CXXXVI gives the absolute number of men found with the different chest circumferences in the different distribution zones. It also gives for each zone per 1,000 men the number having each of the classes of chest circumference. It shows also what proportion of sizes of each 1,000 blouses distributed should be sent to each of the distribution zones in order to meet the size requirements of men of these zones. Thus Table CXXXVI-B states that to zone 1 there should be distributed in every 1,000 blouses 285 of chest size 90-94, 382 of chest size 85-89, 189 of chest size 80-84. On the other hand, to zone 11 there should be sent 363 blouses of chest size 90-94, 324 of chest size 85-89, and only 124 of chest size 80-84. To zone 12 there should be sent only 8 blouses of size 100-104, whereas to zone 11, 23 per 1,000 blouses of size 100-104 should be sent. To zone 4 there should be sent 30 blouses per 1,000 of size 75-79, whereas to zone 11 there should be sent only 10 such.

Table CXXXVI-C states that in distributing 1,000 blouses of size 60-64, 512, or over half of them, should go to zone 2; 268, or over one-fourth, should go to zone 7, the remaining one-fourth should be distributed as indicated, but none at all should be sent to zones 9, 10, 12, and 13. Of 1,000 blouses of size 65-69, one-third of all should be sent to zone 7; 278, or over one-fourth, to zone 2; the remainder will be variously distributed as indicated, but only 1 or 2 should be sent to zones 4, 11, and 12. Of 1,000 blouses of size 75-79, 284 should be sent to zone 7; another one-fourth, precisely 265, should be sent to zone 2; 130 should be sent to zone 5; but only 9 should be sent to zone 11, and 3 to zone 12. Similarly the tables give the proper distribution for all of the different sizes of blouses.

The sizes of breeches are determined primarily by waist circumference. Distribution by waist circumference is shown in Table CXXXVII. This table gives the absolute frequency by zones of occurrence of the different waist circumference in the 101,576 men measured. The table indicates the proper proportion of the different sizes of breeches in a shipment of 1,000 to any zone. Thus, in a shipment of 1,000 breeches to zone 1, 4 should be of waist circumference 60-64, 60 of waist circumference 65-69, 283 of waist 70-74, 368 of waist 75-79, 185 of waist 80-84, 67 of waist 85-89, 22 of waist 90-94, 7 of waist 95-99, and 3 of waist 100-104. Similar data are given for each zone.

Table CXXXVII-C shows the proper distribution to the different zones of 1,000 breeches of different waist circumference sizes. Thus of 1,000 breeches of waist 60-64, 331, or about one-third of all, should be sent to zone 7; 309 to zone 2; 9 to zone 9, etc. It may be pointed out, however, that there is reason for thinking that the men measured may not constitute the real proportion of recruits drawn from the different zones. If the total number of men measured in the various zones be divided by the total number of men drafted from these different zones, as given in the report of the Provost Marshal General, there will be obtained for each zone the proportion of drafted men who were measured at demobilization.

Table LXXIII gives the distribution of different colored races measured in the various zones. This table, for many reasons made clear in the last sections, must not be taken as an actual relative frequency of the different colored races in these zones. It appears that the most colored men were measured from zone 5, including the Southeastern States. The next largest proportion is in zone 9, including Louisiana and Mississippi, although an equally large number was measured from zone 4. An attempt was made to distinguish the mulattoes, quadroons, and sambos, but it can not be hoped that this attempt succeeded. A large proportion of sambos, or three-fourths blacks, were measured from zone 9, Louisiana and Mississippi, and a smaller proportion from zone 5, the Southeastern States. On the other hand, more mulattoes and quadroons were measured from zone 5 than from zone 9.

The distribution of blouse and breeches groups for white and colored troops taken separately are shown in Tables CXXIX-CXXXII.

G. DISTRIBUTION OF EYE COLOR.

Eye color is of importance as a rough index of race. Thus the so-called Nordic race, which has its home in northwestern Europe, is characterized by clear blue eyes. Nearly all other peoples have brown eyes. Hybrids between blue and brown eyed people have light brown or blue eyes with brown spots. Table 130 shows that absolutely the largest number of clear blue eyes was observed from zone 7, but there were more eyes observed from this zone than from any other. There were fewest clear blue eyes from zone 12, but there were fewer eyes examined from this zone than from any other. The absolute numbers, therefore, are not very significant. More important is the proportion of different types found in the different zones.

Table 130-B gives also the proportion of different eye colors in the different zones. Taking the figures as they stand, it appears that the largest proportion of clear blue eyes is found in zone 13 (the Pacific and northern Rocky Mountain States). Next largest percentage is in zone 11, the central Rocky Mountain States, the Dakotas and Nebraska. Third comes zone 7 (42 per cent blue-eyed); this territory has a large proportion of Scandinavians. The smallest rate for clear blue eyes (15 per cent) is found in zone 5, which includes the Southeastern States with their large proportion of colored population. In this zone, moreover, there is an exceptionally large proportion (42 per cent) of persons found with blue eyes having brown spots. It seems possible that the proportion of blue eyes with brown spots found is due to special (and justifiable) precaution of the anthropologist in charge at Camp Gordon in warning his recorders to look for brown spots in apparently blue eyes. If we combine clear blue with blue with brown spots, then the proportion of such eyes in the whole population is about 62 per cent. In zone 13 it is 65 per cent; in zone 11, 70 per cent; in zone 7, 69 per cent; in zone 5, 57 per cent; in zone 4, 53 per cent, which is the lowest proportion of clear blue and blue with brown spots found in any zone. Of light brown eyes the highest rate as given is 45 per cent in zone 9, including Mississippi and Louisiana, of which the population is over one-third colored. Very high rates are found also in zone 5, the Southwest; zone 4, Virginia and Maryland; zone 10, Texas. Low rates are found in zone 11, the central Rocky Mountain States; and zone 7, the Central States, including Minnesota, Wisconsin, and Iowa. Of the dark-brown eyes, the largest rate is found in zone 12, Arizona and New Mexico, and this doubtless is due to the influence of the Indian race here. Next is zone 10, and next zone 8, where the Indian rate is high. Low rates are found in zone 5 of the Southeast, zone 1, New England, and zone 11, the central Rocky Mountain States.

(a) *Clear blue eyes.*—The significance of these results will be clearer from a study of Table CXXXVIII, which gives the proportion of eye color by States. Table 123 gives the distribution of clear blue eye color by States. The States are arranged in descending order of the proportion of clear blue eyes observed. At the top of the list stands Alaska, with a rate

of 54 per cent; next Wisconsin, also with about 54 per cent. This is the State in which in certain sections one-fourth of the inhabitants are Scandinavians. Next comes the State of Maine with 53 per cent; the largest foreign element in Maine is French Canadian, about 13 per cent in one section; otherwise the immigrants are chiefly English Canadians; there are few representatives of south-eastern Europe. Vermont stands next with 51 per cent. Since Maine and Vermont contain a large proportion of French Canadians, it seems probable that the proportion of blue eyes is high among them. Next stands Minnesota with a high Scandinavian population, and then comes Oregon with many Scandinavians and Germans. Massachusetts follows with 49 per cent clear blue eyes. This also has a large representation of French Canadians and Irish. Next comes Michigan and then the State of Washington, both with many representatives from northwestern Europe. At the bottom of the list stands Florida, with only 4 per cent of clear blue eyes among the population. This population includes Negroes, mulattoes, and a considerable number of Cubans and West Indians, some probably who have received their brown eye color from Negro stock. It is perhaps not strange that this State, with its dense Negro population and with its former Spanish blood and its proximity to Cuba, should be the darkest of all the States in respect to eye color. Next to the bottom stands Georgia, which is geographically adjacent to Florida. The numbers of Nevada may be excluded, since there are only two individuals under consideration. This is followed by Alabama, Tennessee, South Carolina, Louisiana, Kentucky, Missouri, North Carolina, and Mississippi, all but one Southern States. The proportion of clear blue eye is, therefore, smallest in those States which have a large proportion of Negro population. Consequently, in general terms, the proportion of clear blue eyes diminishes with latitude. This is to be explained on the ground that blue eye color rose in northern Europe, and that immigrants from northern Europe settled the northern parts of our country; and, also, that the percentage of the Negro population there is small (see Plate XXX, Fig. 7, p. 295).

TABLE 123.—*Absolute and relative numbers of veterans with clear blue eyes, by States of nativity in order of incidence, demobilization, 1919.*

State.	Number of cases.	Ratio.	State.	Number of cases.	Ratio.
Alaska.....	7	538.46	New York.....	3,845	416.13
Wisconsin.....	1,441	538.29	Arkansas.....	1,064	412.08
Maine.....	365	525.94	Colorado.....	93	409.69
Vermont.....	229	512.30	Pennsylvania.....	4,381	401.89
Minnesota.....	969	496.67	California.....	109	391.31
Oregon.....	529	494.39	Wyoming.....	31	387.50
Massachusetts.....	2,365	493.22	District of Columbia.....	87	376.62
Michigan.....	1,821	488.46	Texas.....	1,511	345.45
Washington.....	996	486.92	Maryland.....	387	338.88
Utah.....	51	485.71	Arizona.....	43	330.77
New Hampshire.....	201	485.51	Indiana.....	1,265	319.85
Idaho.....	77	469.51	Virginia.....	614	318.13
Connecticut.....	464	465.39	New Mexico.....	69	300.00
Illinois.....	3,112	463.92	Mississippi.....	582	276.88
Rhode Island.....	186	461.54	North Carolina.....	479	263.91
Montana.....	122	458.65	Missouri.....	651	228.66
North Dakota.....	158	441.34	Kentucky.....	565	192.56
Oklahoma.....	1,008	435.23	Louisiana.....	362	174.12
New Jersey.....	1,374	430.99	South Carolina.....	128	154.40
Nebraska.....	353	428.92	Tennessee.....	426	151.33
West Virginia.....	726	427.82	Alabama.....	246	127.33
Ohio.....	3,027	426.70	Nevada.....	2	111.11
Kansas.....	433	426.60	Georgia.....	330	96.97
South Dakota.....	177	425.48	Florida.....	97	94.73
Delaware.....	127	423.33			
Iowa.....	679	421.74	Total.....	38,354	374.69

(b) *Blue eyes with brown spots.*—The distribution of eye color "blue with brown spots" is given in Table 124. In some ways this affords a remarkable reversal of the order of the States shown in Table 123, for here such States as Tennessee, Kentucky, Missouri, Alabama, Florida, and Georgia, stand at the top of the list, constituting from 42 to 52 per cent of the population. It is impossible to say, however, how much of this large proportion of blue with brown spots found is due to special effort to find it on the part of the observers. The lowest proportion of blue with brown spots is found in certain of the New England States; in Rhode Island only 10 per cent; Massachusetts, 11 per cent; Vermont, 11 per cent; Maine, 13 per cent; New York and Connecticut follow with less than 14 per cent. The proportion of blue and brown spots found in Louisiana is small, 15 per cent, which may in part be accounted for by the fact that men from this State were observed at Camp Shelby, where another anthropologist was in charge, who was perhaps less careful to instruct his observers to note the presence of brown spots upon the blue iris. However, it must be admitted that the proportion of blueness of iris found in men from Louisiana is low and it seems probable that not only the colored population, but also the South French blood, which settled there, has had its influence in depressing the total amount of blue eye color found in that State.

TABLE 124.—*Absolute and relative numbers of veterans with blue eyes with brown spots, by States of nativity in order of incidence, demobilization, 1919.*

State.	Number of cases.	Ratio.	State.	Number of cases.	Ratio.
Tennessee.....	1,463	519.72	Wyoming.....	15	187.50
Kentucky.....	1,510	514.65	Wisconsin.....	474	177.06
Missouri.....	1,420	498.77	Oklahoma.....	408	176.16
Alabama.....	881	456.00	Virginia.....	339	175.65
Florida.....	443	432.62	Utah.....	18	171.42
Georgia.....	1,433	421.10	Michigan.....	626	167.92
Indiana.....	1,616	408.60	New Mexico.....	38	165.21
South Carolina.....	296	357.05	Pennsylvania.....	1,795	164.67
North Dakota.....	101	282.12	Washington.....	332	163.95
Iowa.....	451	280.12	Arkansas.....	423	163.82
Nevada.....	5	277.77	Montana.....	41	154.13
South Dakota.....	114	274.03	Alaska.....	2	153.85
Nebraska.....	218	264.88	Oregon.....	164	153.27
Minnesota.....	485	248.59	Louisiana.....	315	151.51
Arizona.....	32	246.15	New Jersey.....	477	149.62
Kansas.....	248	244.33	District of Columbia.....	34	147.19
Delaware.....	71	236.66	New Hampshire.....	59	142.51
California.....	108	223.60	Idaho.....	23	140.24
Mississippi.....	435	206.95	Connecticut.....	138	138.42
Texas.....	904	206.68	New York.....	1,247	134.95
Illinois.....	1,363	203.19	Maine.....	90	129.68
Colorado.....	46	202.64	Vermont.....	49	109.62
North Carolina.....	366	201.65	Massachusetts.....	521	108.66
West Virginia.....	335	197.41	Rhode Island.....	41	101.74
Maryland.....	222	194.40			
Ohio.....	1,336	188.33	Total.....	23,571	229.79

(c) *Brown eyes.*—Considering dark brown eye color, we find that Louisiana stands at the very head of the table with 48 per cent of her soldiers placed in that category; a relatively low proportion (19 per cent) from Louisiana were found with light brown eyes. In the table (126) of dark brown eyes, next to Louisiana, stand North Carolina, Virginia, District of Columbia, Georgia, Mississippi, Florida, and South Carolina; and here again the Southern States have an excess of dark brown eyes in the population, due to the colored race. The Southern States for the most part stand near the bottom of the list of

light brown eyes, although Louisiana has a median position, with a rate of 19 per cent. Of dark brown eyes, Maine shows the smallest rate, 8.6 per cent; Vermont slightly more, 9.2 per cent; Wisconsin, Idaho, Minnesota, all have less than 11 per cent. New York stands far above the average in the proportion of dark brown eyes found in the population; Pennsylvania is slightly below the average, and Illinois and Michigan are far below the average, with only 15 per cent.

TABLE 125.—*Absolute and relative numbers of veterans with light brown eyes, by States of nativity in order of incidence, demobilization, 1919.*

State.	Number of cases.	Ratio.	State.	Number of cases.	Ratio.
Nevada.....	7	388.89	Ohio.....	404	174.43
Idaho.....	46	280.49	Colorado.....	84	173.91
Wyoming.....	30	250.00	Connecticut.....	194	171.96
New Hampshire.....	101	243.98	Massachusetts.....	191	167.25
Montana.....	64	240.60	New Mexico.....	167	164.53
New Mexico.....	54	234.78	Maine.....	38	162.01
Maine.....	162	223.43	Utah.....	442	157.02
Utah.....	24	228.57	Vermont.....	251	155.90
Vermont.....	89	221.48	Pennsylvania.....	128	154.40
Pennsylvania.....	2,409	220.99	Rhode Island.....	63	151.44
Rhode Island.....	89	220.84	Massachusetts.....	124	150.67
Massachusetts.....	1,043	217.52	New Jersey.....	381	147.56
New Jersey.....	666	208.59	Colorado.....	140	143.51
Colorado.....	47	207.08	Ohio.....	140	142.56
Ohio.....	1,387	195.52	Michigan.....	974	141.82
Michigan.....	728	193.27	Connecticut.....	140	138.17
Connecticut.....	192	192.57	West Virginia.....	31	134.20
West Virginia.....	317	186.80	Louisiana.....	243	125.90
Louisiana.....	387	186.14	New York.....	210	115.70
New York.....	1,716	185.71	Illinois.....	450	113.78
Arizona.....	24	184.61	Wisconsin.....	29	96.67
Illinois.....	1,221	182.02	Texas.....	290	95.43
Wisconsin.....	493	180.42	Mississippi.....	268	93.43
Texas.....	787	179.92	Washington.....	1	76.02
Mississippi.....	375	178.40			
Washington.....	361	178.27			
			Total.....	17,955	175.03

TABLE 126.—*Absolute and relative numbers of veterans with dark brown eyes, by States of nativity in order of incidence, demobilization, 1919.*

State.	Number of cases.	Ratio.	State.	Number of cases.	Ratio.
Louisiana.....	1,006	483.98	Montana.....	1,297	182.88
North Carolina.....	734	404.41	Iowa.....	41	180.61
Virginia.....	729	374.61	North Dakota.....	180	177.57
District of Columbia.....	79	341.99	Utah.....	471	165.43
Georgia.....	1,138	334.41	New Hampshire.....	334	164.94
Mississippi.....	694	320.13	Minnesota.....	462	164.12
Florida.....	328	320.31	Idaho.....	13	162.30
South Carolina.....	257	310.01	Wisconsin.....	157	154.68
New Mexico.....	68	295.65	Vermont.....	730	154.12
Maryland.....	331	289.84	Maine.....	126	153.09
Arkansas.....	701	271.50	Delaware.....	508	151.19
Alabama.....	515	266.56	Alaska.....	905	148.33
Texas.....	1,145	261.77	Idaho.....	538	144.31
New York.....	2,384	258.01	Montana.....	60	144.23
Delaware.....	72	240.00	Iowa.....	38	142.86
Alaska.....	3	230.77	North Dakota.....	220	136.65
Arizona.....	30	221.11	Utah.....	41	114.52
Nevada.....	4	222.22	New Hampshire.....	12	114.23
Oklahoma.....	498	209.84	Minnesota.....	47	113.53
Rhode Island.....	84	206.44	Idaho.....	211	103.15
Pennsylvania.....	2,257	207.05	Wisconsin.....	17	103.06
New Jersey.....	653	204.83	Vermont.....	273	101.98
California.....	97	200.83	Maine.....	41	91.72
Connecticut.....	196	198.60		60	86.45
Kentucky.....	545	185.75			
West Virginia.....	311	183.27			
			Total.....	21,324	112.28

(d) *Eye color in eight European races.*—Table 127 shows the absolute and proportional occurrence of eye color in each of the eight races, of each of which more than 1,000 men were observed. According to this table the Irish show the largest percentage of clear blue eyes, the Scotch second, followed by the Polish, English, German, French, Hebrew, and Italian. If we combine clear blue and blue with brown spots, the highest proportion of blue eyes still remains with the Irish, 73 per cent; next come the Scotch with 71 per cent; next the Polish and English, each about 66 per cent; then come the German with 65 per cent, French with 49 per cent, Hebrews 37 per cent, and Italian 20 per cent. Dark brown eyes naturally run for the most part in inverse order. Italians stand at the head with 51 per cent; Hebrews next with 38 per cent, French 25 per cent, Germans 15 per cent, English 15 per cent, Scotch 14 per cent, Polish 13 per cent, and Irish 11 per cent.

TABLE 127.—*Comparative frequency distribution of eye color in each of eight European races, demobilization.*

SECTION A: ABSOLUTE NUMBERS.

Race.	Total.	Clear blue.	Blue and brown spots.	Light brown.	Dark brown.
English.....	4,194	1,852	920	794	628
Scotch.....	2,049	978	484	310	277
Irish.....	6,144	3,279	1,224	964	677
German.....	7,059	3,008	1,572	1,400	1,079
French.....	1,429	490	212	376	351
Italian.....	3,486	389	319	999	1,779
Polish.....	2,399	1,124	480	485	310
Hebrew.....	1,685	389	232	426	638
	28,445	11,509	5,443	5,754	5,739
	225				
Total.....	28,670				

SECTION B: RACE DISTRIBUTION PER 1,000 OF EACH EYE COLOR.

Race.	Total.	Clear blue.	Blue and brown spots.	Light brown.	Dark brown.
English.....	147.44	160.92	169.03	137.99	109.43
Scotch.....	72.03	84.98	88.92	53.88	48.27
Irish.....	216.00	284.91	224.88	167.54	117.97
German.....	248.16	261.36	288.81	243.31	188.02
French.....	50.24	42.58	38.95	65.35	61.17
Italian.....	122.55	33.80	58.61	173.62	309.98
Polish.....	84.34	97.66	88.19	84.29	54.02
Hebrew.....	59.24	33.80	42.62	74.04	111.17
Total.....	1,000.00	1,000.01	1,000.01	1,000.02	1,000.03

SECTION C: EYE-COLOR DISTRIBUTION PER 1,000 OF EACH RACE.

Race.	Total.	Clear blue.	Blue and brown spots.	Light brown.	Dark brown.	Total.
English.....	147.44	441.57	219.37	189.32	149.74	1,000
Scotch.....	72.03	477.29	236.21	151.30	135.20	1,000
Irish.....	216.00	533.70	199.21	156.89	110.20	1,000
German.....	248.16	426.14	222.69	198.32	152.85	1,000
French.....	50.24	342.90	148.35	263.12	245.62	1,000
Italian.....	122.55	111.59	91.51	286.56	510.32	1,000
Polish.....	84.34	468.50	200.08	202.18	129.23	1,000
Hebrew.....	59.24	230.86	137.69	252.81	378.63	1,000

(e) *Comparison with Civil War data.*—These results may be compared with those given by Baxter and Gould for Civil War recruits. According to Baxter,¹ the examination of 9,649 Englishmen gave a ratio of 71 per cent for blue or gray eyes combined with light hair, and 29 per cent for dark or hazel eyes and dark hair. Assuming that the examiners of recruits did not distinguish between clear blue eyes and those with small brown spots, the ratio of 71 per cent in Civil War times is to be contrasted with 66 per cent among the English at demobilization of the troops of the World War.

The statistics of Baxter for 28,995 Irishmen give a proportion of blue or gray eyes combined with light hair of 70 per cent, to be compared with 73 per cent of our statistics. Baxter finds in an examination of 29,600 Germans a ratio of blue or gray eyes and light hair of 69 per cent; our statistics give 65 per cent. There are copied from Gould² (pp. 196–201) in our Tables 128 and 129 data concerning the color of the eyes of United States soldiers by States and of volunteers by nativity.

TABLE 128.—*Color of eyes: Proportional numbers for different States in 1865 (Gould,² p. 200).*

State of enlistment.	Blue.	Gray.	Hazel.	Dark.	Black.	Total.
Maine.....	458	171	193	70	106	1,000
New Hampshire.....	494	193	168	75	70	1,000
Vermont.....	555	148	82	98	117	1,000
Massachusetts.....	506	184	173	76	61	1,000
Connecticut.....	476	228	124	103	69	1,000
New York.....	467	255	75	140	63	1,000
Pennsylvania.....	319	356	142	150	33	1,000
West Virginia.....	430	258	84	126	102	1,000
Kentucky.....	466	220	91	97	126	1,000
Ohio.....	393	293	120	112	82	1,000
Indiana.....	422	258	139	94	87	1,000
Illinois.....	447	245	121	106	81	1,000
Michigan.....	522	224	93	85	76	1,000
Wisconsin.....	533	202	106	93	66	1,000
Iowa.....	462	239	129	86	84	1,000
Missouri.....	460	245	115	107	73	1,000
Total.....	449	243	128	104	76	1,000

TABLE 129.—*Color of eyes: Proportional numbers for different nativities in United States in 1865 (Gould,² p. 201).*

Nativity.	Blue.	Gray.	Hazel.	Dark.	Black.	Total.
Six New England States.....	499	175	150	83	93	1,000
New York, New Jersey, Pennsylvania.....	415	280	119	126	60	1,000
Ohio, Indiana.....	417	266	127	102	88	1,000
Michigan, Wisconsin, Illinois.....	449	237	121	96	97	1,000
Slave States ^a	432	249	112	110	97	1,000
Kentucky and Tennessee.....	464	221	105	94	116	1,000
Free States west of the Mississippi.....	396	284	159	84	77	1,000
Slave States west of the Mississippi.....	435	243	128	96	98	1,000
British America exclusive of Canada.....	464	203	194	78	61	1,000
Canada.....	432	218	154	107	89	1,000
England.....	472	238	142	94	54	1,000
Scotland.....	478	254	129	83	56	1,000
Ireland.....	505	274	119	69	33	1,000
France, Belgium, and Switzerland.....	328	225	192	151	104	1,000
Germany.....	445	262	107	141	45	1,000
Scandinavia.....	684	172	63	60	21	1,000
Spain, Portugal, and Spanish America.....	239	185	164	197	215	1,000
Miscellaneous.....	349	250	149	158	94	1,000
Total.....	449	243	128	104	76	1,000

^a Not including Kentucky and Tennessee and Slave States west of the Mississippi.

A comparison of the proportions of the population having different colored eyes may be made between Civil War times and those of the demobilization in the recent war. The order of proportion of blue eye color in the Civil War in the different States is as follows: Vermont, 555; Wisconsin, 533; Michigan, 522; Massachusetts, 506; New Hampshire, 494; Connecticut, 476; New York, 467; Kentucky, 466; Iowa, 462; Missouri, 460; Maine, 458; Illinois, 447; West Virginia, 430; Indiana, 422; Ohio, 393; Pennsylvania, 319. The average for the States named is 449 in Civil War times, as contrasted with 375 in the World War. This suggests a marked decrease in the proportion of blue eyes, namely, from 45 to 37 per cent. However, it is to be remembered that the Southern States were not included in the Civil War statistics, and these are just the States that show the smallest proportion of clear blue eyes. The inclusion of such States would inevitably tend to lower the average in the World War statistics. Indeed, if we compare the States which are mentioned both in the Civil War records and in those of the World War we find some cases of marked agreement. Thus Wisconsin was 533; is 539, per 1,000, blue-eyed; Vermont was 555, and has become darker, 512; Massachusetts was 506, has become a trifle darker, 493; Michigan has fallen from 522 to 488; New Hampshire from 494 to 486; Connecticut from 476 to 465; Illinois has increased from 447 to 463, due, no doubt, to the coming in of Scandinavians in recent decades. West Virginia has remained nearly constant at 430 then and 428 now. Ohio was 393, is 427; New York was 467, is 416; a great decrease, due to the immigration from the south and east of Europe. Pennsylvania, on the other hand, has increased enormously from 319 to 401, the meaning of which is not perfectly clear, but is it possibly due to the coming in of large numbers of blue-eyed Poles and Lithuanians. Kentucky was 466 and is 193, which indicates that the recruits from Kentucky to the Northern Army in Civil War times were a highly selected lot of Nordics from the mountain regions and largely excluded Negroes. Indiana has fallen from 447 to 320, again a marked decline.

Since the categories are not the same in 1866 and 1919, it is difficult to compare the darker eyes. It is clear that the West Virginians, however, had a prevalence of dark eye color which is hardly recognized to-day. In general, persons who have much pigment in the iris are more numerous in the United States to-day than they were 55 years ago. It is possible to compare some of the races described in Gould's book with those examined in 1919. Among the English the proportion of blue eyes was 472, is now 442; among the Scotch, then 478, now 477; among the Irish, then 505, now 534; among the French, Belgians, and Swiss, then 328, now 343; among the Germans, then 445, now 426. If we add together the "dark" and the "black" eye colors of Gould, we have a total for the English of 148, as opposed to 150 of our "dark browns"; for the Scotch, 139, as opposed to 135 in 1919; for the Irish, 102, as opposed to 110 at the later date; for the French, 255, as opposed to 246; for the Germans 186, as opposed to 153. It is clear that the dark and black are nearly equivalent to our dark brown, and it is probable that Gould's hazel corresponds nearly with our light brown as well as with our blue with brown spots. The comparison is of interest, showing the comparative stability of proportions in racial populations. But there have been great changes in sections of our country due to extensive immigration.

TABLE 130.—Comparative frequency distribution of eye color by Q. M. C. distribution zones, based on nativity of demobilized troops.

SECTION A: ABSOLUTE NUMBERS.

Eye color.	Total.	Zone 1.	Zone 2.	Zone 4.	Zone 5.	Zone 7.	Zone 8.	Zone 9.	Zone 10.	Zone 11.	Zone 12.	Zone 13.
Clear blue.....	32,345	1,996	7,421	1,050	1,764	12,002	3,175	952	1,452	774	112	1,647
Blue with brown spots.....	23,947	1,174	4,395	623	4,820	7,864	2,198	767	865	479	70	692
Light brown.....	23,585	1,236	5,644	1,060	4,082	5,607	1,679	1,848	1,243	348	78	760
Dark brown.....	10,528	496	2,548	425	916	3,137	1,133	519	579	185	98	492
Number measured.....	90,405	4,902	20,008	3,158	11,562	28,610	8,185	4,086	4,139	1,786	358	3,591
Not measured.....	11,928											
Total.....	102,333											

SECTION B: EYE-COLOR DISTRIBUTION PER 1,000 OF EACH ZONE.

Eye color.....	Total.....	Zone 1.	Zone 2.	Zone 4.	Zone 5.	Zone 7.	Zone 8.	Zone 9.	Zone 10.	Zone 11.	Zone 12.	Zone 13.
Clear blue.....	357.78	407.18	370.90	332.49	152.31	419.50	387.90	232.99	350.81	433.37	312.85	458.65
Blue with brown spots.....	264.89	239.49	219.66	197.28	416.16	274.87	268.54	187.71	208.99	268.19	195.53	192.70
Light brown.....	260.88	252.14	282.08	335.66	352.44	195.98	205.13	452.28	500.31	194.85	217.88	211.64
Dark brown.....	116.45	101.18	127.35	134.58	79.09	109.65	138.42	127.02	139.89	103.58	273.74	137.01
Total.....	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00

H. DISTRIBUTION OF HAIR COLOR.

(a) *General discussion.*—The directions given to the anthropologists called for the use of the following terms in describing hair color: Flaxen, light brown, medium brown, dark brown, red, red and black. The last was explained to mean the presence of melanic pigment mixed with the red as opposed to pure red. The terms used by Gould are light, brown, dark, black, red, sandy, and gray. There were 4,000, or 1 per cent of all, in Gould's statistics returned as gray. In our statistics the term gray was not used, as the color before graying was to be recorded. Tables 131–135 show the proportion of cases of the different hair colors in the different States. In each table the States are arranged in descending order of frequency of the stated hair color.

(b) *Flaxen hair.*—Table 131 gives the list of States in descending order of the population having flaxen hair. From this table it appears that there are proportionally more persons with flaxen hair in Oregon than any other State, 28 per cent; Montana follows with 23 per cent; and Utah with 14 per cent. Minnesota and South Dakota have about 10 per cent each, and this result is largely due to the Scandinavian population. Then follow Alaska, Iowa, and Michigan. At the opposite end of the table stand the Southern Atlantic and Gulf States, with their large Negro and mulatto populations; Florida, Alabama, and Georgia, each with less than 1 per cent; South Carolina, Louisiana, Mississippi, North Carolina, and Kentucky with 2 per cent or less. In the middle of the series lie the New England States and the more densely populated States of the Mississippi Valley, such as Ohio, New Hampshire, Connecticut, Indiana, Wisconsin, Massachusetts, Vermont, Maine, Illinois, Pennsylvania, New Jersey, Rhode Island, Maryland, and New York. One may be quite sure that the presence of flaxen hair is indicative of Nordic blood, and one draws the conclusion that there is a larger proportion of this in Oregon, Montana, and Utah than in the other States. The relative absence of light hair in the Southern States is to be attributed to the colored part of the population (see Plate XXX, Figs. 1, 8, p. 295).

TABLE 131.—*Absolute and relative number of veterans with flaxen hair, by States of nativity in order of incidence, demobilization, 1919.*

States.	Number of cases.	Ratios.	States.	Number of cases.	Ratios.
Oregon.....	302	282.24	Illinois.....	329	49.06
Montana.....	62	233.08	Pennsylvania.....	529	48.53
Utah.....	15	142.86	New Mexico.....	11	47.83
Minnesota.....	195	99.95	New Jersey.....	152	47.68
South Dakota.....	40	96.15	Rhode Island.....	19	47.15
Alaska.....	1	76.92	Maryland.....	53	46.41
Iowa.....	122	75.78	Oklahoma.....	103	44.47
Michigan.....	280	75.11	California.....	20	41.41
Texas.....	310	70.87	New York.....	347	37.55
North Dakota.....	25	69.83	Virginia.....	67	34.72
Idaho.....	11	67.07	District of Columbia.....	8	34.63
Ohio.....	472	66.54	Missouri.....	97	24.07
Wyoming.....	5	62.50	Kentucky.....	63	21.47
Nebraska.....	51	61.97	North Carolina.....	38	20.94
Colorado.....	14	61.67	Mississippi.....	43	20.46
Kansas.....	62	61.06	Arizona.....	2	15.38
New Hampshire.....	25	60.39	Tennessee.....	40	14.21
Connecticut.....	60	60.18	Louisiana.....	29	13.96
Indiana.....	237	59.92	South Carolina.....	9	10.86
Wisconsin.....	157	58.65	Delaware.....	3	10.00
Massachusetts.....	281	58.60	Georgia.....	29	8.52
Vermont.....	26	58.17	Alabama.....	16	8.28
Maine.....	38	54.76	Florida.....	8	7.81
Washington.....	107	52.84			
West Virginia.....	88	51.86	Total.....	5,132	50.03
Arkansas.....	131	50.74			

TABLE 132.—*Absolute and relative number of veterans with red hair, by States of nativity in order of incidence, demobilization, 1919.*

States.	Number of cases.	Ratios.	States.	Number of cases.	Ratios.
Montana.....	14	52.63	Pennsylvania.....	136	12.48
Oregon.....	32	29.91	Indiana.....	49	12.39
Wyoming.....	2	25.00	Idaho.....	2	12.20
Minnesota.....	40	20.50	Oklahoma.....	28	12.09
New Hampshire.....	8	19.32	Arkansas.....	31	12.01
Utah.....	2	19.05	Missouri.....	34	11.94
North Dakota.....	6	16.76	Vermont.....	5	11.19
Maryland.....	19	16.64	Michigan.....	40	10.73
Florida.....	17	16.60	Virginia.....	20	10.36
Washington.....	33	16.30	Alabama.....	20	10.35
Illinois.....	109	16.25	California.....	5	10.35
Connecticut.....	16	16.05	South Dakota.....	4	9.62
New Jersey.....	50	15.68	Ohio.....	67	9.44
Texas.....	68	15.55	Iowa.....	15	9.32
New York.....	138	14.94	North Carolina.....	14	7.71
Rhode Island.....	6	14.89	Wisconsin.....	20	7.47
Massachusetts.....	67	13.97	South Carolina.....	6	7.24
Tennessee.....	39	13.85	Mississippi.....	15	7.14
Kansas.....	14	13.79	Delaware.....	2	6.67
West Virginia.....	23	13.55	Louisiana.....	13	6.25
Nebraska.....	11	13.37	District of Columbia.....	1	4.33
Kentucky.....	39	13.29	Maine.....	3	4.32
Colorado.....	3	13.22			
Georgia.....	43	12.64	Total.....	1,329	12.96

(c) *Dark brown hair.*—Turning to the dark brown hair, we find that the Southern States are at the head of the list, North Carolina and Louisiana at the very top, and Virginia, Mississippi, Maryland, South Carolina, and Georgia stand above the average in percentage of population with dark brown hair. On the other hand, there is less of this in Montana and Oregon relatively than in any other States (see Plate XXX, Fig. 5, p. 295).

(d) *Red hair.*—Red hair was so relatively uncommon that it becomes almost futile to compare the proportions secured. On the face of the returns there are proportionately more red heads in Montana than in any other State, and Oregon comes second—that is, there is a close correlation between the proportion of flaxen and of red hair. However, Maine stands near the middle of the series for flaxen hair and at the bottom of the series for red, indicating that the association is not absolute. The Southern States tend to lie at the bottom of the list of the rates of red hair. Thus Louisiana, Mississippi, South Carolina, North Carolina, Alabama, and Virginia are markedly below the mean of the whole population. On the other hand, Florida stands relatively high at 1.6 per cent (mean of United States, 1.3 per cent). (See Plate XXX, Fig. 1, p. 295).

TABLE 133.—*Absolute and relative number of veterans with light brown hair, by States of nativity in order of incidence, demobilization, 1919.*

States.	Number of cases.	Ratios.	States.	Number of cases.	Ratios.
Alaska.....	5	384.62	Mississippi.....	411	195.53
Nevada.....	6	333.33	Arizona.....	25	192.31
Wisconsin.....	880	328.73	Kentucky.....	563	191.89
Michigan.....	1,190	319.21	New York.....	1,765	191.02
Ohio.....	2,183	307.72	Arkansas.....	490	189.78
Minnesota.....	587	300.87	Delaware.....	55	183.33
Illinois.....	2,000	298.15	Maryland.....	201	176.01
North Dakota.....	102	284.92	Tennessee.....	488	173.36
Utah.....	29	276.19	District of Columbia.....	40	173.16
Wyoming.....	22	275.00	Maine.....	118	170.03
Idaho.....	45	274.39	Texas.....	742	169.64
Washington.....	550	271.60	Montana.....	45	169.17
Nebraska.....	219	266.10	Massachusetts.....	804	167.67
Iowa.....	422	262.11	New Hampshire.....	67	161.84
California.....	126	260.87	Connecticut.....	160	160.48
South Dakota.....	107	257.21	Virginia.....	309	160.10
Kansas.....	261	257.14	Alabama.....	287	148.55
Indiana.....	995	251.58	Florida.....	152	148.44
Missouri.....	646	226.91	South Carolina.....	110	132.69
New Jersey.....	723	226.79	Georgia.....	449	131.94
West Virginia.....	382	225.10	Louisiana.....	270	129.87
Oregon.....	234	218.69	Rhode Island.....	52	129.03
Colorado.....	49	215.86	North Carolina.....	228	125.62
Pennsylvania.....	2,329	213.65	New Mexico.....	28	121.74
Vermont.....	90	201.34			
Oklahoma.....	465	200.78	Total.....	22,506	219.40

TABLE 134.—*Absolute and relative number of veterans with medium brown hair, by States of nativity in order of incidence, demobilization, 1919.*

States.	Number of cases.	Ratios.	States.	Number of cases.	Ratios.
Alaska.....	7	538.46	Arkansas.....	471	182.42
Tennessee.....	1,255	445.83	Mississippi.....	381	181.26
Montana.....	116	436.09	California.....	87	180.12
Alabama.....	782	404.76	Delaware.....	54	180.00
Oregon.....	414	386.92	Massachusetts.....	849	177.06
Kentucky.....	1,083	369.12	Michigan.....	646	173.28
Missouri.....	1,003	352.30	Utah.....	18	171.43
Georgia.....	1,183	347.63	Ohio.....	1,176	165.77
Florida.....	351	342.77	Vermont.....	73	163.31
South Carolina.....	261	314.84	Maryland.....	186	162.87
Indiana.....	1,241	313.78	Texas.....	712	162.78
Wisconsin.....	686	256.26	Arizona.....	20	153.85
Illinois.....	1,649	245.83	North Carolina.....	278	153.17
Wyoming.....	19	237.50	Idaho.....	25	152.44
Iowa.....	376	233.54	Pennsylvania.....	1,598	145.67
Washington.....	450	222.22	West Virginia.....	244	143.78
South Dakota.....	92	221.15	Virginia.....	275	142.49
Minnesota.....	430	220.40	New Jersey.....	431	135.19
North Dakota.....	78	217.88	New Mexico.....	31	134.78
Maine.....	147	211.82	New York.....	1,224	132.47
Rhode Island.....	84	208.44	Louisiana.....	237	114.00
New Hampshire.....	85	205.31	District of Columbia.....	23	99.57
Nebraska.....	162	196.84	Nevada.....	1	55.56
Colorado.....	44	193.83			
Oklahoma.....	441	190.41	Total.....	21,656	213.12
Connecticut.....	187	187.56			

TABLE 135.—*Absolute and relative number of veterans with dark brown hair, by States of nativity in order of incidence, demobilization, 1919.*

States.	Number of cases.	Ratios.	States.	Number of cases.	Ratios.
North Carolina.....	1,207	665.01	Arkansas.....	1,146	443.84
Louisiana.....	1,360	654.16	California.....	213	440.99
New Mexico.....	144	626.09	Colorado.....	100	440.53
District of Columbia.....	143	619.05	Nebraska.....	350	425.27
Nevada.....	11	611.11	Ohio.....	3,007	423.88
Virginia.....	1,143	592.23	Kansas.....	425	418.72
New York.....	5,212	564.07	Alabama.....	776	401.66
Delaware.....	168	560.00	Washington.....	811	400.49
Arizona.....	72	553.85	Michigan.....	1,484	398.07
Mississippi.....	1,159	551.38	Iowa.....	622	386.34
Rhode Island.....	219	543.42	North Dakota.....	134	374.30
Maryland.....	604	528.90	Utah.....	39	371.43
Pennsylvania.....	5,703	523.16	South Dakota.....	154	370.19
New Jersey.....	1,067	522.90	Kentucky.....	1,081	368.44
Massachusetts.....	2,498	520.96	Indiana.....	1,343	339.57
Connecticut.....	518	519.56	Missouri.....	962	337.90
Maine.....	360	518.73	Illinois.....	2,238	333.63
West Virginia.....	571	513.26	Tennessee.....	930	330.37
New Hampshire.....	205	495.17	Minnesota.....	629	322.40
South Carolina.....	400	482.51	Wisconsin.....	799	298.47
Vermont.....	214	478.75	Wyoming.....	23	287.50
Georgia.....	1,619	475.76	Montana.....	6	22.56
Idaho.....	78	475.61	Oregon.....	21	19.63
Texas.....	2,044	467.31			
Florida.....	477	465.82			
Oklahoma.....	1,057	456.39	Total.....	46,446	452.79

(e) *Comparison with Civil War recruits.*—A comparison of the proportion of kind of hair color found in the different States in 1919 with that found in corresponding States in 1866 will be of interest. Assuming that the light hair of Gould's statistics corresponds with the flaxen hair of the statistics of 1919, then we have for the whole territory considered in 1866 a rate of 235 per 1,000 of hair colors belonging to the category of light, and, in 1919, 50 per 1,000 belonging to the category of light. On the face of it, this is an enormous reduction in the proportion of flaxen hair as compared with the light hair of half a century earlier. Fifty years ago the State with the largest percentage of light hair color was Kentucky, with 381 per 1,000; in 1919 the proportion of flaxen hair in Kentucky was 21 per 1,000, and of light brown hair 192 per 1,000, or together 213 per 1,000; in any case an enormous decrease of light hair in the population. This is probably due to the fact that the recruits from Kentucky during the Civil War were drawn especially from the mountain regions and contained few or no colored men, whereas in the World War they were uniformly from the whole State and included colored as well as white.

In the series of light hair in the Civil War we find West Virginia standing second, with 311 per 1,000; in 1919 there are 52 per 1,000 with flaxen hair and 225 per 1,000 with light brown hair; a total of 277 with light hair. This is a marked reduction in the proportion of light hair in this State, due no doubt to the inclusion of many colored men in the present series. Next in the Civil War series of light hair stands Indiana, with a ratio of 294 per 1,000. In the World War this State had a ratio for flaxen hair of 60 per 1,000 and of light-brown hair of 252 per 1,000, or together 312 per 1,000. This indicates no great change in the proportion of light hair in this State. In 1866, of men from Missouri, 291 per 1,000 had light hair; in 1919, 34 per 1,000 of the men from this State were stated to have flaxen hair and 227 per 1,000 light-brown hair; a total of 261 per 1,000, a slight decrease during 50 years. In 1866 the ratio

for light-brown hair for Illinois was 286; in 1919 for flaxen hair it is 49, for light-brown hair 298; a total of 347, apparently an increase in the proportion of blonds in this State, probably due to the immigration of Scandinavians and Germans. Similarly the proportion of blonds has probably risen in Ohio and Wisconsin, remained stationary or fallen in Massachusetts, and increased somewhat in Pennsylvania from 204 to 262. The apparent increases of the lighter colors of hair in Vermont, New York, Connecticut, and Maine may very likely be due to the fact that the categories were not quantitatively distinguished either for Civil War recruits or World War troops, and hence the limits were not drawn uniformly.

In regard to the distribution of light hair color by races, we find that of World War troops there is a larger proportion of flaxen hair in the Polish than in any of the other eight races considered. It is to be noted that Scandinavians were not included in the study, as there were relatively few of them. The proportion of flaxen and light-brown hair together in 1919 is 374 per 1,000 in Germans; for light hair color in 1866 it was 290. In the Civil War soldiers, as in those of the World War, the proportion of light hair stands highest in Germans, if we omit Scandinavians and Polish from consideration. Third in position of World War troops in proportion of light hair are the English, and this position is the same that they occupied in the Civil War. Next in both Civil War and World War series stand the Scotch, then come the Irish and French. The Hebrews come next in the World War series; last come the Italians, with 6 per 1,000 flaxen hair, or 65 per 1,000 of flaxen and light-brown hair together. This proportion agrees pretty well with the proportion of Spanish and Portuguese recruits in the Civil War of 42 per 1,000.

TABLE 136.—Comparative frequency distribution of hair color in each of 8 races, demobilization, 1919.

SECTION A: ABSOLUTE NUMBERS.

Race.	Total.	Flaxen.	Light brown.	Medium brown.	Dark brown.	Clear red.	Red and black.
English.....	4, 196	231	969	950	1, 826	58	133
Scotch.....	2, 045	108	468	459	863	41	106
Irish.....	6, 137	232	1, 157	1, 140	3, 138	156	314
German.....	7, 067	484	2, 163	1, 467	2, 711	48	194
French.....	1, 434	39	199	237	885	11	63
Italian.....	3, 488	21	206	278	2, 636	6	341
Polish.....	2, 402	182	801	470	863	17	69
Hebrew.....	1, 686	27	186	188	1, 131	15	139
Number measured.....	28, 455	1, 324	6, 169	5, 198	14, 053	352	1, 359
Not measured.....	215						
Total.....	28, 670						

SECTION B: HAIR-COLOR DISTRIBUTION PER 1,000 OF EACH RACE.

Race.	Total.	Flaxen.	Light brown.	Medium brown.	Dark brown.	Clear red.	Red and black.	Total.
English.....	147. 46	55. 05	235. 70	228. 55	435. 20	13. 82	31. 69	1, 000
Scotch.....	71. 87	52. 81	228. 85	224. 44	422. 00	20. 05	51. 83	1, 000
Irish.....	215. 67	37. 80	188. 54	185. 77	511. 32	25. 42	51. 17	1, 000
German.....	248. 36	68. 49	306. 08	207. 60	348. 61	6. 79	27. 45	1, 000
French.....	50. 40	27. 19	134. 77	165. 27	617. 19	7. 67	43. 93	1, 000
Italian.....	122. 58	6. 02	59. 06	79. 70	755. 73	1. 72	97. 76	1, 000
Polish.....	84. 41	75. 77	333. 47	195. 67	359. 28	7. 08	28. 72	1, 000
Hebrew.....	59. 25	16. 01	110. 31	111. 51	670. 81	8. 90	82. 44	1, 000

Clear red hair is perhaps the most satisfactory color to serve as a basis for comparison between Civil War and World War troops. In the Civil War the rate for Scotland was higher than that for any other country, namely, 27 per 1,000, and Ireland came next with 23 per 1,000. In the World War series Ireland stands first with 25 per 1,000, and the Scotch second with 20 per 1,000. It is probable that the more recent Scotch immigrants have been drawn from a different part of Scotland than the earlier one. Third in the Civil War series stands England with a rate of 22, whereas for England in the World War series the rate is 14. Next in the Civil War series comes Germany with a rate of 19, but the rate is 6.8 in the case of World War troops, and this rate is exceeded by Hebrews, 9 per 1,000; French, 8 per 1,000, and by the Polish, 7 per 1,000. The rate for the French, Belgians, and Swiss together in the Civil War was 16 per 1,000. The smallest ratio of red hair is found in troops of Italian origin, namely, 1.7 per 1,000. In the case of Civil War troops the smallest ratio was in the Spanish and Portuguese, 3 per 1,000. Red hair seems to be getting rarer in all European stocks.

(f) *By Quartermaster distribution zones.*—Table 137 gives the distribution of the various hair colors in the Quartermaster's distribution zones. The rate for flaxen hair reaches a maximum in zone 11, including the Dakotas, Nebraska, and the three central Mountain States, 75 per 1,000. The next highest rate is in zone 7, surrounding the Great Lakes; next in zone 1, the New England States except Connecticut; next zone 13, the Pacific and northern Rocky Mountain States. The zone with the smallest proportion of flaxen hair, 12 per cent, is zone 5, including the Southeastern States. Just above in order stand zone 9, zone 4, and zone 12, including Arizona and New Mexico, 37 per 1,000. Of clear red hair the largest proportion is found in zone 13, Pacific and northern Rocky Mountain States, 17 per 1,000. Next is zone 10, 16 per 1,000; then comes zone 11, the Dakotas and central Rocky Mountain States, 14 per 1,000; and zone 2, the Middle Atlantic States, also 14 per 1,000. The smallest rate is found in zone 12, from which no case is recorded; and the next is zone 9, the States of Louisiana and Mississippi.

(g) *Hair color in eight European races.*—Table 136-B gives the relative proportion of different classes of hair pigmentation for each of the eight races. This table shows that among the Irish the clear red hair forms a larger proportion of the total than in the case of any other race. Similarly, flaxen hair forms a larger proportion of all hair colors among the Polish than it does among the Germans or any other of the eight races. This table shows strikingly the small amounts of flaxen, light brown, and medium brown hair color among the Italians and the large percentage of dark browns among them.

The table brings out strongly that the Poles in America, probably largely from a restricted area of Polish territory in Europe, are more nearly Nordic in their blue eyes and light hair than are the English, who have suffered so large an admixture of other races. As far as hair color goes, Poles are blonder than the Scotch or the English. It is noteworthy also that among the Scotch, Irish, and Polish, the proportion of clear blue eyes far exceeds the total of flaxen and light brown and clear red hair together. In fact, among the Irish the clear blue eye constitutes 534 per 1,000, whereas the sum of flaxen, light

DISTRIBUTION, HAIR, AND EYE COLOR DEM.—1919

STATES OF NATIVITY

HAIR

FLAXEN AND RED

EYES

CLEAR BLUE AND BLUE WITH BROWN SPOTS


LIGHT AND MEDIUM BROWN

LIGHT BROWN


DARK BROWN

DARK BROWN

7
LINES OF EQUAL PROP. OF FLAXEN HAIR



8
LINES OF EQUAL PROP. OF CLEAR BLUE EYES



SECTION II.

HEIGHT, WEIGHT, AND CHEST CIRCUMFERENCE OF RECRUITS IN RELATION TO VARIOUS DISEASES AND DEFECTS.

I. INTRODUCTORY.

The following study is based upon the physical measurements of defective men from among approximately 2,000,000 men sent to mobilization camps in the United States in connection with the selective drafts of 1917 and 1918. Tabulations have been made separately for those in the first and second million men. The first million include men sent to mobilization camps between September, 1917, and the 1st of May, 1918; the second million of the 1,666,867 who were entrained for mobilization camps between the early part of May, 1918, and November 11, 1918. The second million includes a large (though unknown) proportion of men of the second registration, or of those who had reached the age of 21 subsequent to the first registration.

In studying the results, it must be noted, first of all, that the men measured had already been examined and selected by local boards. They represent the cases accepted by local boards. Presumably all who were rejected on physical grounds fall outside certain limits of acceptance designated in the physical examination standards. On the other hand, some men, whose physical dimensions lay outside the ordinary limits of acceptance, nevertheless got to camp under various broad interpretations of the standards, and a few were sent through accident.

II. STANDARDS OF MEASUREMENTS OF DRAFTED MEN.

(a) *Stature*.—The changing military standards for stature during the period of the draft have been referred to in detail in an earlier chapter. Always men under 60 inches and over 78 inches were to be rejected; but apparently some were sent to camp who were outside the regulation limits, because of exceptional qualifications in other respects.

The mean stature of the first million recruits, including defectives, sent to camp, is 67.49; the standard deviation, or measure of variability, 2.71 inches. The mean is not that of adult males in general, but that of a selected lot, from whom the shortest and tallest has been eliminated. Not until the measurement of the men rejected by the local boards shall have been tabulated will we be able to estimate the true mean stature of young adult American males.

(b) *Weight*.—From the beginning of the selective draft stress was laid upon securing for the Army men of proper weight. Experience indicates that men who are below a certain standard of weight are unable, ordinarily, to carry a heavy pack, and that those over a certain weight are too unwieldy for rapid movement.

The standards of the War Department in the years before the draft provided a minimum weight for all branches of the service of 128 pounds. But it was provided that men 64 inches in height might be accepted who weighed only 120 pounds, if otherwise sound and apparently healthy. It was necessary to obtain special permission from The Adjutant General to enlist a man who weighed less than 120 pounds. The maximum weight was placed at 190 pounds for Infantry, Engineers, Coast Artillery, and Field Artillery, and 165 pounds for Cavalry. At the beginning of the draft local boards were instructed that the minimum weight was 118 pounds and the maximum 211 pounds. But it was provided that, "when the applicant is active, has firm muscles, and is evidently vigorous and healthy" a weight of 8 pounds below the minimum would be accepted for men 61 to 64 inches; of 24 pounds below the minimum for men 73 inches and upward, and for intermediates permissible variations below the standard were given. The regulations further state: "Variations in weight above the standard would not disqualify unless sufficient to constitute obesity."

III. PHYSICAL-EXAMINATION STANDARDS.

(a) *Stature and weight.*—The physical-examination standards for local boards of November, 1917, gave a table (Table 138) showing the relations between standard accepted measurements and the permissible variations from the standard.

TABLE 138.—Standards of height, weight, chest circumference, and mobility of chest, adopted for draft recruits, United States Army, 1917.^a

Column A.				The following variations from the standard shown in Column A are permissible when the applicant is <i>active</i> , has firm muscles, and is <i>evidently vigorous and healthy</i> .			
Standard accepted measurements.							
Stature.	Weight.	Chest measurement: Expiration mobility.		Stature.	Weight.	Chest measurement: Expiration mobility.	
<i>Inches.</i>	<i>Pounds.</i>	<i>Inches.</i>	<i>Inches.</i>	<i>Inches.</i>	<i>Pounds.</i>	<i>Inches.</i>	<i>Inches.</i>
61	118	31	2	61	110	30	2
62	120	31	2	62	110	30	2
63	124	31	2	63	112	30	2
64	128	32	2	64	113	30	2
65	130	32	2	65	114	30	2
66	132	32½	2	66	116	30½	2
67	134	33	2	67	118	30½	2
68	141	33½	2½	68	121	30½	2
69	148	33½	2½	69	124	31	2
70	155	34	2½	70	128	31½	2
71	162	34½	2½	71	133	31½	2
72	169	34½	3	72	138	32½	2½
73	176	35½	3	73	143	32½	2½
74	183	36½	3	74	148	33½	2½
75	190	36½	3½	75	155	34½	2½
76	197	37½	3½	76	161	34½	2½
77	204	37½	3½	77	168	35½	3
78	211	38½	4	78	175	35½	3

^a Selective Service Regulations, Nov. 8, 1917.

It was moreover specified:

Variations in weight above the standard are not disqualifying, unless sufficient to constitute such well-marked obesity as to interfere permanently with normal physical activity.

The standards for local boards of January, 1918, reduced the minimum height from 61 to 60 inches and raised the minimum weight to 120 pounds, but in other

respects did not vary from the table of November, 1917. It was further specified:

Reject registrants whose weight is less than 100 pounds, unless it is plainly due to some recent illness and otherwise the registrants have no disqualifying defect.

Registrants whose weight is more than 100 pounds and less than 114 pounds and who have no other disqualifying defect are to be referred to the Medical Advisory Boards.

Registrants underweight in proportion to their height (see table), unless it is plainly due to some temporary cause, are referred to the Medical Advisory Board. When this underweight can reasonably be explained and the registrant is otherwise physically fit, *accept*.

Registrants with overweight are to be *accepted* unless the obesity interferes with normal physical activity. Refer all doubtful cases to the Medical Advisory Board.

For the Medical Advisory Boards there were issued in March, 1918, standards similar to those furnished the local boards in January. It is moreover stated:

Registrants who weigh less than 114 pounds shall not be accepted for general military service unless in the opinion of the Medical Advisory Board it is a remediable defect.

Registrants who weigh more than 120 pounds, but less than the prescribed weight for the height indicated in the table of measurements of height and weight, may be accepted when in the opinion of the Advisory Board the defect is remediable by camp life. If, however, in the opinion of the Advisory Board the defect is not remediable these registrants, if otherwise physically and mentally fit, shall be accepted for special and limited military service. (Group C.)

From the foregoing extracts from the standards we see that though the weights for each height and the minimum and maximum weights are clearly stated, yet examining boards were permitted considerable latitude in rejecting men whose weight lay outside the standards, and there is internal evidence that boards exercised the discretion thus given to them. For example, in Table I there are recorded over 10,000 men who were under the minimum weight of 114 pounds, and nearly 4,700 men who weighed 200 pounds, which was too great a weight for men even of the maximum stature of 78 inches, of whom there were, indeed, only about 550.

(b) *Chest circumference*.—The Army Regulations require that the circumference of the chest of recruits shall be measured at the time of the physical examination. Ordinarily the circumference of the chest is measured while fully deflated and then when fully expanded. The difference between the two measurements is known as mobility.

The local boards were directed in the first of the physical examination standards (1917) that "all chest measurements are to be taken on a level just above the nipple." Standard chest measurements at expiration for each inch of stature are given in Table 138. The same standards were continued throughout the year. It was prescribed, "All chest measurements to be taken on a level just above the nipple and with the tape horizontal." In January, 1918, there was added to the table as a standard measurement, "Height, 60 inches; weight, 120 pounds; chest at expiration, 31 inches; mobility, 2 inches."

It was further specified: "Registrants whose chest measurements do not come within the limits of the table and who have no disqualifying defect are to be referred to the Medical Advisory Board. Examiners were moreover warned that "Measurements should be taken with the greatest care."

Instructions to the Medical Advisory Boards of February, 1918, repeated these regulations for the local boards and added:

A registrant who appears not to be able to expand the chest $2\frac{1}{2}$ to 3 inches, respectively, as per table, should be examined especially to ascertain if the failure of adequate chest expansion is due

to ignorance and lack of practice. If in the opinion of the Advisory Board the lack of the prescribed expansion is remediable by camp life and the registrant is otherwise physically and mentally fit he shall be accepted. If, however, in the opinion of the Advisory Board the defect of expansion is not remediable and the registrant is otherwise physically and mentally fit he shall be accepted for special and limited military service. (Group C.)

The growing precision and emphasis in physical examination requirements of later date indicate a realization by the board responsible for the standards that the chest circumference was not always taken adequately by local boards. During the earlier period under consideration when examination at camps were made by regimental medical officers, the instructions given to them was that *weight, height, and chest measurements will be copied from data on physical forms (No. 14 P. M. G. O.) furnished by the local boards, except in those cases referred to the specialist for retaking of weight, height, and chest measurements*, in which case the specialist will note his findings in the proper place on record card. Subsequently, however, when the examinations were conducted at each camp by a central examining board, it was the custom for each such board to retake and record the weight, height, and chest measurements.

In the tables referred to in the following sections there are considered only the measurements of chest at expiration, which is certainly somewhat less than the chest circumference at rest. The measurement of the chest circumference at rest was not taken by the medical examiners. The chest circumference at expiration is taken as most nearly representing the circumference of the chest at rest. It may be here mentioned that the average chest circumference found at demobilization, when the chest circumference was taken while at rest, is 34.96 for whites and 34.63 for colored, or probably not far from 34.9 for the whole population. The chest circumference at expiration for the 873,000 men examined by camp boards is given as 33.22; part of the excess of the men at demobilization is to be attributed to exercise and Army training which are adapted to produce an enlargement of the chest. About three-fourths of an inch, however, of the greater size at demobilization is due to the fact that, as stated, the chest was measured at demobilization in a quiescent condition, whereas in the case of recruits it was measured with the lungs deflated as far as possible. As stated, for the entire 873,000 men measured in the early part of the draft, the chest circumference is found on the average to be 33.22 inches at expiration. The standard deviation of this chest circumference is 2.01 inches.

IV. THE DIMENSIONS OF MEN WITH SPECIFIC DEFECTS AND DISEASES.

We now pass to a detailed consideration of the three physical measurements in men with the different classes of defects and diseases and the interpretation of the peculiarities that these dimensions show.

1. PULMONARY TUBERCULOSIS.

There are included in our statistics 10,701 men found at mobilization camps to have pulmonary tuberculosis.

(a) *Stature*.—The average stature of such men is 68.07 inches, which is 0.58 inch greater than the average height of the first million men as shown at the bottom of Table I. The standard deviation in stature of these men is 2.74, which is 0.03 more than the standard deviation of all statures as given in

Table I. That is to say, in respect to stature men with pulmonary tuberculosis are not a random sample of the population, but on the average are selected from the taller men. The significance of tall stature of men with pulmonary tuberculosis is probably not that the organism induces extra growth, but that the tall races of men are less resistant to the *Bacillus tuberculosis* than are the shorter races of stockier build. That the taller races are more susceptible to tuberculosis of the lungs is indicated by a study of Dublin and Baker.²⁴ They show that the rate of mortality from pulmonary tuberculosis is: Among persons born in England, Scotland, and Wales and living in Pennsylvania, 150 per 100,000; living in New York State, 215 per 100,000. Of persons born in Ireland the respective rates are 343 and 589. For persons born in Italy the corresponding rates are 82 and 112, and for persons born in Russia (largely Russian Jews) 107 and 115. This observation then supports the view that pulmonary tuberculosis affects particularly taller races.

Of the 6,048 men found with pulmonary tuberculosis in the second million examined at mobilization camps, the mean stature is 68.12, which is even taller than in the case of the first million. This second group includes more young men, of the age of 21. The standard deviation of stature of men with pulmonary tuberculosis among men of the second million is 2.76 inches, which is 0.06 inch larger than for the first million men.

The distribution of statures in the population of men found with pulmonary tuberculosis is shown in Plate XXXI. This shows at a glance that the modal stature is over one-half an inch greater in this group than in the population at large, and that, on the whole, men with tuberculosis form a group characterized by tall stature.

(b) *Weight*.—Of the 10,701 men found with pulmonary tuberculosis at mobilization camps among the first and second million, the average weight is 130.44 pounds. This is about 11 pounds below the average. This deficiency in weight is the more remarkable inasmuch as the men with tuberculosis are an exceptionally tall lot of men, over half an inch taller than the average. The index of build is important in this connection. As shown in Table 189, the index of build for pulmonary tuberculosis is 28.15, the lowest index, except that of underweight cases, of any class of defects. The reduced weight of men found with pulmonary tuberculosis is in accordance with general experience, since loss of weight is one of the most marked symptoms of active tuberculosis. That the loss of mean weight is not greater is due to the fact that the more advanced cases of active pulmonary tuberculosis were eliminated by local boards and are not included in our statistical tables. It is only the residuum that was sent to camp and there diagnosed as having pulmonary tuberculosis, which is considered in our tables.

The standard deviation of the mean weight is 14.95 pounds for the first million, 14.36 for the second million, and 14.74 for the two combined. This is about 2.75 to 3 pounds below the standard deviation for the whole population. This small standard deviation is partly in consequence of the reduced mean weight, but largely because the men with pulmonary tuberculosis practically all show loss of weight, and relatively few of them show a deviation in the positive direction. They are mostly men of low mean weight, and show comparatively little variation therefrom.

The relation between the distribution of weights of the population of men with pulmonary tuberculosis and the population of recruits in general is graphically shown in Plate XXXIV. This curve brings out strikingly the great weight deficiency of men with pulmonary tuberculosis, and this is the more striking in view of the fact that they have a stature that is above the average. The modal weight is about 10 pounds below the average, and there is almost an entire absence of the greater weights, above 185 pounds.

(c) *Chest circumference.*—In the 10,649 men found having pulmonary tuberculosis at mobilization camps the average chest circumference is 32.09 inches, or 1.13 inch less than the average for the whole population examined. For the first million men the average chest circumference is 32.33 inches; for the second million 31.90 inches. That for the second million is nearly half an inch less than that for the first million. This is a somewhat remarkable result in view of the fact that the men of the second million are taller than those of the first and indicates that the men with tuberculosis in the second million were much slenderer than those of the first million. These facts show that, as ordinary observation confirms, persons with pulmonary tuberculosis tend to have relatively small chest circumference despite their great stature. The low variability suggests that the small chest circumference is not necessarily the consequence of tuberculosis, for if it were, we should have persons with large chest circumference who were beginning to show signs of pulmonary tuberculosis, and those with small chest circumference in whom the disease had progressed far. Consequently were the small chest circumference merely caused by pulmonary tuberculosis, variable chest circumference would be expected. On the other hand, the low variability suggests that the small chest circumference is a constitutional trait; that is, those in whom the chest developed inadequately are apt to acquire active symptoms of tuberculosis, or, to put it in another way, persons with a tuberculous diathesis are characterized from youth on by small development of the chest, as well as by tall stature.

The relation between the distribution of chest circumference in the population found with pulmonary tuberculosis and the population of recruits in general is shown graphically in Plate XXXVII. It appears that the chest girth of the population with tuberculosis is far below that of the population of recruits in general.

(d) *Robustness.*—The index of build, as determined by using the second power of the height as a divisor, for the group of men with pulmonary tuberculosis is 28.15, which is 2.82 units below the average of the United States. This, as stated above, is the smallest index of build of any of the groups of defects, except that of underweight.

Pignet's ²⁰ index of robustness for men with pulmonary tuberculosis is 30.27. This brings the group into Pignet's class of very weak constitution. For each inch of the average height there are 1.92 pounds of weight as compared with the normal 2.097, and 0.472 inch of chest measure (expiration) as compared with the normal 0.492. In summary, the average tuberculous subject is tall, narrow chested, and underweight.

TABLE 139.—Correlation between height and weight in recruits with tuberculosis (pulmonary), first (P₁) and second (P₂) million draft recruits.

Height, in inches.	Total	Weight, in pounds.																							
		89 and under.	90-94	95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204
58 and under...	9	1											1												
59	14		1								3	3	2												
60	25	1	2								3	5	3	1											
61	35		3								6	8	6	4											
62	122		2								26	38	25	14											
63	240		4								68	119	111	4											
64	805		2								123	209	102	54											
65	783		1								166	338	184	134											
66	1,196		1								215	406	255	222											
67	1,444		1								283	466	333	283											
68	1,594										255	466	333	283											
69	1,486										270	406	255	222											
70	1,242										270	406	255	222											
71	868										331	406	255	222											
72	568										51	130	101	80											
73	263										21	49	27	11											
74	135										11	11	11	3											
75	53										7	7	7	4											
76	23										4	4	4	2											
77	5																								
78	8																								
79	1																								
80 and over...	1																								
Total.	10,701	3	9	167	372	716	1,186	1,462	1,468	1,460	1,142	913	642	435	235	167	133	47	29	13	9	5	8	2	

P₁—Number of cases, 4,853.
 Height: Mean, 68.01 inches, standard deviation, 2.70±0.02
 Weight: Mean, 131.77 pounds, standard deviation, 14.95±0.11 pounds.
 Correlation: 0.4551±0.0076.

P₂—Number of cases, 9,048.
 Height: Mean, 68.12 inches; standard deviation, 2.78±0.02
 Weight: Mean, 129.42 pounds; standard deviation, 14.36±0.09 pounds.
 Correlation: 0.5533±0.0060.

standard deviation, 2.74±0.01
 Weight: Mean, 130.44 pounds; standard deviation, 14.74±0.07 pounds.
 Correlation: 0.4734±0.0080.

TABLE 140.—(Correlation between height and chest circumference (expiration) in recruits with tuberculosis (pulmonary), first (P_1) and second (P_2) million draft

Height, in inches.	Total.	Chest, in inches.															
		28 and under.	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43 and over.
38 and under	9		1	3	3	2	1	2	1		1						
39	12	6	7	11	17	11	11	3	2		1						
40	34	12	17	25	38	26	21	11	1								
41	64	22	32	41	50	41	33	18	13	4							
42	121	41	59	88	102	104	61	33	19	5							
43	236	82	122	188	223	223	151	107	24	11							
44	502	147	240	388	479	479	212	111	44	20	6						
45	782	222	392	600	818	818	247	123	65	32	6						
46	1,192	333	599	921	1,211	1,211	374	171	94	41	14						
47	1,435	413	744	1,100	1,435	1,435	448	225	123	53	14						
48	1,579	514	909	1,264	1,654	1,654	514	265	141	89	36						
49	1,498	477	844	1,176	1,498	1,498	477	248	141	90	36						
50	1,237	377	644	943	1,237	1,237	377	200	127	42	11						
51	884	266	444	644	884	884	266	135	80	30	11						
52	566	177	299	444	566	566	177	114	44	20	3						
53	282	88	144	211	282	282	88	54	28	10	3						
54	135	44	77	111	135	135	44	28	14	6	3						
55	53	17	28	44	53	53	17	11	4	2	1						
56	23	7	11	17	23	23	7	6	3	1	1						
57	5	1	1	1	5	5	1	1	2	1	1						
58	1	1	1	1	1	1	1	1	1	1	1						
59	1	1	1	1	1	1	1	1	1	1	1						
60	1	1	1	1	1	1	1	1	1	1	1						
61	1	1	1	1	1	1	1	1	1	1	1						
62	1	1	1	1	1	1	1	1	1	1	1						
63	1	1	1	1	1	1	1	1	1	1	1						
64	1	1	1	1	1	1	1	1	1	1	1						
65	1	1	1	1	1	1	1	1	1	1	1						
66	1	1	1	1	1	1	1	1	1	1	1						
67	1	1	1	1	1	1	1	1	1	1	1						
68	1	1	1	1	1	1	1	1	1	1	1						
69	1	1	1	1	1	1	1	1	1	1	1						
70	1	1	1	1	1	1	1	1	1	1	1						
71	1	1	1	1	1	1	1	1	1	1	1						
72	1	1	1	1	1	1	1	1	1	1	1						
73	1	1	1	1	1	1	1	1	1	1	1						
74	1	1	1	1	1	1	1	1	1	1	1						
75	1	1	1	1	1	1	1	1	1	1	1						
76	1	1	1	1	1	1	1	1	1	1	1						
77	1	1	1	1	1	1	1	1	1	1	1						
78	1	1	1	1	1	1	1	1	1	1	1						
79	1	1	1	1	1	1	1	1	1	1	1						
Total	10,649	126	628	1,368	2,026	2,562	1,918	1,223	617	290	89	22	14	2	1	1	2

P_1 —Number of cases: 4,627.
 Height: Mean, 63.02 inches; standard deviation, 2.89 \pm 0.02 inches.
 Mean, 32.33 inches; stand-
 dard deviation, 2.73 \pm 0.01
 Mean, 32.08 inches; stand-

TABLE 139. — *Correlation between height and weight in recruits with tuberculosis (pulmonary), first (P₁) and second (P₂) million draft recruits.*

Height, in inches.	Total.	Weight, in pounds.																								P ₁ — Number of cases: 4,633. Height: Mean, 68.01 inches; standard deviation, 2.70±0.02 Weight: Mean, 131.77 pounds; standard deviation, 14.95± 0.11 pounds. Correlation: 0.4551±0.0078.												P ₂ — Number of cases: 6,048. Height: Mean, 68.12 inches; standard deviation, 2.76±0.02 Weight: Mean, 129.42 pounds; standard deviation, 14.36± 0.09 pounds. Correlation: 0.5533±0.0060.												P ₁ and P ₂ — Number of cases: 10,701. Height: Mean, 68.07 inches; standard deviation, 2.74±0.01 Weight: Mean, 130.44 pounds; standard deviation, 14.74± 0.07 pounds. Correlation: 0.4754±0.0060.																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																	
		89 and under.	90- 94	95- 99	100- 104	105- 109	110- 114	115- 119	120- 124	125- 129	130- 134	135- 139	140- 144	145- 149	150- 154	155- 159	160- 164	165- 169	170- 174	175- 179	180- 184	185- 189	190- 194	195- 199	200- 204	89 and under.	90- 94	95- 99	100- 104	105- 109	110- 114	115- 119	120- 124	125- 129	130- 134	135- 139	140- 144	145- 149	150- 154	155- 159	160- 164	165- 169	170- 174	175- 179	180- 184	185- 189	190- 194	195- 199	200- 204																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																		
58 and under	9	1										1	1				1																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																		

TABLE 140.—Correlation between height and chest circumference (expiration) in recruits with tuberculosis (pulmonary), first (P_1) and second (P_2) million draft registrants.

Height, in inches		Chest, in inches.														
Total.	28 and under.	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43 and over.
55 and under	9	1														
56	12	3	4	3	2	1				1						
57	34	6	7	4	5	2	3									
58	64	2	11	17	11	5	3		1							
59	121	6	29	25	18	21	8									
60	236	2	32	41	50	33	18									
61	782	14	60	89	102	61	38									
62	1,192	20	74	158	179	107	87									
63	1,435	13	92	210	223	212	111									
64	1,579	14	100	211	318	317	171									
65	1,469	17	96	206	330	328	171									
66	1,237	7	64	176	275	285	178									
67	684	6	26	70	219	248	172									
68	596	2	4	32	122	200	141									
69	282	2	4	13	63	135	114									
70	185	2	5	15	30	56	54									
71	123	2	12	16	16	30	18									
72	23	2	3	8	11	6	8									
73	3	1	2	2	3	6	3									
74	1				2	2	2									
75																
76																
77																
78																
79																
Total.	10,649	126	628	1,358	2,026	2,283	1,915	1,233	617	290	89	22	14	2	1	2

P_1 —
 Number of cases: 10,649.
 Height: Mean, 68.07 inches; standard deviation, 2.73 ± 0.01 inches.
 Chest circumference (expiration): Mean, 32.00 inches; standard deviation, 1.85 ± 0.01 inches.
 Correlation: 0.3412 ± 0.0082 .

P_2 —
 Number of cases: 10,649.
 Height: Mean, 68.07 inches; standard deviation, 2.73 ± 0.01 inches.
 Chest circumference (expiration): Mean, 32.00 inches; standard deviation, 1.85 ± 0.01 inches.
 Correlation: 0.3412 ± 0.0082 .

2. SIMPLE GOITER.

The dimensions of men who show simple goiter are of very great interest, because goiter is a disease of the thyroid gland and the secretions of this gland are believed to have important relations to the growth of the body. It is commonly accepted that persons with a thyroid that is especially active during early years of development tend to a large stature, whereas those with less active thyroid secretions remain relatively short. The tall groups are relatively slender, the short groups relatively stout. It is to be noted, however, that goiter usually first makes its appearance after adolescence, when the form of the body is already established, and hence its influence is less than though the disturbance of the thyroid gland occurred at an earlier age.

(a) *Stature*.—The mean stature of 7,099 men found at mobilization camps among the first and second million to have goiter is 67.94 inches, which is 0.45 inch greater than the average stature of the first million men, as indicated in Table I. Thus the men with hypertrophied thyroid gland show themselves to be nearly half an inch taller than the average. Since enlargement of the thyroid gland, in its early stage at least, may possibly be accompanied by an excessive secretion, the tall stature of the goitrous cases may be in part due to this excessive secretion. On the other hand, it must be recognized that the persons affected with goiter belong especially to the taller races in the United States. As pointed out in another publication,⁹ goiter is found especially in the extreme northwest, in Washington and Oregon, and in the region of the Great Lakes. The extreme northwest is certainly characterized by tall stature, and in the States of Wisconsin and Minnesota, both States with a high proportion of goiter, there are many representatives of the Scandinavians and Germans, who belong to the taller races. So it is impossible to ascribe the exceptional height of men found with goiter exclusively, if at all, to the hypersecretion of the thyroid gland. The standard deviation in stature of the goitrous cases is 2.58, which is slightly less than the standard deviation of all statures (2.71) shown in Table I. This restricted variability of stature indicates that the goitrous population is selected for high stature more than the population as a whole, and this is because the goitrous localities contain a rather homogeneous population of tall men as compared with the population of the United States as a whole. Indeed a comparison of the distribution of statures in men with simple goiter, as shown in Table 141, with the distribution in Table I (which shows the distribution of statures for the unselected population), reveals a marked deficiency in the lower statures and a corresponding excess in the tall statures. The mean height of men found with goiter in the second million is practically the same as the first million—namely, 67.95—so that there was no important change in the stature of the men selected for this disease in the two periods of examination.

The relation between the distribution of height of men with simple goiter and its distribution in the population at large is shown in Plate XXXI. It appears at a glance that the men with goiter are markedly taller than the population at large. This is probably because such tall men have come to inhabit the goitrous districts, or that the taller races, such as Scandinavians, are more often affected.

(b) *Weight*.—Of the 7,099 men in whom simple goiter was found at mobilization camps among the first and second million recruits, the average weight is 142.36 pounds. The average weight is only 0.82 pound above the average for the whole population, which is about six times the probable error. The index of build is 30.84, which is slightly less than that of the first million men as a whole. The population with goiter is a tall and slender one. The slight deficiency in build is, however, probably no greater than the deficiency in build that characterizes tall men in general.

The standard deviation of the mean weight is 16.50 pounds for the first and second million men. This standard deviation is 0.92 pound less than the standard deviation of the whole population of the first million which is not a very significant difference. The weight and standard deviation for the second million are not significantly different from those of the first million.

The relation between the distribution of weights in the population of men with simple goiter and that of recruits in general is shown in Plate XXXIV. The graph shows that the population with simple goiter is a heavy population as compared with the population of recruits in general. This is, however, associated with the great stature of the population with simple goiter, the significance of which has been referred to in the preceding paragraph.

(c) *Chest circumference*.—In the 7,085 men found with goiter among the the first and second million the average chest circumference is 33.11 inches, or 0.11 below the average of the whole population, as shown in Table II. The standard deviation of this dimension is 1.95, or about 0.06 inch below the average of the whole population. The average chest circumference for this group in the second million men is 33.13, which is slightly greater than for both million men, and is slightly less for the first million, 33.04.

The relation between the distribution of chest circumference in the population found with simple goiter and that of the recruits in general is shown graphically in Plate XXXVII. There is no great difference between the two distributions, though there is a slight inferiority in chest girth in the case of the goitrous population, and this is more striking in view of the large stature of this population.

(d) *Robustness*.—The index of build of men with simple goiter is 30.84, or 0.23 unit below that of the average for the United States. Pignet's index of robustness for this group is 21.94, which places them in the class of medium constitution. For each inch of the average height there are 2.10 pounds of weight, as compared with the normal 2.097; and 0.487 inch of chest measurement (expiration), as compared with the normal 0.492.

TABLE 142.—Correlation between height and chest circumference (expiration) in recruits with goiter (simple), first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Chest, in inches.												Total.		
	28 and under.	29	30	31	32	33	34	35	36	37	38	39		40	41
58 and under	4														
59	10														
60	9														
61	23														
62	10														
63	164	4	10	16	40	15	20	7	3	1	1	1			
64	307	7	22	31	50	30	33	24	19	4	3	3			
65	527	2	6	54	85	125	68	90	83	35	9	4	4		
66	571	2	14	40	62	133	168	126	83	19	10	6	2		
67	1,037	5	21	79	143	226	228	179	95	60	15	4	1		
68	1,338	1	22	56	122	222	280	211	123	67	19	4	1		
69	1,021	3	11	41	115	219	192	173	127	73	38	11	7		
70	800	2	3	33	81	162	158	108	119	69	43	10	3		
71	512		15	41	111	101	104	70	53	22	14	3	1		
72	305		6	15	52	57	65	41	23	13	8	3	1		
73	143		1	5	18	33	33	21	21	7	7	1	2		
74	61			2	8	9	12	16	6	3	2	2	1		
75	26				2	4	2	2	1	1	1	1	1		
76	14														
77	7														
78	6														
79	1														
Total..	7,085	26	109	438	881	1,333	1,454	1,255	808	444	202	87	38	7	3

P_1 —Number of cases, 1,509
Height: Mean, 57.94 inches; standard deviation 2.51 ± 0.03 inches.
Chest circumference (expiration) Mean, 33.04 inches; standard deviation, 1.94 ± 0.02 inches.
Correlation: 0.2162 ± 0.0151 .

P_2 and P_3 —
Height: Mean, 57.94 inches; standard deviation 2.59 ± 0.02 inches.
Chest circumference (expiration) Mean, 33.13 inches; standard deviation, 1.94 ± 0.02 inches.
Correlation: 0.2162 ± 0.0151 .

Standard deviation, 2.53 ± 0.01
Mean, 33.11 inches; standard deviation, 1.94 ± 0.02 inches.

3. EXOPHTHALMIC GOITER.

Exophthalmic goiter is the name applied to a set of symptoms that accompanies hypersecretion of the thyroid gland. It is characterized by some swelling of the gland itself, which, however, may be very slight, and it induces a rapid heart beat, cardiac hyperirritability, and protrusion of the eyeballs in advanced cases. The geographical distribution indicates that the exogenous causes that induce it are similar to those which induce simple goiter.

(a) *Stature*.—The average stature of 2,620 men found in mobilization camps with exophthalmic goiter, among the first and second million men examined, is 67.97 inches, which is 0.48 inch greater than the average of the first million men, as indicated in Table I. This excess of stature is about the same as for simple goiter and is to be explained on similar grounds, especially on the ground of high racial stature found in the population subject to it or inhabiting the goitrous districts. The average stature of the 439 cases of goiter found among the first million men is 67.94 and for the 2,181 in the second million, 67.97. The standard deviation of men with exophthalmic goiter is 2.65 inches, or 0.06 inch less than the standard deviation of all the first million men measured. The low standard deviation is due to the fact that exophthalmic goiter is especially prevalent in regions occupied by tall men.

A comparison of the statures of men with exophthalmic goiter as compared with the population at large is given in Plate XXXI. This shows that on the whole the selected population with exophthalmic goiter is strikingly taller than the population at large. This is probably because tall persons either have come to inhabit the regions especially subject to exophthalmic goiter or are more subject to the disease.

(b) *Weight*.—Of the 2,620 men found at mobilization camps, among the first and second million, with exophthalmic goiter the mean weight was 138.82 pounds, or 2.72 pounds below the average for the first million men. This difference indicates that exophthalmic goiter has some influence upon the weight. The standard weight associated with 68 inches of stature is 142.6 pounds. The men found with exophthalmic goiter were, therefore, 3.72 pounds below the average of men of their stature. This difference indicates that patients with exophthalmic goiter are slenderer than the men of their size; and it is not improbable that this reduction in weight is due to the disease. It is noteworthy that the correlation between stature and weight is 0.516 for simple goiter, and only 0.476 for exophthalmic, indicating that stature and weight are more closely associated in simple goiter than in exophthalmic. The index of build for men of exophthalmic goiter is 30.05, as contrasted with the index of 30.84 for men of simple goiter. This leads to the conclusion that men with exophthalmic goiter are of slender build, probably in consequence of the disease itself.

The relation between the distribution of weights in the population with exophthalmic goiter and that of the population of recruits in general is shown graphically in Plate XXXIV. The mode is about 132 pounds, which is 5 pounds below the mode of recruits in general. In view of the fact that persons

with exophthalmic goiter have on the whole a stature greater than the average, this suggests that a large proportion of persons afflicted with the disease are underweight because of the effects of the disease, and this would seem to be an explanation of the marked excess of persons with exophthalmic goiter having weights between 132 and 112 pounds. The irregularity in the curve at 142 pounds, or 5 pounds above the average, is possibly due to some error in recording or in tabulating. The cases are too few to give satisfactory averages.

(c) *Chest circumference.*—Of the 2,622 men found with exophthalmic goiter among the first and second million, the average chest circumference is 32.85 inches, or 0.37 inch less than the average of the whole population, as shown in Table II, and this despite the fact that the men with exophthalmic goiter are taller than the average. The relation of chest circumference to height is 0.483, which is less than 0.487 in the case of simple goiter and much less than 0.492 in the population as a whole. This again leads to the conclusion that men with exophthalmic goiter are a slender, small-chested type. The standard deviation of chest circumference is 1.98, which is a relatively small standard deviation. The small size of this standard deviation is partly due to the small absolute size of the chest, but in part is probably due to the effect of the disease itself.

The relation between the distribution of chest circumferences in the population with exophthalmic goiter and in the population of recruits in general is shown graphically in Plate XXXVII. This shows a marked deficiency in chest girth of the population with exophthalmic goiter, despite the fact that it is, on the whole, above the average in stature, and supports the conclusion that exophthalmic goiter results in malnutrition.

(d) *Robustness.*—The index of build of men with exophthalmic goiter is 30.05, which is 0.79 unit less than that of the group with simple goiter and 1.02 units less than the average for the United States. Pignet's index of robustness for this group is 24.28. This index places men with exophthalmic goiter in Pignet's group of medium constitution. For each inch of the average height there are 2.04 pounds of weight as compared with the normal 2.097, and 0.483 inch of chest as compared with the normal 0.492.

TABLE 144.—Correlation between height and chest circumference (expiration) in recruits with exophthalmic goiter, first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Total.	Chest, in inches.															
		28 and under	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43 and over.
58 and under	1																
59	5																
60	6																
61	10																
62	28																
63	71																
64	120																
65	187																
66	304																
67	378																
68	397																
69	382																
70	306																
71	210																
72	105																
73	63																
74	23																
75	9																
76	6																
77	2																
78																	
79																	
80 and over	1																
Total	2,622	6	69	209	380	540	503	392	287	138	57	23	11	3	3		1

P_1 —Number of cases, 432.
 Height: Mean, 67.64 inches, standard deviation, 2.53 ± 0.09 inches.
 Chest circumferences (expiration): 33.01 inches; standard deviation, 1.91 ± 0.04 inches.
 Correlation: 0.2469 ± 0.0072 .

P_2 —Number of cases, 432.
 Height: Mean, 67.64 inches, standard deviation, 2.57 ± 0.03 inches.
 Chest circumferences (expiration): 32.82 inches; standard deviation, 1.91 ± 0.02 inches.
 Correlation: 0.2469 ± 0.0072 .

Standard deviation, 2.65 ± 0.02 inches.
 Mean, 32.85 inches; 305.

TABLE 145.—Correlation between height and weight in recruits with myopia, first (P_1) and second (P_2) million draft recruits.

Height, in inches.		Weight, in pounds.																														
		89 and under.	90-94	95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204	205-209	210-214	215-219	220-224			
58 and under.	3						2		1																							
	10				3	2																										
	59				1		1	3		2																						
	60				1		1	3		2																						
	32				2		10	4	4																							
	58				2	5	8	12	8	5	3	2	1																			
	62						4	4																								
	63						8	14	16	14	3	4	3	4																		
	108						18	20	36	30	16	11	12	7	2																	
	135						20	20	36	30	16	11	12	7	2																	
	154						24	25	45	33	49	39	26	17	10	5	4															
	185						1	4	9	24	23	36	41	32	24	12	14	8														
	200																															
	209																															
	348																															
	350																															
	386																															
	396																															
	235																															
	209																															
	270																															
	322																															
71																																
122																																
86																																
73																																
24																																
17																																
1																																
75																																
6																																
76																																
2																																
Total	2,420	1	14	40	101	143	210	255	316	282	246	194	172	134	97	57	51	30	22	15	13	10	10	2	3	1	1					

P_1 —Number of cases, 778.
Height: Mean, 67.23 inches; standard deviation, 2.83 ± 0.05 inches.

Weight: Mean, 140.23 pounds; standard deviation, 18.07 ± 0.31 pounds.
Correlation: 0.5121 ± 0.0178 .

P_2 —Number of cases, 1,642.
Height: Mean, 67.01 inches; standard deviation, 2.77 ± 0.03 inches.

Weight: Mean, 138.75 pounds; standard deviation, 18.61 ± 0.22 pounds.
Correlation: 0.4906 ± 0.0128 .

P_1 and P_2 —Number of cases, 2,420.
Height: Mean, 67.08 inches; standard deviation, 2.79 ± 0.03 inches.

Weight: Mean, 139.23 pounds; standard deviation, 18.45 ± 0.18 pounds.
Correlation: 0.4912 ± 0.0104 .

TABLE 146.—Correlation between height and chest circumference (expiration) in recruits with myopia, first (P₁) and second (P₂) million draft recruits.

Height, in inches.	Total.	Chest, in inches.														
		28 and under.	29	30	31	32	33	34	35	36	37	38	39	40	41	42
58 and under.	3															
59	10															
60	11															
61	32	1	1	2	2	3	4	1	2	3	1	1	1			
62	58	1	5	6	9	10	11	7	9	2	1	1	1			
63	108	2	8	15	16	15	12	13	9	3	4	1	1			
64	185	3	17	35	40	36	26	26	21	6	2	1	1			
65	304	7	31	66	54	51	62	44	24	16	2	1	1			
66	308	2	28	61	64	67	66	38	30	18	5	2	2			
67	347	1	5	29	61	62	70	45	34	26	11	9	3	1		
68	349	2	6	19	51	62	63	30	44	23	9	4	3	1		
69	255	2	6	10	31	45	45	20	37	13	8	4	2	1		
70	209		1	7	24	41	43	24	20	8	3	2	1	1		
71	132		1	4	10	23	22	13	17	14	1	2	1	1		
72	24		2	4	8	16	11	5	4	2	2	1	2	2		
73	17				1	4	5	3	4	1	1	1	1	1		
74	4					1	1	1								
75	3															
76	2															
77																
Total.	2,417	17	55	192	356	436	464	357	262	145	68	32	24	6	2	1

P₁—Number of cases: 776.
 Height: Mean, 67.23 inches; standard deviation, 2.93
 ±0.05 inches.
 Chest circumference (expiration): Mean, 33.13 inches;
 standard deviation, 2.12 ± 0.04 inches.
 Correlation: 0.2177 ± 0.0291.

P₂ and
 inches; standard deviation, 2.70
 (expiration): Mean, 32.66 inches;
 2.12 ± 0.02 inches.
 .01660.

4. MYOPIA.

Myopia, or short-sightedness, is a constitutional hereditary defect of the lens; not that all myopics are born so, but that there is in many persons a constitutional tendency for the eye to become myopic under the influence of bad conditions of life, especially such as lead to abuse of the eyes.

(a) *Stature*.—The mean stature of 2,420 men found at mobilization camps in the first and second million with myopia is 67.08 inches, or 0.41 inch below the average stature of Table 1 for the first million. The mean stature for the myopics among the first million men was 67.23, and the second, 67.01. This group and that of hyperopics constitutes the shortest groups associated with the various defects other than that of astigmatics. The reason for the short stature of men with myopia is not difficult to infer. They were not rendered short because of eyesight, but many of them belong to races which have an especial tendency toward developing the myopic condition in the environment in which they prefer to live. As shown in "Defects Found in Drafted Men," 1920, defective vision in general has an especially high rate in the eastern manufacturing sections of the country, which include many French-Canadian and Polish Jews. Errors of refraction, of which myopia is the most common of the specified types, occur especially in urban districts and reach a maximum in New York City, a city characterized by a large proportion of Polish Jews. Indeed, it is well known that this race, which is one of the shortest races in the United States, is especially liable to this defect. Thus in the British report upon physical examinations of men of military age, 1917-1918 (Ministry of National Service²⁵, Vol. 1, p. 107), it is said that the very large Jewish population of Leeds helps to swell the number of cases of myopia, etc. We may conclude, therefore, that the short stature of persons with myopia is due in part to the high incidence of this defect in persons of short race.

Plate XXXIII gives a comparison between the statures of men found with myopia and the population at large. It is apparent that the population with myopia consists of a group of short men, some of whom are Russian and Polish Jews, who have a tendency toward myopia and short stature.

(b) *Weight*.—Of 2,420 men found with myopia at mobilization camps among the first and second million examined, the average weight was 139.23 pounds, or 2.31 pounds below the average. For the first million men the weight is 140.23 pounds and for the second million 138.75 pounds. This low weight of men with myopia is, of course, associated with their low mean stature. They are light in weight as a whole, not because myopia affects the weight, but because the myopics are commoner among certain small races than in the population at large. The standard deviation of weight in men with myopia was for the first million 18.07, or 0.65 above the average of the whole first million. The standard deviation of weight in men with myopia among the second million is 18.61, which tends to raise the excess of the standard deviation. The high standard deviation (or index of variability) of the weight of the myopics is, like the high standard deviations in respect to stature, due to the fact that the myopics constitute a marked deviation from the normal distribution inasmuch as it is weighted with excess of men of short stature.

The relation between the distribution of weights of the population with myopia and that of the population of recruits in general is shown graphically in Plate XXXVI. From this graph it appears that the population with myopia is characterized by small weight as, indeed, it is by small stature. This result merely supports the conclusion reached above that men with myopia include a racial group of small persons.

(c) *Chest circumference.*—Of 776 men found with myopia at mobilization camps among the first million, the average chest circumference at expiration is 33.13 inches. In the 1,641 men among the second million the average chest circumference is 32.89 inches. For the two groups together, 2,417 men, the mean chest circumference is 32.97. This average is somewhat less than the average chest circumference of the first million men, 33.22; the smaller chest circumference of the myopic men is doubtless to be attributed to the large proportion of smaller men found among them. That the chest circumference is only slightly less than the average is due to the fact that just these shorter men have a relatively high chest circumference, in accordance with the generally greater robustness of shorter men.

The standard deviation of chest circumference is 2.12 for the first million men and 2.12 ± 0.02 for the second million men. The standard deviation for the myopic men among the first and second million combined is 2.12 ± 0.02 . Thus the standard deviation is considerably greater than the average, which is to be explained on the same ground as the greater standard deviation of stature and weight, namely, on account of the excess of small men with absolutely small chest circumference.

In general, then, the conclusion to be drawn concerning the dimensions of myopic men is that myopia is especially characteristic of certain small races (especially the Polish and Russian Jews).

The relation between the distribution of chest circumferences in the population with myopia and the population of recruits in general is shown graphically in Plate XXXVII. Here we see that the chest girth for the population with myopia is slightly less than that of recruits in general, which is no doubt due to the fact that the population with myopia contains an excess of individuals of small races.

(d) *Robustness.*—The index of build of men with myopia is 30.95, which is 0.13 below the average for the United States. Pignet's index is 21.52. The men of this group belong in the class with medium constitution. For each inch of the average height there are 2.08 pounds of weight, as compared with the normal 2.097; and 0.492 inch of chest measure (expiration), as compared with the normal 0.492.

5. HYPEROPIA.

(a) *Stature.*—The average stature of 188 men found with hyperopia at mobilization camps among the first million is 67.28 inches; among the second million, 67.03 inches for 781 men; or for the total of 969 the mean stature is 67.08 inches, which is 0.41 inch below the average of all. This indicates that the hyperopic group contains an excess of short men. This is probably, as in the case of myopia, due less to any influence that hyperopia has upon growth than to the circumstance that hyperopia occurs in men that belong to the short races.

TABLE 147.—Correlation between height and weight in recruits with hyperopia, first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Total.	Weight, in pounds.																							
		89 and under.	90-94	95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204
58 and under.	3																								
59	3																								
60	6																								
61	17																								
62	21																								
63	33																								
64	70																								
65	100																								
66	133																								
67	134																								
68	150																								
69	116																								
70	86																								
71	48																								
72	29																								
73	9																								
74	5																								
75	1																								
76	1																								
Total.....	969				2	7	38	50	77	114	118	120	133	94	71	37	35	28	17	13	4	2	3	3	3

P_1 —Number of cases: 188.
Height: Mean, 67.28 inches; standard deviation, 2.65 ± 0.09 inches.

Weight: Mean, 139.13 pounds; standard deviation, 17.23 ± 0.60 pounds.
Correlation: 0.4146 ± 0.0407 .

P_2 —Number of cases: 781.
Height: Mean, 67.08 inches; standard deviation, 2.73 ± 0.06 inches.

Weight: Mean, 138.98 pounds; standard deviation, 16.10 ± 0.27 pounds.
Correlation: 0.4596 ± 0.0190 .

P_1 & P_2 —Number of cases: 969.
Height: Mean, 67.08 inches; standard deviation, 2.72 ± 0.04 inches.

Weight: Mean, 138.96 pounds; standard deviation, 16.29 ± 0.26 pounds.
Correlation: 0.4511 ± 0.0173 .

TABLE 148.—Correlation between height and chest circumference (expiration) in recruits with hyperopia, first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Total.	Chest, in inches.																43 and over.
		28 and under.	29	30	31	32	33	34	35	36	37	38	39	40	41	42		
46 and under.....	5																	
46.....	1																	
47.....	6																	
48.....	17																	
49.....	21																	
50.....	38																	
51.....	70																	
52.....	100																	
53.....	132																	
54.....	184																	
55.....	150																	
56.....	116																	
57.....	88																	
58.....	48																	
59.....	29																	
60.....	9																	
61.....	5																	
62.....	1																	
Total.....	988		29	68	123	182	208	157	113	63	18	10	6	1			2	
Total.....																		

Standard deviation, 2.45 ± 0.09
Mean, 33.26 inches; stand-

ard deviation, 2.74 ± 0.06
Mean, 33.00 inches; stand-

ard deviation, 2.79 ± 0.04
Mean, 33.06 inches; stand-

The standard deviation in stature of hyperopics is 2.65 for the first million men and 2.73 for the second million men, or 2.72 ± 0.04 for the two combined. This is only slightly greater than the standard deviation of stature for the whole of the first million men; the difference is much less than the probable error. Apparently, even though the mean stature of the hyperopics is slightly depressed, they conform closely to the average distribution of frequencies and hence possess the average variability in stature, of the population in general. The normal variability oscillates about a low mode.

(b) *Weight*.—Of the 188 men found with hyperopia in mobilization camps among the first million men, the mean weight is 139.13 pounds, or 2.41 pounds below the mean weight of the first million. This deficiency in weight is doubtless associated with the small size of hyperopics. The mean weight for the hyperopics among the second million is 138.98 pounds. For the two groups together it is 138.96 pounds; 2.58 pounds below the mean weight of the whole of the first million. This low weight is again doubtless associated with the small mean stature. The variability of hyperopics is indicated by the standard deviation of 16.29 ± 0.25 , which is more than 1 pound below the standard deviation for the whole of the first million men. This markedly low standard deviation for weight indicates that we have in hyperopics a fairly homogeneous group of men of slightly less than normal weight.

The relation between the distribution of weights in the population found with hyperopia and the population of recruits in general is shown in Plate XXXIV. As the number of persons in this population is small, the irregularity of the curve of distribution is probably not significant. On the whole the curve of weights of persons with hyperopia falls below that of the population in general.

(c) *Chest circumference*.—In 188 men found with hyperopia at mobilization camps among the first million, the average chest circumference is 33.26 inches, or 0.4 inch above the average chest circumference of the whole first million. In the 781 men found with hyperopia in the second million, the average chest circumference is 33.00 inches. For the 969 men in both, the chest circumference is 33.05 inches, or 0.17 inch below the mean chest circumference for the whole of the first million men. This relatively small mean chest circumference is doubtless associated with the generally small size of men with hyperopia. The standard deviation of chest circumference of men with hyperopia among the first million is 2.03, for the second million 1.96 ± 0.03 ; for the total 968, 1.98 ± 0.03 ; a variability which again is slightly, but hardly less significantly, than the standard deviation of the first million men, which is 2.01.

We may conclude that the hyperopics, like the myopics, include an especially large proportion of short men; in fact, they constitute more nearly a distinct lot of short men than the myopics. It is probable that this also is a matter of race.

The distribution of chest circumference in the population with hyperopia and the population of recruits in general is shown graphically in Plate XXXVII. It appears that the chest girth is slightly less than that of the population of recruits in general, which is probably associated with the smaller average size of the population with hyperopia.

(d) *Robustness*.—Men with hyperopia have an index of build of 30.88, or slightly less than that of men with myopia, and 0.19 less than the average of the United States. Pignet's index is 21.44, which places them in the medium group. For each inch of the average height there are 2.07 pounds of weight as compared with the normal 2.097, and 0.493 inch of chest measurement expirations, as compared with the normal 0.492.

6. ASTIGMATISM.

(a) *Stature*.—The average stature of 517 men found with astigmatism at mobilization camps among the first million is 66.95 inches; for the 1,075 among the second million it is 67.13; for the two groups combined, 1,592 men had the mean height of 67.07, which is 0.42 below the mean stature of the whole population of the first million men. The stature of astigmatics among the first million is 0.54 inch below the average stature of men of the first million. This is certainly a significant difference. Indeed if one compares in Table 184 relative distribution of statures in the line labeled at the left "Astigmatism" with the bottom line of the table, it will be seen that the short statures, 62–66 inches, inclusive, are uniformly in excess, whereas the taller statures, 68 inches upward, are for the most part in deficiency. However, there are relatively few astigmatics among the very short men, 61 inches and under (except a few cases 58 inches and under). This deficiency in frequencies of statures 61 and 59 inches strikingly separates astigmatics from the myopics, which have an excess in these stature classes. The excess of myopics in the lower stature classes does not extend above 65 inches, whereas in the astigmatics the excess extends to 66 inches. Astigmatics form a group that is as short on the average as the myopics, but it does not include so many of the very short men. The standard deviation for the astigmatics found among the 2,000,000 men is 2.71, which is probably not significantly less than the standard deviation of myopics of 2.79 inches. This indicates that though the astigmatics are a short people they do not include so many of an extremely short race as do the myopics.

There are several possible explanations of this extraordinary deficiency in stature of men with astigmatism found in mobilization camps. First, the hypothesis may be entertained that astigmatism is especially common in cities and that the population in cities contains men of inferior nutrition and consequently shorter stature than those of rural districts. This hypothesis may be tested by comparing the statures of men of eastern manufacturing sections with those of the population at large. For the eastern manufacturing group the mean stature is 66.77 inches; for the population as a whole, 67.49. But it has been already pointed out that this deficiency of eastern manufacturing sections can not be ascribed merely to conditions of life in these sections, but doubtless to the fact that shorter races, immigrated from Europe, have remained in these sections. The stature of people from Chicago is 67.09, which is only 0.04 inch below the average of the whole country, and from Denver is 67.67, which is slightly greater than the average of the whole country. Recruits from St. Paul and Minneapolis average still higher, 67.83. It is clearly not urbanity, but race, that chiefly determines the smaller stature of

some cities. The association of astigmatic persons with cities is to be ascribed rather to the short races living therein than to the fact that conditions of life in cities may be bad for the eyes. Perhaps one may say that peoples with hereditary tendency toward astigmatism are more apt to develop the tendency in cities than when they live in rural districts.

The deficiency in stature in men found with astigmatism may be due to racial factors. It is indeed well known that defects of vision, including astigmatism, are exceptionally frequent in recruits coming from New York city ("Defects Found in Drafted Men," p. 366). The rate for errors of refraction is given for New York city as 68.8 per 1,000. It was, however, still greater in Boston, 73.6. The high rate of errors of refraction of the classified cases of which astigmatism, next to myopia, is the largest item, is, as pointed out, probably due to the exceptionally large number of Hebrews in the cities. However, astigmatism is less predominantly found among the Hebrews than myopia, and that is probably why Boston exceeds New York city in the proportion of errors of refraction. Possibly there are other short races which are peculiarly subject to astigmatism (as, for example, South Italians, French Canadians, and Portuguese) which may occur in greater proportion in Boston than in New York city.

We may conclude, therefore, that the association of short stature with astigmatism is an association of two independent traits which are both racial characteristics.

(b) *Weight*.—In 517 men found with astigmatism at demobilization camps among the first million, the average weight is 138.59 pounds; for 1,075 men in the second million, 139.43 pounds; and for 1,592 men in both groups together, 139.16, or 2.38 pounds below the average for the whole of the first million men. This deficiency is, of course, associated with generally smaller size of the men found to have astigmatism. The standard deviation of this weight is for the first million men 17.25 pounds; for the second million 16.87 pounds; and for both together it is 17.00 ± 0.20 , which is 0.42 of a pound below the average of the whole of the first million men, a difference which is not very significant, being only a little more than twice the probable error. It is, however, in line with the low standard deviation found in men with eye defects, indicating one or more short racial groups.

The relation between the distribution of weights in the population found with astigmatism and that of the population of recruits in general is shown in Plate XXXIV. It appears from this graph that the population with astigmatism has a weight that is below the average of the population in general, a condition which is associated with the small stature of many of them. The mode of the astigmatic population is 2 or 3 pounds less than that of the population at large and stands much higher than the average population. This indicates that astigmatics are less variable in weight than the average, although it appears that they are more variable in stature than the population at large. The conclusion is justified, that in the population with astigmatism there is an excess of small persons, doubtless belonging to one or more small races.

TABLE 149.—Correlation between height and weight in recruits with astigmatism, first (P_1) and second (P_2) million draft recruits.

Height in inches.	Total.	Weight in pounds.																									
		89 and under.	90- 94	95- 99	100- 104	105- 109	110- 114	115- 119	120- 124	125- 129	130- 134	135- 139	140- 144	145- 149	150- 154	155- 159	160- 164	165- 169	170- 174	175- 179	180- 184	185- 189	190- 194	195- 199	200- 204	205- 209	210- 214
58 and under.	2																										
59.	2																										
60.	5																										
61.	14																										
62.	41																										
63.	88																										
64.	129																										
65.	163																										
66.	230																										
67.	220																										
68.	228																										
69.	171																										
70.	131																										
71.	90																										
72.	43																										
73.	21																										
74.	9																										
75.	3																										
76.	1																										
77.																											
78.																											
Total.	1,592				8	14	64	83	134	161	216	213	172	144	113	85	58	40	35	19	8	10	7	3	3	1	1

P_1 —Number of cases: 517.
 Height: Mean, 66.95 inches; standard deviation, 2.77 ± 0.06 inches.
 Weight: Mean, 138.59 pounds; standard deviation, 17.25 ± 0.36 pounds.
 Correlation: 0.5452 ± 0.0208 .

P_2 —Number of cases: 1,075.
 Height: Mean, 67.13 inches; standard deviation, 2.68 ± 0.04 inches.
 Weight: Mean, 139.43 pounds; standard deviation, 16.87 ± 0.25 pounds.
 Correlation: 0.4121 ± 0.0171 .

P_1 and P_2 —Number of cases: 1,592.
 Height: Mean, 67.07 inches; standard deviation, 2.71 ± 0.03 inches.
 Weight: Mean, 139.16 pounds; standard deviation, 17.00 ± 0.20 pounds.
 Correlation: 0.4573 ± 0.0134 .

TABLE 150.—Correlation between height and chest circumference (expiration) in recruits with astigmatism, first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Total.	Chest, in inches.															
		28 and under.	29	30	31	32	33	34	35	36	37	38	39	40	41	42	
58 and under.	2						1			1							
59	2				1			1									
60	5					1	1	2	1								
61	14		1	1	2	5		4		1							
62	41		1	1	8	12	8	6	2	1	1						
63	88		2	12	16	14	20	11	10	2				1			
64	129		6	16	18	26	28	19	6	6	1	2	1				
65	162		5	14	31	35	30	20	17	6	3		1				
66	230		2	23	40	48	47	36	15	10	5	1	1	1	1		
67	220		3	14	25	60	36	36	20	14	5	5	1			1	
68	226		2	9	36	49	42	31	31	20	3		2	1			
69	170		4	3	18	25	48	26	21	12	7	2	2	2			
70	130			4	18	26	30	18	12	6	10	4	1	1			
71	90		2	6	10	20	11	17	7	9	4	1	2	1			
72	44			1	1	5	11	14	6	4	1	1					
73	20					2	5	7	3	2				1			
74	9			1		1		1	2	2	1						
75	3							1						1			
76	1											1					
77																	
78	1							1									
Total..	1,587	28	105	224	329	319	251	154	96	41	17	13	8	1		1	

 P_1 —Number of cases: 517.Height: Mean, 66.95 inches; standard deviation, 2.77 ± 0.06 inches.Chest circumference (expiration): Mean, 33.06 inches; standard deviation, 2.02 ± 0.04 inches.Correlation: 0.2515 ± 0.0278 . P_2 —Number of cases: 1,070.Height: Mean, 67.13 inches; standard deviation, 2.68 ± 0.04 inches.Chest circumference (expiration): Mean, 33.01 inches; standard deviation, 2.01 ± 0.03 inches.Correlation: 0.1641 ± 0.0201 . P_1 and P_2 —Number of cases: 1,587.Height: Mean, 67.07 inches; standard deviation, 2.71 ± 0.03 inches.Chest circumference (expiration): Mean, 33.03 inches; standard deviation, 2.01 ± 0.02 inches.Correlation: 0.1928 ± 0.0163 .

(c) *Chest circumference*.—Of the 517 men found with astigmatism at mobilization camps among the first million, the average chest circumference is 33.06 inches; that of the 1,070 astigmatics found in the second million is 33.01; and for 1,587 men in both together it is 33.03, or 0.19 inch less than the average mean chest circumference of the first million men. This small chest circumference is associated with the low average stature and weight. The standard deviation of the chest circumference is for astigmatics among the first million men, 2.02; for the second million, 2.01; and for both groups together, 2.01 ± 0.02 . This is the same as the standard deviation in chest circumference for the whole of the first million men and indicates that the astigmatics form, on the whole, quite as homogeneous a group as the population at large, although a group slightly below the average in size.

The relation between the distribution of chest circumference in the population found with astigmatism and the population of recruits in general is shown graphically in Plate XXXVII. It appears that the population with astigmatism has, on the average, a small chest circumference, which is no doubt associated with their prevailing small height and weight, owing to the fact that this part of this population contains an excess of small races.

(d) *Robustness*.—The index of build of men with astigmatism is 30.94, or only 0.13 unit below the average of the United States. This index of robustness (Pignet²⁰) is 21.38, which is close to that of men with hyperopia. For each inch of the average height there are 2.08 pounds of weight, as compared with the normal 2.097, and 0.493 inch of chest measurement (expiration), as compared with the normal 0.492.

It will be observed that the foregoing three groups of men with errors of refraction have all an index of build and robustness slightly inferior to the average of the United States. This inferiority is to be ascribed less to any influence of errors in refraction upon the body than to the fact that errors of refraction are especially marked in certain races, especially Polish and Russian Jews, who are physically less well developed than the average.

7. HYPERTROPHIC TONSILLITIS.

Enlarged tonsils of such degree as to warrant record were found in 23,732 men at mobilization camps among the first million, and 28,299 among the second million draft recruits.

(a) *Stature*.—The average stature of men found among the first million to be affected with hypertrophic tonsillitis is 67.47 inches, which is 0.02 inch below the average stature of the whole population. The average stature of men found in the second million to have hypertrophied tonsils is 67.48. For the two combined, 52,031 men, the average is 67.48, which is practically the mean stature. We may conclude that, so far as stature is concerned, men with hypertrophic tonsils are typical of the whole population. This indicates that there is probably no race that is especially subject to this disease, and that apparently it has not affected the body nutrition, and hence the development. The standard deviation of height in the two groups is 2.71 and 2.74, respectively, and for the two combined, 2.73. The index of variability is practically the same as for the population as a whole, which confirms the conclusion that hypertrophic tonsils are fairly uniformly distributed through the population, so far as stature is concerned.

The distribution of statures in the population with hypertrophic tonsillitis as compared with the whole population of recruits is indicated graphically in Plate XXXIII. The distribution of statures nearly coincides in the two groups, but there are more men slightly above mediocre stature than below in the tonsillitis population than in that at large.

(b) *Weight*.—Of 23,732 men found with hypertrophic tonsils among the first million at mobilization camps the average weight is 142.19, and among 28,299 men in the second million 141.46. Taking both groups together, 52,031, we have a mean weight of 141.79, which is 0.25 above the average weight of the whole of the first million men examined. This is a real difference, though not a large one. The standard deviation in weight is for the first million men, 17.77 pounds; for the second million, 17.84 pounds; and for the two combined, 17.80 ± 0.04 . This is an excess of 0.38 pound over the average for the whole population of the first million men, a difference which is about nine times the probable error, and hence is significant. This indicates that in respect to weight, men with hypertrophic tonsils are more variable than the average population and suggests that the group includes an excess of men whose weight is above and a group whose weight is below the average. By comparing the distribution of weights in the hypertrophic tonsil group with that of the totals in the last line of Table I we find that the commonest weight for both the total and the hypertrophic tonsil group is 137 pounds and that, though there is a larger proportion of men in the modal group among those with large tonsils

TABLE 151.—Correlation between height and weight in recruits with tonsillitis (hypertrophic), first (P_1) and second (P_2) million draft recruits.

Height, in inches.

54 and under.
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80 and over
Total

P_1 Number of cases: 22,732.
 Height: Mean, 67.47 inches, standard deviation, 2.71 \pm 0.01 inches.
 Weight: Mean, 142.19 pounds, standard deviation, 17.77 \pm 0.05 pounds.
 Correlation: 0.4938 \pm 0.0084.

P_2 Number of cases: 28,290.
 Height: Mean, 67.48 inches; standard deviation, 2.74 \pm 0.01 inches.
 Weight: Mean, 141.46 pounds; standard deviation, 17.34 \pm 0.05 pounds.
 Correlation: 0.5001 \pm 0.0080.

P_1 and P_2 —
 Number of cases: 51,021.
 Height: Mean, 67.48 inches; standard deviation, 2.73 \pm 0.01 inches.
 Weight: Mean, 141.79 pounds, standard deviation, 17.80 \pm 0.04 pounds.
 Correlation: 0.4792 \pm 0.0080.

TABLE 152.—Correlation between height and chest circumference (expiration) in recruits with tonsillitis (hypertrophic), first (P_1) and second (P_2) million draft recruits.

Height, in inches	Chest, in inches.															
	28 and under.	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43 and over.
Total.	35	1,057	3,108	6,384	9,616	10,435	8,790	5,887	3,340	1,629	791	554	84	43	21	37
58 and under.	35	1	5	4	10	5	7	2	1	1	1	1	1	1	1	1
59	30	12	27	37	37	47	32	15	8	6	2	6	1	1	1	1
60	1	3	61	80	105	90	71	28	16	14	2	10	8	1	1	1
61	490	36	113	195	238	175	125	62	24	24	16	23	14	1	1	1
62	82	40	133	330	415	352	278	160	74	67	25	48	4	1	1	1
63	1,958	115	190	543	683	606	443	223	147	114	53	51	5	1	1	1
64	3,208	133	280	757	923	994	712	487	258	182	100	82	11	6	1	1
65	5,094	24	100	908	1,206	1,318	970	615	445	258	120	90	18	10	1	1
66	6,398	156	503	908	1,435	1,567	1,333	855	557	249	110	80	14	8	1	1
67	7,597	158	491	886	1,435	1,567	1,333	855	557	249	110	80	14	8	1	1
68	7,652	8	128	390	890	1,435	1,333	855	557	249	110	80	14	8	1	1
69	8,747	74	329	685	1,227	1,304	1,094	671	394	213	88	48	10	3	1	1
70	4,917	63	229	685	1,227	1,304	1,094	671	394	213	88	48	10	3	1	1
71	3,250	20	103	454	812	1,017	1,017	655	319	153	77	35	2	1	1	1
72	1,906	1	38	256	491	648	655	319	153	77	47	17	2	1	1	1
73	939	7	15	66	101	193	189	153	77	47	26	16	2	1	1	1
74	397	6	6	21	43	68	87	75	45	26	16	2	1	1	1	1
75	142	2	3	3	16	21	35	28	6	10	7	2	1	1	1	1
76	55	1	1	2	4	2	12	6	1	5	3	1	1	1	1	1
77	22	1	1	1	2	2	2	2	1	3	1	1	1	1	1	1
78	11	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1
79	10	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1
80 and over.	1	1	2	2	3	1	2	4	1	1	1	1	1	1	1	1
Total.	51,985	1,057	3,108	6,384	9,616	10,435	8,790	5,887	3,340	1,629	791	554	84	43	21	37

P_1 —Number of cases: 23,712.
 Height: Mean, 67.47 inches; standard deviation, 2.70±0.01 inches.
 Chest circumference (expiration): Mean, 33.29 inches; standard deviation, 2.03±0.01 inches.
 Correlation: 0.2284±0.0042.

P_2 —Number of cases: 23,712.
 Height: Mean, 67.47 inches; standard deviation, 2.73±0.01 inches.
 Chest circumference (expiration): Mean, 33.18 inches; standard deviation, 2.03±0.01 inches.
 Correlation: 0.2284±0.0042.

than in the total, yet men of 177 pounds are likewise in excess among those with enlarged tonsils, and the same is true of all weights above 192 pounds. There is, therefore, a clear excess of very heavy men with hypertrophic tonsils, and this accounts at once for the high mean weight and the high standard deviation of such men. That inflamed and enlarged tonsils should be more prevalent in heavy (though not tall men) is a point which should attract the attention of the physiologist and pathologist and be of help in understanding the causes of this condition. As shown in "Defects Found in Drafted Men," (1920, p. 132), the States with the highest ratio of hypertrophic tonsils are West Virginia, Virginia, and Pennsylvania, containing a large proportion of tall men, especially the mountaineers of the first two named States. Men from these States were examined at one camp where special attention was paid to infections of the head and throat, and it seems probable that there were thus brought into the total an exceptionally large number of tall men recorded with tonsillitis. Another center of high incidence of tonsillitis comprised the States of Mississippi, Arkansas, Oklahoma, and the contiguous States of Louisiana and Alabama. Southern whites are known to show a high ratio of this disease. The southern agricultural whites at least are above the average in stature, and this again contributes to the result. Finally, exceptionally high rates for tonsillitis (more than double the average) were found in the mining, Indian, and Scotch sections of the country, in all of which the average weight is high. Tonsillitis may possibly be associated with conditions in the mining groups, but the same explanation would not hold in the case of groups occupying Indian reservations and the Scotch. The large amount of tonsillitis found in New Mexico, Colorado, and California may perhaps be associated with the large amount of tuberculosis found in these States, due to the immigration thither of persons with this disease, but that there is a causal relation between the two diseases must not be hastily concluded, both because the defect rate for tonsillitis in Arizona, in which the rate for tuberculosis is highest, is below the average, but also because men with tuberculosis have a weight far below the average, while those with tonsillitis have a weight slightly above the average.

The relation between the distribution of weights of the population with hypertrophic tonsillitis and that of recruits in general is shown in Plate XXXVI. The graph brings out strikingly the fact that the population with hypertrophic tonsillitis differs in weight, as indeed in stature, in no important respect from the population at large.

(c) *Chest circumference.* —In the 23,712 men found with hypertrophic tonsils at mobilization camps among the first million, the average chest circumference is 33.29 inches, or 0.07 inch above the average of the first million. The average chest circumference for the 28,273 men with tonsillitis among the second million is 33.08, and the average for the two lots, 51,985 men, is 33.18 inches, which is close to the average for the whole of the first million men examined (33.22). Despite the slight excess of weight of these men, therefore, we have a slight deficiency of chest circumference. It is doubtful, however, if this is significant. The standard deviation of chest circumference of men with tonsillitis among the first million was 2.03 ± 0.01 ; for the second million, 2.10 ± 0.01 ; and for the two groups it is 2.07 ± 0.004 , which is 0.06 above the standard deviation

of chest circumference for the whole. This indicates a slight lack of homogeneity in the chest circumference, suggestive of possibly two groups. There is a very slight excess in the proportion of men of 35 inches upward with hypertrophic tonsillitis, and a corresponding slight deficiency of men 32 inches and under.

The relation between the distribution of chest circumference in the population with hypertrophic tonsillitis and the population of recruits in general is shown graphically in Plate XXXVII. The two curves nearly coincide, as is the case also in height and weight, indicating that the population with hypertrophic tonsillitis is nearly a random sample of the whole population.

(d) *Robustness*.—The index of build of men with hypertrophic tonsillitis is 31.14, which is 0.07 above the average of the United States. Pignet's index is 20.85. Pignet's index places the men with hypertrophic tonsillitis in the class with good constitution. For each inch of the average height there are 2.10 pounds of weight, as compared with the normal 2.097, and 0.492 inch of chest measurement (expiration), as compared with the normal 0.492.

8. TACHYCARDIA, SIMPLE.

Exceptionally rapid heart beat without other indications of organic disease was assigned to this category.

(a) *Stature*.—Of the 447 men with this defect among the first million the average stature is 67.73 inches, and in the 1,700 men found with the defect among the second million it is 67.76 inches. Of both groups together, 2,147 men, the mean stature is 67.76, which is 0.27 inch above the average stature of the whole of the first million men. The average stature of men found with tachycardia among the first million men is 0.24 inch above the average of the whole. This excess in stature of men with tachycardia is of the same order as the excess stature of men with exophthalmic goiter, with which some cases of simple tachycardia are probably associated. As shown in "Defects Found in Drafted Men" (p. 137), the highest rate for tachycardia is found in the State of Michigan. High rates are found also in South Dakota, Washington, and Wisconsin. These are all States occupied by men of exceptionally tall stature, and they have, therefore, influenced the average stature of men found with tachycardia. Tachycardia is indeed found especially among the Scandinavian, German, and Finn sections, which are those in the central Northern States in the Great Lakes region. It seems clear that the tall stature of some of the men with tachycardia is due to thyroid disturbance, which is again due to the fact that some races of men of prevalingly tall stature are especially predisposed to goiter or have settled in the geographic districts in which goiter is induced. The standard deviation of stature of men found with simple tachycardia among the first million is 2.71 inches, among the second million 2.66. For both groups it is 2.68 ± 0.03 . The small standard deviation of the tachycardia group is possibly significant, indicating that there has been something of a selection of tall men and that the tendency to tachycardia is not uniformly distributed through all statures. This is shown also in Table 184 through a comparison of the rates in the line "Simple tachycardia" with the total rates at the bottom of the table. Here we see that the rates for tachycardia are abnormally high in men with stature of 69 inches and over, and abnormally low in men with stature of less than 69 inches.

TABLE 154.—Correlation between height and chest circumference (expiration) in recruits with tachycardia, first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Total.	Chest, in inches.													43 and over.	
		28 and under.	29	30	31	32	33	34	35	36	37	38	39	40		41
28 and under.	2	2	1					1								
29	2	2	2		2	2	4	1								
30	7	3	2	5	3	13	4	13	1							
31	16	6	6	10	13	31	11	21	3							
32	26	8	16	29	24	41	15	36	11							
33	67	23	39	70	52	83	45	77	22							
34	109	37	62	110	82	124	66	111	35							
35	187	64	104	187	133	200	107	186	60							
36	265	87	144	259	188	282	145	261	93							
37	309	99	171	311	214	335	165	306	110							
38	330	109	190	330	247	372	183	338	120							
39	350	117	208	350	272	406	202	368	133							
40	374	130	230	374	311	446	223	390	146							
41	399	150	260	399	330	477	257	418	160							
42	424	171	289	424	354	511	277	452	174							
43	449	190	310	449	374	540	292	481	186							
44	474	211	339	474	399	565	310	506	199							
45	499	230	360	499	424	590	329	531	211							
46	524	250	389	524	449	615	348	556	224							
47	549	271	419	549	474	640	367	581	236							
48	574	290	449	574	499	665	386	606	249							
49	599	310	479	599	524	690	405	631	261							
50	624	330	509	624	549	715	424	656	274							
51	649	350	539	649	574	740	443	681	286							
52	674	370	569	674	599	765	462	706	299							
53	699	390	599	699	624	790	481	731	311							
54	724	410	629	724	649	815	500	756	324							
55	749	430	659	749	674	840	519	781	336							
56	774	450	689	774	699	865	538	806	349							
57	799	470	719	799	724	890	557	831	361							
58	824	490	749	824	749	915	576	856	374							
59	849	510	779	849	774	940	595	881	386							
60	874	530	809	874	799	965	614	906	399							
61	899	550	839	899	824	990	633	931	411							
62	924	570	869	924	849	1,015	652	956	424							
63	949	590	899	949	874	1,040	671	981	436							
64	974	610	929	974	899	1,065	690	1,006	449							
65	999	630	959	999	924	1,090	709	1,031	461							
66	1,024	650	989	1,024	949	1,115	728	1,056	474							
67	1,049	670	1,019	1,049	974	1,140	747	1,081	486							
68	1,074	690	1,049	1,074	999	1,165	766	1,106	499							
69	1,099	710	1,079	1,099	1,024	1,190	785	1,131	511							
70	1,124	730	1,109	1,124	1,049	1,215	804	1,156	524							
71	1,149	750	1,139	1,149	1,074	1,240	823	1,181	536							
72	1,174	770	1,169	1,174	1,099	1,265	842	1,206	549							
73	1,199	790	1,199	1,199	1,124	1,290	861	1,231	561							
74	1,224	810	1,229	1,224	1,149	1,315	880	1,256	574							
75	1,249	830	1,259	1,249	1,174	1,340	899	1,281	586							
76	1,274	850	1,289	1,274	1,199	1,365	918	1,306	599							
77	1,299	870	1,319	1,299	1,224	1,390	937	1,331	611							
Total	2,143	79	168	330	411	447	311	209	104	37	24	9	6	3	4	1
deviation, 2.89±0.03 can, 32.81 inches;																

deviation, 2.72 ± 0.06
 mean, 32.79 inches.

deviation, 2.66 ± 0.03
 mean, 32.81 inches.

deviation, 2.68 ± 0.03
 mean, 32.81 inches.

The relation between the distribution of statures in the population with simple tachycardia and that of the population of drafted men in general is shown in Plate XXXIII. The graph shows at a glance that the population with simple tachycardia consists of men strikingly taller than the average. There are relatively fewer men with statures from 61 to 68 inches and relatively more men with statures 69 to 76 inches. The mode is shifted from 67½ to 69 inches. This shows that men with simple tachycardia are prevaillingly tall men. This result is, as stated, probably not due to the influence of tallness, but to the fact that simple tachycardia is in some cases associated with disturbances of the thyroid gland, and this in turn by conditions in those sections that are inhabited by tall races, largely the Scandinavians. However, the possibility that great size of the body may be responsible for rapid heart beat, apart from thyroid disturbance, must not be overlooked.

(b) *Weight*.—Of 447 men found with simple tachycardia in the first million examined at camps the average weight is 137.06 pounds, which is 4.48 below the average for the whole first million men. Of tachycardia cases among the second million, 1,700 men, the mean weight is 137.45 pounds; and for both combined, 2,147 men, it is 137.37 pounds, which is 4.17 pounds below that of all the first million men. This marked deficiency in weight, despite tall stature, must certainly be significant and suggests an insufficiency in metabolism. The standard deviation in weight of tachycardia cases in the first million men is 17.36; in the second million men, 17.63; and for both lots together, 17.57 ± 0.18 . The difference from the standard deviation for the average of the whole first million is only 0.15, or about once the probable error, so that the difference is probably not a significant one, and the group of tall but slender men, who are especially liable to tachycardia, constitutes a group which has nearly the same distribution about the mode as has the whole population.

The relation between the distribution of weight of the population found with simple tachycardia and the population of recruits in general is shown graphically in Plate XXXVI. This indicates that the population with tachycardia is below average weight. The irregularity in the curve is probably due to the small number of cases. This deficiency in weight of the population with tachycardia is the more striking in view of the fact that persons with the disease are on the whole taller than the average. The result is probably due to an insufficiency of nutrition caused by the condition itself.

(c) *Chest circumference*.—Of 447 men found with simple tachycardia at mobilization camps among the first million the average chest circumference is 32.79 inches, or 0.43 inch less than the average chest circumference of the whole first million men. In the 1,696 men found among the second million the average chest circumference is 32.81, and for the two lots together, 2,143 men, the average chest circumference is 32.81, which is 0.41 inch below the average. This low mean chest circumference of men with tachycardia is associated with their low weight. The standard deviation of chest circumference was for men of the first million 2.03, and for the second million 2.05, and for the two combined 2.04 ± 0.02 . This is only 0.03 inch above the average for the whole first million, a difference which is probably not significant, indicating that the chest circumference of the slender men was not more variable around the new mode than the population in general.

The relation between the distribution of chest girth in the population found with simple tachycardia and that of the population of recruits in general is shown in Plate XXXVIII. One sees that the population with simple tachycardia has a chest circumference which is below the average, corresponding with the low average weight, despite the high average stature. The slender form is probably due to the disturbance of nutrition consequent upon the disease.

(d) *Robustness*.—Men with simple tachycardia have an index of build of 29.92, which is 1.15 below the average index of build of recruits. Pignet's index is 24.50. It places such men among the worst of the groups with medium constitution. It appears, then, that men with simple tachycardia have inferior constitution. For each inch of the average height there are 2.03 pounds of weight, as compared with the normal 2.097, and 0.484 inch of chest measurement (expiration), as compared with the normal 0.492.

9. CARDIAC HYPERTROPHY.

(a) *Stature*.—An enlargement of the heart sufficient to warrant recording was found among the first million men at mobilization camps in 503 cases, the average stature being 67.68 inches, or 0.19 inch above the average of the stature of the first million men. For the 840 cases found among the second million the average stature is 67.79, and for the two groups, 1,343 men, 67.75, or 0.26 inch above the mean of the whole first million men. The excess in stature of men with cardiac hypertrophy is a little less than twice the probable error of the standard deviation of the height of the population and is possibly significant. It indicates that men of large stature had enlarged hearts, probably in part because the larger bodies throw more work upon the heart, which has to enlarge to meet the functional demand made upon it. At least it is probable that one class of cases of enlarged hearts belong to this category. The standard deviation of men with enlarged hearts of the first million is 2.86; among men of the second million it is 2.64; and for both groups together 2.73 ± 0.04 . The standard deviation of stature in the cases of cardiac hypertrophy is thus 0.02 inch more than the average for the whole of the first million men. The mode has moved to a higher level than found in the whole population, yet the distribution around this mode is typical of the whole population. The details of distribution of statures of men with cardiac hypertrophy are given in Table 155.

The relation between the distribution of stature in the population with cardiac hypertrophy and of drafted men in general is shown graphically in Plate XXXIII. It appears at once that men with cardiac hypertrophy are a taller group than that of the general population. This is shown by the deficiency of short men and the excess of tall men, especially of men from 69 to 74 inches. It is shown also by the fact that the mode is one-half inch above the average.

(b) *Weight*.—Of the 503 men found with cardiac hypertrophy among the first million examined at mobilization camps, the average weight is 139.23 pounds, or 2.31 pounds less than the mean weight of the whole of the first million men. The mean weight of 840 men with enlarged hearts found in the second million is 141.24, and of both lots, 1,343 men, 140.49. This is about 1 pound less than the average weight of the whole of the first million men.

TABLE 155.—Correlation between height and weight in recruits with cardiac hypertrophy, first (P_1) and second (P_2) million draft recruits.

		Weight, in pounds.																								
Height, in inches.																										
Total.																										
89 and under.	90-94	95-100	105-110	110-114	115-119	120-124	125-129	130-134	135-140	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204	205-210	210-214	215-219	220-224
58 and under.	1																									
59	7			1	1	1	1	1	1																	
60	2																									
61	6				1	1	1	1	1																	
62	14			1	1	2	4	6	9																	
63	42		1	2	3	6	10	6	7																	
64	74		1	2	7	12	15	12	23																	
65	129		2	11	8	10	22	23	15	16	10	3	4	5	3	2	2	1								
66	164		3	4	3	9	22	24	25	28	17	11	13	12	8	4	1	2								
67	177			1	4	5	15	23	29	35	27	22	14	9	6	3	5	2								
68	199				1	5	14	25	31	35	27	24	14	7	11	7	2	1								
69	167				1	1	3	13	17	20	24	26	17	14	11	7	3	2								
70	161					1	2	8	17	20	25	14	13	9	11	4	3	2								
71	93							3	15	5	5	8	7	4	4	8	2	1								
72	60							6	5	3	5	7	4	1	1	1	1									
73	28							2	3	2	3	1	1	1	2	1	1									
74	9																									
75	6																									
76	2																									
77	1																									
78	1																									
Total.....	1,343	7	12	34	53	101	153	157	182	159	140	95	80	60	42	25	14	10	6	3	2	3	1	2	1	1

P_1 —Number of cases: 503.
Height: Mean, 67.68 inches; standard deviation, 2.86±0.06 inches.

P_2 —Number of cases: 840.
Height: Mean, 67.79 inches; standard deviation, 2.64±0.04 inches.
Weight: Mean, 139.23 pounds; standard deviation, 16.76±0.36 pounds.
Correlation: 0.4576±0.0238.

P_1 —Number of cases: 840.
Height: Mean, 67.79 inches; standard deviation, 2.64±0.04 inches.
Weight: Mean, 141.24 pounds; standard deviation, 16.86±0.05 pounds.
Correlation: 0.4044±0.0195.

P_2 —Number of cases: 1,343.
Height: Mean, 67.75 inches; standard deviation, 2.73±0.04 inches.
Weight: Mean, 140.49 pounds; standard deviation, 16.85±0.22 pounds.
Correlation: 0.4252±0.0151.

TABLE 158.—Correlation between height and chest circumference (expiration) in recruits with cardiac hypertrophy, first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Total.	Chest, in inches.																43 and over.
		28 and under.	29	30	31	32	33	34	35	36	37	38	39	40	41	42		
58 and under.	1						1	1										
59	7		1				1											
60	2																	
61	6						3	3										
62	14		3	3			9	10	2	2	1							
63	42	1	3	9	21	12	20	14	3	2								
64	74		4	10	16	21	25	15	15	7	4							
65	129		6	15	23	28	36	24	16	10	6	1						
66	163	5	3	15	25	30	36	27	23	11	7	1						
67	176		3	14	28	30	36	32	26	11	7	1						
68	200		4	17	21	33	46	32	25	11	7	1						
69	167		6	17	21	33	38	29	16	10	6	1						
70	166	1	6	19	21	35	36	29	15	10	6	1						
71	48			6	10	21	19	17	10	6	4	1						
72	40			3	3	11	13	12	6	7	3	1						
73	97			2	2	4	4	6	2	3	2	1						
74	4							1	1	1								
75	6							1	1									
76	2																	
77	1																	
78	1																	
Total	1,330	7	32	91	181	248	276	233	135	80	34	6	8	3		2	1	

P_1 —Number of cases: 800.
 Height: Mean, 67.57 inches; standard deviation, 2.57 ± 0.05 inches.
 P_2 —
 Height: Mean, 67.57 inches; standard deviation, 2.57 ± 0.05 inches.
 Chest: Mean, 33.58 inches; standard deviation, 2.63 ± 0.04 inches.
 Correlation coefficient, $r = 0.04$ inches.

These men, then, are taller than the average and of slightly less weight. Their index of build is 30.61, as contrasted with 31.07, which is the index of robustness of the whole of the first million men. In other words, men with cardiac hypertrophy are prevailingly tall and slender. The standard deviation of the weight for the first million is 16.75 pounds, or about 0.67 less than the standard deviation in weight of the whole population of the first million men. For cases of cardiac hypertrophy among the second million the standard deviation in weight is 16.86, and for the two groups together it is 16.85 ± 0.22 . This is a standard deviation of 0.57 pound less than the average for the whole first million. It appears that men with cardiac hypertrophy are not only a slender group, but that they are less variable about this lower weight mode than the population in general. This suggests that either slender men are most apt on this account to have hypertrophied hearts or else, more probably, that the conditions which have led to enlarged hearts in these tall men have resulted in an abnormal diminution in weight.

The relation between the distribution of weights of the population found with cardiac hypertrophy and the population of recruits in general is shown in Plate XXXVI. On the whole this population is characterized by less than average weight and this despite the fact that the population contains more tall persons than the population at large. The principal mode is the same as for the population at large.

(c) *Chest circumference*.—Of the 500 men found with cardiac hypertrophy among the first million men examined at mobilization camps the average chest circumference is 32.88 inches, or 0.34 below the mean chest circumference of the first million men. For 839 men in the second million the chest circumference is 33.03. For 1,339 men in the two groups it is 32.97, or 0.25 below the mean chest circumference of the whole of the first million men. This low chest circumference is associated with low weight and confirms the conclusion that men with hypertrophied hearts are tall and slender people. The standard deviation of chest circumference is for men with enlarged hearts, among the first million, 2.02; among the second million, 1.99; and for both together, 2 ± 0.03 . This is very close to the standard deviation of the whole of the first million men and suggests that while the mean chest circumference is low yet the variations around this mode are those typical of the whole population. This result leads to the conclusion that the hypertrophied heart has caused a symmetrical reduction in chest circumference and weight in that part of the population which has been affected.

The relation between the distribution of chest girth in the population found with cardiac hypertrophy and the population of recruits in general is shown graphically in Plate XXXVII. It is obvious that the population with cardiac hypertrophy has on the whole a smaller chest circumference than the population in general and this is probably associated with the reduced weight which they show, probably as a consequence of the defect.

(d) *Robustness*.—Men with cardiac hypertrophy have an index of build of 30.61, or 0.46 below the average for the United States. Pignet's index is 22.66. Thus they are placed in the group with medium constitution. For each inch of average height there are 2.07 pounds of weight, as compared with the normal 2.097, and 0.487 inch of chest measurement (expiration), as compared with the normal 0.492.

10. MITRAL INSUFFICIENCY.

(a) *Stature*.—The average stature of 4,257 men found to have mitral insufficiency at mobilization camps out of the first million examined is 67.86 inches, or 0.37 inch above the mean stature of the first million men. The mean stature of 4,603 cases with mitral insufficiency out of the second million men is 67.82 inches; for both groups, 8,860 men, it is 67.84 inches, or 0.35 inch above the mean stature. It is clear that mitral insufficiency is found especially in tall men. If we examine the distribution of endocarditis and valvular diseases of the heart as given in "Defects Found in Drafted Men" (p. 133), we find that the highest rate occurs in the States of Washington, Utah, Michigan, Maryland, and others, including several States with exceptionally tall men. However, in Texas, in which the average stature is exceptionally high, the ratio of valvular diseases found is below the average. The standard deviation of stature of men found with mitral insufficiency out of the first million is 2.73; out of the second million, and for the combined group, it is the same. This standard deviation is not significantly different from that of the population at large. Thus the men with mitral insufficiency constitute a group with a high mode but with essentially the same distribution about that mode as a normal population. The causes then which have lifted the mode have acted similarly and in essentially uniform fashion upon "the run" of the population.

The relation between the distribution of stature in the population with mitral insufficiency and of drafted men in general is shown in Plate XXXIII. Here, as in cardiac hypertrophy, it is obvious that men with mitral insufficiency constitute a group of tall persons. This is shown by the regular deficiency of men below the mode in stature, by the regular excess of men above the mode and by the fact that the mode is $\frac{1}{2}$ inch above the modal stature of the population of drafted men.

(b) *Weight*.—The mean weight of 4,257 men found to have mitral insufficiency in mobilization camps of the first million examined is 139.11; in 4,603 from the second million, 138.87. The average of the total 8,860 cases is 138.99, which is 2.55 pounds below the mean weight for the whole population of the first million. This places men with mitral insufficiency below the average of the population. The index of build of men with mitral insufficiency is 30.20, which is decidedly less than that of the average for the whole first million men, 31.07. It appears then that men with mitral insufficiency are on the average tall and slender men, the same type of men we have seen to be affected with cardiac hypertrophy. Cardiac hypertrophy and mitral insufficiency are in a way correlated, for if the valves of the heart are inadequate then the muscles of the heart must make good the deficiency and this hyperactivity leads to increase in size of the muscles of the heart. The hydrostatic problem that the heart has to meet is increased by the increase in stature of the man.

The relation between the distribution of weights in the population found with mitral insufficiency and the population of recruits in general is shown in Plate XXXVI. This graph shows a small but constant inferiority in weight of persons found with mitral insufficiency and this despite the fact that they

TABLE 157.—Correlation between height and weight in recruits with mitral insufficiency, first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Total.	Weight, in pounds.																									
		80 and under.	90-94	95-100	105-110	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-200	205-210	215-220	225-230	
58 and under	8	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
59	9	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
60	30	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
61	52	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
62	115	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
63	276	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
64	459	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
65	732	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
66	1,034	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
67	1,244	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
68	1,350	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
69	1,464	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
70	1,584	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
71	1,716	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
72	1,860	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
73	2,016	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
74	2,184	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
75	2,364	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
76	2,556	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
77	2,760	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
78	2,976	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
Total	8,880	8	44	112	273	523	778	923	1,083	1,127	1,008	830	686	484	346	200	132	81	53	36	22	20	22	3	1	3	1

P_1 —Number of cases: 4,257.
Height: Mean, 67.86 inches, standard deviation, 2.73±0.02
Weight: Mean, 139.11 pounds, standard deviation, 16.94±0.12 pounds.
Correlation: 0.4890±0.0078

P_2 —Number of cases: 4,603.
Height: Mean, 67.83 inches, standard deviation, 2.73±0.02
Weight: Mean, 138.67 pounds, standard deviation, 16.94±0.12 pounds.
Correlation: 0.5029±0.0074

P_1 —Number of cases: 4,603.
Height: Mean, 67.83 inches, standard deviation, 2.73±0.02
Weight: Mean, 138.67 pounds, standard deviation, 16.94±0.12 pounds.
Correlation: 0.5029±0.0074

standard deviation, 2.73±0.01
ds; standard deviation, 16.79±

TABLE 158.—Correlation between height and chest circumference (expiration) in recruits with mitral insufficiency, first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Chest, in inches.																
	28 and under.	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43 and over.	
58 and under.....	8	2	2	2	2	2	1	3	1								
59.....	9	1	1	1	1	2	5	2	2								
60.....	29	3	4	4	7	10	3	4	4								
61.....	51	2	11	12	25	24	10	4	2								
62.....	115	4	16	14	57	50	23	11	6								
63.....	275	6	37	35	92	71	46	42	7	3	1	2					
64.....	459	6	54	105	177	143	99	37	26	3	2	2					
65.....	780	4	98	101	229	199	118	80	23	11	6	2					
66.....	1,032	5	113	200	272	253	169	108	57	34	9	7					
67.....	1,258	6	110	184	293	272	214	147	71	32	12	1					
68.....	1,376	13	85	156	222	231	194	98	66	36	16	4					
69.....	1,137	3	87	138	222	231	194	98	66	36	16	4					
70.....	978	1	47	115	164	209	164	110	73	37	13	7					
71.....	619	6	28	65	120	118	127	76	42	15	8	4					
72.....	398	3	12	41	64	78	87	68	35	9	5	3					
73.....	205	2	6	17	36	40	39	25	20	12	4	4					
74.....	100	2	2	6	10	18	26	18	13	4	3	3					
75.....	43	1	1	3	6	7	9	6	3	4	2	1					
76.....	19				1	2	8	3	2	4	2	1					
77.....	4					1	1	1	1	1							
78.....	5					1	3	1									
Total.....	8,830	57	310	733	1,271	1,801	1,713	1,356	823	459	199	83	33	5	2	1	4

Number of cases: 4,590.
 Height Mean, 67.63 inches; standard deviation, 2.73±0.02 inches.
 Chest circumference (expiration): Mean, 32.66 inches; standard deviation, 2.05±0.01 inches.
 Correlation: 0.2666±0.0091.

Standard deviation, 2.73±0.02
 Mean, 32.66 inches; stand-

Standard deviation, 2.73±0.01
 Mean, 32.76 inches; stand-

are men on the whole of a stature above the average. This result indicates that the population with mitral insufficiency is undernourished, probably in consequence of the valvular defect.

(c) *Chest circumference.*—The average chest circumference of 4,240 men found with mitral insufficiency in the first million is 32.86, and in 4,590 men in the second million 32.65. The average for the 8,830 in both groups is 32.75, which is 0.47 inch less than the average chest circumference. This small chest circumference is associated with the slender build which is, as we have seen, characteristic of the group with mitral insufficiency. The standard deviation of chest circumference for the combined groups is 2.0, which is essentially the same as that of the whole population. It appears then that, so far as chest circumference goes, if the mode has been diminished, the distribution about the mode is about the same as the mode of the whole population. It seems probable, therefore, that tall and short persons are affected in equal degree, so that the reduction in chest circumference of that part of the population with mitral insufficiency has affected them in equal proportion.

The relation between the distribution of chest girth in the population found with mitral insufficiency and the population of recruits in general is shown graphically in Plate XXXVIII. It appears at once that the population with mitral insufficiency has a chest girth strikingly below that of the population in general—a fact which is associated with their low average weight, despite the high average stature. This result is therefore probably due to malnutrition in consequence of the disease.

(d) *Robustness.*—Men with mitral insufficiency have an index of build of 30.20, or 0.87 below the average for the United States. Pignet's index is 24.12. Thus they fall into the group with medium constitution. For each inch of the average height there are 2.05 pounds of weight, as compared with the normal 2.097, and 0.483 inch of chest measurement (expiration), as compared with the normal 0.492.

11. MITRAL STENOSIS.

(a) *Stature.*—Of 1,521, in the first million men, affected with mitral stenosis, the mean height is 67.71 inches, which is 0.22 inch above the average stature for the first million men. The mean stature for 991 men in the second million, 67.50, is somewhat less than for the first million. For the 2,512 men in the two groups it is 67.63 inches, or 0.14 inch above the average. The standard deviation of stature of men with mitral stenosis is 2.72 for the first million, and 2.73 for the second, and 2.72 ± 0.03 for the two groups, which is about the same as the standard deviation of the whole population of the first million given in Table I.

The relation between the distribution of stature in the population with mitral stenosis and that of drafted men in general is shown in Plate XXXIII. This graph indicates that the population with mitral stenosis contains on the whole a slightly greater stature than the population of drafted men in general. However, the contrast is much less than the case of either mitral insufficiency or cardiac hypertrophy. The mode for the population with mitral stenosis is the same as that of the drafted men in general.

(b) *Weight*.—The weight of 1,521 men with mitral stenosis among the first million is 137.46; and for the 991 men among the second million, 135.93; and for the 2,512 in both groups, 136.85 pounds, which is 4.69 pounds below the average of the first million men. The standard deviation is extraordinarily low, being 15.24 for the first million men; 16.16 for the second million; and 15.63 ± 0.15 for the two groups, which is strikingly below the standard deviation for the population in general. This means that tall, slender men are prevalently affected with mitral stenosis. The reduced weight is not merely a consequence of the mitral stenosis, for if it were the standard deviation would be large. Rather the men with mitral stenosis are a selected lot of the population characterized by their tall and slender form.

The relation between the distribution of weights in the population found with mitral stenosis and that of the population of recruits in general is shown in Plate XXXVI. This graph shows clearly that the population with mitral stenosis is inferior in weight on the average to the population in general and this despite the fact that they are on the average slightly taller than the population of recruits in general. This reduction in weight is therefore probably due to imperfect development resulting from the disease.

(c) *Chest circumference*.—Of the 1,516 men found with mitral stenosis at mobilization camps among the first million men, the average chest circumference is 32.77 inches, which is 0.45 inch less than the average of the whole population, and of the 991 men found in the second million the average chest circumference is 32.47. Of 2,507 men in the two groups together the average is 32.65, which is 0.57 inch less than the average for the first million as shown in Table I. This small chest circumference accords with the evidence derived from weight that men with mitral stenosis are tall and slender.

The standard deviation of chest circumference is 1.89 for the two groups, which is 0.12 less than the standard deviation of the chest circumference of the population of Table II. This accords also with the small standard deviation for weight and suggests the conclusion that men with mitral stenosis are not a random sample of the population, but are (in part) a selected group, characterized by tall stature, small weight, and narrow chest circumference, and that their peculiarities are associated constitutionally, to at least a certain extent, with a diseased or defective condition of the valves.

(d) *Robustness*.—Men with mitral stenosis have an index of build of 29.93, or 1.14 below the average of the United States. This is the lowest index of build of the groups with heart defects excepting the group with simple tachycardia. Pignet's index of robustness is 24.81, which places it in the lower part of the medium group. For each inch of the average height there are 2.02 pounds of weight, as compared with the normal 2.097, and 0.483 inch of chest measurement (expiration), as compared with the normal 0.492.

TABLE 159.—Correlation between height and weight in recruits with mitral stenosis, first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Total.	Weight, in pounds.																							
		89 and under.	90-94	95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204
58 and under.	4																								
59	7			1							1	1	2	2											
60	8				1						2														
61	16				2																				
62	43																								
63	85																								
64	140			1	3	7	13	24	15	31	18	13	10	2	1										
65	212				1	7	12	17	23	37	34	34	16	9	5										
66	320				3	6	10	29	40	63	50	52	32	14	16	3	2								
67	367				2	4	10	27	49	48	70	49	40	25	15	9	6	10	3	1	1				
68	369				2	4	8	19	38	41	55	50	47	33	29	21	9	9	2	1	2				
69	321				2	1	2	8	19	36	45	59	38	30	33	16	14	2	4	3	2				
70	262						3	2	11	21	26	50	45	33	27	13	10	4	11	4	2				
71	186						1		7	3	17	24	21	21	25	21	14	3	3	2	2				
72	91								6	6	5	14	9	13	12	9	7	3	3	2	2				
73	53								1	3	3	4	4	12	6	4	6	6	2	1	1	1			
74	18										1														
75	4																								
76	1																								
77	1																								
78	2																								
79	2																								
Total	2,512			2	17	35	74	169	248	304	339	354	271	208	177	103	71	59	37	19	8	9	3	2	3

P_1	Number of cases: 1,521. Height: Mean, 67.71 inches; standard deviation, 2.72±0.03 inches. Weight: Mean, 137.46 pounds; standard deviation, 15.24±0.19 pounds. Correlation: 0.4631±0.0133.	P_2	Number of cases: 991. Height: Mean, 67.50 inches; standard deviation, 2.73±0.04 inches. Weight: Mean, 135.93 pounds; standard deviation, 16.16±0.24 pounds. Correlation: 0.5105±0.0158.
P_1 and P_2	Number of cases: 2,512. Height: Mean, 67.63 inches; standard deviation, 2.72±0.03 inches. Weight: Mean, 136.85 pounds; standard deviation, 15.83±0.15 pounds. Correlation: 0.4931±0.0102.		

TABLE 160.—Correlation between height and chest circumference (expiration) in recruits with mitral stenosis, first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Total.	Chest, in inches.														Total.
		28 and under.	29	30	31	32	33	34	35	36	37	38	39	40	41	42
58 and under	4	1														
59	7		1		1											
60	8		1	2	3											
61	16		5	4	4	1	2	2	1	1						
62	43		10	10	10	10	5	7	1	2						
63	88	3	13	13	22	23	24	8		1						
64	140		8	16	26	34	24	24	7							
65	211		9	13	41	63	64	29	12							
66	320	2	13	24	63	81	64	41	20	7						
67	367	4	16	31	60	79	60	62	32	16						
68	356	1	9	28	46	61	79	65	37	17						
69	321	1	8	24	43	61	64	42	39	16						
70	260		10	11	26	55	45	31	27	18						
71	186			7	24	36	41	28	26	16						
72	91		1	2	12	17	21	16	8	9						
73	52		1	3	6	8	8	10	9	3						
74	18					1	2	3	6	1						
75	4															
76	1															
77	1															
78	2															
79	2															
Total	2,507	14	95	186	369	528	463	390	224	120	34	20	4			

P_1 and P_2 —
 Number of cases: 921
 Height: Mean, 67.50 inches; standard deviation, 2.79±0.04
 Chest circumference (expiration): Mean, 32.47 inches; standard deviation, 1.95±0.03 inch.
 Correlation: 0.2669±0.0030.

 P_1 and P_2 —
 Number of cases: 2,507
 Height: Mean, 67.68 inches; standard deviation, 2.73±0.08
 Chest circumference (expiration): Mean, 32.85 inches; standard deviation, 1.99±0.02 inch.
 Correlation: 0.2626±0.0127.

12. VALVULAR DISEASES OF THE HEART (UNCLASSIFIED).

(a) *Stature*.—The mean stature of men found at mobilization camps, in the 3,419 men in the first and second million draft recruits, with unclassified valvular disease of the heart, is 67.60 inches, or 0.11 inch greater than the population in Table I. The standard deviation of the height of these men with unclassified valvular diseases of the heart is 2.67, which is practically the same as the variability of the whole population as shown in Table I.

The relation of distribution of statures in the population with valvular diseases of the heart as compared with the whole population of drafted men is shown in the graph on Plate XXXII. While the two curves of distribution are intertwined to a considerable extent, yet it is clear that there are certain elements of the population with valvular diseases of the heart that are above average stature. Thus there is a clear excess of such diseases in men 69 to 72 inches tall. However, the mode in the population with valvular diseases of the heart lies at 67 inches, or $\frac{1}{4}$ inch below that of the population of drafted men in general.

(b) *Weight*.—Of 909 men found with unclassified valvular diseases of the heart among the first million at mobilization camps, the average weight is 138.49 pounds, or 3.05 pounds below the average of the population in Table I; for the 2,510 in the second million it is 136.78; and for 3,419 men in both groups 137.24, being 4.30 pounds below the mean weight for the first million men. The standard deviation in weight for the first million is 16.49 pounds, or 0.93 pound below the standard deviation of the population in Table I; for the second million it is 17.40; for the two combined it is 17.35 ± 0.14 . This is less than the standard deviation for the whole of the first million as given in Table I, but as the difference is only equal to one-half of the probable error it is probably not very significant.

The relation between the distribution of weights in the population with unclassified valvular diseases of the heart and the population of recruits in general is shown graphically in Plate XXXV. It appears at once that the affected population has a weight clearly below the average and this despite the fact that the statures are practically the same as the average. We have, therefore, evidence of a lack of nutrition in the population with unclassified valvular diseases of the heart, no doubt partly due to the disease itself.

(c) *Chest circumference*.—Of 906 men found with unclassified valvular diseases of the heart among the first million at mobilization camps, the average chest circumference is 32.77 inches, or 0.45 inch less than the population in Table II; for the 2,500 such men found in the second million the chest circumference is 32.49 inches; and for the 3,406 men in both groups combined it is 32.56, which is 0.66 less than the mean chest circumference of the average for the first million men. The standard deviation of chest circumference of those in the first million men is 1.88, or 0.13 below the standard deviation of the whole population in Table II; for the second million it is 2.01 ± 0.02 ; and for the two groups combined 1.98 ± 0.02 .

From these measurements we find that men with unclassified valvular diseases of the heart are tall men with smaller chest circumference and with somewhat less variability than the population as a whole.

TABLE 161.—Correlation between height and weight in recruits with valvular disease of heart (unclassified), first (P_1) and second (P_2) million draft recruits.

[illegible]

TABLE 162.—Correlation between height and chest circumference (expiration) in recruits with valvular disease of heart (unclassified), first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Chest, in inches.																
	28 and under.	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43 and over.	
34 and under.																	
35	3		1		2	4		2	1								
36	8		4	2	6	2		1									
37	15		7	5	8	2		3	1								
38	21		10	10	12	16		6									
39	46		26	22	26	32		12	1								
40	104	2	4	6	38	32	24	17	2								
41	189	7	15	23	73	65	34	23	10								
42	347	8	20	51	73	78	49	33	17								
43	513	13	30	62	93	89	67	45	9								
44	684	23	45	88	118	107	82	56	2								
45	488	3	15	44	86	76	55	31	1								
46	460	3	13	26	67	76	55	30	3								
47	350	2	4	22	38	81	78	40	6								
48	297	2	5	10	28	54	45	31	3								
49	139	1	1	10	12	26	28	16	4								
50	56	2	1	5	3	7	13	8	2								
51	25			2	5	3	4	1	1								
52	6			1	1		1										
53	4																
54	1																
55	2																
Total.	3,406	36	118	331	538	687	668	477	286	142	63	20	14	3	1	1	

P_1 —Number of cases: 906.
 Height: Mean, 67.53 inches; standard deviation, 2.67±0.04 inch.
 Chest circumference (expiration): Mean, 32.77 inches; standard deviation, 1.80±0.03 inch.
 Correlation: 0.2445±0.0211.

P_2 —Number of cases: 2,500.
 Height: Mean, 67.63 inches; standard deviation, 2.67±0.03 inch.
 Chest circumference (expiration): Mean, 32.49 inches; standard deviation, 2.01±0.02 inch.
 Correlation: 0.1896±0.0130.

P_1 and P_2 —Number of cases: 3,406.
 Height: Mean, 67.60 inches; standard deviation, 2.67±0.02 inch.
 Chest circumference (expiration): Mean, 32.56 inches; standard deviation, 1.94±0.02 inch.
 Correlation: 0.2020±0.0111.

Thus in the four categories of heart defects—cardiac hypertrophy, mitral insufficiency, mitral stenosis, and unclassified valvular diseases of the heart—we see that the stature of the affected population is clearly in excess of the average of the whole population. What is the significance of this excess of persons showing valvular diseases of the heart? The first suggestion that occurs to one is that the heart as a pump has to raise fluid about 2 feet above its own level and has to force it through a complicated system of capillaries that occurs in all parts of the body. The taller the individual the more work does the heart have to do and the more back pressure there is upon the valves, both in carrying the fluid to a higher level and in forcing it through a greater number of capillaries. Because of the extra work involved in pumping the blood in persons of large stature, when the muscles or valves of the heart become diseased or crippled as the result of any cause, then the valves or the muscles may become insufficient and show organic disturbance, sooner than in shorter men.

The relation between the distribution of chest girths in men found with unclassified valvular diseases of the heart and in recruits in general is shown graphically in Plate XXXVIII. This graph shows strikingly the abnormally small chest girth of the populations found with unclassified valvular diseases of the heart. This result is associated with the low average weight in this part of the population, despite their average stature. The resulting slender build is no doubt largely the effect of malnutrition consequent upon the disease.

(d) *Robustness*.—Men with unclassified valvular diseases of the heart have an index build of 30.04, or 1.03 below the average of the United States. Pignet's index is 24.78, which places them in the group with medium constitution. For each inch of the average height there are 2.03 pounds of weight, as compared with the average 2.097, and 0.482 inches of chest measurement (expiration), as compared with the normal 0.492.

13. VARICOSE VEINS AND VARICOCELE.

(a) *Stature*.—The average stature of men found at mobilization camps, among 1,409 men in the first million, is 68.34 inches, with varicose veins, which is 0.85 inch above the average of the first million men, as indicated in Table I. The average stature of 2,014 men with this defect found among the second million is 68.49; and for the 3,423 men in the two groups is 68.43, or 0.94 inch above the average height of the whole population. The standard deviation of stature of men with varicose veins among the first million is 2.70; among the second million, 2.77; and among the two groups combined, 2.74 ± 0.02 inches. This is essentially the same as the variability of the statures of the whole population as shown in Table I.

The average stature of 3,453 men among the first million at mobilization camps with varicocele is 68.32 inches, which is 0.83 inch above the average stature of the population in Table I. For the 2,396 men in the second million the average stature is 68.44, and for the 5,849 men in the two groups together 68.37, which is 0.88 inch above the mean stature of the whole population. The standard deviation of the mean stature of men with varicocele among the first million is 2.78; among the second million, 2.71; and for the two groups together, 2.75 ± 0.02 inches, which is somewhat higher than the average for the

whole population, but not significantly so. What is clear in the stature of men having the two defects mentioned is that they are strikingly tall.

The relation between the distribution of stature of the population with varicose veins and the population of recruits in general is shown graphically in Plate XXXII. It appears at once that the population with varicose veins is characterized by great stature. There is a marked deficiency of men below modal stature and a marked excess of men above. The modal stature for the population with varicose veins is at 68 inches, or 0.5 inch above the population of drafted men in general. As in the case of hemorrhoids, so here the mode has a relatively high frequency, indicating relatively small variability in the population with varicose veins and enforcing the conclusion that men with varicose veins are those afflicted primarily because of their tall stature.

The relation between the distribution of statures of men with varicocele as compared with the population of recruits in general is shown graphically in Plate XXXII. Here we see, as in the case of the population with varicose veins, that the population is one of tall men. There is a marked deficiency of men with stature below the average and a marked excess of men with stature above the average. Also the mode is at 68 inches, or 0.5 inch above that of recruits in general, and the fact that it is strikingly higher than the mode of recruits in general indicates a relatively small variability in stature of men with varicocele and enforces the conclusion that men with this defect are affected primarily because of their great stature.

(b) *Weight*.—In 1,409 men found with varicose veins among the first million at mobilization camps the average weight is 146.43 pounds, or 4.89 above the average of the population of Table II. For the 2,014 among the second million the average weight is 146.45, and for the 3,423 men in both lots it is 146.44, or 4.90 above the mean weight of the whole population. This abnormally great weight is in part associated with the great height, nearly an inch above the average, found in these men. By comparing Table 163, showing the relation of stature to weight in men with varicose veins, with Table I, showing the relation of stature to weight among the whole of the first million men, it appears that men with varicose veins are heavy for their height. Thus the mean weight of men 68 inches tall in the whole population is 142.61 pounds, while the mean weight of men 68 inches tall who have varicose veins is 145.52 pounds, or 2.91 above the average of the whole population.

The standard deviation in weight of men found with varicose veins is for the first million 18.39, or 0.97 pounds above the standard deviation in weight of the population in Table I. For the second million the standard deviation in weight is 18.62, and for the two groups together, 18.53 ± 0.15 . This is 1.11 pounds above the standard deviation and over seven times the probable error. It is with one exception the largest standard deviation found. This measures the remarkable variability in weight of men with varicose veins and suggests that this defect is found not merely in a particular stature-weight class, but that it is found in a considerable range of stature classes all of which comprise abnormally stout.

In 3,453 men found with varicocele among the first million examined at mobilization camps the average weight is 141.88 pounds, or 0.34 pound above the average of the population of Table I; for the 2,396 men in the second million the average weight is 141.55; and for the 5,849 in both groups combined it is

141.75, which is 0.26 pound above the average of the first million as shown in Table I. The standard deviation for varicocele in the first million men is 16.68 pounds, or 0.74 below that of the whole population. For varicocele in the second million the standard deviation is 16.18, and for both groups together it is 16.47 ± 0.10 . This is 0.95 pound below the standard deviation of the average population of the first million, as shown in Table I. This low standard deviation is, therefore, in striking contrast with that of varicose veins, and indicates that men affected with varicocele constitute probably a special type and this special type includes exceptionally tall men, though only of average weight; hence men exceptionally tall and slender.

The relation between the distribution of weights in the population with varicose veins and in the population of recruits in general is shown graphically in Plate XXXV. It appears at once that the population with varicose veins is a heavy population, as it is also a tall population. Hence it appears that persons with varicose veins are prevailingly larger persons than the population in general.

The relation between the distribution of weight of persons with varicocele and of recruits in general is shown graphically also in Plate XXXV. It appears that on the whole the population with varicocele is slightly heavier than that of recruits in general, a result which is sufficiently accounted for by the clear excess in stature of the population with varicocele.

(c) *Chest circumference*.—In 1,412 men found among the first million men examined at mobilization camps with varicose veins the average chest circumference is 33.70 inches, or 0.48 inch above the average chest circumference of men of Table II; for 2,014 men in the second million the average chest circumference is 33.64, and for the 3,426 men in both groups together, 33.67. This is 0.45 inch above the average mean chest circumference, which is correlated with the great weight of men found with varicose veins. The standard deviation of chest circumference is for the first million 2.14, or 0.13 above the standard deviation of the population of Table II. For the second million and the two groups combined it is the same (2.14). Men with varicose veins are accordingly not only taller than the average, but have a greater chest circumference and are more variable in this respect than the average of the population, indicating that the defect is found not only in a particular chest circumference-stature class, but that it is found in a considerable range of height classes all of which have large chests just because they are abnormally stout.

Varicocele was found in 3,441 men among the first million examined at mobilization camps. In them the average chest circumference is 33.24 inches, or 0.02 above the average of the whole population of Table II. For the 2,395 men in the second million the average chest circumference is 32.79, and for the 5,835 men in both groups the average is 33.06, or 0.16 below the average for the first million as shown in Table II. The standard deviation of chest circumference of men of the first million is 1.95 inches, or 0.06 inch below the standard deviation of the whole population of Table II. For the cases of varicocele found among the second million the standard deviation in chest circumference is 1.95, and for the two groups together 1.97 ± 0.01 . This is 0.04 inch below the standard deviation for the average of the first million as shown in Table II, and this difference is probably a significant one. Owing to the fact that men showing varicocele are taller than the average, the slight

deficiency of chest circumference indicates that they are not stout, as is confirmed also by their weight. Their reduced variability suggests that the selected tall men having varicocele belong for the most part to a race of such men.

To sum up, it appears that both varicose veins and varicocele are associated with excess stature and that this result is probably primarily a hydrostatic one. The blood vessels of the lower part of the body have to support columns of fluid which are longer in taller men. It is to be expected that veins will give way more commonly where the hydrostatic pressure is greater than where it is less.

From the large size of the standard deviation of weight associated with varicose veins, it seems probable that varicose veins, though found prevalently in heavy men, are found also in some slender men of very tall stature, and in some prevalently short men of great weight, so that both weight and stature are concerned in the production of varicose veins. In the case of varicocele, on the other hand, the hypothesis seems to be favored that chiefly tall men, prevalently of average or slightly less than average robustness, show the defect.

The relation between the distribution of chest girths in the population found with varicose veins and that of the population of recruits in general is shown graphically in Plate XXXVIII. Here there is a clear excess of persons with large chest circumference which is no doubt associated with the generally large size of persons with varicose veins and suggests that the defect has little influence on nutrition, or vice versa.

The relation between the distribution of chest girths in the population with varicocele and the population of recruits in general is also shown graphically in plate XXXVIII. It appears that there is no very important difference between the two populations, though there is a slight, but fairly constant, deficiency in chest girths in the population with varicocele, and this despite the fact that that population contains an excess of tall and heavy men. It appears then that the population with varicocele is characterized by slenderness of build.

(d) *Robustness*.—Men with varicose veins have an index of build of 31.28, or 0.21 unit above the average of the United States. Pignet's index is 19.90. This places them in the group with good constitutions.

Men with varicocele have an index of build of 30.33, or 0.74 unit below the average of the United States. Pignet's index is 23.43. This places them in the group with medium constitution. One notes then that men with varicocele are strikingly inferior in build and robustness to those with varicose veins.

For the men with varicose veins for each inch of the average height there are 2.14 pounds of weight, as compared with the normal 2.097, and 0.492 inch of chest measurement (expiration), as compared with the normal 0.492; while in men with varicocele the weight per inch is 2.07 pounds and the chest measurement 0.484 inch. Thus both sets of men are abnormally tall, but while those with varicose veins are of normal chest and overweight those with varicocele are small chested and underweight.

TABLE 163.—Correlation between height and weight in recruits with varicose veins, first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Total.	Weight, in pounds.																												
		89 and under.	90-94	95-100	101-106	107-110	111-115	116-120	121-125	126-130	131-135	136-140	141-145	146-150	151-155	156-160	161-165	166-170	171-175	176-180	181-185	186-190	191-195	196-200	201-205	206-210	211-215	216-220	221-225	226-230
58 and under.	1																													
59	3																													
60	6																													
61	20	1																												
62	25																													
63	51																													
64	152																													
65	219																													
66	329																													
67	425																													
68	536																													
69	475																													
70	421																													
71	312																													
72	253																													
73	107																													
74	80																													
75	30																													
76	13																													
77	3																													
78	3																													
79	1																													
Total.	3,423	2	2	4	15	45	100	157	270	322	365	425	418	312	242	209	160	115	83	53	42	26	10	28	10	3	2	2	2	1

P_1 —Number of cases: 1,409.
Height: Mean, 68.54 inches; standard deviation, 2.70±0.03
Weight: Mean, 146.43 pounds; standard deviation, 18.39±0.25 pounds.
Correlation: 0.4833±0.0138.

P_2 —Number of cases: 2,014.
Height: Mean, 68.49 inches; standard deviation, 2.77±0.03
Weight: Mean, 146.45 pounds; standard deviation, 18.62±0.24 pounds.
Correlation: 0.4608±0.0118.

P_1 and P_2 —Number of cases: 3,423.
Height: Mean, 68.43 inches; standard deviation, 2.74±0.02
Weight: Mean, 146.44 pounds; standard deviation, 18.63±0.15 pounds.
Correlation: 0.4606±0.0090.

TABLE 164.—(Correlation between height and chest circumference (expiration) in recruits with varicose veins, first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Total.	Chest, in inches.															
		28 and under.	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43 and over.
58 and under	1								1								
59	3			1	1		1	1									
60	6			3	4	2	2	2	5	2							
61	20		1	5	6	6	5	5	1								
62	25		1	8	7	8	8	6	10	3							
63	51		4	9	26	20	28	18	14	14	3	3	1				
64	152	1	2	14	31	40	38	34	25	15	8	5	2				
65	219	3	4	17	45	60	69	54	36	24	13	6	1	2			
66	329		4	14	41	64	97	81	53	41	15	9	5				
67	425	1	8	25	59	80	116	95	64	40	24	10	3	1			
68	538	3	2	16	47	71	106	80	68	43	17	14	6	2			
69	475	1	2	13	34	57	85	57	39	21	17	5	1				
70	421		3	9	15	36	55	39	21	18	9	9	1				
71	312	4	3	5	14	24	42	26	39	21	24	11	6	1			
72	233			1	4	8	18	12	21	12	9	2	4				1
73	107		1	1	2	3	12	7	14	5	1	2	1	2			
74	60						5	4	10	1	2	1	2	1			
75	30				2	1	3		4	2							
76	13						3										
77	2																
78	3							2		2							
79	1								1			1					
Total	3,426	13	33	138	339	491	690	616	460	323	160	93	53	8	6		3

 P_1 —

Number of cases: 1,412.

Height: Mean, 68.35 inches; standard deviation, 2.70 ± 0.03 inch.Chest circumference (expiration): Mean, 33.70 inches; standard deviation, 2.14 ± 0.03 inch.Correlation: 0.2066 ± 0.0172 . P_2 —

Number of cases: 2,014.

Height: Mean, 68.49 inches; standard deviation, 2.77 ± 0.03 inch.Chest circumference (expiration): Mean, 35.64 inches; standard deviation, 2.14 ± 0.02 inch.Correlation: 0.2062 ± 0.0144 . P and P_2 —

Number of cases: 3,426.

Height: Mean, 68.43 inches; standard deviation, 2.75 ± 0.02 inch.Chest circumference (expiration): Mean, 33.67 inches; standard deviation, 2.14 ± 0.02 inch.Correlation: 0.2073 ± 0.0110 .

TABLE 165.—Correlation between height and weight in recruits with varicocele, first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Total.	Weight, in pounds.																									
		89 and under.	90-94	95-100	101-105	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204	205-210	215-219	
58 and under.	3																										
59	22																										
60	14																										
61	32																										
62	67																										
63	109																										
64	163																										
65	269																										
66	566																										
67	730																										
68	935																										
69	974																										
70	737																										
71	640																										
72	331																										
73	189																										
74	102																										
75	44																										
76	23																										
77	3																										
78	4																										
79																											
Total	5,849		1	25	32	147	214	333	580	694	759	701	637	532	403	244	170	130	91	51	31	22	8	13	1	1	

P_1 : Number of cases: 2,453
 Height: Mean, 68.32 inches; standard deviation, 2.78 ± 0.02 inch.
 Weight: Mean, 141.88 pounds; standard deviation, 16.88 ± 0.14 pound.
 Correlation: 0.4995 ± 0.0046 .

P_2 : Number of cases: 2,396.
 Height: Mean, 68.44 inches; standard deviation, 2.71 ± 0.03 inch.
 Weight: Mean, 141.55 pounds; standard deviation, 16.18 ± 0.16 pound.
 Correlation: 0.4854 ± 0.0105 .

S.E.D.
 standard deviation, 2.73 ± 0.02
 ds; standard deviation, $16.47 \pm$

TABLE 166.—Correlation between height and chest circumference (expiration) in recruits with varicocele, first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Total.	Chest, in inches.														
		28 and under.	29	30	31	32	33	34	35	36	37	38	39	40	41	42
58 and under.....	3	1														
59.....	16		1	1	1	2	3	3	5	2			1			
60.....	14		1	1	2	3	3	3	1							
61.....	32		1	2	6	6	7	4	5	1						
62.....	56		8	8	14	14	8	4	3	2						
63.....	109		14	14	29	20	12	16	7	1						
64.....	193		11	22	37	45	31	21	13	2						
65.....	369		10	40	63	76	80	47	23	18						
66.....	556	4	24	62	97	101	105	77	66	37						
67.....	730		17	51	112	148	156	119	87	51						
68.....	932	2	18	57	127	168	202	180	117	61						
69.....	873		14	56	88	171	179	166	117	55						
70.....	736		11	19	71	132	176	131	92	30						
71.....	540	3	2	20	61	83	114	113	79	38						
72.....	331		7	7	27	47	64	61	54	20						
73.....	199		5	5	20	26	38	22	20	10						
74.....	101		1	3	7	22	14	21	16	7						
75.....	44				1	7	8	6	9	2						
76.....	23				2	1	3	5	2	4						
77.....	2					1	1	1	1							
78.....	3															
79.....	4															
Total.....	5,836	12	132	368	757	1,075	1,204	1,011	646	367	156	64	38	3	2	1

P_1 —Number of cases: 3,441.
 Height: Mean, 68.33 inches; standard deviation, 2.75 ± 0.02 inch.
 Chest circumference (expiration): Mean, 33.24 inches; standard deviation, 1.95 ± 0.02 inch.
 Correlation: 0.2575 ± 0.0107 .

P_2 —Number of cases: 2,395.
 Height: Mean, 68.44 inches; standard deviation, 2.71 ± 0.03 inch.
 Chest circumference (expiration): Mean, 32.79 inches; standard deviation, 1.95 ± 0.02 inch.
 Correlation: 0.1836 ± 0.0133 .

P_1 and P_2 —Number of cases: 5,836.
 Height: Mean, 68.38 inches; standard deviation, 2.74 ± 0.02 inch.
 Chest circumference (expiration): Mean, 33.06 inches; standard deviation, 1.97 ± 0.01 inch.
 Correlation: 0.2237 ± 0.0084 .

14. HEMORRHOIDS.

(a) *Stature*.—The average stature of 1,027 men among the first million found at mobilization camps to have hemorrhoids is 67.82 inches, or 0.33 inch above the average of the stature of the population of Table I; for the 797 men in the second million the average is 67.77 inches; and for the 1,824 men in both groups combined 67.80 inches, which is 0.31 inch above the mean stature for the whole of the first million men as shown in Table I.

The standard deviation of stature of men found with hemorrhoids is for the first million 2.68, which is 0.03 less than the standard deviation of the whole population of Table I; for the second million it is 2.91; and for both lots combined it is 2.78 ± 0.03 , a value which differs from the standard deviation of the first million by a little more than twice the probable error.

Men found with hemorrhoids are therefore a somewhat selected lot, being taller than the average. This excess stature is probably one of the causes of hemorrhoids, just as it is of varicose veins and varicocele. Since the variability of the population with hemorrhoids is the same as that of the general population, we may conclude that the men with hemorrhoids constitute a normally distributed part of the population, only distributed about a higher mean stature.

The relation between the distribution of stature in the population with hemorrhoids and the population of recruits in general is shown in Plate XXXII. It appears at once that the population with hemorrhoids consists of men taller than the average. This is indicated both by the constant deficiency of short men 60 to 67 inches tall and the constant excess of tall men 68 to 76 inches tall. The mode in the distribution curve of the population with hemorrhoids is at 68 inches, or one-half inch above that of recruits in general. Moreover, this mode is relatively high and acute, enforcing the lesson that the population with hemorrhoids is affected with this condition largely because of their tall stature.

(b) *Weight*.—The average weight of the 1,027 men found with hemorrhoids among the first million examined at mobilization camps is 141.44 pounds, or 0.10 below the average of the population of Table I; for 797 men in the second million the mean weight is 139.06; and for the 1,824 men in both groups it is 140.39 (Table 167), which is 1.15 below the average of the first million as shown in Table I. This low average weight is associated with abnormally high stature, so that men with hemorrhoids are a tall and slender group. The standard deviation for the first million is 16.78, or 0.64 below the standard deviation of Table I; for the second million it is 16.75; and for both together it is 16.76 pounds, which is 0.66 pound below the standard deviation of the first million men as indicated in Table I. This result indicates that the population with hemorrhoids is a specially selected population, selected tall and slender men, and that this build is in some way causally related to hemorrhoids and has not been induced merely by the hemorrhoids.

The relation between the distribution of weights of the population found with hemorrhoids and that of the whole population of recruits is shown graphically in Plate XXXV. The flattening at the top of the curve is possibly due to the small number of cases.

TABLE 167.—Correlation between height and weight in recruits with hemorrhoids, first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Total.	Weight, in pounds.																								
		89 and under: 84	90-94	95-100	100-105	110-115	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-170	175-179	180-184	185-189	190-194	195-199	200-204	205-209	210-214	215-219	220-224
54 and under.	2																									
56	4																									
58	5																									
60	11																									
61	11																									
62	27																									
63	56																									
64	100																									
65	159																									
66	201																									
67	253																									
68	301																									
69	337																									
70	381																									
71	427																									
72	478																									
73	525																									
74	570																									
75	619																									
76	674																									
77	731																									
78	791																									
79 and over.	851																									
Total.	1,824	2	4	14	46	92	133	202	225	219	219	197	140	108	75	51	38	21	15	5	3	4	9	1	1	1

P_1 —Number of cases: 1,027.
Height: Mean, 67.52 inches; standard deviation, 2.58±0.04 inch.

Weight: Mean, 141.44 pounds; standard deviation, 16.78±0.25 pound.
Correlation: 0.5115±0.0155.

P_2 —Number of cases: 797.
Height: Mean, 67.77 inches; standard deviation, 2.91±0.05 inch.

Weight: Mean, 139.06 pounds; standard deviation, 16.73±0.28 pound.
Correlation: 0.5285±0.0172.

P_1 and P_2 —Number of cases: 1,824.
Height: Mean, 67.60 inches; standard deviation, 2.78±0.03 inch.

Weight: Mean, 140.39 pounds; standard deviation, 16.76±0.19 pound.
Correlation: 0.5219±0.0115.

TABLE 168.—Correlation between height and chest circumference (expiration) in recruits with hemorrhoids, first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Chest, in inches.															Total.
	28 and under.	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43 and over.
28 and under.	2															
29	4	1														
30	6	3	1													
31	11	3	4	1												
32	37	11	17	21	2											
33	96	33	44	34	10	1										
34	159	77	88	54	34	9	1									
35	201	139	117	84	46	38	30	17	2							
36	252	199	168	109	61	43	44	20	12	1						
37	299	237	196	126	81	54	44	29	22	6	1					
38	347	281	228	168	109	61	44	30	13	4	1					
39	381	311	258	208	138	84	45	34	13	3	2	1				
40	426	356	294	244	184	114	54	37	8	4	1	1				
41	478	408	344	284	214	144	64	44	11	3	1	1				
42	529	459	394	334	264	194	74	54	16	2	2	1				
43	579	509	444	384	314	244	84	64	21	3	2	1				
44	629	559	494	434	364	294	94	74	26	4	2	1				
45	679	609	544	484	414	344	104	84	31	5	2	1				
46	729	659	594	534	464	394	114	94	36	6	2	1				
47	779	709	644	584	514	444	124	104	41	7	2	1				
48	829	759	694	634	564	494	134	114	46	8	2	1				
49	879	809	744	684	614	544	144	124	51	9	2	1				
50 and over.	929	859	794	734	664	594	154	134	56	10	2	1				
Total.	1,819	1,519	1,219	919	619	319	119	49	29	19	14	7	1			

P_1 —Number of cases: 1,024.
 Height: Mean, 67.32 inches, standard deviation, 2.68
 ± 0.04 inch.
 Chest circumference (expiration): Mean, 32.22 inches;
 standard deviation, 1.87 ± 0.03 inch.
 Correlation: 0.2280 ± 0.0006 .

P_2 and
 inches; standard deviation, 2.91
 (expiration): Mean, 32.94 inches;
 standard deviation, 1.90 ± 0.03 inch.
 Correlation: 0.2238.

Standard deviation,
 : Mean, 33.10
 ± 0.03 inch.

(c) *Chest circumference.*—The average chest circumference found in the 1,024 men with hemorrhoids among the first million at mobilization camps is 33.22 inches, or the same as the average of the whole population of Table II. In the 795 men with the defect among the second million the mean chest circumference is 32.94, and for the 1,918 men in both groups together it is 33.10, which is 0.12 inch below the average for the first million men, as shown in Table II. Since these men, however, are taller than the average, we may say that the smaller chest circumference means that the men are tall and slender.

The standard deviation for chest circumference for men with hemorrhoids for the first million is 1.87, or 0.14 less than the standard deviation of the whole population of Table II. For the second million it is 1.89. Combining these with the first million we have a standard deviation of 1.88, or 0.13 less than the standard deviation of the first million, shown in Table II. This again indicates that the men with hemorrhoids constitute in respect to chest circumference also a selected class, and that their tall, slender form is antecedent to the incidence of hemorrhoids.

The relation between the distribution of chest girths in the population found with hemorrhoids and the population in general is shown graphically in Plate XXXVIII. Though there is no very striking difference between the two distributions, yet there is a slight excess of men undersize, which is associated with a slight deficiency in weight found in the same population, despite the fact that they are of slightly greater height than the average.

(d) *Robustness.*—The group of men with hemorrhoids has an index of build of 30.54, or 0.53 below the average for the United States. Pignet's index of robustness is 22.50. This places them in the group with medium constitution, and they are thus shown to be between the group with varicose veins and varicocele in physical development.

For each inch of the average height there are 2.07 pounds of weight, as compared with the average 2.097, and 0.488 inch of chest measurement (expiration), as compared with the normal 0.492.

15. ASTHMA.

(a) *Stature.*—The average stature of 614 men with asthma in the first million men examined at mobilization camps was 67.22 inches, or 0.27 less than the average of the whole population in Table I. In 967 men in the second million men the average is 67.26 inches, and for the 1,581 men in both lots together it is 67.24 inches (Table 169), which is 0.25 inch below the mean stature of the first million men. Men with asthma are of slightly less than mean average stature and this is probably indicative of their environmental or racial selection or both. It appears that asthma is much commoner in the Northern States than in the Southern and the Northern States contain a larger proportion of short men. In the mountain regions of Tennessee and Kentucky, where there are very tall men, asthma is relatively uncommon. The short stature of men found with asthma is not due to the disease itself, but to the fact that the larger part of the population is found in that environment of the country in which the causative factors for asthma are especially found.

(b) *Weight.*—In 614 men found with asthma among the first million at mobilization camps the average weight is 139.38 pounds (Table 169), or 2.16

pounds below the average of the population of Table I; for the 967 men in the second million the mean weight is 138.78 pounds, and for the 1,581 men in both groups together it is 139.01 pounds, or 2.53 pounds below the mean weight for the whole of the first million. This low weight is only in part accounted for by the low average stature, since the average weight for a stature of 67.24 inches is 141.02 pounds, while for asthmatics it is 139.01 pounds. The standard deviation for the first million is 17.28 pounds, or 0.14 pound below the standard deviation in weight of the population of Table I. The standard deviation for the second million is 18.35, an extraordinary increase over the standard deviation for the first million men. The average of the two lots is 17.94 ± 0.22 , which is 0.52 above the standard deviation of the entire first million men, as shown in Table I. This result suggests the tentative conclusion that asthma is partly responsible for the small size; that it reduces the weight.

The relation between the distribution of weights in the population found with asthma and that of the population of recruits in general is shown graphically in Plate XXXIV. It appears from the graph that there is an excess of men underweight in the population with asthma, but this is associated with the deficiency in average stature of such men. The mode in the distribution of weights of asthmatics agrees with that of the population at large—about 137 pounds. It may be, however, that there is a deficiency of build among the asthmatics which is determined by the disease itself. In any case there is a marked deficiency of men between 142 and 169 pounds of weight.

(c) *Chest circumference.*—The average chest circumference of the 612 men found with asthma among the first million is 33.57, or 0.35 inch above the average chest circumference of the population of Table I; for the 967 men in the second million (Table 170) it is 33.19; and for the 1,579 men in both combined (Table 170) it is 33.34. This is 0.12 inch above the average chest circumference of all recruits. Since the average stature of men with asthma is less than the average of the whole population studied, and since they are below the average in weight, this large average chest circumference would seem to be in some way determined by the disease. This conclusion is confirmed by the circumstance that the standard deviation for chest circumference for the two combined is 2.12, or 0.11 above the average, an excess which is about four times the probable error. This high variability suggests that the enlarged chest circumference of asthmatic men has been superimposed upon both large and small men, doubtless in consequence of the disease. We may conclude, then, that just the tendency to violent inhalations and expansions of the chest are responsible for the extraordinary development of the chest even in the relatively short and slender asthmatics.

The relation between the distribution of the chest circumference (expiration) in the population found with asthma among the draft recruits and in the population of recruits in general is shown in Plate XXXVII. It appears from this graph that the curve, although somewhat irregular, is moved to the right, showing a greater chest circumference (expiration). The apparent mode is, however, between 32 and 33 inches, or about one-half an inch to the left of the mode of the population of the recruits in general. This shifting of the mode to the left is expected from the small size of asthmatics. It represents the part of the asthmatic population whose chest is not yet abnormally enlarged.

TABLE 169.—*Correlation between height and weight in recruits with asthma, first (P_1) and second (P_2) million draft recruits.*

Height, in inches.	Total.	Weight, in pounds.																									
		80 and under.	90-94	95-100	100-105	110-115	115-120	120-125	130-134	135-140	140-145	145-150	150-155	155-160	160-165	165-170	170-175	175-180	180-185	185-190	190-195	200-205	210-215	215-220	220-225	225-230	230-235 and over.
56 and under.	2																										
59.	6																										
60.	7																										
61.	15																										
62.	21																										
63.	72																										
64.	112																										
65.	178																										
66.	196																										
67.	242																										
68.	281																										
69.	136																										
70.	40																										
71.	83																										
72.	27																										
73.	7																										
74.	3																										
75.	4																										
76.																											
77.																											
78.																											
79.																											
Total.	1,581																										

P_1 —
Number of cases: 614.
Height: Mean, 67.22 inches; standard deviation, 2.77 ± 0.05 inch.
Weight: Mean, 139.38 pounds; standard deviation, 17.28 ± 0.23 pound.
Correlation: 0.3833 ± 0.0282 .

P_2 —
Number of cases: 997.
Height: Mean, 67.26 inches; standard deviation, 2.67 ± 0.04 inch.
Weight: Mean, 138.78 pounds; standard deviation, 18.35 ± 0.28 pound.
Correlation: 0.4226 ± 0.0178 .

P_1 and P_2 —
Number of cases: 1,581.
Height: Mean, 67.24 inches; standard deviation, 2.71 ± 0.03 inch.
Weight: Mean, 139.01 pounds; standard deviation, 17.94 ± 0.25 pound.
Correlation: 0.4069 ± 0.0142 .

TABLE 170.—Correlation between height and chest circumference (expiration) in recruits with asthma, first (P_1) and second (P_2) million draft recruits.

Height in inches.										Chest, in inches.										Total.	28 and under.		P ₁ and P ₂																
																					28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43 and over.	P ₁ and P ₂		
58 and under.....																				2	2																	Number of cases: 967.	Number of cases: 1,579.
59.....																				6	2																	Height: Mean, 67.23 inches; standard deviation, 2.77 ± 0.05 inch.	Height: Mean, 67.25 inches; standard deviation, 2.71 ± 0.03 inch.
60.....																				7	1																		
61.....																				15	2																		
62.....																				21	3																		
63.....																				32	4																		
64.....																				72	3																		
65.....																				111	6																		
66.....																				178	31																		
67.....																				195	17																		
68.....																				242	6																		
69.....																				231	2																		
70.....																				186	4																		
71.....																				138	1																		
72.....																				93	7																		
73.....																				27	1																		
74.....																				7	5																		
75.....																				3	2																		
76.....																				7	2																		
77.....																				4	7																		
78.....																				1	7																		
79.....																				1	7																		
Total.....																				1,579	6	27	74	175	313	298	254	213	110	52	27	18	6	3	1	2			

Number of cases: 612.
Height: Mean, 67.23 inches; standard deviation, 2.77 ± 0.05 inch.

Chest circumference (expiration): Mean, 33.57 inches; standard deviation, 2.11 ± 0.04 inch.
Correlation: 0.1274 ± 0.0298.

Number of cases: 967.
Height: Mean, 67.25 inches; standard deviation, 2.67 ± 0.04 inch.

Chest circumference (expiration): Mean, 33.19 inches; standard deviation, 2.11 ± 0.03 inch.
Correlation: 0.1628 ± 0.0211.

Number of cases: 1,579.
Height: Mean, 67.25 inches; standard deviation, 2.71 ± 0.03 inch.

Chest circumference (expiration): Mean, 33.34 inches; standard deviation, 2.12 ± 0.03 inch.
Correlation: 0.1477 ± 0.0166.

P_1 —Number of cases: 612.
 Height: Mean, 67.23 inches; standard deviation, 2.77 \pm 0.05 inch.
 Chest circumference (expiration): Mean, 33.57 inches; standard deviation, 2.11 \pm 0.04 inch.
 Correlation: 0.1274 \pm 0.0288.

P_2 —Number of cases: 997.
 Height: Mean, 67.26 inches; standard deviation, 2.67 \pm 0.04 inch.
 Chest circumference (expiration): Mean, 33.19 inches; standard deviation, 2.11 \pm 0.03 inch.
 Correlation: 0.1628 \pm 0.0211.

P_1 and P_2 —Number of cases: 1,579.
 Height: Mean, 67.25 inches; standard deviation, 2.71 \pm 0.03 inch.
 Chest circumference (expiration): Mean, 33.34 inches; standard deviation, 2.12 \pm 0.03 inch.
 Correlation: 0.1477 \pm 0.0166.

(d) *Robustness*.—Men with asthma have an index of build of 30.75, which is 0.32 below the average of the United States. Pignet's index is 21.09, which places them in the class with good constitution. For each inch of the average height there are 2.07 pounds of weight, as compared with the normal 2.097 and 0.496 inch of chest measurement (expiration), as compared with the average 0.492.

16. DEFECTIVE AND DEFICIENT TEETH.

(a) *Stature*.—The average stature of the population found with defective and deficient teeth among the 5,166 men in the first million at mobilization camps is 67.26 inches, or 0.23 inch below the average; for 12,817 men in the second million (Table 171) the average stature is 67.26; and for the 17,983 in both together, 67.26, or 0.23 below the average stature for the first million. It appears that men with defective and deficient teeth are strikingly shorter than the average. It does not follow that the short stature is due to the bad teeth.

The standard deviation of stature of men found with defective and deficient teeth is for the first million 2.68, which is only 0.03 less than the standard deviation of the whole population in Table I; for the second million it is 2.69; and for the two combined it is 2.69 ± 0.01 , which is 0.02 below the average standard deviation for the first million. The difference is very slight, but so far as it goes, it suggests that the small stature of men with defective and deficient teeth is due rather to a racial characteristic than to any direct influence upon stature by the teeth. The study, "Defects Found in Drafted Men,"* shows that there is an exceptionally low rate for defective and deficient teeth among the white agriculturists of the South, among the mountain whites, native whites of Scotch origin, and areas having a large proportion of Scandinavians, Germans, and Austrians. On the other hand, the rate is high in the eastern manufacturing, commuting, and maritime groups, and especially in those sections containing French Canadians. Thus, in general, the defective teeth are found in small proportions in those parts of the country occupied by tall men and in large proportions in those parts of the country occupied by short men. It seems probable that we have to do here with a varying racial resistance to dental caries.

The relation between the distribution of stature in men with defective and deficient teeth and that of the population of recruits at large is shown graphically in Plate XXXII. One sees at a glance that men with defective and deficient teeth are somewhat shorter on the whole than the population of recruits in general. This is shown by the uniform excess of men 62 to 67 inches in stature and the uniform deficiency of men 68 to 72 inches tall. The modal stature of men with defective and deficient teeth is 67 inches, or one-half inch below the mode of the population of recruits in general; this indicates that the population with defective and deficient teeth is shorter than recruits in general, probably racially shorter, for reasons given above.

(b) *Weight*.—The average weight of the 5,166 men found with defective and deficient teeth among the first million at mobilization camps is 139.18 pounds, or 2.36 below the average of the population; for the 12,817 men in the second million it is 137.97 pounds; and for the 17,983 men in both groups together (Table 171) 138.32 pounds, or 3.22 pounds below the mean weight of the first million. This deficiency in weight is only in part accounted for by the low

TABLE 171.—Correlation between height and weight in recruits with defective and deficient teeth and dental caries, first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Total.	Weight, in pounds.																									
		80 and under	90- 94	100- 104	105- 109	110- 114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155- 159	160- 164	165- 169	170- 174	175- 179	180- 184	185- 190	190- 194	195- 200	200- 204	205-210-215		
58 and under	19	2																									
59	30	2	1																								
60	72	1	1	7	12	10	32	33	16	11	10	9	3	1	2	1	1	1	1								
61	153	1	1	7	27	64	137	133	49	24	49	36	18	17	6	1											
62	243	1	1	13	35	68	144	170	137	75	139	85	51	31	16	6											
63	796	1	1	6	49	98	306	215	187	138	277	170	115	69	42	23	18	9	3	2	3	2					
64	1,210	3	1	17	112	229	595	422	280	278	523	371	273	177	114	74	41	30	15	7	2	2	1				
65	1,949	5	27	16	29	100	181	322	369	374	589	377	261	163	89	42	23	20	8	4	2	5					
66	2,416	10	34	10	53	167	246	383	367	389	577	340	210	139	105	46	33	21	11	7	9	4	4	5			
67	2,722	1	4	11	34	88	189	328	322	410	596	310	246	160	115	78	55	33	24	12	12	4	4	5			
68	2,594	1	1	3	17	40	118	176	235	301	506	281	203	176	154	85	55	33	32	16	13	5	2	9			
69	2,150	1	1	3	5	23	41	80	131	177	211	202	130	136	112	75	67	38	22	14	10	3	3	1			
70	1,407	1	1	1	1	1	6	10	28	44	48	77	90	73	59	43	34	34	22	14	10	3	3	1			
71	865	1	1	1	1	1	1	6	10	14	23	33	38	40	31	25	18	6	9	7	6	1	5	1			
72	589	1	1	1	1	1	1	1	1	2	3	7	11	10	11	12	12	5	3	2	5	2	1	2			
73	278	1	1	1	1	1	1	1	1	2	3	3	2	2	7	3	3	3	2	4	4	3	2	1			
74	101	1	1	1	1	1	1	1	1	2	3	1	1	1	2	4	1	1	1								
75	45	1	1	1	1	1	1	1	1	1	1	1	1	1	2	4	1	1	1								
76	12	1	1	1	1	1	1	1	1	1	1	1	1	1	2	4	1	1	1								
77	9	1	1	1	1	1	1	1	1	1	1	1	1	1	2	4	1	1	1								
78	5	1	1	1	1	1	1	1	1	1	1	1	1	1	2	4	1	1	1								
79	5	1	1	1	1	1	1	1	1	1	1	1	1	1	2	4	1	1	1								
80 and over	1	1	1	1	1	1	1	1	1	1	1	1	1	1	2	4	1	1	1								
Total	17,983	5	6	19	102	244	646	1,183	1,651	1,651	2,012	1,674	1,308	955	641	454	294	172	130	80	45	30	29	11	5	1	

P_1 —Number of cases: 5,168.
Height: Mean, 67.26 inches; standard deviation, 2.69±0.01
inch.
Weight: Mean, 139.18 pounds; standard deviation, 16.94±
0.11 pound.
Correlation: 0.6107±0.0099.

P_2 —Number of cases: 12,817.
Height: Mean, 67.26 inches; standard deviation, 2.69±0.01
inch.
Weight: Mean, 137.97 pounds; standard deviation, 16.90±
0.07 pound.
Correlation: 0.5054±0.0044.

standard deviation, 2.69±0.01
Weight: Mean, 136.22 pounds; standard deviation, 16.90±
0.06 pound.
Correlation: 0.5057±0.0057.

TABLE 172.—Correlation between height and chest circumference (expiration) in recruits with defective and deficient teeth and dental caries, first (P_1) and second (P_2) million draft recruits.

Height in inches.		Chest, in inches.														Total.	
		28 and under.	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43 and over.
58 and under.....	17	1	2	4	2	4	2	2	5	1	1	1	1				
59.....	30	3	6	9	2	16	13	1	5	2							
60.....	73	3	5	32	14	31	33	6	4								
61.....	154	7	21	48	71	80	65	32	11	5	1	1	1				
62.....	344	14	41	107	158	166	141	79	44	18	7	3	3	1			
63.....	763	11	37	124	220	264	272	155	89	35	15	6	2	2			
64.....	1,044	20	80	161	304	427	373	272	170	81	29	14	10	1	1		
65.....	2,409	15	59	205	385	521	456	365	239	98	40	22	13	2	1	1	
66.....	2,712	8	49	172	368	529	571	468	285	147	69	28	12	6	1	3	
67.....	2,590	6	32	131	298	534	557	459	294	158	67	35	9	8	4	2	
68.....	2,156	11	32	100	234	384	441	404	262	152	71	23	8	3	1	1	
69.....	1,498	4	11	59	114	246	312	270	216	151	82	16	14	1	2		
70.....	588	1	9	42	80	154	179	194	163	82	46	10	7	3	1		
71.....	277		1	19	38	84	114	126	96	58	31	9	3	2			
72.....	101			8	19	26	43	61	53	32	21	10	3	1			
73.....	45			1	3	11	21	31	10	15	8	4	2				
74.....	11			1	2	5	9	11	7	3	6	1					
75.....	9				1	3	1	2	2	3	1						
76.....	5							1	1	2							
77.....										1							
78.....																	
79.....																	
Total.....	17,932	106	389	1,224	2,343	3,510	3,567	2,932	1,962	1,044	483	188	91	28	13	9	3

P_1 —Number of cases: 5,150.
Height: Mean, 67.26 inches; standard deviation, 2.67 ± 0.02 inch.
Chest circumference (expiration): Mean, 33.25 inches; standard deviation, 1.94 ± 0.01 inch.
Correlation: 0.2713 ± 0.0087 .

P_2 —Number of cases: 12,782.
Height: Mean, 67.26 inches; standard deviation, 2.69 ± 0.01 inch.
Chest circumference (expiration): Mean, 32.89 inches; standard deviation, 2.02 ± 0.01 inch.
Correlation: 0.2496 ± 0.0086 .

P_1 and P_2 —Number of cases: 17,932.
Height: Mean, 67.26 inches; standard deviation, 2.69 ± 0.01 inch.
Chest circumference (expiration): Mean, 33.00 inches; standard deviation, 2.00 ± 0.01 inch.
Correlation: 0.2551 ± 0.0047 .

average stature of the group, since men with a height of 67.24 inches are expected to have an average weight of 141.02 pounds. There is thus a deficiency in weight of men with defective and deficient teeth even when regard is taken for their short stature.

The standard deviation in weight for both groups combined is 16.89 ± 0.06 , or 0.53 pound below the average of the first million. This low standard deviation indicates that defective and deficient teeth are found predominantly in men belonging to a short and slender race. It is to be noted that the highest State rates for defective and deficient teeth are found in Vermont, New Hampshire, Rhode Island, Massachusetts, and Maine, all of which have about three times the average rate. Now these are just the States occupied by an excess of French Canadian groups that have a rate of 40.01 for defective and deficient teeth, which is by far the largest ratio of any of the groups. At the same time this group is characterized by exceptionally low stature.

The relation between the distribution of weight of the population found with defective and deficient teeth and that of the population of recruits in general is shown graphically in Plate XXXV. It appears at once that the population with defective and deficient teeth is on the whole characterized by having a weight inferior to the average; but they are, however, prevalingly short persons, so that there is little evidence that they are badly nourished on account of the defective teeth.

(c) *Chest circumference.*—The average chest circumference in the 5,150 men found with defective and deficient teeth among the first million at mobilization camps is 33.25, or 0.03 above the average of the whole population. For the 12,782 men in the second million it is 32.89, and for the 17,932 men in the two groups combined (Table 172) it is 33.00, or 0.22 inch less than the average for the whole of the first million. The standard deviation of chest circumference for the two groups is 2.00, or 0.01 below the standard deviation for the first million. This is not a significant difference.

The relation between the distribution of chest girths in the population found with defective and deficient teeth and recruits in general is shown graphically in Plate XXXVIII. This shows that, on the whole, persons with defective and deficient teeth have a smaller chest girth than the average, as indeed they have a smaller weight and stature. On the whole, they contain an excess of men of small size, belonging to small races.

(d) *Robustness.*—Men with defective and deficient teeth have an index of build of 30.33, which is 0.74 below the average for the United States. Pignet's index is 22.31, which places them in the class with the medium constitution. For each inch of the average height there are 2.06 pounds of weight, as compared with the normal 2.097, and 0.491 inch of chest measurement (expiration), as compared with the normal 0.492.

17. HERNIA.

(a) *Stature.*—The average stature of 13,822 men with hernia found among the first million men at mobilization camps is 67.40, which is only 0.09 inch below the mean stature of the population of Table I; for the 20,398 men in the second million it is 67.47; for the 34,220 men in both combined (Table 173),

67.44, or 0.05 less than the average for the first million. The standard deviation of stature of men with hernia is for both groups 2.76 ± 0.01 , which is 0.05 inch above the average for the first million as shown in Table I. One may conclude, therefore, that hernia occurs in the different statures in about the same proportion as the different statures occur in the whole population.

The relation between the distribution of stature in the population with hernia and that of the population of drafted men in general is shown graphically in Plate XXXII. This curve indicates that men with hernia are not far from a fair sample of the whole population in respect to stature. There is, however, a slight excess of men shorter than the average. This is shown by the deficiency in the population with hernia between 67 and 70 inches, which overbalances the shift of the modal point from $\frac{1}{2}$ to $\frac{1}{4}$ inch to the right. The excess of short men is, however, not at all marked.

(b) *Weight*.—In 13,822 men found with hernia among the first million at mobilization camps, the average weight is 141.69 pounds, or 0.15 pound above the average; for the 20,398 men in the second million the weight is 140.91 pounds, and for the 34,220 men in both groups combined (Table 173), 141.23, which is 0.31 pound below the average of the first million. Since the men with hernia are slightly below the average stature, this result in the case of such men shows about normal build. The standard deviation of weight for both groups combined is 17.17, or 0.25 pound below the standard deviation in weight of the population of Table I. This indicates that hernia is especially apt to affect persons who are slightly under weight, although stature has practically nothing to do with its occurrence. This result might have been anticipated since it is just the men who are below normal vigor, as indicated partly by underweight, who, whatever their size, are most apt to show the effects of a strain in the abdominal muscles and the ligaments of the inguinal region.

The relation between the distribution of weights in the population found with hernia and that of the population of recruits in general is shown graphically in Plate XXXV. It appears that there is no important difference in the distribution of weights in the two populations, as indeed we have found there is no important difference in stature.

(c) *Chest circumference*.—The average chest circumference in the 13,822 men found with hernia among the first million at mobilization camps is 33.23 inches, or 0.01 inch greater than the average chest circumference for the whole population of Table II; for the 20,398 men in the second million it is 33.04; and for the 34,220 men in both groups combined (Table 174) it is 33.11, which is 0.11 less than the average of the first million as shown in Table II. The standard deviation of chest circumference for both groups combined is 2.00 ± 0.01 , which is practically that of all of the first million men, as shown in Table II. It appears, then, that men in whom hernia is found have slightly less average weight and chest circumference than men of their height, which is almost exactly the average.

TABLE 173.—Correlation between height and chest circumference (expiration) in recruits with hernia, first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Total.	Chest, in inches.														49 and over.	
		24 and under.	28	30	31	32	33	34	35	36	37	38	39	40	41		42
58 and under.....	46																
59.....	99																
60.....	118																
61.....	340																
62.....	639																
63.....	1,283																
64.....	2,233																
65.....	3,312																
66.....	4,420																
67.....	4,858																
68.....	5,007																
69.....	4,227																
70.....	3,245																
71.....	2,245																
72.....	1,130																
73.....	1,207																
74.....	281																
75.....	109																
76.....	49																
77.....	17																
78.....	11																
79.....	7																
80 and over.....	6																
Total.....	34,220	110	659	2,052	4,365	6,444	6,940	5,776	3,886	2,198	1,084	407	235	53	25	5	9

P_1 —Number of cases: 13,822.
 Height: Mean, 67.40 inches; standard deviation, 2.74 ± 0.01 inch.
 Chest circumference (expiration): Mean, 33.23 inches; standard deviation, 2.60 ± 0.01 inch.
 Correlation: 0.2515 ± 0.0084 .

P_2 —Number of cases: 13,822.
 Height: Mean, 67.40 inches; standard deviation, 2.74 ± 0.01 inch.
 Chest circumference (expiration): Mean, 33.11 inches; standard deviation, 2.60 ± 0.01 inch.
 Correlation: 0.2523 ± 0.0084 .

TABLE 175.—Correlation between height and weight in recruits with enlarged inguinal rings, first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Total.	Weights, in pounds.															
		89 and under.	90-94	95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164
58 and under.	31																
59	76																
60	144																
61	400																
62	176																
63	1,749																
64	1,609																
65	2,839																
66	4,066																
67	5,541																
68	6,301																
69	6,318																
70	5,357																
71	4,753																
72	1,568																
73	778																
74	334																
75	133																
76	48																
77	15																
78	9																
79	6																
80 and over.	3																
Total.	43,619																

P_1 —Number of cases: 20,142.
 Height: Mean, 67.54 inches; standard deviation, 2.69±0.01 inch.
 Weight: Mean, 140.17 pounds; standard deviation, 16.64±0.06 pound.
 Correlation: 0.5174±0.0035.

P_2 —Number of cases: 23,477.
 Height: Mean, 67.40 inches; standard deviation, 2.71±0.01 inch.
 Weight: Mean, 140.00 pounds; standard deviation, 16.46±0.05 pound.
 Correlation: 0.5077±0.0033.

P_1 and P_2 —Number of cases: 43,619.
 Height: Mean, 67.46 inches; standard deviation, 2.70±0.01 inch.
 Weight: Mean, 140.08 pounds; standard deviation, 16.54±0.04 pound.
 Correlation: 0.5115±0.0024.

TABLE 176.—Correlation between height and chest circumference (expiration) in recruits with enlarged inguinal rings, first (P_1) and second (P_2) million draft recruits.

Height, in inches.		Chest, in inches.														Total.	
		28 and under.	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43 and over.
58 and under.....	31	3	4	7	6	6	2	1	1	1
59.....	75	5	6	12	14	18	8	10	1
60.....	144	6	20	24	39	28	14	12
61.....	400	11	57	75	95	66	48	29	11	3
62.....	748	5	39	131	181	128	77	48	23	10	7
63.....	1,609	14	65	296	358	319	194	111	57	21	6	2
64.....	2,839	16	104	503	636	553	403	187	110	37	15	7	1
65.....	4,082	16	116	571	656	884	628	337	148	76	26	8	2
66.....	5,538	13	119	442	827	1,112	865	525	264	98	40	23
67.....	6,497	16	118	438	895	1,402	1,078	600	355	146	58	27	5	2
68.....	6,619	17	90	367	780	1,267	1,163	800	421	183	78	46	5	4
69.....	5,373	6	69	253	569	1,152	972	690	411	184	68	37	5	2
70.....	4,060	4	30	133	390	721	750	574	331	156	59	30	7	2
71.....	2,759	4	24	80	216	454	551	414	249	106	44	25	5	1
72.....	1,787	3	16	48	109	166	152	134	99	86	28	17
73.....	133	1	5	16	25	73	68	63	44	23	12	6
74.....	48	2	5	16	25	22	23	18	14	5	2
75.....	16	3	5	6	8	11	3	7	3	1
76.....	2	4	8	3
77.....	1	1	2	3
78.....
79.....
80 and over.....	3
Total.....	43,625	114	810	2,747	5,566	8,457	9,027	7,346	4,913	2,681	1,165	472	244	32	14	3	4

P_1 —Number of cases: 20,161.
 Height: Mean, 67.55 inches; standard deviation, 2.70±0.01 inch.
 Chest circumference (expiration): Mean, 33.03 inches; standard deviation, 1.92±0.01 inch.
 Correlation: 0.2410±0.0045.

P_2 —Number of cases: 23,464.
 Height: Mean, 67.40 inches; standard deviation, 2.71±0.01 inch.
 Chest circumference (expiration): Mean, 33.09 inches; standard deviation, 1.97±0.01 inch.
 Correlation: 0.2237±0.0042.

P_1 and P_2 —Number of cases: 43,625.
 Height: Mean, 67.47 inches; standard deviation, 2.71±0.01 inch.
 Chest circumference (expiration): Mean, 33.06 inches; standard deviation, 1.95±0.004 inch.
 Correlation: 0.2310±0.0031.

The relation between the distribution of chest girths in the men found with hernia and the population of recruits in general is shown graphically in Plate XXXVIII. There is very little difference in the two curves, but there is an indication of a slight deficiency in chest girth in men with hernia despite the fact that in stature they are fair samples of the whole population. This deficiency in chest girth is possibly due to the condition of malnutrition which favored the hernia.

(d) *Robustness*.—Men with hernia show an index of build of 31.05, or 0.02 under the average of the United States. Pignet's index is 21.17, which places them in the class with good constitution. For each inch of the average height there are 2.09 pounds of weight, as compared with the normal 2.097, and 0.491 inch of chest measurement (expiration), as compared with the averages 0.492.

18. ENLARGED INGUINAL RINGS.

(a) *Stature*.—The mean stature of 20,142 men found with enlarged inguinal rings among the first million at mobilization camps is 67.54 inches, or 0.05 above the average for the population of Table 175; for the 23,477 men in the second million it is 67.40; and for the 43,619 men in both combined 67.46, or 0.03 below the mean height of the first million men. This is an insignificant difference. The standard deviation of stature for both groups is 2.70, which is again almost exactly the standard deviation for the first million. It appears, therefore, that recruits showing enlarged inguinal rings are typical in their stature of the whole population of recruits; just as are those who show well-developed hernia.

(b) *Weight*.—The average weight of 20,142 men found with enlarged inguinal rings among the first million at mobilization camps (Table 175) is 140.17 pounds, or 1.37 below the average of the population of Table I; for the 23,477 men in the second million it is 140.00 pounds; and for the 43,619 men in both groups combined it is 140.08, or 1.46 pounds below the average. The standard deviation is 16.54 ± 0.04 , or 0.88 pound below the standard deviation for the first million as shown in Table I. This indicates that, as in the case of hernia, so in the case of enlarged inguinal rings, the defect is found prevalingly in slender persons. It is because they are slender that they have enlarged inguinal rings rather than that the weight is reduced because they have enlarged inguinal rings.

(c) *Chest circumference*.—The average chest circumference of the 20,161 men found with enlarged inguinal rings among the first million is 33.03 inches, or 0.19 below the average chest circumference of the population studied; for the 23,464 men in the second million it is 33.09; and for the 43,625 men in both combined (Table 176) 33.06, which is 0.16 below the average chest circumference for the first million as shown in Table II. This result indicates again that men with enlarged inguinal rings are slender. The standard deviation for the two groups combined is 1.95 ± 0.004 . This small standard deviation combined with the small chest circumference and low weight indicates that the men with enlarged inguinal rings belong prevalingly to a race of average stature, but that is underweight and abnormally slender.

(d) *Robustness*.—The index of build is 30.78, or 0.29 below normal. Pignet's index is 21.89. The pounds of weight to each inch of average height are 2.09, and the inches of chest measurement (expiration) 0.490.

19. FLAT-FOOT.

(a) *Stature*.—The average stature of 175,358 men with flat-foot among the first million is 67.30 inches, or 0.19 below the average stature of the population of Table I. For the 94,990 men in the second million (Table 177) the mean stature is 67.28, and for the 270,348 men in both groups combined it is 67.30, or 0.19 below the average for the first million as shown in Table I. The standard deviation for the two combined is 2.70 ± 0.003 , or 0.01 below the standard deviation for the total of the first million as shown in Table I. Thus men with flat-foot are shorter and less variable in stature than the population at large. This suggests that we have here to do with a prevalence of flat-foot in the short races.

(b) *Weight*.—The average weight of 175,358 men found with flat-foot among the first million at mobilization camps is 143.24, or 1.70 pounds above the average of the population of Table I. For 94,990 men in the second million (Table 177) it is 143.31, and for 270,348 men in both groups combined it is 143.26, or 1.72 pounds above the average of the first million as shown in Table I. This high mean weight is present despite the fact that the average stature of men found with flat-foot is slightly below the average for the whole population. The standard deviation for weight of men with flat-foot for the two groups combined is 18.41 ± 0.02 , or 0.99 pound above the average for the first million shown in Table I. This result shows that men with flat-foot are relatively heavy, and that all physical types of men who become heavy may gain flat-foot.

(c) *Chest circumference*.—The number of cases of flat-foot were so many and the preliminary inspection indicated that the chest circumference deviated so slightly from the normal that, on account of lack of funds, it was decided not to do the work required to make out the table of relation of height to chest circumference for men with flat-foot.

(d) *Robustness*.—The index of build of men with flat-foot is 31.63, or 0.56 above the average of the United States. The chest circumference for men with flat-foot was not calculated, so their index of robustness can not be determined. There are 2.13 pounds of weight for each inch of average height, as compared with the average 2.097.

TABLE 177.—Correlation between height and weight in recruits with flat-foot, first (P_1) and second (P_2) million draft recruits.

	Number of cases, 175,386.	standard deviation, 2.70 ± 0.003
Height: Mean, 67.30 inches;	standard deviation, 2.73 ± 0.004	
Weight: Mean, 143.24 pounds;	standard deviation, 18.97	
Correlation: 0.4786 ± 0.0012.		

20. DEFECTIVE PHYSICAL DEVELOPMENT.

This term is a vague one used often by the examining boards to avoid recording a more specific diagnosis. It is frequently applied to persons who are far under the normal degree of robustness and also to many cases of malformation of the trunk, such as flat chest or curved spine.

(a) *Stature*.—The average stature of 758 men found with defective physical development among the first million examined at mobilization camps is 66.34 inches, or 1.15 inches less than the average stature of the population in Table I. For the 534 men in the second million it is 66.91, and for 1,292 men in both groups together (Table 178) 66.57, or 0.92 inch below the mean stature of the whole of the first million as given in Table I. We have here a very striking inferiority in stature of the men with "defective physical development." And there is reason for thinking that many persons who were below the standard minimum stature were on that account given the diagnosis "defective physical development." The standard deviation in stature of men placed in this category is for the two groups 3.84 ± 0.05 , which is the largest standard deviation of stature shown in Table 187. This simply means that the diagnosis has a very scattered application through the whole range of the human statures. It is applied, as we have seen, prevailing to very short persons, but also to tall persons who are very thin, flat chested, or otherwise malformed. A comparison of the range of different statures of men with defective physical development and of the total defective population shows clearly the significance of this high variability. For whereas in the population as a whole there is a larger proportion of men with the stature of 67 inches (14.6 per cent) than of any other inch class; yet, of men diagnosed as having defective physical development, there were in this stature class only 9.8 per cent. Instead of the proportion in the classes at each side of the mean diminishing as in the normal frequency curve, in this selected class the numbers actually increase, being 11.1 per cent for men 66 inches tall and 11.9 per cent for men 68 inches tall. The proportion of men 59 inches tall is nearly 25 times the proportion of such men in the whole population. There are nearly 11 times as many men 60 inches tall in this special group as in the whole population. Also there are disproportionately high ratios for statures 71 inches and above. Of men 79 inches tall, there are nearly 15 times as many in the defective physical development group as in the population at large. It is the extremes, then, that were prevailingly diagnosed as of defective physical development.

(b) *Weight*.—The average weight of the 758 men found with defective physical development among the first million at mobilization camps (Table 178) is 128.94 pounds, or 12.60 pounds below the average of the population of Table I. For 534 men in the second million the average weight is 123.43 pounds, and for 1,292 in both groups it is 125.51, which is 16.03 pounds below the average of the first million. This exceptionally low weight is only in part accounted for by the low average stature of this group. The standard deviation of the groups is 18.57 pounds ± 0.25 , which is 1.15 pounds above the average standard deviation for the whole first million. These figures show clearly that the group of defective physical development includes men belonging to races of various sizes, victims

TABLE 178.—Correlation between height and weight in recruits with defective physical development, first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Total.	Weight, in pounds.																							
		89 and under.	90-94	95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204
58 and under.....	2				2																				
59.....	68			6	10	13	11	12	3	4	2	1	3	1			1	1							
60.....	57			5	8	7	7	11	5	3	3	3	1	3			1	1							
61.....	37			1	4	13	1	6	3		4	1	2			1	1								
62.....	43			1	8	11	3	3	3	3	7	1				2	1								
63.....	59			8	13	17	10	3	2	3	1	1				1									
64.....	69			5	4	14	13	8	8	5	6	1	3												
65.....	123			2	10	21	30	12	13	8	9	7	3	5	2										
66.....	144			1	11	22	29	16	17	9	7	6	9	4	4										
67.....	126			1	6	8	19	22	11	15	12	7	11	3	5	2									
68.....	154				2	9	21	31	21	18	13	10	11	8	10	2									
69.....	126				1	2	2	11	24	16	8	14	8	10	2	7									
70.....	106				2	1	9	16	9	5	13	15	6	11	5	3									
71.....	78						3	9	10	16	9	9	9	5	3										
72.....	45						2	2	6	5	7	7	3	4	1	1									
73.....	25							1	2	1	1	2	2	2	2	1									
74.....	11															1									
75.....	4															1									
76.....	5															1									
77.....	2																								
78.....	3																								
79.....	4																								
80 and over.....	1																								
Total.....	1,292			30	81	138	169	176	127	103	111	90	77	60	32	24	20	19	12	7	6	1	3	4	2

P_1 —Number of cases: 758.
 Height: Mean, 66.34 inches; standard deviation, 4.01±0.07 inch.
 Weight: Mean, 128.94 pounds; standard deviation, 18.14±0.31 pound.
 Correlation, 0.4600±0.0193.

P_2 —Number of cases: 534.
 Height: Mean, 66.91 inches; standard deviation, 3.56±0.07 inch.
 Weight: Mean, 123.43 pounds; standard deviation, 18.96±0.36 pound.
 Correlation, 0.5008±0.0219.

P_1 and P_2 —Number of cases: 1,292.
 Height: Mean, 66.57 inches; standard deviation, 3.84±0.05 inch.
 Weight: Mean, 125.51 pounds; standard deviation, 18.57±0.25 pound.
 Correlation: 0.4644±0.0147.

TABLE 179.—Correlation between height and chest circumference (expiration) in recruits with defective physical development, first (P_1) and second (P_2) million draft recruits.

	Height, in inches.					Chest, in inches.													Total.
	28 and under.	29	30	31	32	33	34	35	36	37	38	39	40						
56 and under.	2	1	1	1	1	1	1	1	1	1	1	1	1	1					
59	68	10	13	10	12	12	9	9	1	1	1	1	1	1					
60	56	7	10	9	9	8	6	6	1	1	1	1	1	1					
61	37	1	7	7	7	7	4	2	2	1	1	1	1	1					
62	42	6	10	9	7	4	2	2	2	1	1	1	1	1					
63	60	16	15	11	9	6	2	2	2	1	1	1	1	1					
64	67	3	13	20	11	8	2	2	2	1	1	1	1	1					
65	124	17	27	24	20	10	9	6	3	2	1	1	1	1					
66	142	20	27	31	26	18	16	13	7	5	3	2	1	1					
67	127	3	10	17	25	18	15	13	7	5	3	2	1	1					
68	154	3	10	28	37	31	18	15	9	8	5	3	2	1					
69	124	2	6	28	25	23	18	13	9	8	5	3	2	1					
70	105	2	4	16	16	25	17	10	8	6	5	3	2	1					
71	76	2	3	16	15	25	17	12	8	6	5	3	2	1					
72	43	1	2	4	11	16	8	12	7	4	3	2	1	1					
73	25	1	1	4	4	8	3	8	4	3	2	1	1	1					
74	11	1	1	1	3	2	2	2	1	1	1	1	1	1					
75	4	1	1	1	1	1	1	1	1	1	1	1	1	1					
76	5	1	1	1	1	1	1	1	1	1	1	1	1	1					
77	2	1	1	1	1	1	1	1	1	1	1	1	1	1					
78	3	1	1	1	1	1	1	1	1	1	1	1	1	1					
79	4	1	1	1	1	1	1	1	1	1	1	1	1	1					
80 and over.	1	1	1	1	1	1	1	1	1	1	1	1	1	1					
Total.	1,284	126	217	251	239	166	106	69	33	21	13	11	2	2					

P_1 —Number of cases: 752.
Height: Mean, 66.34 inches; standard deviation, 4.02±0.07 inch.

Chest circumference (expiration): Mean, 32.15 inches; standard deviation, 2.21±0.04 inch.
Correlation: 0.1792±0.0228.

P_2 —Number of cases: 832.
Height: Mean, 66.50 inches; standard deviation, 3.56±0.07 inch.

Chest circumference (expiration): Mean, 31.43 inches; standard deviation, 2.21±0.04 inch.
Correlation: 0.1792±0.0228.

Standard deviation, 3.84±0.06
Mean, 31.53 inches; standard deviation, 2.21±0.04 inch.

of various environmental conditions which have prevented full physical development or the achievement of the physical standards set for military service. A comparison of the distribution of weights of these men and the distribution of weights in the whole populations shows that there is an extraordinary deficiency of heavy men. Thus the ratios for men over 137 pounds are about half the corresponding normal ratios, from 152 upward about one-third the normal ratios. On the other hand, there are proportionately nearly nine times as many men of 102 pounds found in this group as in the population at large, and of men 95-99 pounds there are 11 times as many. By comparing these ratios with those of height, we see that men with defective physical development were prevailingly exceedingly slender men.

(c) *Chest circumference.*—The chest circumference in the 752 men found with defective physical development among the first million (Table 179) is 32.15 inches, or 1.07 inches less than the average chest circumference of the whole population of Table II. For the 532 men in the second million the mean is 31.43, and for 1,284 men in the two combined 31.85, or 1.37 less than the average of the total first million. This low mean chest circumference is correlated with the low weight of prevailingly slender men. The standard deviation of the chest circumference in the two groups is 2.18 ± 0.03 , or 0.17 inch above the standard deviation of the first million. It appears, then, that the group with defective physical development contains very short and very tall men, all under weight and all of prevailingly small chest circumference and showing a marked variability as contrasted with the population at large. We are not here dealing with a racial trait, but with a mixture of races and of causes having this in common, that they result in men who, in form and proportions, deviate far from military standards.

(d) *Robustness.*—Men with defective physical development have an index of build of 28.32, or 2.75 below the average of the United States. Their index of robustness is 29.94, or 9.06 below the average of the United States, placing them in the class of weak constitution. For each inch of the average height there are 1.89 pounds of weight as compared with the normal 2.097 pounds, and 0.479 inch of chest measurement (expiration), as compared with the normal 0.492.

21. UNDERWEIGHT.

The requirements of weight for each unit of height are given in Table 138, p. 297, copied from the physical examination standards.

(a) *Stature.*—The average stature of 2,686 men found to be underweight among the first million at mobilization camps is 66.22 inches, or 1.27 less than the average stature of the population of Table I. For 9,943 in the second million men the average stature is 65.30; for 12,129 men, both lots (Table 180), it is 65.50, or 1.99 inches below the mean height. It will be recalled that local boards, during most of the draft period, were instructed not to send to camp men under 61 (later 60) inches in height. It appears, however, from Table 180 that 241 men 59 inches and under were examined at camp and recorded as being underweight. The low average stature is of course due to the fact that weight and stature are closely correlated and the "underweight" is frequently one who has less than average stature. However, the proportion of men 74-77 inches tall was larger than in the population as a whole, showing that there was an exceptionally large number of very tall men who were below the appropriate

weight for their stature. Of men 61 inches tall, those rejected for underweight were five times the normal proportion of this stature. Of men 60 inches tall, there were about eight times the normal proportion rejected for underweight, and similarly for the shorter groups. The standard deviation of stature of men found at camps to be underweight for both groups is 3.36 ± 0.01 , which is 0.65 inch above the standard deviation for the whole population of the first million. This high variability is clearly due to the fact that underweight, while found especially in the short men, was found also in the very tall men. Consequently underweight men are a very variable group with respect to stature.

(b) *Weight*.—The average weight of 2,686 men diagnosed as underweight among the first million at mobilization camps (Table 180) is 114.67 pounds, or 26.87 pounds below the average of the population of Table I. For 9,443 men in the second million it is 109.88; for 12,129 men in both groups together, 110.94, or 30.60 pounds below the average for the first million. The standard deviation for the groups combined is 9.89 ± 0.04 pounds, which is the lowest standard deviation of weight found in the first million men. This result was, of course, to have been anticipated, since we have in this group one selected for a single feature, namely, weight. It constitutes, therefore, so far as weight goes, a very homogeneous lot, but not so homogeneous as would be the case were only the small men considered. The small standard deviation, moreover, combined with the prevailing causes of underweight, indicates that the majority of men concerned belong to the small races.

(c) *Chest circumference*.—The average chest circumference of the 2,708 men found to be underweight among the first million at mobilization camps is 30.94 inches, or 2.29 inches less than the average. For the 9,424 men among the second million (Table 181) the chest circumference is 30.32; and for 12,132 men in both lots (Table 181) it is 30.46 or 2.76 inches less than the mean of the whole population. The standard deviation of chest circumference for the two lots is 1.53 ± 0.01 inches, or 0.48 inch less than the average standard deviation in chest circumference for the first million. It appears, then, that the underweight group is characterized by extremes of statures and by slenderness of body, by small chest circumference, and by relatively slight variability in respect to slenderness. The slight variability in chest circumference is, however, partly due to the small average chest circumference. However, if we divide the standard deviation by the mean we find for this, the coefficient of variability, a ratio of 0.56, which is much less than that for the population at large, 0.91. This indicates that the chest circumference of underweight men is not only absolutely but also relatively smaller than that of the population at large. The men of this class had, therefore, an exaggerated and relatively uniform slenderness of build.

(d) *Robustness*.—The index of build of men classified as underweight is 25.86, or 5.21 below the average, and the lowest index of the whole United States. Pignet's index is 37.36, or 16.48 below the average of the whole country, placing them in the class of bad constitution. For each inch of the average height there are 1.69 pounds of weight, as compared with the normal 2.097, and 0.465 inch of chest measurement (expiration), as compared with the normal 0.492.

TABLE 180.—Correlation between height and weight in recruits underweight, first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Total.	Weight, in pounds.																							
		89 and under.	90-94	95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204
58 and under.....	76	6	4	16	18	15	11	5		1															
59.....	59	4	4	33	40	43	24	15		2															
60.....	165	4	12	76	98	125	56	19	4	3															
61.....	401	8	27	99	110	244	119	23	2																
62.....	626	2	23	130	285	363	104	37																	
63.....	956	4	19	118	323	514	217	50	1																
1,248	1,248	5	19	118	323	514	217	50	1																
64.....	1,469	3	16	81	337	535	378	109	5																
1,594	1,594	4	12	74	260	568	480	192	3																
65.....	1,352	1	8	48	169	472	434	227	20																
1,197	1,197	3	3	24	104	305	408	263	71	13															
66.....	861	1		3	37	182	284	177	94	33	1														
67.....	881	1		3	37	182	284	177	94	33	1														
68.....	659	2	1	3	16	68	159	116	121	54	22	7													
69.....	476				4	23	69	116	121	54	22	7													
70.....	334					9	22	16	64	41	32	13	2												
71.....	283					1	8	2	25	47	34	22	14	11											
72.....	160						1		11	20	16	6	12	15	7										
73.....	120								3	7	4	5	2	2	5	2									
74.....	63								1		1	4	2	6	3	2									
75.....	38																								
76.....	10																								
77.....																									
78.....																									
79.....	1																								
Total.....	12,129	43	129	705	1,815	3,468	2,774	1,517	651	439	217	156	95	54	24	13	10	13	2	2	1				1

P_1 —Number of cases: 2,698.
 Height: Mean, 66.22 inches; standard deviation, 3.51±0.03 inch.
 Weight: Mean, 114.67 pounds; standard deviation, 11.61±0.11 pound.
 Correlation: 0.7339±0.0060.

P_2 —Number of cases: 9,443.
 Height: Mean, 66.30 inches; standard deviation, 3.29±0.02 inch.
 Weight: Mean, 109.88 pounds; standard deviation, 9.07±0.05 pound.
 Correlation: 0.6873±0.0037.

P_1 and P_2 —Number of cases: 12,129.
 Height: Mean, 66.50 inches; standard deviation, 3.36±0.01 inch.
 Weight: Mean, 110.94 pounds; standard deviation, 9.89±0.04 pound.
 Correlation: 0.6970±0.0031.

TABLE 181.—Correlation between height and chest circumference (expiration) in recruits underweight, first (P_1) and second (P_2) million draft recruits.

Height, in inches.		Chest, in inches.												
Total.		28 and under.	29	30	31	32	33	34	35	36	37	38	39	
58 and under.	76	9	14	22	18	11	1	1	2					
59	168	8	35	48	31	23	17	13			1			
60	399	34	79	112	90	56	14	7						
61	622	57	142	181	150	66	19							
62	951	102	220	278	211	96	32	9	1					
63	1,252	113	277	410	274	121	45	9						
64	1,470	140	314	472	315	153	43	20	11		1			
65	1,385	161	343	478	371	166	53	16	4			3		
66	1,598	98	278	431	346	159	52	12	8					
67	1,387	101	220	383	274	149	41	13	10			1		
68	1,196	54	149	248	221	119	40	14	8	1				
69	662	33	95	185	175	107	34	18	9	2				
70	477	21	53	121	121	94	40	14	9		1			
71	333	9	30	57	63	81	29	17	10					
72	291	5	14	46	71	65	51	27	7					
73	159		9	22	40	40	32	9	3					
74	119		2	11	20	36	26	17	5					
75	63		3	3	7	14	18	9	6					
76	38	1	2	6	8	4	6	5	3					
77	9			2	1		2	3	1					
78														
79	2						1							
Total.....	12,132	946	2,279	3,516	2,837	1,559	586	236	99	15	7	5	37	

P_1 —Number of cases: 2,708.
 Height: Mean, 66.30 inches; standard deviation, 3.51 ± 0.03 inch.
 Chest circumference (expiration): Mean, 30.94 inches; standard deviation, 1.72 ± 0.02 inch.
 Correlation: 0.2843 ± 0.0119 .

P_2 —Number of cases: 9,424.
 Height: Mean, 65.30 inches; standard deviation, 3.29 ± 0.02 inch.
 Chest circumference (expiration): Mean, 30.32 inches; standard deviation, 1.44 ± 0.01 inch.
 Correlation: 0.2312 ± 0.0066 .

P_1 and P_2 —Number of cases: 12,132.
 Height: Mean, 65.50 inches; standard deviation, 3.36 ± 0.01 inch.
 Chest circumference (expiration): Mean, 30.46 inches; standard deviation, 1.53 ± 0.01 inch.
 Correlation: 0.2459 ± 0.0058 .

22. OVERWEIGHT AND OBESITY.

Table 138 specifies the standard weights for each height and the minimum weight for each height which will permit of acceptance. There was a maximum weight for each stature, and this defined the overweight and obese men. The overweight men, however, reached such extremes that it was not feasible to tabulate all of the classes of weight.

(a) *Stature*.—The mode of stature of recruits as far as tabulated stands at 69 inches or $1\frac{1}{2}$ inches above the mode of the whole population as shown in Table I. The group is also clearly a more variable one than the population as a whole.

(b) *Weight*.—The average weight of the 271 men found with overweight and obesity among the first million at mobilization camps is not calculated because, by the method of tabulating, more than half of the men placed in this class were grouped in the category "200 pounds and over." For the same reason the standard deviation was not calculated.

(c) *Chest circumference*.—The average chest circumference of the 271 men found with overweight and obesity among the first million at mobilization camps is 36.92 inches, or 3.70 inch above the average chest circumference of the average male population at large. Owing to the fact that in tabulating chest circumference, 39 inches and over were massed into one class, the standard deviation of chest circumference has not been calculated.

(d) *Robustness*.—As stated above, the weight of men classified as overweight or obese was grouped in many cases as 200 pounds and over, and as a result the average weight could not be accurately determined. Hence the indices of build and robustness could not be calculated.

23. CRYPTORCHIDISM, HYPOSPADIA, ANORCHISM, AND MONORCHISM.

This group is a heterogeneous one, comprising some cases of accidental mutilation and others of imperfect development of the genitalia, owing to their retention of an infantile condition.

(a) *Stature*.—The average stature of 1,808 men found with one of these defects among the first million is 67.34 inches, which is only 0.15 inch less than the average stature of the whole population of Table I. For 3,140 men in the second million the stature is 67.49, and for 4,948 men, both lots together (Table 182), 67.44, or 0.05 inch below the mean height for the first million. The standard deviation for the two lots is 2.81 ± 0.02 , which is 0.10 above the standard deviation of the whole first million. It appears, therefore, that in respect to stature, persons with the named defects, though these are of an infantile or undeveloped nature, are typical of the whole population. They are, however, slightly more variable in stature than the rest of the population, and this seems to be due to the fact that there is an excess in this group of very short men under 60 inches and of men 70–75 inches tall, and a corresponding deficiency in the middle statures of 67 inches. This indicates that there is a slight association with the effects due to internally secreting glands, which influence both stature and the development of the genitalia.

TABLE 182.—Correlation between height and weight in recruits with cryptorchidism, anorchism, monorchism, and hypospadias, first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Total.	Weight, in pounds.																								
		89 and under.	90-94	95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204	205-209
54 and under.	14																									
59	6																									
60	31																									
61	51																									
62	96																									
63	202																									
64	287																									
65	476																									
66	634																									
67	723																									
68	965																									
69	591																									
70	462																									
71	333																									
72	172																									
73	81																									
74	42																									
75	21																									
76	6																									
77	3																									
78	1																									
79	1																									
Total.....	4,946	2	2	8	28	51	158	267	414	522	544	553	618	425	341	306	224	133	101	50	56	31	28	14	23	4

Number of cases: 1,908.
Height: Mean, 67.34 inches; standard deviation, 2.40±0.03 inch.
Weight: Mean, 140.81 pounds; standard deviation, 18.61±0.21 pound.
Correlation: 0.5198±0.0116.

P₁.....

Number of cases: 3,140.
Height: Mean, 67.49 inches; standard deviation, 2.81±0.02 inch.
Weight: Mean, 139.93 pounds; standard deviation, 17.49±0.16 pound.
Correlation: 0.4666±0.0094.

P₂.....

Number of cases: 4,948.
Height: Mean, 67.44 inches; standard deviation, 2.81±0.02 inch.
Weight: Mean, 140.25 pounds; standard deviation, 17.91±0.12 pound.
Correlation: 0.4867±0.0073.

P_1 —
Number of cases: 1,808.
Height: Mean, 67.34 inches; standard deviation, 2.80±0.03 inch.

Weight: Mean, 140.81 pounds; standard deviation, 18.81±0.21 pound.
Correlation: 0.5186±0.0116.

P_2 —
Number of cases: 3,140.
Height: Mean, 67.49 inches; standard deviation, 2.81±0.02 inch.

Weight: Mean, 139.93 pounds; standard deviation, 17.48±0.15 pound.
Correlation: 0.4666±0.0094.

P_1 and P_2 —
Number of cases: 4,948.
Height: Mean, 67.44 inches; standard deviation, 2.81±0.02 inch.

Weight: Mean, 140.25 pounds; standard deviation, 17.91±0.12 pound.
Correlation: 0.4967±0.0073.

TABLE 163.—Correlation between height and chest circumference (expiration) in recruits with cryptorchidism, anorchism, monorchism, and hypospadias, first (P_1) and second (P_2) million draft recruits.

Height, in inches.	Total.	Chest, in inches.														43 and over.
		28 and under.	29	30	31	32	33	34	35	36	37	38	39	40	41	42
55 and under	14															
56	6		1													
57	31	2	5	10	6	3	4	1	1	1						
58	51		10	20	10	12	4	4	4	2						
59	90	3	15	36	37	19	14	11	11	1						
60	202	3	12	35	60	55	25	46	26	9	1					
61	290	3	19	51	75	113	41	66	17	6						
62	474	1	13	46	106	138	121	90	25	24	10					
63	653		15	49	106	145	134	108	66	28	10					
64	723		18	60	141	165	123	119	64	34	13					
65	829		16	72	161	172	125	104	63	37	14					
66	922		9	11	27	43	72	69	41	32	11					
67	1000		1	11	16	26	33	42	15	17	6					
68	1171						12	21	14	9	3					
69	1311						8	10	6	5	1					
70	1422						5	4	4	1						
71	1511															
72	1611															
73	1711															
74	1811															
75	1911															
76	2011															
77	2111															
78	2211															
79	2311															
Total	4,943	19	122	378	679	990	926	832	463	287	130	60	56	9	3	4

P_1 —Number of cases: 1,898.
 Height: Mean, 67.54 inches; standard deviation, 2.50 ± 0.03 inch.
 Chest circumference (expiration): Mean 33.18 inches; standard deviation, 2.05 ± 0.02 inch.
 Correlation: 0.2299 ± 0.0150 .

P_2 and P_3 —Number of cases: 4,943.
 Height: Mean, 67.44 inches; standard deviation, 2.51 ± 0.02 inch.
 Chest circumference (expiration): Mean, 33.05 inches; standard deviation, 2.10 ± 0.01 inch.
 Correlation: 0.2107 ± 0.0092 .

(b) *Weight*.—Of the 1,808 men found with these defects among the first million, the average weight is 140.81 pounds, or 0.73 below the average of the population of Table I. For 3,140 men in the second million, the average is 139.93, and for 4,948 men in both groups (Table 182), 140.25, which is 1.29 pounds below the average weight of the whole of the first million. The standard deviation in weight for both lots is 17.91 ± 0.12 , or 0.49 pounds above the standard deviation in weight of the whole population. This result indicates that the group is a rather heterogeneous one so far as weight goes, but characterized on the whole by slightly less than normal weight, despite the fact that the average stature is practically normal. Men with these defects are therefore on the whole slightly slenderer than the average population. The high standard deviation indicates that the defect is more apt to be found in lighter and heavier men than in men of more nearly normal weight. This accords again with the view that these defects are associated with glandular disturbances which are well known to influence weight.

(c) *Chest circumference*.—The average chest circumference in the 1,808 men found with these defects among the first million at mobilization camps (Table 183) is 33.18, or 0.04 inch below the average of the population studied. For the 3,135 men found with these defects in the second million the average chest circumference is 32.95 inches, or for 4,943 men in both lots together 33.03, or 0.19 inch below the mean chest circumference of the first million men.

The standard deviation of the chest circumference for both lots is 2.10 ± 0.01 . This is practically the same as the standard deviation for the whole population. We conclude, therefore, that the part of the population with the named defect is very like the population at large, except that it is slightly underweight and slender and that this condition affects different parts of the normal frequency distribution nearly uniformly, so that there is no marked selection of a particular class.

(d) *Robustness*.—The index of build of men with cryptorchidism, hypospadias, anorchism, and monorchism is 30.84, or 0.23 less than the average of the whole United States. Pignet's index is 21.83, or 0.95 above the average of the United States, which places them in the class of good constitution. For each inch of the average height there are 2.08 pounds of weight, as compared with the normal 2.097, and 0.490 inch of measurement (expiration), as compared with the normal 0.492.

V. SUMMARY: BODILY DIMENSIONS IN RELATION TO DISEASES.^a

The foregoing sections have revealed the fact that populations selected because of the possession of some common disease or defect have in many cases proportions which deviate widely from those of the population of recruits in general.

The findings in this respect are summarized in Tables 184–192, and in Plates XXXIX–XLI. In Plate XXXIX the deviations in stature from the average are given for the populations detected with each of 23 defects. This figure shows that the greatest deviation above the average stature is found in that population which has varicose veins; next that which has varicocele;

^a It will be noted that in what follows, the averages of height, weight, and chest at expiration are those taken only from men showing the various defects and diseases referred to in Table 187. For the average of height, weight, and chest circumference of the whole population, reference has to be made to Tables I, II, and III.

next that characterized by pulmonary tuberculosis; next the two forms of goiter, and then certain forms of valvular disease of the heart, tachycardia, and hemorrhoids. On the other hand, striking deviations below the average in stature are found in populations classified as underweight, defective physical development, or as possessing astigmatism, myopia, hyperopia, asthma, defective and deficient teeth, and flat feet. The reasons for the deviations in these representative populations are treated in the corresponding sections above.

Plate XL gives the deviations in weight of various populations, characterized by having particular diseases or defects, from the average weight found in the entire population of recruits. Here, far more than in height, most of the deviations are below the normal. That is because almost all of the diseases and defects tend to interfere with bodily functioning and to reduce the weight. In the case of varicose veins, however, the defect itself is probably largely induced by excessive stature, and so we find persons with this defect to be on the average far above the mean weight of the whole population. In the case of simple goiter, the excess of weight found in the population is merely associated with the excess of stature that this population shows. The "build" is not abnormal. (See Table 189.) On the other hand, in pulmonary tuberculosis and various valvular diseases of the heart there is clear evidence that deficiency of weight is determined by the diseases. In the case of the population with defective and deficient teeth, the reduction in weight is possibly influenced by inadequate nutrition. Other populations whose weight is below the average are those characterized by eye defects, but these are populations composed to an unusual extent of persons belonging to races characterized by short stature.

Plate XLI gives the distribution in chest circumference of the populations characterized by different defects and diseases from the mean chest circumference of the whole population of recruits. Here, again, most of the deviations are in deficiency. In the case of varicose veins the population is characterized by great build, excessive weight, and thus also of excessive chest girth. In the case of the population characterized by asthma there is reason for thinking that the excess chest circumference is induced by the disease itself. Passing to the populations characterized by abnormally small chest measurements, we find, in addition to the groups of underweight and defective physical development, the group characterized by pulmonary tuberculosis, and, following that, various groups characterized by organic and functional diseases of the heart. Here also are the populations with errors of refraction whose small chest measurement is correlated with general small size on account of the small races which form so large a part of these populations.

Plate XLII is drawn up in a similar manner to Plate XIV, page 177. Here an attempt is made to show the interrelation of stature, weight, and chest circumference (expiration) as associated with certain diseases or defects.

Passing downward the first heavy horizontal line shows the average stature of the first million draft recruits, while the second and third shows the quotients of the average weight and chest circumference (expiration) divided by the height. It is apparent at once that the average stature of the men with certain diseases or defects is above that of the population of recruits in general. Included in this number are defects of the veins, namely, varicose veins, varicocele, and hemorrhoids; tuberculosis; organic and functional cardiac conditions, namely, mitral insufficiency, simple tachycardia, cardiac hypertrophy,

mitral stenosis, and valvular diseases of the heart unclassified; and, finally, exophthalmic goiter and simple goiter. Only one of these conditions, varicose veins, shows both a proportional weight and chest circumference (expiration) above the average. Here the proportional weight stands well up above, while that for the chest circumference (expiration) reaches the average line. Simple goiter also has a proportional weight slightly above the average, but the proportional chest circumference (expiration) is below it. For all of the other conditions with excessive stature the proportional weight and chest circumference (expiration) are well below the average, and it is apparent that the men with these diseases or defects are on the average tall, slender, and small-chested. This is most marked in cases of tuberculosis. For men with hypertrophied tonsils the stature, the proportional weight, and chest circumference (expiration) are practically the same as the average of the population of recruits in general. On the other hand, the proportional weight and chest circumference (expiration) of recruits with hernia and relaxed inguinal rings are below the average, and the same is true of recruits with congenital genital defects, as well as of those with defective and deficient teeth.

The build of the asthmatic cases is of considerable interest, since it is apparent that the stature is considerably below the average, as is also the proportional weight, but the greater proportional chest circumference (expiration) is much above the average. This latter condition is due no doubt to the effects of the disease itself. The three refractive errors, hyperopia, myopia, and astigmatism, have proportional weight below the average, with proportional chest circumference (expiration) slightly above.

In figure 2 of Plate XLII the weight is taken as the controlling factor, while in the second and third sections below there is shown the quotient of the weight divided by the height, and the weight divided by the chest circumference (expiration). As shown in figure 1, simple goiter affects weight less than exophthalmic goiter; consequently the quotient of the weight divided by the chest circumference (expiration) is greater for the patient with simple goiter than for those with exophthalmic goiter. On the other hand, since the chest circumference (expiration) for asthmatics has increased while the proportional weight has decreased, the quotient of the weight divided by the chest circumference (expiration) is much reduced.

In figure 3 the chest circumference (expiration) is taken as the controlling factor, while in the second and third sections below there is shown the quotient of the chest circumference (expiration) divided by the height and the weight divided by the chest circumference (expiration). It is again apparent here that men with varicose veins have a well-developed chest, are above the average in stature, and have great proportional weight. It is also apparent that for asthmatics the chest circumference (expiration) has increased out of proportion to the stature and weight. Further study of Plate XLII will reveal many interesting facts showing the interrelation of stature, weight, and chest circumference (expiration) as associated with the special diseases or defects considered.

Table 189 summarizes the relations of index of build and index of robustness (Pignet) ²⁰ associated with the various diseases. The heavy build of many recruits with errors of refraction is striking; they belong to stocky races. The

dependence of flat-foot and varicose veins on build is fairly clear. It is noteworthy that recruits with varicose veins stand at the top of the list for robustness. That recruits with asthma stand so high is due to their large chest girth. The shape of recruits with defective development of the genitalia is probably due to the influence of the sex glands on development. The heart conditions are associated with a low average robustness, as indeed also a slender build.

The variability of the stature of recruits with various diseases presents many points of interest (Table 190). In general it appears that, when the aberrant stature that is associated with a disease is so associated because tall or short races are especially apt to be affected by the disease, the variability is low. Thus, recruits with goiter have low stature-variability. But goiter appears prevailing in the Northwestern States which are inhabited by tall "Nordics." We have seen also that short races are especially apt to have defective and deficient teeth; and so the stature of the class shows less variability than the average. On the other hand, the great variability in stature of recruits with myopia is due, as Plate XXXIII shows, to the fact that there are two groups in the lot—a group of racially short stature (largely Polish Jews) and of other recruits of average stature. Likewise cardiac hypertrophy comprises persons of normal stature and also a group of especially tall persons. On the other hand, underweight occurs in tall and short races and is due to a diversity of causes, and the resulting group is very variable in stature. The high variability of the group of cryptorchidism, etc., is partly due to the heterogeneity of the group.






The variability of the weight of recruits with the various defects and diseases is shown in Table 191. This table combined with Table 190 shows that men with varicose veins are of varied races, but generally tall and heavy. Thus stature and weight are not caused by the condition of the veins; for, so far as stature goes, the group is less variable than the average; and as for weight it is only a little more variable (as measured by the coefficient of variation). A tolerably uniformly tall and heavy lot of men have become victims of varicose veins; the disease is induced in part by the build. Varicocele is likewise found in tall and gaunt men of the Nordic type, and such defectives are tolerably uniform in this respect.

In other cases the variability of weight is due to the composite constitution of the group. Thus, as has already been pointed out, the myopics are composed both of the average population and also a special lightweight (and short) group. The asthmatics seem to comprise a group of normal weight and one of underweight (probably due to the disease in its advanced stages). Men with flat-foot are of somewhat less than average stature, very heavy on the average, but comprising some small and light men.

The clearest case of an uniformly low variability induced by disease is that of pulmonary tuberculosis. A group of abnormally tall persons of average variability in stature shows an abnormally and extraordinarily uniform low weight. Low weight is one of the principal symptoms of the disease. Again, mitral stenosis is found in men of average stature but far below average weight; in them stature is not affected, but weight is abnormally low, and the group is remarkably uniform in this respect.

HEIGHT DISTRIBUTION BY SPECIAL DISEASES OR DEFECTS (M & P.)

RATIOS PER 1000




FINE LINE CURVE DENOTES AVERAGE FOR U. S.

PLATE XXXIII.

WEIGHT DISTRIBUTION BY SPECIAL DISEASES OR DEFECTS (M & F)

RATIOS PER 1000

2



FINE LINE CURVE DENOTES AVERAGE FOR M. & F.
PLATE XXXIV.

WEIGHT DISTRIBUTION BY SPECIAL DISEASES OR DEFECTS (M & N)
 RATIO PER 1000

2

⁴
 VERUOCILE

⁶
 VERUOL

³
 VERUOCILE VERUOL

⁵
 DEFECTIVE & DEFICIENT TEETH

THE LINE CURVE DENOTES AVERAGE FOR U. S.
 PLATE XXXV.

WEIGHT DISTRIBUTION BY SPECIAL DISEASES OR DEFECTS (R & R)
 RATIOS PER 1000

2

1

4

3


2

5
 AVERAGE

FINE LINE CURVE DENOTES AVERAGE FOR U. S.
 PLATE XXXVI


CHEST (EXP.) DISTRIBUTION BY SPECIAL DISEASES OR DEFECTS (R₁ & R₂)
RATIOS PER 1000

4
ASTHMA




6
HYPEROPIA

7
MYOPIA



9
CARDIAC HYPERTROPHY



FINE LINE CURVE DENOTES AVERAGE FOR U. S.

PLATE XXXVII.


CHEST (EXP.) DISTRIBUTION BY SPECIAL DISEASES OR DEFECTS (P_1 & P_2)

RATIOS PER 1000

■

■

5
HEMONORRHOES




■

4

■

9
ASTHMA

8
DYSPEPTIC & DEFICIENT TESTIN




■

■

FINE LINE CURVE DENOTES AVERAGE FOR U.S.
PLATE XXXVIII.

HEIGHT MEASUREMENT, SPECIAL DISEASES (P&P) IN RELATION TO THE AVERAGE, INDUCED MEN (P.)



**WEIGHT MEASUREMENT, SPECIAL DISEASES (P&P;₂)
IN RELATION TO THE AVERAGE, INDUCTED MEN (P)**





PLATE XL.

**CHEST MEASUREMENT (EXPIRATION), SPECIAL DISEASES (P₁ & P₂)
IN RELATION TO THE AVERAGE, INDUCED MEN (P₁)**



**WEIGHT MEASUREMENT, SPECIAL DISEASES (R&P₂)
IN RELATION TO THE AVERAGE, INDUCED MEN (P)**




TABLE 184. -- *Height distribution by special diseases or defects, for first and second million draft recruits.*

SECTION A: ABSOLUTE NUMBERS.

Disease.	Total.	Height, in inches.																			80 and over.			
		58 and under.	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76		77	78	79
Tuberculosis, pulmonary.....	10,701	9	14	35	64	122	240	505	783	1,198	1,444	1,584	1,496	1,242	888	598	283	135	53	23	5	8	1	1
Gout, simple.....	7,069	4	10	9	33	70	165	308	529	871	1,089	1,143	1,021	801	515	305	143	61	26	14	7	6	1	1
Exophthalmic goiter.....	2,620	1	5	6	10	28	71	126	187	304	378	398	382	305	210	105	63	23	9	4	2	2	1	1
Asthma.....	1,581	2	6	7	15	21	72	112	178	196	242	231	186	138	93	40	27	7	3	4	1	1	1	1
Strabismus.....	1,592	2	2	5	14	41	88	129	163	220	228	228	171	131	80	43	21	9	3	1	1	1	1	1
Myopia.....	999	3	3	6	17	21	38	70	100	133	134	150	116	86	48	29	9	5	1	1	1	1	1	1
Hyperopia.....	2,420	3	10	11	32	58	108	185	304	509	348	350	238	209	132	86	24	17	1	1	1	1	1	1
Tonsillitis, hypertrophic.....	52,031	35	82	205	462	999	1,982	3,211	5,004	6,392	7,571	7,555	6,748	4,925	3,250	1,913	942	400	143	57	22	11	10	1
Cardiac hypertrophy.....	1,343	1	7	2	6	14	42	74	129	177	190	167	161	93	60	28	9	6	2	1	1	1	1	1
Mitral insufficiency.....	8,960	8	9	30	52	115	276	459	732	1,034	1,244	1,380	1,140	984	620	399	205	101	44	19	4	5	2	2
Tachycardia, simple.....	2,512	4	7	8	16	43	85	140	212	320	367	399	320	262	186	91	53	18	11	3	1	1	1	1
Valvular disease of heart.....	3,419	2	2	7	16	46	67	109	187	265	292	309	330	243	238	139	57	25	9	4	1	1	1	1
Hemorrhoids.....	1,824	2	4	5	11	27	56	100	159	253	301	237	181	127	78	45	20	9	4	1	1	1	1	1
Varicose veins.....	3,423	1	3	6	25	51	152	219	329	426	536	475	421	312	233	107	60	30	13	3	3	4	1	1
Varicocele.....	5,849	3	22	14	32	67	109	193	369	556	730	695	574	421	312	233	107	60	30	13	3	3	4	1
Defective and deficient teeth.....	17,983	19	30	73	155	343	798	1,210	1,949	2,416	2,722	2,694	2,159	1,497	965	599	278	101	45	12	9	5	5	5
Hernia.....	34,324	47	71	120	341	642	1,288	2,249	4,433	8,871	5,017	4,242	3,249	2,135	1,209	607	283	109	40	17	11	7	6	6
Inguinal rings, enlarged.....	43,619	31	75	144	400	749	1,609	2,839	5,541	6,501	5,018	4,383	3,087	2,785	1,598	778	334	133	48	15	9	6	3	3
Flat-foot.....	270,348	135	530	1,026	2,770	5,644	11,124	18,297	27,639	33,843	40,273	39,628	32,135	24,515	15,334	8,591	4,055	1,645	621	290	113	73	63	4
Defective physical development.....	1,292	2	68	57	37	59	60	123	144	126	154	126	105	78	45	25	11	4	5	2	3	4	1	1
Underweight.....	12,129	76	165	401	926	1,248	1,469	1,594	1,382	1,197	861	659	476	334	263	160	120	63	38	10	1	1	1	1
Cryptorchidism, hypospadias, anorchism, and monorchism.....	4,948	14	6	31	51	95	202	287	476	634	695	723	591	482	383	172	81	42	21	6	3	1	1	1
Total.....	463,033	407	1,139	2,224	5,231	10,186	19,830	32,480	48,791	63,299	71,786	71,949	59,636	45,559	29,424	16,901	8,214	3,547	1,398	630	218	147	107	20

SECTION B: RATIOS PER 1,000.

Disease.	Total.	Height, in inches.													Total.										
		59	60	61	62	63	64	65	66	67	68	69	70	71		72	73	74	75	76	77	78	79	80 and over.	
Tuberculosis, pulmonary.....	10,701	0.84	1.31	3.27	5.96	11.40	22.43	47.19	73.17	111.90	148.03	180.12	203.16	220.06	234.98	248.06	260.45	272.02	282.75	292.61	301.67	310.00	317.60	324.50	330.70
Gout, simple.....	7,099	.56	1.41	1.27	4.65	9.86	25.24	43.10	74.52	122.09	148.18	161.03	173.83	186.58	199.28	211.93	224.53	237.08	249.58	262.03	274.43	286.78	299.08	311.33	323.53
Exophthalmic goiter.....	2,630	.38	1.29	2.49	3.52	10.69	27.10	48.09	71.37	116.03	144.27	161.19	175.80	190.11	204.13	217.85	231.28	244.40	257.11	269.42	281.33	292.94	304.25	315.26	326.00
Asthma.....	1,681	.27	1.27	1.91	2.49	9.49	25.75	46.54	70.84	112.97	153.07	168.11	181.66	195.66	209.11	222.01	234.36	246.15	257.38	268.05	278.16	287.71	296.71	305.16	313.96
Angina pectoris.....	1,892	.16	1.26	3.14	8.79	25.75	46.54	70.84	112.97	153.07	168.11	181.66	195.66	209.11	222.01	234.36	246.15	257.38	268.05	278.16	287.71	296.71	305.16	313.96	
Myopia.....	2,909	.31	2.10	3.19	17.04	21.67	44.63	76.45	123.62	157.25	188.29	216.71	242.63	267.01	289.88	311.20	332.09	352.56	372.61	392.24	411.46	430.27	448.60	467.43	
Hypertension.....	2,436	.14	2.13	4.55	13.22	21.67	44.63	76.45	123.62	157.25	188.29	216.71	242.63	267.01	289.88	311.20	332.09	352.56	372.61	392.24	411.46	430.27	448.60	467.43	
Tonsillitis, hypertrophic.....	52,031	.24	1.56	3.96	9.46	19.20	37.71	61.71	90.17	122.55	153.85	181.44	206.33	228.62	249.31	268.40	286.89	304.78	322.07	338.76	354.85	370.34	385.23	399.52	
Cardiac hypertrophy.....	1,343	.74	1.02	3.39	4.87	10.98	31.15	53.10	82.62	116.10	144.41	168.48	193.33	218.06	242.67	267.15	291.50	315.72	339.82	363.80	387.67	411.43	435.08	458.62	
Mitral insufficiency.....	6,860	.90	1.74	2.38	7.53	17.12	33.64	65.77	94.39	127.67	158.00	186.41	213.82	241.23	268.64	296.05	323.46	350.87	378.28	405.69	433.10	460.51	487.92	515.33	
Altrial stenosis.....	2,312	.68	2.22	3.18	7.53	17.12	33.64	65.77	94.39	127.67	158.00	186.41	213.82	241.23	268.64	296.05	323.46	350.87	378.28	405.69	433.10	460.51	487.92	515.33	
Rheumatism, simple.....	2,147	.86	2.34	3.39	7.53	17.12	33.64	65.77	94.39	127.67	158.00	186.41	213.82	241.23	268.64	296.05	323.46	350.87	378.28	405.69	433.10	460.51	487.92	515.33	
Valvular diseases of heart.....	3,149	.90	2.18	3.18	7.53	17.12	33.64	65.77	94.39	127.67	158.00	186.41	213.82	241.23	268.64	296.05	323.46	350.87	378.28	405.69	433.10	460.51	487.92	515.33	
Gonorrhoea.....	1,634	.10	2.18	3.18	7.53	17.12	33.64	65.77	94.39	127.67	158.00	186.41	213.82	241.23	268.64	296.05	323.46	350.87	378.28	405.69	433.10	460.51	487.92	515.33	
Varicose veins.....	2,846	.20	2.18	3.18	7.53	17.12	33.64	65.77	94.39	127.67	158.00	186.41	213.82	241.23	268.64	296.05	323.46	350.87	378.28	405.69	433.10	460.51	487.92	515.33	
Diabetes.....	17,083	.51	1.67	3.59	8.93	19.07	33.71	67.29	103.38	138.35	173.31	208.27	243.23	278.19	313.15	348.11	383.07	418.03	452.99	487.95	522.91	557.87	592.83	627.79	
Chorea.....	43,619	.17	3.30	8.93	19.07	33.71	67.29	103.38	138.35	173.31	208.27	243.23	278.19	313.15	348.11	383.07	418.03	452.99	487.95	522.91	557.87	592.83	627.79		
Chorea, chorea.....	37,324	.17	3.30	8.93	19.07	33.71	67.29	103.38	138.35	173.31	208.27	243.23	278.19	313.15	348.11	383.07	418.03	452.99	487.95	522.91	557.87	592.83	627.79		
Chorea, chorea.....	43,619	.17	3.30	8.93	19.07	33.71	67.29	103.38	138.35	173.31	208.27	243.23	278.19	313.15	348.11	383.07	418.03	452.99	487.95	522.91	557.87	592.83	627.79		
Chorea, chorea.....	270,348	.16	1.96	3.12	28.64	30.28	45.67	63.41	95.20	111.46	97.82	119.29	97.82	119.29	97.82	119.29	97.82	119.29	97.82	119.29	97.82	119.29	97.82	119.29	
Chorea, chorea.....	1,292	.16	1.96	3.12	28.64	30.28	45.67	63.41	95.20	111.46	97.82	119.29	97.82	119.29	97.82	119.29	97.82	119.29	97.82	119.29	97.82	119.29	97.82	119.29	
Chorea, chorea.....	12,179	6.27	13.60	33.04	61.61	78.62	102.80	121.11	131.42	113.64	98.66	70.99	64.33	38.24	27.54	18.19	13.19	8.99	6.19	3.13	1.83	0.89	0.08	0.08	
Chorea, chorea.....	4,948	2.63	1.21	6.27	10.31	19.40	40.82	58.00	96.20	126.13	140.46	146.12	119.44	97.41	87.30	34.76	8.40	4.24	1.21	0.61	0.20	0.00	0.00	0.00	
Total.....	493,033	.83	2.31	4.51	10.61	20.60	40.26	65.88	96.90	126.39	145.00	145.73	120.97	92.41	66.66	34.28	10.66	7.19	2.02	1.28	.44	.30	.22	.04	

TABLE 185.—Weight distribution by special diseases or defects, for first and second million draft recruits.

SECTION A: ABSOLUTE NUMBERS.

Disease.	Total.	Weight, in pounds.																							
		90- under.	94	99	100- 104	105- 109	110- 114	115- 119	120- 124	125- 129	130- 134	135- 139	140- 144	145- 149	150- 154	155- 159	160- 164	165- 169	170- 174	175- 179	180- 184	185- 189	190- 194	195- 199	200- 204
Tuberculosis, pulmonary.....	10,701	3	9	46	167	372	718	1,186	1,462	1,498	1,460	1,142	913	642	435	285	167	133	47	29	13	9	5	8	2
Galer, simple.....	7,096	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Galer, exophthalmic.....	2,620	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Asthma.....	1,581	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Asigmatism.....	1,562	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Hyperopia.....	2,426	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Myopia.....	52,031	3	1	13	68	401	1,286	2,442	3,862	5,097	6,014	6,354	6,122	5,139	4,343	3,153	2,319	1,730	1,145	810	564	316	262	202	355
Tonsillitis, hypertrophic.....	1,343	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Cardiac hypertrophy.....	8,860	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Mitral insufficiency.....	2,512	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Atrial stenosis.....	2,512	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Tachycardia, simple.....	2,147	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Valvular diseases of heart (others).....	3,419	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Hemorrhoids.....	1,824	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Varicose veins.....	3,423	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Varicocele.....	5,849	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Defective and deficient teeth.....	17,963	5	6	19	102	218	347	1,153	1,333	1,651	2,096	2,267	2,012	1,974	1,823	1,621	1,434	1,274	1,139	911	724	581	448	364	15
Hernia.....	34,324	2	5	19	135	314	627	1,603	2,470	3,191	3,961	4,499	5,231	5,631	5,983	6,269	6,568	6,823	7,074	7,284	7,464	7,624	7,774	7,914	8,044
Inguinal rings, enlarged.....	43,619	1	1	16	143	364	1,140	2,293	3,463	4,688	5,955	7,263	8,512	9,763	10,983	12,183	13,363	14,543	15,723	16,903	18,083	19,263	20,443	21,623	22,803
Flat-foot.....	270,348	3	9	90	769	1,941	5,966	11,428	17,957	24,041	29,381	32,061	31,134	27,651	23,670	17,830	13,708	10,300	6,857	4,756	3,571	1,963	1,486	1,110	2,456
Defective physical develop- ment.....	1,292	1	30	81	138	199	176	127	103	111	90	77	60	32	24	20	19	12	7	6	1	3	4	2	1
Underweight.....	12,126	43	126	705	1,815	3,468	2,774	1,517	651	439	217	156	96	54	24	13	10	13	2	2	1	1	1	1	1
Cryptorchidism, hypospadi- anorchism, and monorchism.....	4,948	2	2	8	28	51	158	267	414	522	544	558	618	425	381	306	224	133	101	50	56	31	28	14	27
Total.....	493,053	66	163	966	3,534	7,796	15,036	24,438	35,672	46,298	54,362	58,110	55,620	48,111	40,151	29,892	22,354	16,463	10,971	7,367	5,362	3,037	2,287	1,645	3,302

SECTION B: RATIOS PER 1,000.

TABLE 186.—*Chest circumference (expiration) distribution by special disease or defects, for first and second million draft recruits*

SECTION A: ABSOLUTE NUMBERS.

Disease.	Total	Chest, in inches.													43 and over.	
		29 and under.	30	31	32	33	34	35	36	37	38	39	40	41		42
Tuberculosis, pulmonary.....	10,640	126	628	2,026	2,393	1,918	1,253	617	290	89	23	14	2	1	2	2
Gout, simple.....	7,065	26	100	438	881	1,454	1,265	808	444	202	87	38	7	1	2	1
Exophthalmic goiter.....	2,622	6	60	209	640	503	392	297	139	57	23	11	3	3	1	2
Asthma.....	1,579	6	27	74	176	313	296	254	213	110	52	27	18	6	1	2
Asigmatism.....	1,387	28	105	224	320	319	251	154	96	41	17	13	8	1	1	2
Hyperopia.....	958	20	68	123	182	205	157	113	63	18	10	6	1	2	1	2
Myopia.....	2,417	17	192	356	426	464	357	282	143	68	33	24	6	2	1	2
Tonsillitis, hypertrophic.....	51,985	131	3,067	6,364	9,616	10,435	8,769	5,887	3,369	1,620	781	554	94	43	21	37
Cardiac hypertrophy.....	1,339	7	32	91	248	276	232	138	80	34	6	3	5	2	2	1
Mitral insufficiency.....	8,630	57	310	733	1,271	1,801	1,336	823	459	199	83	33	5	3	1	4
Mitral stenosis.....	2,607	14	98	168	328	448	380	224	120	34	20	4	6	3	4	1
Tachycardia, simple.....	2,143	79	108	330	411	447	311	209	104	37	24	7	1	1	2	3
Valvular disease of heart.....	3,408	36	118	331	667	688	477	256	142	53	20	9	3	1	1	2
Hemorrhoids.....	1,619	6	32	95	132	372	333	228	88	48	14	53	6	6	2	2
Varicose veins.....	3,428	13	33	138	339	690	616	460	223	190	93	63	3	3	1	3
Varicore.....	5,856	12	132	308	757	1,075	1,204	1,011	646	357	156	64	38	3	2	1
Defective and deciduous teeth.....	17,932	106	399	1,224	2,343	3,897	2,962	1,962	1,044	453	188	91	28	13	9	3
Hernia.....	34,220	110	659	2,062	6,444	6,940	5,776	3,688	2,186	1,034	407	236	53	25	6	9
Inguinal rings, enlarged.....	43,625	114	810	2,747	8,457	9,027	7,346	4,913	2,681	1,196	472	244	32	14	3	4
.....	1,294	30	126	217	251	106	69	33	21	13	11	2	1	1	1	1
.....	12,132	946	2,276	2,837	1,556	886	256	96	15	7	5	37	1	1	1	1
monorchism.....	4,943	19	122	328	679	926	582	493	287	130	60	56	9	3	4	6
Total.....	222,334	1,782	7,309	17,836	30,654	42,681	34,622	22,790	12,626	5,737	2,477	1,518	270	123	56	78

SECTION B: RATIOS PER 1,000.

Chest, In inches.

Disease.	Total.	28 and under.	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43 and over.	Total.
Tuberculosis, pulmonary.	10,649	11.33	58.97	130.34	190.25	215.33	190.11	115.70	57.94	27.23	8.36	2.07	1.31	0.19	0.00	0.28	0.19	1,000
Galer, simple.	7,065	3.67	13.38	61.82	124.35	189.14	205.22	177.13	114.04	62.67	28.51	12.28	5.36	1.99	0.14	0.28	0.19	1,000
Exophthalmic goiter.	2,622	2.26	20.32	78.71	144.68	205.85	191.84	149.50	102.46	62.63	21.74	8.77	4.20	1.14	1.14	0.28	0.19	1,000
Asthma.	1,579	3.90	17.04	46.87	110.83	198.23	188.73	160.86	134.90	69.66	22.98	17.10	11.40	3.80	1.90	0.63	1.27	1,000
Strabismus.	1,587	17.64	66.16	141.15	207.31	201.01	158.16	97.04	60.46	26.83	10.71	6.19	5.04	0.63	0.63	2.07	1,000
Hyperopia.	968	7.03	20.66	70.25	127.07	189.02	211.78	167.70	108.40	65.08	18.60	10.33	6.20	1.03	0.83	0.41	2.07	1,000
Myopia.	2,417	22.78	78.44	147.29	180.39	191.97	166.68	113.24	69.90	28.13	13.24	9.83	2.48	0.83	0.41	0.71	1,000
Tonsillitis, hypertrophic.	51,985	2.52	20.33	60.94	122.80	185.21	200.73	173.26	103.06	66.19	31.34	15.02	10.06	1.62	0.83	0.40	0.71	1,000
Cardiac hypertrophy.	1,339	6.23	23.90	67.96	135.18	185.21	194.08	151.30	93.20	51.98	22.54	4.48	2.74	2.24	1.40	0.75	0.75	1,000
Mitral insufficiency.	8,830	6.46	33.11	83.01	143.94	203.96	194.08	151.30	93.20	51.98	22.54	4.48	2.74	2.24	1.40	0.75	0.75	1,000
Mitral stenosis.	2,507	5.58	37.89	74.19	159.15	210.61	196.65	155.98	89.35	47.87	13.66	7.98	1.60	0.87	0.23	0.11	0.45	1,000
Tachycardia, simple.	2,143	36.86	78.39	153.99	191.79	208.59	145.12	97.53	48.53	17.77	11.20	4.20	2.80	1.40	0.87	0.45	1,000
Valvular disease of heart.	3,406	10.57	34.64	97.18	163.83	201.70	196.12	140.05	83.97	41.99	18.60	5.87	4.11	0.88	0.29	0.29	0.29	1,000
Hemorrhoids.	1,819	3.79	17.59	52.23	114.90	205.61	183.07	153.07	125.34	94.28	28.39	7.70	3.85	0.55	0.29	0.29	1.10	1,000
Varicose veins.	3,426	3.79	9.63	40.23	98.95	143.32	201.40	178.80	134.27	94.28	28.39	7.70	3.85	0.55	0.29	0.29	1.10	1,000
	17,852	2.06	22.62	63.08	126.71	184.20	206.31	173.24	110.69	62.89	28.73	10.97	6.51	1.51	0.34	0.17	0.17	1,000
	34,220	5.91	21.69	68.28	130.66	195.74	198.92	164.62	110.53	68.22	28.94	10.48	6.07	1.96	0.72	0.50	0.17	1,000
	43,625	3.21	19.26	59.66	128.14	188.31	202.81	168.79	113.62	64.23	30.22	11.69	6.87	1.55	0.73	0.18	0.18	1,000
	1,294	2.61	18.57	62.97	127.59	193.96	206.92	168.39	112.62	61.46	27.39	10.82	6.99	1.73	0.82	0.07	0.09	1,000
	12,132	77.96	187.83	286.81	233.84	128.50	48.13	19.45	8.16	1.24	0.38	0.41	3.05	1.56	1,000
	4,943	3.84	24.68	66.36	137.37	198.26	187.34	168.32	99.74	58.06	28.30	13.96	11.33	1.82	0.61	0.81	1.21	1,000
Total.	222,334	8.01	33.42	80.22	137.87	188.21	191.97	155.72	102.64	56.79	25.89	11.14	6.83	1.21	0.56	0.26	0.35	1,000

TABLE 186.—Chest circumference (expiration) distribution by special diseases or defects, for first and second million draft recruits

SECTION A: ABSOLUTE NUMBERS.

Disease	Total.	Chest, in inches.															43 and over.
		29 and under.	29	30	31	32	33	34	35	36	37	38	39	40	41	42	
.....	10,449	126	628	1,386	2,028	2,293	1,918	1,233	617	200	89	22	14	2	1	2
.....	7,045	26	109	438	2,681	1,333	1,454	1,255	806	444	202	87	38	7	1	2
.....	2,622	6	69	209	390	1,640	1,603	1,282	287	138	57	23	11	3	3	1
.....	1,579	6	27	74	176	313	298	254	213	110	53	27	18	4	3	2
.....	1,547	28	105	224	329	319	251	154	96	41	17	13	6	1
.....	968	20	68	123	182	205	157	113	63	18	10	6	2
.....	2,417	17	65	182	335	436	464	357	262	145	66	32	24	6	2
.....	51,985	131	1,057	3,166	6,394	9,616	10,435	8,769	6,687	3,380	1,629	781	554	84	43	21	37
.....	1,329	7	32	91	181	248	276	232	138	80	34	6	8	3	2	1
.....	4,820	57	310	733	1,271	1,801	1,713	1,338	823	459	199	83	33	5	2	4
.....	2,507	14	95	185	320	411	447	390	224	120	34	20	4	1
.....	2,143	79	168	330	447	477	411	209	104	37	20	14	3	1	1
.....	3,408	118	331	558	687	668	477	285	142	63	30	14	3	1	1
.....	1,819	6	32	95	209	374	372	333	228	98	46	14	7	1	2
.....	3,426	13	33	138	259	491	660	616	460	223	160	93	33	8	6	3
.....	5,806	12	132	368	757	1,075	1,204	1,011	646	367	186	64	38	3	2	3
.....	17,653	106	889	1,224	2,343	3,510	3,897	2,952	1,962	1,044	483	188	91	28	13	9	3
.....	34,220	110	659	1,632	3,365	5,444	6,940	5,778	3,893	2,196	1,034	497	236	53	25	5	9
.....	45,625	114	810	2,717	5,686	8,457	9,027	7,346	4,913	2,681	1,196	472	244	32	14	3	4
.....	1,264	30	126	217	406	457	466	323	21	33	21	13	11	2
.....	12,132	946	2,279	3,516	2,837	1,559	866	286	99	15	7	5	37
monstrous,	4,943	19	122	328	679	980	926	832	493	287	130	69	56	9	3	4	6
Total.....	222,334	1,782	7,209	17,636	30,654	41,946	42,681	34,622	22,799	12,636	4,767	2,477	1,513	270	123	56	78

SECTION B: RATIOS PER 1,000.

Chest, in inches.

Disease.	Total.	28 and under.	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43 and over.	Total.
10,049	11.83	58.97	130.34	190.26	215.33	180.11	115.79	57.94	27.23	8.36	2.07	1.31	0.19	0.09	0.09	0.28	0.19	1,000
7,095	3.67	15.38	61.82	124.35	188.14	205.22	177.13	114.04	62.67	28.51	12.28	5.36	2.99	1.14	0.14	0.28	0.38	1,000
2,622	2.29	26.32	79.71	144.83	205.95	191.84	146.40	102.40	62.63	31.74	8.77	4.20	1.14	0.14	0.14	0.28	0.38	1,000
1,579	3.80	17.10	46.87	110.83	198.23	188.73	160.88	134.90	66.60	25.83	17.10	11.40	3.80	1.90	0.90	0.63	1.27	1,000
1,587	17.64	46.16	141.15	207.31	201.01	188.16	97.04	60.40	25.83	10.71	8.19	3.04	1.04	0.63	0.63	2.07	1,000
2,417	7.03	22.06	70.23	127.07	188.02	211.78	162.19	116.74	65.06	18.60	10.33	6.20	1.03	0.63	0.63	0.63	2.07	1,000
51,985	22.76	78.44	147.29	190.39	191.77	191.77	147.70	108.40	60.99	28.13	13.24	9.83	2.48	0.83	0.83	0.41	0.41	1,000
1,339	2.52	20.33	60.94	122.80	194.98	200.73	168.68	113.24	65.19	31.34	15.02	10.66	1.62	0.83	0.83	0.41	0.41	1,000
8,830	6.46	35.11	83.01	145.94	233.96	208.12	172.26	123.06	69.75	25.39	4.48	5.97	2.24	0.75	0.75	0.45	0.45	1,000
2,577	5.58	37.89	74.19	159.15	210.61	198.65	145.12	93.20	51.98	22.54	9.40	3.74	0.57	0.57	0.57	0.45	0.45	1,000
2,143	38.86	78.39	153.99	191.79	208.69	145.12	93.20	51.98	22.54	9.40	3.74	0.57	0.57	0.57	0.45	0.45	1,000
3,406	10.57	34.64	97.18	183.83	201.70	198.12	140.08	97.83	48.53	17.27	7.98	1.20	2.80	1.40	1.40	1.87	1.87	1,000
1,819	3.30	17.59	52.23	114.90	205.61	194.13	163.07	125.94	41.09	18.50	5.97	4.11	0.88	0.28	0.28	0.28	0.28	1,000
3,436	3.79	9.63	40.23	98.90	143.32	201.40	173.94	134.27	53.88	26.38	7.70	16.47	2.35	1.75	1.75	1.10	1.10	1,000
5,836	2.06	22.62	63.08	126.71	184.20	200.31	173.94	110.09	62.69	26.73	10.97	6.51	0.51	0.51	0.51	0.88	0.88	1,000
17,982	6.91	21.69	68.20	130.69	195.74	198.92	164.62	110.53	58.22	26.94	10.48	8.07	1.56	0.72	0.72	0.50	0.50	1,000
34,220	3.21	18.26	56.96	128.14	188.31	202.81	168.79	113.62	64.23	30.22	11.80	6.87	1.56	0.72	0.72	0.50	0.50	1,000
45,625	2.61	13.57	62.97	127.59	193.96	208.92	168.39	113.62	61.46	27.39	10.82	8.07	1.56	0.72	0.72	0.50	0.50	1,000
1,294	28.36	98.13	196.00	233.84	198.14	198.14	152.28	92.65	25.74	10.36	4.41	3.06	1.56	0.87	0.87	0.09	0.09	1,000
12,132	77.98	187.85	290.81	233.84	198.14	198.14	152.28	92.65	25.74	10.36	4.41	3.06	1.56	0.87	0.87	0.09	0.09	1,000
4,943	3.84	24.68	65.36	137.37	198.28	187.34	168.32	99.74	58.06	26.30	13.96	11.83	1.82	0.61	0.61	1.21	1.21	1,000
Total.....	222,334	32.42	80.22	137.87	186.21	191.97	155.72	102.64	58.79	26.89	11.14	6.83	1.21	0.56	0.56	0.25	0.35	1,000

TABLE 187.—Summary of means, standard deviations, and probable errors, coefficients of variability and probable errors, of recruits found with specified diseases and defects; also correlations between pairs of dimensions for first and second million recruits. (From Tables 139-183)

[Height and chest in inches and weight in pounds.]

					Probable error of stand- ard devia- tion.	Coeffi- cient of varia- tion.	Probable error of coeffi- cient of varia- tion.	Correla- tion.	Probable error of correla- tion.
Pulmonary tuber- culosis.....	10,701	First and second.	68.07	2.736	±0.013	0.04019	0.00019	0.4754	±0.0050
			130.44	14.740	±.088	.11300	.00053		
	4,653	First.....	68.01	2.702	±.019	.03673	.00020	.4554	±.0078
			131.77	14.850	±.108	.11346	.00079		
	6,048	Second...	68.12	2.762	±.017	.04054	.00024	.5533	±.0080
			126.42	14.858	±.090	.11094	.00067		
Simple goiter.....	10,649	First and second.	68.07	2.731	±0.013	.04011	.00019	.2412	±.0062
			32.09	1.848	±.009	.05758	.00029		
	4,627	First.....	68.02	2.693	±.019	.03669	.00021	.2391	±.0093
			32.33	1.875	±.013	.05799	.00035		
	6,022	Second...	68.12	2.759	±.017	.04050	.00024	.2499	±.0081
			31.90	1.893	±.011	.05658	.00030		
Exophthalmic goiter	7,069	First and second.	67.94	2.578	±.015	.03794	.00020	.5180	±.0059
			142.36	16.498	±.093	.11586	.00094		
	1,813	First.....	67.94	2.544	±.028	.03744	.00049	.4861	±.0121
			142.39	16.287	±.182	.11789	.00151		
	5,286	Second...	67.95	2.579	±.017	.03812	.00020	.5200	±.0067
			142.35	16.573	±.109	.11642	.00108		
Myopia.....	7,065	First and second.	67.94	2.579	±.015	.03796	.00020	.2616	±.0075
			33.11	1.950	±.011	.05999	.00032		
	1,809	First.....	67.94	2.544	±.029	.03744	.00033	.2411	±.0151
			33.04	1.893	±.011	.05996	.00035		
	5,276	Second...	67.94	2.580	±.017	.03812	.00020	.2700	±.0066
			33.13	1.953	±.013	.05995	.00040		
Hyperopia.....	2,620	First and second.	67.97	2.647	±.025	.03994	.00028	.4765	±.0402
			136.82	16.430	±.153	.11300	.00112		
	430	First.....	67.94	2.535	±.068	.03771	.00083	.4676	±.0246
			136.39	16.430	±.180	.11590	.00270		
	2,181	Second...	67.97	2.669	±.027	.03997	.00032	.4758	±.0401
			136.39	16.430	±.083	.11867	.00142		
Astigmatism.....	2,622	First and second.	67.97	2.649	±.025	.03997	.00027	.2440	±.0124
			32.35	1.978	±.018	.06015	.00056		
	430	First.....	67.94	2.535	±.068	.03731	.00083	.2439	±.0302
			33.01	1.914	±.044	.05796	.00117		
	2,183	Second...	67.97	2.672	±.027	.03991	.00039	.2454	±.0136
			32.82	1.967	±.020	.06005	.00061		
Myopia.....	2,420	First and second.	67.06	2.787	±.027	.04135	.00089	.4912	±.0104
			126.28	16.430	±.179	.13253	.00126		
	778	First.....	67.23	2.827	±.048	.04205	.00099	.5121	±.0178
			140.23	18.069	±.309	.12885	.00215		
	1,642	Second...	67.01	2.785	±.033	.04136	.00047	.4806	±.0129
			128.75	16.611	±.111	.13413	.00152		
Hyperopia.....	2,417	First and second.	67.06	2.791	±.027	.04146	.00099	.2065	±.0131
			32.97	2.119	±.021	.06427	.00068		
	776	First.....	67.23	2.831	±.049	.04131	.00099	.2177	±.0331
			33.13	2.116	±.036	.06387	.00103		
	1,641	Second...	67.01	2.780	±.033	.04119	.00047	.2028	±.0140
			32.69	2.117	±.025	.06411	.00070		
Astigmatism.....	969	First and second.	67.06	2.719	±.042	.04063	.00081	.4511	±.0178
			136.96	16.299	±.250	.11722	.00167		
	188	First.....	67.29	2.650	±.082	.03899	.00145	.4145	±.0407
			139.13	17.228	±.600	.12883	.00436		
	781	Second...	67.03	2.723	±.047	.04077	.00069	.4896	±.0190
			136.96	16.096	±.275	.11581	.00196		
Astigmatism.....	968	First and second.	67.06	2.726	±.042	.04064	.00081	.2398	±.0204
			33.06	1.977	±.030	.05962	.00092		
	188	First.....	67.29	2.650	±.082	.03899	.00145	.2840	±.0451
			33.26	2.026	±.071	.06091	.00218		
	780	Second...	67.03	2.742	±.047	.04091	.00089	.2317	±.0229
			33.00	1.982	±.034	.05945	.00104		
Astigmatism.....	1,892	First and second.	67.07	2.711	±.032	.04062	.00047	.4573	±.0124
			126.16	17.000	±.208	.12216	.00143		
	517	First.....	66.95	2.767	±.058	.04123	.00085	.5432	±.0208
			139.50	17.245	±.262	.12843	.00265		
	1,075	Second...	67.13	2.682	±.039	.03995	.00050	.4121	±.0171
			130.43	16.968	±.246	.12098	.00181		
Astigmatism.....	1,567	First and second.	67.07	2.712	±.033	.04044	.00041	.1926	±.0163
			33.08	2.014	±.024	.05977	.00074		
	517	First.....	66.95	2.767	±.058	.04133	.00085	.2513	±.0278
			33.06	2.019	±.042	.05977	.00127		
	1,070	Second...	67.13	2.684	±.039	.03999	.00050	.1641	±.0201
			33.01	2.011	±.029	.05992	.00077		

TABLE 187.—Summary of means, standard deviations, and probable errors, coefficients of variability and probable errors, of recruits found with specified diseases and defects; also correlations between pairs of dimensions for first and second million recruits. (From Tables 139-183.)—Continued.

[Height and chest in inches and weight in pounds.]

Disease.	N	Probable error of standard deviation.	Coefficient of variation.	Probable error of coefficient of variation.	Correlation.	Probable error of correlation.
Hypertrophic tonsils		± .006	.04041	.00008		
		± .087	.12556	.00025	.4762	± .0026
		± .008	.04014	.00013		
		± .055	.12501	.00042	.4638	± .0034
		± .006	.04065	.00011		
		± .050	.12613	.00034	.5001	± .0030
		± .006	.04068	.00008		
		± .004	.03243	.00012	.2065	± .0028
		± .008	.04006	.00012		
		± .006	.06101	.00013		± .0042
Tachycardia.....		± .008	.04065	.00011		
		± .006	.06342	.00016		± .0036
		± .026	.03043	.00042		
		± .181	.12791	.00136	.3757	± .0125
		± .061	.03906	.00039		
		± .392	.12666	.00230		± .0253
		± .031	.03630	.00046		
		± .204	.12629	.00148	.3523	± .0143
		± .028	.03949	.00040		
		± .021	.06224	.00061	.1789	± .0141
Cardiac hypertrophy		± .061	.04016	.00039		
		± .046	.06186	.00134	.2597	± .0296
		± .031	.03632	.00046		
		± .024	.06233	.00069	.1548	± .0300
		± .086	.04022	.00052		
		± .219	.11976	.00156	.4252	± .0151
		± .061	.04229	.00065		
		± .356	.12028	.00254	.4576	± .0238
		± .043	.03993	.00066		
		± .055	.11836	.00197	.4044	± .0196
Mitral insufficiency		± .036	.04021	.00052		
		± .026	.06075	.00079	.1948	± .0177
		± .061	.04237	.00065		
		± .043	.06153	.00127	.2633	± .0281
		± .043	.03884	.00066		
		± .033	.06023	.00093	.1487	± .0228
		± .014	.04027	.00022		
		± .065	.12081	.00061		± .0064
		± .020	.04020	.00029		
		± .122	.11949	.00067	.4860	± .0079
Mitral stenosis.....		± .019	.04033	.00028		
		± .119	.12201	.00085	.5029	± .0077
		± .014	.04024	.00022		
		± .010	.06107	.00032	.2338	± .0068
		± .020	.04020	.00029		
		± .014	.05913	.00043	.1972	± .0100
		± .019	.04026	.00028		
		± .014	.06279	.00033	.2886	± .0051
		± .026	.04028	.00038		
		± .149	.11426	.00109	.4951	± .0102
Valvular disease of heart (unclassified).		± .033	.04011	.00049		
		± .187	.11087	.00135	.4831	± .0133
		± .041	.04046	.00090		
		± .245	.11838	.00181	.5105	± .0158
		± .026	.04027	.00038		
		± .018	.06776	.00047	.2326	± .0127
		± .033	.04010	.00049		
		± .023	.06600	.00061	.2109	± .0166
		± .041	.04046	.00090		
		± .030	.06609	.00094	.2589	± .0200
Valvular disease of heart (unclassified).		± .022	.03948	.00024		
		± .142	.12641	.00100	.4546	± .0092
		± .042	.03952	.00063		
		± .261	.11808	.00190	.5023	± .0167
		± .023	.03946	.00038		
		± .166	.12720	.00119	.4459	± .0108
		± .022	.03951			
		± .016	.03078	.00046	.2020	± .0111
		± .042	.03946	.00063		
		± .030	.05749	.00090	.2445	± .0211
Valvular disease of heart (unclassified).		± .026	.03951	.00036		
		± .019	.06177	.00047	.1896	± .0130

TABLE 187.—Summary of means, standard deviations, and probable errors, coefficients of variability and probable errors, of recruits found with specified diseases and defects; also correlations between pairs of dimensions for first and second million recruits. (From Tables 139-183)—Continued

(Height and chest in inches and weight in pounds.)

						Probable error of stand- ard devia- tion.	Coeffi- cient of varia- tion.	Probable error of coeffi- cient of varia- tion.	Correla- tion.	Probable error of correla- tion.
						2	± .022	.04007	.00033	± .0090
						8	± .151	.12532	.00131	± .0136
						6	± .034	.03945	.00497	± .0115
						9	± .234	.12558	.00159	± .0110
						2	± .030	.04047	.00042	± .0172
						5	± .198	.12718	.00135	± .0144
						5	± .022	.04011	.00033	± .0067
						8	± .017	.06350	.00046	± .0086
						6	± .034	.03955	.00051	± .0106
						7	± .027	.06341	.00076	± .0084
						2	± .029	.04047	.00042	± .0107
						8	± .023	.06356	.00064	± .0133
						3	± .017	.04027	.00025	± .0115
						4	± .103	.11622	.00073	± .0155
						9	± .023	.04058	.00033	± .0172
						6	± .136	.11754	.00084	± .0178
						5	± .026	.03967	.00038	± .0142
						8	± .158	.11429	.00111	± .0166
						8	± .017	.04004	.00025	± .0133
						5	± .012	.03944	.00039	± .0107
						4	± .022	.04030	.00033	± .0133
						1	± .016	.05869	.00041	± .0133
						2	± .026	.03963	.00038	± .0133
						4	± .019	.05959	.00056	± .0133
Hemorrhoids.....	1,824	First and second.	t....	67.80	2.782	± .031	.04103	.00045	.5219	± .0115
			t....	140.39	16.757	± .187	.11936	.00134		
	1,027	First.....	t....	67.82	2.681	± .040	.03953	.00080	.5115	± .0155
			t....	141.44	16.784	± .250	.11867	.00177		
	797	Second.....	t....	67.77	2.906	± .049	.04288	.00067	.5285	± .0172
			t....	139.06	16.747	± .283	.12043	.00202		
Asthma.....	1,819	First and second.	t....	67.80	2.783	± .031	.04105	.00045	.5202	± .0150
			t....	33.10	1.884	± .021	.05869	.00067		
	1,024	First.....	t....	67.82	2.680	± .040	.03952	.00080	.5230	± .0208
			t....	33.22	1.869	± .028	.05826	.00075		
	796	Second.....	t....	67.77	2.910	± .049	.04294	.00067	.5169	± .0228
			t....	32.94	1.892	± .032	.05744	.00084		
Asthma.....	1,681	First and second.	t....	67.24	2.710	± .033	.04030	.00047	.4069	± .0142
			t....	139.01	17.945	± .215	.12902	.00155		
	614	First.....	t....	67.82	2.770	± .053	.04121	.00077	.3633	± .0232
			t....	139.38	17.280	± .333	.12398	.00260		
	967	Second.....	t....	67.26	2.670	± .041	.03970	.00061	.4226	± .0178
			t....	139.78	18.351	± .282	.13223	.00217		
Dental caries, defective and deficient teeth.	1,579	First and second.	t....	67.25	2.710	± .033	.04030	.00047	.1477	± .0166
			t....	33.34	2.120	± .025	.05359	.00071		
	612	First.....	t....	67.25	2.771	± .053	.04122	.00077	.1274	± .0288
			t....	33.57	2.114	± .041	.05297	.00116		
	967	Second.....	t....	67.26	2.670	± .041	.03970	.00061	.1628	± .0211
			t....	33.19	2.112	± .032	.05363	.00092		
Hemorrhoids.....	17,983	First and second.	Height....	67.26	2.689	± .010	.03998	.00010	.5067	± .0087
			Weight....	138.32	16.889	± .080	.12210	.00043		
	5,166	First.....	Height....	67.26	2.676	± .019	.03979	.00027	.5107	± .0080
			Weight....	139.18	16.839	± .112	.12089	.00081		
	12,817	Second.....	Height....	67.26	2.694	± .011	.04006	.00016	.5064	± .0044
			Weight....	137.97	16.900	± .071	.12249	.00051		
Hemorrhoids.....	17,982	First and second.	Height....	67.26	2.686	± .010	.03998	.00010	.2551	± .0047
			Chest....	33.00	2.004	± .007	.06073	.00021		
	5,150	First.....	Height....	67.26	2.674	± .018	.03976	.00027	.2713	± .0087
			Chest....	33.25	1.943	± .013	.05844	.00059		
	12,782	Second.....	Height....	67.26	2.690	± .011	.03999	.00013	.2493	± .0056
			Chest....	32.89	2.018	± .009	.06136	.00026		
Hemorrhoids.....	34,324	First and second.	Height....	67.44	2.762	± .007	.04095	.00010	.5188	± .0134
			Weight....	141.23	17.187	± .044	.12155	.00030		
	13,870	First.....	Height....	67.40	2.743	± .011	.04070	.00016	.5285	± .0054
			Weight....	141.69	17.221	± .070	.12154	.00048		
	20,454	Second ..	Height....	67.47	2.774	± .009	.04111	.00013	.5130	± .0047
			Weight....	140.91	17.122	± .011	.12151	.00040		
Hemorrhoids.....	34,220	First and second.	Height....	67.44	2.760	± .007	.04093	.00010	.2426	± .0027
			Chest....	33.11	2.002	± .006	.06047	.00015		
	13,822	First.....	Height....	67.40	2.741	± .011	.04067	.00016	.2515	± .0041
			Chest....	33.23	1.991	± .008	.05992	.00020		
	20,398	Second ..	Height....	67.47	2.772	± .010	.04108	.00042	.2372	± .0035
			Chest....	33.04	2.005	± .007	.06098	.00044		

TABLE 187.—Summary of means, standard deviations, and probable errors, coefficients of variability and probable errors, of recruits found with specified diseases and defects; also correlations between pairs of dimensions for first and second million recruits (From Tables 139-183)—Continued.

[Height and chest in inches and weight in pounds.]

Disease.	Number measured.	First or second million.		Stand-ard deviation.	Probable error of stand-ard deviation.	Coeffi-ent of varia-tion.	Probable error of coeffi-ent of varia-tion.	Correla-tion.	Probable error of correla-tion.	
Enlarged inguinal rings.	43,619	{First and second.	6	2.702	± .006	.04006	.00009	.5115	± .0024	
			6	16.543	± .038	.11810	.00028			
	20,142	First.....	4	2.695	± .009	.03990	.00010	.5174	± .0035	
			7	16.637	± .056	.11899	.00040			
	23,477	Second...	0	2.708	± .008	.04015	.00013	.5077	± .0033	
			0	16.462	± .051	.11759	.00038			
	43,625	{First and second.	7	2.706	± .006	.04025	.00009	.2310	± .0031	
			6	1.945	± .004	.05883	.00017			
Flat-foot.....	20,161	First.....	5	2.701	± .009	.03999	.00013	.2410	± .0045	
			3	1.916	± .006	.05801	.00019			
	23,464	Second ..	0	2.708	± .008	.04018	.00013	.2237	± .0042	
			9	1.959	± .006	.05960	.00020			
	270,348	{First and second.	0	2.699	± .003	.04010	.00004	.4721	± .0010	
			6	18.413	± .017	.12853	.00012			
	175,358	First.....	0	2.687	± .059	.03993	.00004	.4786	± .0012	
			4	18.102	± .021	.12838	.00014			
Defective physical development.	94,990	Second...	6	2.723	± .004	.04047	.00005	.4610	± .0017	
			1	18.975	± .030	.13241	.00020			
	1,292	{First and second.	7	3.842	± .051	.05771	.00066	.4644	± .0157	
			1	18.568	± .246	.14794	.00186			
	758	First.....	4	4.012	± .070	.09048	.00105	.4600	± .0193	
			4	18.144	± .315	.14072	.00255			
	534	Second...	1	3.561	± .074	.06322	.00103	.5008	± .0219	
			3	18.951	± .391	.15362	.00289			
Underweight.....	1,284	{First and second.	7	3.841	± .051	.05770	.00067	.1997	± .0181	
			6	2.180	± .029	.05845	.00080			
	752	First....	4	4.015	± .070	.06256	.00104	.1792	± .0238	
			5	2.206	± .038	.06862	.00105			
	532	Second...	0	3.556	± .074	.06315	.00103	.2483	± .0274	
			3	2.071	± .043	.06689	.00124			
	12,129	{First and second.	Height.....	65.50	3.360	± .015	.05130	.00021	.6970	± .0031
			Weight.....	110.94	9.893	± .043	.08917	.00039		
Cryptorchidism, monorchism, anorchism, hypospadi.	2,686	First.....	Height.....	66.22	3.507	± .032	.05296	.00046	.7339	± .0060
			Weight.....	114.67	11.514	± .107	.10128	.00093		
	9,443	Second....	Height.....	65.30	3.289	± .016	.05037	.00025	.6873	± .0037
			Weight.....	109.88	9.070	± .046	.07566	.00035		
	12,132	{First and second.	Height.....	65.50	3.357	± .015	.05125	.00021	.2469	± .0058
			Chest.....	30.46	1.531	± .007	.05028	.00021		
	2,708	First.....	Height.....	66.20	3.509	± .032	.05301	.00046	.2843	± .0019
			Chest.....	30.94	1.720	± .016	.05559	.00046		
	9,424	Second....	Height.....	65.30	3.285	± .016	.05031	.00025	.2312	± .0066
			Chest.....	30.32	1.442	± .007	.04756	.00021		
	4,948	{First and second.	Height.....	67.44	2.811	± .019	.04168	.00027	.4867	± .0073
			Weight.....	140.25	17.908	± .121	.12769	.00087		
	1,806	First.....	Height.....	67.34	2.803	± .031	.04162	.00045	.5186	± .0116
			Weight.....	140.81	18.608	± .209	.13214	.00146		
	3,140	Second....	Height.....	67.49	2.814	± .024	.04170	.00034	.4666	± .0094
			Weight.....	139.93	17.483	± .149	.12494	.00108		
	4,943	{First and second.	Height.....	67.44	2.812	± .019	.04170	.00027	.2107	± .0092
			Chest.....	33.03	2.102	± .014	.06364	.00040		
	1,806	First.....	Height.....	67.34	2.803	± .031	.04162	.00045	.2299	± .0150
			Chest.....	33.18	2.050	± .023	.06178	.00067		
	3,135	Second....	Height.....	67.49	2.816	± .024	.04172	.00034	.2029	± .0115
			Chest.....	32.95	2.126	± .018	.06452	.00052		

TABLE 188.—*The mean stature and weight of recruits found with the specified diseases and defects among the first two million draft recruits, arranged in descending order of the means.*

Defect.	Mean stature.	Defect.	Mean weight.
	Inches.		Pounds.
Varicose veins.....	68.43	Varicose veins.....	155.44
Varicocele.....	68.37	Flat-foot.....	153.26
Pulmonary tuberculosis.....	68.07	Simple goiter.....	142.36
Exophthalmic goiter.....	67.97	Hypertrophic tonsillitis.....	141.79
Simple goiter.....	67.94	Varicocele.....	141.75
Mitral insufficiency.....	67.84	Hernia.....	141.23
Hemorrhoids.....	67.80	Cardiac hypertrophy.....	140.49
Cardiac hypertrophy.....	67.75	Hemorrhoids.....	140.29
Tachycardia.....	67.76	Cryptorchidism.....	140.25
Mitral stenosis.....	67.63	Enlarged inguinal rings.....	140.08
Valvular diseases of heart.....	67.60	Myopia.....	139.23
Hypertrophic tonsillitis.....	67.48	Astigmatism.....	138.16
Enlarged inguinal rings.....	67.46	Asthma.....	138.01
Hernia.....	67.44	Mitral insufficiency.....	138.99
Cryptorchidism, etc.....	67.44	Hyperopia.....	138.98
Flat-foot.....	67.30	Exophthalmic goiter.....	138.82
Defective teeth.....	67.28	Defective and deficient teeth.....	138.32
Asthma.....	67.24	Tachycardia.....	137.37
Myopia.....	67.08	Valvular diseases of heart.....	137.24
Hyperopia.....	67.08	Mitral stenosis.....	136.85
Astigmatism.....	67.07	Pulmonary tuberculosis.....	130.44
Defective physical development.....	66.57	Defective physical development.....	125.51
Underweight.....	65.60	Underweight.....	110.94
Average, United States, first million.....	67.49	Average, United States, first million.....	141.54

TABLE 189.—*The index of build (weight multiplied by 1,000, divided by the stature squared) and Pignet's index of robustness of recruits found with the specified diseases and defects, arranged in order of standing, first and second million draft recruits.*

Defect.	Index of build.	Defect.	Pignet's index.	Class.
Flat-foot.....	31.63	V.....	19.90	Good.
Varicose veins.....	31.26	H.....	20.86	Average.
Hypertrophic tonsillitis.....	31.14	A.....	21.09	Do.
Hernia.....	31.06	H.....	21.18	Do.
Myopia.....	30.96	A.....	21.39	Do.
Astigmatism.....	30.94	H.....	21.44	Do.
Hyperopia.....	30.86	M.....	21.53	Do.
Cryptorchidism.....	30.84	Ci.....	21.83	Do.
Simple goiter.....	30.84	E.....	21.89	Do.
Enlarged inguinal rings.....	30.78	Si.....	21.94	Do.
Asthma.....	30.75	D.....	22.51	Do.
Cardiac hypertrophy.....	30.61	H.....	22.50	Do.
Defective teeth.....	30.56	Ci.....	22.66	Do.
Hemorrhoids.....	30.54	V.....	23.43	Do.
Varicocele.....	30.53	M.....	24.12	Do.
Mitral insufficiency.....	30.50	E.....	24.26	Do.
Exophthalmic goiter.....	30.06	T.....	24.50	Do.
Valvular diseases of heart (unclassified).....	30.04	V.....	24.78	Do.
Mitral stenosis.....	29.98	M.....	24.81	Do.
Tachycardia, simple.....	29.92	D.....	28.94	Weak.
Defective physical development.....	28.82	P.....	30.27	Do.
Pulmonary tuberculosis.....	28.15	U.....	37.36	Inadequate.
Underweight.....	25.86			

TABLE 190.—*Variability (standard deviation, in inches, and coefficient of variability) of stature, associated with various defects and diseases, first and second million draft recruits.*

Defect.	Standard of deviation.	Defect.	Coefficient of variability.
Defective physical development.....	3.842	D	0.08771
Underweight.....	3.300	U	.05130
Cryptorchidism, etc.....	2.811	C	.04168
Myopia.....	2.787	M	.04155
Hemorrhoids.....	2.783	H	.04103
Hernia.....	2.783	H	.04096
Varicocele.....	2.753	V	.04063
Varicose veins.....	2.742	V	.04042
Pulmonary tuberculosis.....	2.736	P	.04041
Mitral insufficiency.....	2.732	M	.04030
Hypertrophic tonsils.....	2.727	H	.04028
Cardiac hypertrophy.....	2.725	C	.04027
Mitral stenosis.....	2.724	M	.04027
Hyperopia.....	2.719	H	.04022
Astigmatism.....	2.711	A	.04019
Asthma.....	2.710	A	.04010
Enlarged inguinal rings.....	2.702	E	.04007
Flat-foot.....	2.699	F	.04006
Defective teeth.....	2.689	D	.03998
Tachycardia.....	2.675	T	.03948
Valvular diseases of the heart (unclassified).....	2.669	V	.03948
Exophthalmic goiter.....	2.647	E	.03884
Simple goiter.....	2.578	S	.03794
United States first million recruits.....	2.714	U	.04021

TABLE 191.—*Variability (standard deviation, in pounds, and coefficient of variability) of weight, associated with various diseases and defects among first and second million draft recruits, arranged in descending order of size.*

Defect.	Standard of deviation.	Defect.	Coefficient of variability.
Varicose veins.....	18.528	Defective physical development.....	0.14794
Defective physical development.....	18.568	Myopia.....	.13283
Myopia.....	18.452	Asthma.....	.12902
Flat-foot.....	18.413	Flat-foot.....	.12863
Asthma.....	17.945	Tachycardia.....	.12791
Cryptorchidism.....	17.908	Cryptorchidism.....	.12769
Hypertrophic tonsils.....	17.808	Varicose veins.....	.12652
Tachycardia.....	17.571	Valvular diseases of the heart (unclassified).....	.12641
Valvular diseases of the heart (unclassified).....	17.348	Hypertrophic tonsils.....	.12556
Hernia.....	17.167	Astigmatism.....	.12216
Astigmatism.....	17.000	Defective teeth.....	.12210
Defective teeth.....	16.899	Hernia.....	.12155
Cardiac hypertrophy.....	16.845	Mitral insufficiency.....	.12061
Mitral insufficiency.....	16.791	Cardiac hypertrophy.....	.11976
Hemorrhoids.....	16.757	Hemorrhoids.....	.11936
Enlarged inguinal rings.....	16.543	Exophthalmic goiter.....	.11832
Simple goiter.....	16.498	Enlarged inguinal rings.....	.11810
Varicocele.....	16.474	Hyperopia.....	.11732
Exophthalmic goiter.....	16.425	Varicocele.....	.11622
Hyperopia.....	16.299	Simple goiter.....	.11588
Mitral stenosis.....	15.637	Mitral stenosis.....	.11426
Pulmonary tuberculosis.....	14.740	Pulmonary tuberculosis.....	.11320
Underweight.....	9.893	Underweight.....	.08917
United States first million recruits.....	17.420	United States first million recruits.....	.12307

TABLE 192.—*Relative weight (weight divided by the height) and relative chest circumference (chest circumference (expiration) divided by the height and also by the weight) for men found with special diseases or defects in the first and second million draft recruits, 1917-18.*

Special disease.	Number of men measured.	Mean weight.	Mean chest.	Mean chest.
		Mean height.	Mean height.	Mean weight.
		Pounds.	Inch.	Inch.
Average for the United States (P ₁).....	873, 159	2.097	0.492	0.234
Varicose veins.....	3, 426	2.140	.492	.230
Varicocele.....	5, 849	2.070	.484	.233
Tuberculosis, pulmonary.....	10, 701	1.920	.472	.246
Exophthalmic goiter.....	2, 622	2.040	.483	.237
Goiter, simple.....	7, 009	2.100	.487	.233
Mitral insufficiency.....	8, 860	2.060	.483	.236
Hemorrhoids.....	1, 824	2.070	.488	.236
Tachycardia, simple.....	2, 147	2.030	.484	.239
Cardiac hypertrophy.....	1, 343	2.070	.487	.235
Mitral stenosis.....	2, 512	2.020	.483	.239
Valvular disease of heart.....	3, 419	2.080	.482	.237
Tonsillitis, hypertrophic.....	52, 031	2.100	.492	.234
Inguinal rings, enlarged.....	43, 625	2.090	.490	.236
Hernia.....	34, 324	2.090	.491	.234
Anorchism, monorchism, cryptorchidism, and hypospadias.....	4, 948	2.080	.490	.235
Flat-foot.....	270, 348	2.130		
Defective and deficient teeth.....	17, 963	2.060	.491	.239
Asthma.....	1, 581	2.070	.496	.240
Hyperopia.....	969	2.070	.493	.238
Myopia.....	2, 420	2.080	.492	.237
Astigmatism.....	1, 582	2.030	.493	.237
Defective physical development.....	1, 292	1.890	.479	.254
Underweight.....	1, 432	1.690	.465	.275

TABLE 193.—*Table for converting centimeters into inches.*

1 centimeter=0.393704 inch.
 1 decimeter=3.937040 inches.
 1 meter=39.370400 inches.

Centimeters.	Inches.	Centimeters.	Inches.	Centimeters.	Inches.	Centimeters.	Inches.
1.....	0.394	51.....	20.079	101.....	39.764	151.....	59.449
2.....	0.787	52.....	20.473	102.....	40.158	152.....	59.843
3.....	1.181	53.....	20.866	103.....	40.552	153.....	60.237
4.....	1.575	54.....	21.260	104.....	40.945	154.....	60.630
5.....	1.969	55.....	21.654	105.....	41.339	155.....	61.024
6.....	2.362	56.....	22.047	106.....	41.733	156.....	61.418
7.....	2.756	57.....	22.441	107.....	42.126	157.....	61.812
8.....	3.150	58.....	22.835	108.....	42.520	158.....	62.205
9.....	3.543	59.....	23.229	109.....	42.914	159.....	62.599
10.....	3.937	60.....	23.622	110.....	43.307	160.....	62.993
11.....	4.331	61.....	24.016	111.....	43.701	161.....	63.386
12.....	4.724	62.....	24.410	112.....	44.095	162.....	63.780
13.....	5.118	63.....	24.803	113.....	44.489	163.....	64.174
14.....	5.512	64.....	25.197	114.....	44.882	164.....	64.567
15.....	5.906	65.....	25.591	115.....	45.276	165.....	64.961
16.....	6.299	66.....	25.984	116.....	45.670	166.....	65.355
17.....	6.693	67.....	26.378	117.....	46.063	167.....	65.749
18.....	7.087	68.....	26.772	118.....	46.457	168.....	66.142
19.....	7.480	69.....	27.166	119.....	46.851	169.....	66.536
20.....	7.874	70.....	27.559	120.....	47.244	170.....	66.930
21.....	8.268	71.....	27.953	121.....	47.638	171.....	67.323
22.....	8.661	72.....	28.347	122.....	48.032	172.....	67.717
23.....	9.055	73.....	28.740	123.....	48.426	173.....	68.111
24.....	9.449	74.....	29.134	124.....	48.819	174.....	68.504
25.....	9.843	75.....	29.528	125.....	49.213	175.....	68.898
26.....	10.236	76.....	29.922	126.....	49.607	176.....	69.292
27.....	10.630	77.....	30.315	127.....	50.000	177.....	69.686
28.....	11.024	78.....	30.709	128.....	50.394	178.....	70.079
29.....	11.417	79.....	31.103	129.....	50.788	179.....	70.473
30.....	11.811	80.....	31.496	130.....	51.182	180.....	70.867
31.....	12.205	81.....	31.890	131.....	51.575	181.....	71.260
32.....	12.599	82.....	32.284	132.....	51.969	182.....	71.654
33.....	12.992	83.....	32.677	133.....	52.363	183.....	72.048
34.....	13.386	84.....	33.071	134.....	52.756	184.....	72.442
35.....	13.780	85.....	33.465	135.....	53.150	185.....	72.835
36.....	14.173	86.....	33.859	136.....	53.544	186.....	73.229
37.....	14.567	87.....	34.252	137.....	53.937	187.....	73.623
38.....	14.961	88.....	34.646	138.....	54.331	188.....	74.016
39.....	15.354	89.....	35.040	139.....	54.725	189.....	74.410
40.....	15.748	90.....	35.433	140.....	55.119	190.....	74.804
41.....	16.142	91.....	35.827	141.....	55.512	191.....	75.197
42.....	16.536	92.....	36.221	142.....	55.906	192.....	75.591
43.....	16.929	93.....	36.614	143.....	56.300	193.....	75.985
44.....	17.323	94.....	37.008	144.....	56.693	194.....	76.379
45.....	17.717	95.....	37.402	145.....	57.087	195.....	76.772
46.....	18.110	96.....	37.796	146.....	57.481	196.....	77.166
47.....	18.504	97.....	38.189	147.....	57.874	197.....	77.560
48.....	18.898	98.....	38.583	148.....	58.268	198.....	77.953
49.....	19.291	99.....	38.977	149.....	58.662	199.....	78.347
50.....	19.685	100.....	39.370	150.....	59.056	200.....	78.740

TABLE 194.—*Table for converting inches into centimeters.*

1 inch=2.539979 centimeters.
1 foot=30.479748 centimeters.

Inches.	Centi- meters.	Inches.	Centi- meters.	Inches.	Centi- meters.	Inches.	Centi- meters.
1.....	2.540	26.....	66.039	51.....	129.539	76.....	193.038
2.....	5.080	27.....	68.579	52.....	132.079	77.....	195.578
3.....	7.620	28.....	71.119	53.....	134.619	78.....	198.118
4.....	10.160	29.....	73.659	54.....	137.159	79.....	200.658
5.....	12.700	30.....	76.199	55.....	139.699	80.....	203.198
6.....	15.240	31.....	78.739	56.....	142.239	81.....	205.738
7.....	17.780	32.....	81.279	57.....	144.779	82.....	208.278
8.....	20.320	33.....	83.819	58.....	147.319	83.....	210.818
9.....	22.860	34.....	86.359	59.....	149.859	84.....	213.358
10.....	25.400	35.....	88.899	60.....	152.399	85.....	215.898
11.....	27.940	36.....	91.439	61.....	154.939	86.....	218.438
12.....	30.480	37.....	93.979	62.....	157.479	87.....	220.978
13.....	33.020	38.....	96.519	63.....	160.019	88.....	223.518
14.....	35.560	39.....	99.059	64.....	162.559	89.....	226.058
15.....	38.100	40.....	101.599	65.....	165.099	90.....	228.598
16.....	40.640	41.....	104.139	66.....	167.639	91.....	231.138
17.....	43.180	42.....	106.679	67.....	170.179	92.....	233.678
18.....	45.720	43.....	109.219	68.....	172.719	93.....	236.218
19.....	48.260	44.....	111.759	69.....	175.259	94.....	238.758
20.....	50.800	45.....	114.299	70.....	177.799	95.....	241.298
21.....	53.340	46.....	116.839	71.....	180.339	96.....	243.838
22.....	55.880	47.....	119.379	72.....	182.879	97.....	246.378
23.....	58.420	48.....	121.919	73.....	185.418	98.....	248.918
24.....	60.959	49.....	124.459	74.....	187.958	99.....	251.458
25.....	63.499	50.....	126.999	75.....	190.498	100.....	253.998

REFERENCES.

- (1) Baxter, J. H.: Statistics, Medical and Anthropological, of the Provost Marshal General's Bureau. Derived from records of the examination for military service in the armies of the United States during the late War of the Rebellion, of over a million recruits, drafted men, substitutes, and volunteers. Washington, D. C., Government Printing Office, 1875. 2 vols.
- (2) Gould, B. A.: Investigations in the Military and Anthropological Statistics of American Soldiers. For the U. S. Sanitary Commission. New York: Hurd and Houghton, 1869.
- (3) Letter of A. B. Coxe, Lieutenant Colonel, General Staff, Executive Division, July 16, 1918, No. 10905-10, including memo. from Combat Subsection, Military Information 2.
- (4) Manouvrier, L.: Note in *American Journal of Physical Anthropology*. Washington, D. C., 1918, I, 376.
- (5) Martin, R.: *Lehrbuch der Anthropologie*. Jena: Gustav Fisher, 1914.
- (6) Second Report of the Provost Marshal General. Washington, D. C., Government Printing Office, 1918, 31.
- (7) First Report of the Provost Marshal General. Washington, D. C., Government Printing Office, 1917, 44.
- (8) Second report of the Provost Marshal General. Washington, D. C., Government Printing Office, 1918, 463.
- (9) Love, A. G., and Davenport, C. B.: Defects Found in Drafted Men. Prepared under the direction of the Surgeon General, M. W. Ireland, War Department. Washington, D. C., Government Printing Office, 1920.
- (10) Verbal order, S. G. O., October, 1918.
- (11) Joint communication, Provost Marshal General and Surgeon General to Adjutant General, December 9, 1917; indorsement, January 28, 1918. On file, Record Room, S. G. O., No. 413.5.
- (12) Office order No. 68, S. G. O., July 23, 1918. On file, Record Room, S. G. O., No. .024.
- (13) Letter, S. G. O., June 6, 1918; approved June 27, 1918. A. G. O. File No. 320.21, Med. Dept. (Misc. Div.).
- (14) Deniker, J.: *The Races of Man*. Contemporary Science Series. London: The Walter Scott Publishing Company; and New York: Charles Scribner's Sons, 1906, xxiii, 611.
- (15) Harris, J. A., and Benedict, F. G.: *A Biometric Study of Basal Metabolism in Man*. Washington, Carnegie Institution, 1919, Publication No. 279, vi.
- (16) Hoffman, F. L.: *Army Anthropometry and Medical Rejection Statistics*. Newark, N. J., Prudential Press, 1917.
- (17) Livi, R.: *Anthropometria Militarie*. Rome: Presso il Giornale Medico del Regio Esercico, 1905. 2 vols.
- (18) Ammon, O.: *Zur Anthropologie der Badener*. Jena: Gustav Fisher, 1899.
- (19) Love, A. G., and Davenport, C. B.: *Physical Examination of the First Million Draft Recruits, Methods and Results*. Compiled under direction of the Surgeon General, M. W. Ireland. Washington, D. C., Government Printing Office, 1919.
- (20) Pignet: Du coefficient de robusticité. *Le Bulletin Médical*, Paris, 1901, XV, No. 33.
- (21) Butza: Sur la valeur du coefficient de robusticité, Pignet. *La Caducée*, Paris, 1905, v.
- (22) *Manual of Surgical Anatomy*, Medical Department U. S. Army and Navy. Washington, D. C., 1918.
- (23) Papillault, G.: L'homme moyen à Paris. *Bull. Soc. Anthropol.*, Paris, 1902, (5) iii.
- (24) Dublin, Louis, and Baker, G. W.: *Mortality of Race Stocks in Pennsylvania and New York*. Quarterly Publication of American Statistical Association, Concord, N. H., 1920, xvii, March.
- (25) Great Britain, Ministry of National Service, 1917-1919: Report upon the physical examination of men of military age by National Service Medical Board from November 1, 1917, to October 31, 1918. London, His Majesty's Stationery Office, 1920, Vol. I.

ADDITIONAL AUTHORITIES CONSULTED.

- (1) Final Report of the Provost Marshal General to the Secretary of War on the Operations of the Selective Service System to July 15, 1919. Washington, D. C., Government Printing Office, 1920.
- (2) Ripley, W. F.: *The Races of Europe*. New York: D. Appleton and Company, 1899.
- (3) Sergi, G.: *The Mediterranean Race*. Contemporary Science Series. London: The Walter Scott Publishing Company, and New York: Charles Scribner's Sons, 1901, xii.
- (4) Weissenberg, S.: *De Siidrussischen Juden*. *Arch. f. Anthropol.*, Braunschweig, 1895, xxiii.
- (5) Pearson, Karl: *Mathematical Contributions, etc.* London: Dulau and Company, Ltd., 1899.

APPENDIX

TABLE I.—Correlation between height and weight for first million draft recruits.
SECTION A: ABSOLUTE NUMBERS DERIVED FROM SUMMATION OF SECTION.

Height in inches.	Total.	Weight, in pounds.																					
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204
58 and under 59.....	3,124	14	69	111	201	259	252	333	325	310	296	243	209	153	93	124	55	17	15	10	8	15	12
60.....	2,877	10	108	245	367	395	350	270	255	186	165	135	106	92	63	52	24	6	13	4	11	14	7
61.....	7,477	32	240	616	1,138	1,297	1,123	927	643	440	331	210	106	100	75	59	31	23	14	8	22	36	4
62.....	15,644	36	351	975	2,081	2,557	2,221	1,674	1,074	643	456	285	194	100	75	59	31	23	14	8	22	36	4
63.....	30,935	38	396	1,263	3,174	4,644	4,972	4,740	3,846	2,777	1,811	1,211	727	418	306	214	136	78	47	31	30	42	10
64.....	52,547	15	358	1,306	3,812	6,413	7,969	8,104	7,365	5,902	3,974	2,663	1,714	1,018	684	454	255	186	121	83	46	56	37
65.....	81,904	17	276	1,178	3,911	7,859	10,711	12,344	12,147	10,563	7,773	5,459	3,509	2,159	1,403	834	571	401	267	159	112	99	85
66.....	109,964	8	202	778	3,042	7,264	11,780	15,071	16,431	15,824	12,610	8,553	5,378	3,911	2,554	1,744	1,089	719	426	292	203	167	198
67.....	127,844	1	128	497	1,902	5,483	10,196	15,018	18,123	18,872	15,081	10,786	6,900	4,376	2,833	1,945	1,389	908	608	496	356	281	318
68.....	126,987	3	77	251	1,005	3,105	7,320	12,018	16,310	18,872	15,081	10,786	6,900	4,376	2,833	1,945	1,389	908	608	496	356	281	318
69.....	110,508	1	36	100	442	1,453	3,863	7,607	11,994	14,449	11,378	7,482	4,582	2,751	1,803	1,203	824	503	303	207	146	94	683
70.....	83,702	50	65	189	573	1,676	4,676	9,709	14,394	19,636	14,449	11,378	7,482	4,582	2,751	1,803	1,203	824	503	303	207	146	94
71.....	54,357	28	34	45	191	551	1,440	3,016	5,016	6,944	6,524	7,283	7,354	6,308	5,088	3,896	2,508	1,726	1,146	777	517	360	681
72.....	31,370	18	23	41	60	166	335	1,277	2,073	3,152	3,155	3,721	4,285	3,455	2,886	2,018	1,399	892	552	426	256	155	335
73.....	15,186	6	12	17	12	35	127	301	488	665	1,187	1,563	1,926	1,612	1,312	939	860	608	377	252	160	116	228
74.....	6,391	3	4	5	5	6	16	37	88	174	339	503	747	754	612	739	639	400	212	160	116	63	127
75.....	2,620	3	4	5	5	6	8	16	33	97	162	234	316	304	328	328	279	224	182	130	78	32	60
76.....	1,071	2	2	2	2	2	2	11	16	27	55	79	112	109	109	133	137	96	85	49	32	26	25
77.....	360	2	2	2	2	2	2	3	6	19	27	28	23	25	22	40	40	28	33	14	12	14	14
78.....	259	2	2	2	2	2	2	1	10	18	21	21	21	16	18	20	18	22	17	10	8	10	24
79.....	296	2	2	2	2	2	2	13	16	16	26	28	23	16	21	21	18	9	10	12	8	10	24
Total.....	868,445	184	2,356	7,435	21,388	41,503	63,567	84,726	100,964	106,889	100,607	88,057	72,362	53,431	39,797	29,063	18,954	12,629	8,385	5,467	3,907	2,966	4,688

Number of cases: 868,445. Height: Mean, 67.49 inches; standard deviation, 2.71±0.0014 inch. Weight: Mean, 141.54 pounds; standard deviation, 17.42±0.0089 pound. Correlation: 0.4810±0.0006.

TABLE I.—*Correlation between height and weight for first million draft recruits—Continued.*
SECTION B: RATIO PER 1,000 OF THE SEPARATE WEIGHTS SHOWN FOR EACH HEIGHT.

Height, in inches.	Proportion each height per 1,000.	Weight, in pounds.																			Total. ^a			
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189		190-194	195-199	200-204
59	3.63	4.48	22.09	35.53	64.34	82.91	80.67	106.59	104.03	98.23	94.75	77.78	66.90	48.98	29.77	39.69	17.60	5.44	4.80	3.20	2.56	4.80	3.84	1,000—
60	3.32	6.58	37.41	84.38	127.12	161.70	150.19	123.98	88.03	64.43	57.16	46.77	36.72	31.87	21.82	18.01	8.31	2.08	4.50	1.39	3.81	2.56	4.80	1,000—
61	8.61	4.28	32.10	62.33	133.02	163.45	163.19	141.97	97.01	73.30	45.64	28.15	26.71	23.40	10.03	7.89	4.15	3.08	1.87	1.07	3.81	2.56	4.80	1,000—
62	18.01	2.30	22.44	40.83	102.60	153.12	160.72	153.22	124.32	89.77	58.52	39.15	23.50	13.51	8.90	6.92	3.39	2.43	1.66	1.07	3.81	2.56	4.80	1,000—
63	35.62	1.23	6.81	24.90	72.56	122.04	151.65	154.22	140.16	112.32	75.64	50.68	32.62	19.37	13.02	8.64	4.85	3.54	1.82	1.07	3.81	2.56	4.80	1,000—
64	60.51	.29	3.37	13.98	47.75	95.95	130.78	150.71	148.31	128.97	94.91	66.63	42.84	26.36	17.13	11.40	6.97	4.90	3.26	1.94	1.07	3.81	2.56	1,000—
65	94.31	.07	1.84	7.08	27.66	66.06	107.13	138.21	148.41	143.90	114.67	83.21	58.00	35.57	23.23	15.96	9.97	6.84	3.87	2.66	1.69	1.07	3.81	1,000—
66	126.62	.01	1.00	3.89	14.88	42.89	79.75	117.69	141.76	148.07	129.48	104.45	75.09	49.32	31.41	21.53	13.33	8.50	5.46	3.88	2.20	1.69	1.07	1,000—
67	147.21	.02	.59	1.83	7.73	23.89	58.31	92.46	125.48	145.19	139.10	121.45	92.91	64.45	44.88	31.33	18.33	11.99	7.55	5.26	3.46	2.20	1.69	1,000—
68	149.68	.01	.33	.90	4.00	13.15	34.95	68.83	104.62	130.75	140.23	131.24	111.36	82.93	59.61	42.63	24.90	16.32	10.85	7.36	5.19	3.39	2.20	1,000—
69	177.2560	.76	2.26	6.85	20.02	45.51	80.60	115.00	135.93	137.17	126.26	97.01	78.34	53.92	34.93	22.34	15.14	9.26	6.08	3.83	2.20	1,000—
70	96.3852	.63	1.31	3.51	10.14	26.49	55.48	98.12	130.03	133.99	136.59	116.05	93.60	71.49	47.90	31.75	21.08	13.56	9.51	6.02	3.83	1,000—
71	36.1257	.73	1.31	1.91	5.29	14.22	36.72	66.08	101.53	118.62	136.53	125.92	110.14	85.69	64.33	44.60	28.43	17.60	13.58	9.51	6.02	1,000—
72	17.5039	.70	1.12	.79	2.30	8.36	19.81	43.76	78.10	104.82	126.73	126.92	120.35	106.07	81.62	56.59	40.01	24.81	16.58	13.58	9.51	1,000—
73	3.0247	.78	1.41	.78	2.50	6.11	13.77	27.23	53.04	78.70	104.82	126.73	126.92	120.35	106.07	81.62	56.59	40.01	24.81	16.58	13.58	1,000—
74	1.1515	.30	1.01	2.29	3.05	4.22	8.02	12.60	27.02	61.35	80.31	120.61	116.03	125.18	106.40	83.40	69.47	49.62	29.77	25.04	18.15	1,000—
75	1.23	2.80	3.73	3.73	3.73	10.27	14.94	25.21	27.08	61.35	80.31	120.61	116.03	125.18	106.40	83.40	69.47	49.62	29.77	25.04	18.15	1,000—
76	5.56	8.33	22.22	3.98	3.98	13.89	62.78	41.67	77.75	73.76	104.58	101.71	124.18	117.92	89.63	70.36	45.75	29.88	33.33	38.99	1,000—
77	7.72	11.58	7.72	3.98	3.98	38.61	69.49	81.08	104.25	81.08	61.78	69.50	77.22	69.50	77.75	65.64	38.61	30.80	34.75	54.05	1,000—
78	3.38	16.89	40.54	43.92	43.92	54.05	87.84	94.59	77.70	70.95	54.05	70.95	70.95	60.81	30.40	33.78	40.54	27.03	33.78	81.08	1,000—
Total21	2.71	8.56	24.63	47.79	73.20	97.36	115.24	123.08	115.85	101.40	83.32	61.53	45.82	33.47	21.83	14.54	9.66	6.30	4.50	3.42	5.40	1,000+

SECTION C: RATIO PER 1,000 OF THE SEPARATE HEIGHTS SHOWN FOR EACH WEIGHT.

Proportion weight each inch per 1,000.	Height, in inches.																								Total.
	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	Total.		
85-90	0.21	76.00	103.26	173.91	195.65	206.53	51.53	117.15	43.45	5.43	10.30	4.43	21.22	11.88	7.64	2.55	1.27	1.27	0.55	0.69	0.13	0.00	1,000-		
100-104	2.71	29.29	45.84	101.87	131.14	169.87	175.92	154.00	104.64	54.33	32.68	13.45	8.74	4.57	3.09	1.79	1.67	1.27	1.27	0.55	0.69	0.13	0.00	1,000+	
105-109	8.56	15.40	37.96	82.85	97.30	148.40	175.92	182.86	142.23	68.85	46.99	20.67	8.84	2.10	1.92	1.79	1.67	1.27	1.27	0.55	0.69	0.13	0.00	1,000+	
110-114	24.63	17.62	46.22	97.30	111.90	185.36	211.90	200.36	175.02	132.11	74.81	35.01	13.81	4.60	2.45	2.55	2.55	1.27	1.27	0.55	0.69	0.13	0.00	1,000+	
115-119	47.79	35.51	78.22	140.16	178.22	225.36	268.50	260.40	210.40	160.40	115.15	60.78	26.37	8.67	2.61	3.01	3.01	1.27	1.27	0.55	0.69	0.13	0.00	1,000+	
120-124	73.20	53.43	110.94	185.65	238.43	295.65	345.69	345.69	285.04	217.10	160.56	115.18	60.08	25.32	8.51	6.22	6.22	1.27	1.27	0.55	0.69	0.13	0.00	1,000+	
125-129	97.56	74.74	150.58	245.69	318.04	388.26	438.04	438.04	358.04	270.75	200.75	150.75	90.08	35.36	11.51	6.22	6.22	1.27	1.27	0.55	0.69	0.13	0.00	1,000+	
130-134	113.24	85.25	180.64	295.65	373.76	455.22	505.22	505.22	405.22	305.22	225.22	165.22	115.22	45.32	11.51	6.22	6.22	1.27	1.27	0.55	0.69	0.13	0.00	1,000+	
135-139	123.08	95.26	200.64	325.65	413.76	505.22	555.22	555.22	455.22	355.22	275.22	205.22	145.22	45.32	11.51	6.22	6.22	1.27	1.27	0.55	0.69	0.13	0.00	1,000+	
140-144	101.40	77.24	164.64	275.65	363.76	455.22	505.22	505.22	405.22	305.22	225.22	165.22	115.22	45.32	11.51	6.22	6.22	1.27	1.27	0.55	0.69	0.13	0.00	1,000+	
145-149	83.72	63.28	140.64	235.65	323.76	415.22	465.22	465.22	365.22	265.22	185.22	115.22	65.22	11.51	6.22	6.22	6.22	1.27	1.27	0.55	0.69	0.13	0.00	1,000+	
150-154	61.53	45.82	107.24	187.65	275.76	367.22	417.22	417.22	317.22	217.22	137.22	67.22	11.51	6.22	6.22	6.22	6.22	1.27	1.27	0.55	0.69	0.13	0.00	1,000+	
155-159	45.82	33.47	82.24	142.65	230.76	322.22	372.22	372.22	272.22	172.22	92.22	22.22	6.22	6.22	6.22	6.22	6.22	1.27	1.27	0.55	0.69	0.13	0.00	1,000+	
160-164	33.47	23.24	57.24	97.65	152.76	222.22	272.22	272.22	172.22	72.22	12.22	2.22	6.22	6.22	6.22	6.22	6.22	1.27	1.27	0.55	0.69	0.13	0.00	1,000+	
165-169	21.83	15.35	39.35	69.76	114.87	169.98	219.98	219.98	119.98	19.98	1.98	1.98	6.22	6.22	6.22	6.22	6.22	1.27	1.27	0.55	0.69	0.13	0.00	1,000+	
170-174	14.54	10.35	27.35	47.76	82.87	127.98	177.98	177.98	77.98	17.98	1.98	1.98	6.22	6.22	6.22	6.22	6.22	1.27	1.27	0.55	0.69	0.13	0.00	1,000+	
175-179	9.66	6.83	17.83	32.84	57.85	92.96	132.96	132.96	32.96	7.96	1.96	1.96	6.22	6.22	6.22	6.22	6.22	1.27	1.27	0.55	0.69	0.13	0.00	1,000+	
180-184	6.80	4.93	12.93	22.94	42.95	72.96	112.96	112.96	12.96	2.96	1.96	1.96	6.22	6.22	6.22	6.22	6.22	1.27	1.27	0.55	0.69	0.13	0.00	1,000+	
185-189	4.00	2.82	7.82	14.83	26.84	46.85	76.86	76.86	16.86	3.86	1.86	1.86	6.22	6.22	6.22	6.22	6.22	1.27	1.27	0.55	0.69	0.13	0.00	1,000+	
190-194	3.42	2.49	6.49	11.50	20.51	35.52	55.53	55.53	10.53	2.53	1.53	1.53	6.22	6.22	6.22	6.22	6.22	1.27	1.27	0.55	0.69	0.13	0.00	1,000+	
195-199	5.40	3.90	9.90	17.91	31.92	51.93	71.94	71.94	16.94	3.94	1.94	1.94	6.22	6.22	6.22	6.22	6.22	1.27	1.27	0.55	0.69	0.13	0.00	1,000+	

* The maximum or minimum value of the + or - did not exceed 0.03.

TABLE II.—Correlation between height and chest circumference (expiration), first million draft recruits.

SECTION A: ABSOLUTE NUMBERS DERIVED FROM SUMMATION OF SECTIONS.

Height, in inches.	Total.	Chest, in inches.										
		29	30	31	32	33	34	35	36	37	38	39
59.....	3,086	128	241	366	622	586	480	331	172	76	52	32
60.....	2,921	104	331	459	595	530	402	221	124	50	22	23
61.....	7,572	444	910	1,474	1,545	1,336	912	525	246	97	46	37
62.....	15,646	850	1,822	2,884	3,503	2,810	1,941	1,105	541	229	87	76
63.....	31,207	1,407	3,316	5,500	6,678	5,902	4,024	2,328	1,113	522	226	191
64.....	52,923	2,153	4,940	8,920	11,189	8,990	7,419	4,314	2,163	1,030	404	341
65.....	82,426	2,759	6,757	12,514	17,125	16,150	12,547	7,618	3,895	1,806	802	653
66.....	110,816	2,884	7,708	15,679	22,330	22,824	17,708	11,030	5,708	2,616	1,231	1,038
67.....	128,291	2,583	7,549	16,940	24,865	26,558	22,018	15,945	8,946	3,613	1,611	1,348
68.....	130,624	2,053	6,289	14,386	24,163	27,306	23,628	15,945	8,946	4,182	1,991	1,636
69.....	111,123	1,739	4,290	10,779	18,997	23,133	21,563	14,860	8,732	4,051	1,927	1,378
70.....	85,880	362	3,350	6,921	13,261	17,291	16,741	12,250	7,391	3,567	1,666	1,062
71.....	54,609	155	2,611	5,896	10,946	10,833	11,518	8,642	5,306	2,654	1,235	747
72.....	31,323	70	201	3,877	7,731	6,054	6,614	5,252	3,511	1,784	862	480
73.....	15,254	21	79	1,709	4,046	2,798	3,203	2,553	1,905	985	468	210
74.....	6,411	12	29	583	1,645	1,070	1,395	1,176	872	498	237	90
75.....	2,620	2	14	83	223	163	217	516	409	227	124	35
76.....	1,060	2	4	44	103	61	68	209	139	88	38	16
77.....	361	6	1	14	29	39	56	64	61	32	10	5
78.....	256	6	1	8	21	39	56	47	37	27	9	5
79.....	298	6	5	15	36	49	86	43	42	21	11	14
Total.....	873,159	18,083	49,090	103,294	159,379	175,858	152,663	103,414	59,015	28,175	13,151	11,027

Number of cases: 873,159. Height: Mean, 67.49 inches; standard deviation, 2.72±0.0014 inch. Chest circumference (expiration): Mean, 33.22 inches; standard deviation, 2.01±0.0010 inch. Correlation: 0.294±0.0007.

SECTION B: RATIO PER 1,000 OF THE SEPARATE CHEST MEASUREMENTS TO EACH HEIGHT.

Height, in inches.	Chest, in inches.											Total ^a
	29	30	31	32	33	34	35	36	37	38	39	
Proportion each chest per 1,000.												
50	41.48	78.09	118.60	201.56	180.80	155.53	107.26	55.74	24.63	16.65	10.37	1,000+
51	56.15	113.32	157.12	208.70	181.44	157.62	75.70	42.45	17.12	7.53	7.87	1,000+
52	58.64	120.18	164.68	204.05	176.44	150.45	69.34	32.49	12.81	6.08	4.89	1,000+
53	53.64	114.95	151.98	221.04	177.31	122.47	69.73	34.14	14.45	5.49	4.90	1,000+
54	45.09	103.26	136.24	214.00	189.12	128.94	74.63	33.57	16.73	7.24	6.44	1,000+
55	40.68	93.34	126.55	211.43	188.77	140.19	81.52	35.87	19.46	8.77	8.44	1,000+
56	33.47	81.98	108.55	207.77	185.94	149.80	92.42	47.25	21.91	9.73	9.37	1,000+
57	26.03	69.56	131.81	201.51	183.97	159.80	99.54	52.05	23.01	11.11	10.51	1,000+
58	20.13	56.15	127.43	183.99	207.00	171.61	122.06	67.72	32.01	12.52	12.52	1,000+
59	15.72	48.14	111.66	170.86	208.19	192.51	134.00	78.38	36.45	15.24	15.24	1,000+
60	11.55	38.61	97.00	138.10	206.14	193.60	146.32	98.11	42.02	20.24	16.43	1,000+
61	8.81	24.72	71.35	141.57	198.39	210.94	168.25	97.16	48.60	22.63	19.81	1,000+
62	6.63	19.88	56.57	128.34	192.05	209.81	168.60	111.37	56.91	27.35	23.70	1,000+
63	4.92	13.15	46.39	108.09	183.07	203.05	163.44	124.64	65.10	30.62	28.13	1,000+
64	4.58	12.32	33.85	100.61	165.91	216.20	183.44	156.10	77.68	33.97	32.76	1,000+
65	3.28	11.07	31.68	85.12	143.13	203.05	168.94	147.21	86.64	47.83	44.35	1,000+
66	4.58	12.96	40.74	95.37	150.92	200.91	183.51	177.29	88.64	53.18	41.66	1,000+
67	5.54	11.08	38.78	80.34	163.98	198.36	177.29	168.98	103.64	57.70	44.32	1,000+
68	23.44	3.91	31.25	152.34	152.34	152.34	152.34	144.83	103.64	57.70	44.32	1,000+
69	20.13	16.78	50.34	120.81	164.41	187.91	144.29	140.94	70.47	38.92	46.98	1,000+
Total	20.74	36.22	118.30	182.64	201.42	174.83	118.43	67.60	32.27	15.06	12.63	1,000+

SECTION C: RATIO PER 1,000 OF THE SEPARATE HEIGHTS TO EACH CHEST MEASUREMENT.

Chest, in inches.	Height, in inches.											Total ^a
	59	60	61	62	63	64	65	66	67	68	69	
Proportion each chest per 1,000.												
50	20.74	7.07	9.06	24.54	45.98	77.77	119.00	132.50	136.40	142.76	113.45	1,000+
51	56.22	4.91	6.74	18.54	37.12	67.55	100.63	137.64	157.02	164.60	128.12	1,000+
52	118.30	3.54	4.44	14.27	27.92	53.24	86.36	131.79	155.26	164.26	141.22	1,000+
53	182.54	3.90	4.73	9.66	21.96	41.90	70.20	107.45	140.10	156.14	151.00	1,000+
54	201.42	3.33	3.63	7.60	15.96	33.56	56.81	91.82	129.78	151.01	155.27	1,000+
55	174.65	3.14	3.01	5.97	12.71	26.36	48.60	80.87	115.96	144.21	154.75	1,000+
56	118.42	3.20	2.14	5.08	10.66	22.51	41.71	73.66	105.65	135.32	154.17	1,000+
57	67.60	2.91	2.10	4.17	9.17	18.86	36.55	66.00	97.74	136.84	149.90	1,000+
58	32.27	2.70	1.77	3.44	8.13	18.53	36.56	64.10	92.85	128.23	148.42	1,000+
59	15.06	2.95	1.67	3.50	6.62	17.18	35.28	60.99	93.53	122.51	151.40	1,000+
60	12.63	2.90	2.09	3.36	6.90	17.32	30.93	50.22	94.14	122.25	148.26	1,000+
Total	3.53	3.35	8.67	18.15	35.74	60.61	126.92	146.92	149.60	127.38	96.07	1,000+

^a The maximum or minimum value of the + or - did not exceed 0.03.

TABLE I.—Correlation between height and weight for first million draft recruits.
SECTION A: ABSOLUTE NUMBERS DERIVED FROM SUMMATION OF SECTION.

Height in inches.	Total.	Weight, in pounds.																					
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204
58 and under 59.....	3,124	14	69	111	201	259	252	333	325	310	296	243	209	153	93	124	55	17	15	10	8	15	12
60.....	2,857	19	108	245	367	395	350	270	255	186	165	135	106	92	63	59	24	6	13	4	11	14	7
61.....	7,477	32	240	616	1,138	1,209	1,123	977	643	440	331	210	196	100	75	52	31	23	14	8	22	36	4
62.....	15,644	36	351	975	2,081	2,557	2,553	2,221	1,674	1,131	714	456	285	194	139	91	53	38	26	9	18	32	10
63.....	30,935	38	396	1,263	3,174	4,644	4,972	4,740	3,846	2,777	1,811	1,211	777	418	308	214	136	78	47	31	30	42	34
64.....	52,547	15	358	1,308	3,812	6,413	7,969	8,104	7,365	5,902	3,974	2,663	3,509	2,159	1,403	934	255	186	121	83	46	56	37
65.....	81,904	17	276	1,145	3,911	7,859	10,711	12,344	12,147	10,563	7,773	5,459	6,378	3,911	2,554	1,744	1,066	719	426	292	203	167	186
66.....	109,964	8	202	778	3,042	7,264	11,780	15,198	16,431	16,824	12,610	9,151	9,600	6,305	4,015	2,753	1,689	1,086	698	496	356	281	318
67.....	127,844	1	128	497	1,902	5,483	10,196	15,071	18,123	18,930	16,553	13,363	12,077	8,376	5,533	3,946	2,383	1,558	981	694	450	352	520
68.....	129,987	3	77	251	1,005	3,105	7,320	12,018	16,310	18,872	18,081	15,786	12,305	9,162	6,587	4,713	2,751	1,803	1,199	813	574	375	681
69.....	110,508	1	36	100	442	1,453	3,863	7,607	11,594	14,449	15,498	14,503	12,568	9,354	6,308	4,513	2,924	1,870	1,267	775	584	404	663
70.....	83,702	50	65	130	445	1,571	3,676	7,899	11,746	16,626	11,378	11,452	10,568	7,354	5,068	3,688	2,508	1,726	1,146	737	517	360	881
71.....	64,357	28	34	45	191	591	1,551	3,440	6,016	9,626	6,524	7,283	7,354	5,068	3,688	2,508	1,726	1,146	737	517	360	881	
72.....	31,370	18	23	23	41	60	166	446	3,016	4,944	6,524	7,283	7,354	5,068	3,688	2,508	1,726	1,146	737	517	360	881	
73.....	15,198	6	12	12	17	12	35	127	1,152	2,073	3,185	3,721	4,285	3,950	3,455	2,612	2,218	1,399	892	552	426	285	535
74.....	6,391	3	5	5	9	5	16	27	88	174	339	503	747	754	1,829	1,612	1,239	860	608	377	252	210	407
75.....	2,620	3	4	4	5	6	8	16	21	33	97	162	234	316	304	739	639	480	338	212	160	116	228
76.....	1,071	3	2	2	3	4	4	8	11	16	27	29	79	112	109	133	137	96	85	49	32	26	60
77.....	380	2	2	2	2	3	3	3	5	19	15	28	23	25	22	20	40	28	33	14	12	14	25
78.....	259	2	2	2	2	2	2	1	10	18	21	27	21	16	18	20	18	22	17	10	8	9	14
79.....	296	1	1	1	2	5	12	13	16	26	28	23	21	16	21	21	18	9	10	12	8	10	24
Total.....	898,445	184	2,356	7,635	21,388	41,503	63,567	84,726	100,064	106,889	100,607	88,087	72,362	53,431	39,797	29,083	18,984	12,629	8,385	5,467	3,977	2,906	4,688

Number of cases: 898,445. Height: Mean, 67.49 inches; standard deviation, 2.71±0.0014 inch. Weight: Mean, 141.54 pounds; standard deviation, 17.42±0.0089 pound. Correlation: 0.4810±0.0006.

TABLE I.—*Correlation between height and weight for first million draft recruits—Continued.*
SECTION B: RATIO PER 1,000 OF THE SEPARATE WEIGHTS SHOWN FOR EACH HEIGHT.

Height, in inches.	Proportion each height per 1,000.	Weight, in pounds.																							
		93-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204	Total. ^a	
59	3.63	4.48	22.09	35.53	64.34	82.91	80.67	105.59	104.03	96.23	94.75	77.78	66.90	48.98	29.77	39.60	17.60	5.44	4.80	3.20	2.56	4.80	3.84	1,000+	
60	3.32	6.58	37.41	84.87	127.12	136.82	121.23	93.53	88.03	64.43	57.16	46.77	36.72	31.87	21.82	18.01	8.31	2.08	4.50	1.39	3.20	3.81	4.80	2.42	1,000+
61	8.01	4.28	32.10	82.38	152.20	161.70	150.19	123.98	88.00	68.85	44.27	28.09	26.21	13.47	10.03	7.80	4.15	3.08	4.81	1.07	2.94	2.05	1.64	1,000+	
62	18.01	2.30	22.44	62.33	133.02	163.45	163.19	141.97	107.01	72.30	45.64	29.15	18.23	12.40	8.89	5.82	3.39	2.43	1.66	1.58	1.07	2.05	1.64	1,000+	
63	35.62	1.23	12.80	40.83	102.60	150.12	160.72	153.22	124.32	96.77	58.52	39.15	23.50	13.61	9.89	6.92	4.40	2.52	1.36	1.36	1.00	1.07	1.10	1,000+	
64	60.51	2.29	6.81	24.90	72.56	122.04	151.65	150.71	148.31	112.32	75.64	50.68	32.62	19.37	13.02	8.64	6.85	3.54	1.52	1.94	1.77	1.37	1.70	1,000+	
65	94.31	.29	3.37	13.98	47.75	95.95	130.78	130.71	148.41	128.97	94.91	66.63	42.84	26.36	17.13	11.40	9.97	6.54	3.26	2.66	1.94	1.37	1.04	1,000+	
66	126.62	.07	1.84	7.06	27.66	66.09	107.13	138.21	149.41	143.90	114.67	83.21	58.00	35.57	23.23	15.86	9.97	6.50	3.87	3.88	2.66	1.52	1.69	1,000+	
67	147.21	.01	1.00	3.89	14.88	42.89	79.75	117.89	141.76	145.07	120.48	104.63	75.09	49.32	31.41	20.33	13.21	8.50	5.46	5.26	2.78	2.20	2.49	1,000+	
68	149.64	.02	.59	1.83	7.73	23.89	56.31	92.46	125.48	140.19	130.10	121.45	92.91	64.45	44.88	30.61	24.00	10.85	7.55	7.36	5.19	3.39	6.16	1,000+	
69	96.38	.01	.60	.76	2.26	6.85	20.02	45.51	80.60	115.00	135.93	133.17	126.26	97.01	76.34	53.92	47.90	31.75	15.14	9.26	6.96	4.83	8.28	1,000+	
70	62.59	.01	.52	.63	.83	3.51	10.14	26.49	55.48	98.12	130.03	133.98	135.29	116.05	93.60	71.49	64.33	22.34	15.14	13.56	9.51	6.62	12.53	1,000+	
71	36.12	.01	.57	.73	.83	3.51	10.14	26.49	55.48	98.12	130.03	133.98	135.29	116.05	93.60	71.49	64.33	22.34	15.14	13.56	9.51	6.62	12.53	1,000+	
72	17.50	.01	.39	.79	1.12	1.91	5.20	8.36	19.81	43.76	78.10	104.82	126.53	125.92	120.35	106.07	81.62	56.59	40.01	24.81	17.60	13.82	26.78	1,000+	
73	7.36	.01	.47	.78	1.41	2.79	2.60	4.22	13.77	27.23	53.04	78.70	116.88	117.98	126.58	115.63	99.96	75.11	52.89	33.17	25.04	18.15	35.68	1,000+	
74	3.02	.01	1.15	1.53	3.73	3.05	3.05	6.11	8.02	12.60	37.02	61.83	89.31	120.61	116.03	125.19	106.49	85.60	66.47	49.62	29.77	24.05	48.47	1,000+	
75	1.23	.01	2.80	1.53	3.73	3.05	3.05	6.11	8.02	12.60	37.02	61.83	89.31	120.61	116.03	125.19	106.49	85.60	66.47	49.62	29.77	24.05	48.47	1,000+	
76	.41	.01	5.56	7.72	2.78	8.33	22.22	3.98	13.89	52.78	41.67	77.78	63.99	104.25	61.11	111.11	127.92	89.63	79.36	45.75	38.80	33.33	56.02	1,000+	
77	.30	.01	7.72	3.38	6.76	11.58	7.72	3.98	38.61	66.49	81.08	104.25	81.08	61.78	99.50	77.22	69.50	77.78	91.67	38.61	34.75	34.75	54.05	1,000+	
78	.34	.01	3.38	7.72	6.76	16.89	40.54	43.92	54.05	87.84	94.59	77.70	70.95	54.05	70.95	70.95	60.81	30.40	33.78	40.54	27.03	33.78	81.08	1,000+	
Total.	.21	2.71	8.56	24.63	47.79	73.20	97.36	115.24	123.08	115.85	101.40	83.32	61.53	43.82	33.47	21.83	14.54	9.66	6.30	4.50	3.42	5.40	3.42	5.40	

SECTION C: RATIO PER 1,000 OF THE SEPARATE HEIGHTS SHOWN FOR EACH WEIGHT.

Weight in pounds.		Proportion each weight per 1,000.		Height, in inches.																	Total.						
				59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75		76	77	78	79	80	
95-99	0.21	76.09	103.26	173.91	195.65	206.52	81.52	92.39	43.48	5.43	16.30	5.43	21.22	11.88	7.64	2.55	1.27	1.27	1.27	0.85	0.85	0.27	0.13	0.13	1,000-		
100-104	2.71	29.29	45.64	101.87	148.98	168.08	151.95	117.15	85.74	54.33	32.68	15.28	8.74	4.57	3.09	1.61	1.67	1.27	1.27	1.27	0.85	0.85	0.13	0.13	1,000-		
105-109	8.56	14.93	32.96	82.85	131.14	168.87	175.92	154.00	104.64	66.85	33.76	13.45	8.54	4.57	3.09	1.61	1.67	1.27	1.27	1.27	0.85	0.85	0.13	0.13	1,000+		
110-114	24.63	9.40	17.16	53.21	97.30	148.40	178.23	182.88	142.23	88.93	46.99	20.67	13.81	8.67	5.45	2.9	2.25	1.4	1.4	1.4	0.7	0.7	0.12	0.12	1,000+		
115-119	47.79	6.24	9.52	29.13	61.61	111.90	145.40	158.36	175.02	132.11	74.81	35.01	23.37	13.81	8.67	5.45	2.9	2.25	1.4	1.4	1.4	0.7	0.7	0.12	0.12	1,000+	
120-124	73.20	3.96	5.51	17.67	40.16	78.22	125.36	168.50	185.32	160.40	115.15	60.77	35.01	23.37	13.81	8.67	5.45	2.9	2.25	1.4	1.4	1.4	0.7	0.7	0.12	0.12	1,000+
125-129	97.56	3.93	5.19	10.94	26.21	53.95	95.65	145.69	179.39	177.88	141.55	89.78	47.40	30.13	11.51	3.01	1.88	1.19	1.19	1.19	0.65	0.65	0.10	0.10	1,000+		
130-134	115.24	3.25	2.55	6.42	16.73	38.43	73.69	121.37	164.17	181.08	162.96	115.84	67.40	30.13	11.51	3.01	1.88	1.19	1.19	1.19	0.65	0.65	0.10	0.10	1,000+		
135-139	123.08	2.90	1.74	4.12	10.58	25.98	55.22	98.82	145.04	177.10	176.56	135.18	90.06	45.32	19.39	6.22	3.77	1.96	1.96	1.96	1.18	1.18	0.21	0.21	1,000+		
140-144	115.85	2.76	1.53	3.29	7.10	18.00	39.50	77.26	125.34	164.53	176.72	154.03	113.09	64.56	42.26	11.80	5.71	3.37	3.37	3.37	1.5	1.5	0.21	0.21	1,000+		
145-149	101.40	2.76	1.46	2.71	3.63	10.05	22.69	48.49	88.14	132.67	166.90	170.06	146.04	101.63	59.22	18.09	10.32	1.84	1.84	1.84	1.09	1.09	0.31	0.31	1,000-		
150-154	83.52	2.86	1.72	1.87	3.49	7.82	19.05	40.41	73.20	118.00	156.76	171.47	151.97	118.06	73.93	38.05	14.11	5.91	5.91	5.91	2.10	2.10	0.30	0.30	1,000-		
155-159	61.53	2.34	1.58	1.88	3.13	7.36	17.19	35.25	64.18	100.89	146.57	165.51	150.56	127.85	88.82	45.96	20.33	7.64	7.64	7.64	4.58	4.58	0.45	0.45	1,000-		
160-164	45.82	2.34	1.58	1.88	3.13	7.36	17.19	35.25	64.18	100.89	146.57	165.51	150.56	127.85	88.82	45.96	20.33	7.64	7.64	7.64	4.58	4.58	0.45	0.45	1,000-		
165-169	33.47	2.27	1.79	2.03	3.13	7.36	17.19	35.25	64.18	100.89	146.57	165.51	150.56	127.85	88.82	45.96	20.33	7.64	7.64	7.64	4.58	4.58	0.45	0.45	1,000-		
170-174	21.83	2.90	1.27	1.64	2.80	7.18	13.45	30.13	57.82	89.11	123.37	145.14	154.27	137.07	108.47	68.10	38.01	17.72	17.72	17.72	2.22	2.22	1.74	1.74	1,000+		
175-179	14.54	1.35	1.48	1.82	3.01	6.18	14.43	31.84	50.81	85.94	116.99	142.99	151.10	136.67	108.38	72.51	40.31	21.71	21.71	21.71	1.04	1.04	0.34	0.34	1,000+		
180-184	9.66	1.79	1.55	1.67	3.01	5.61	14.43	31.84	50.81	85.94	116.99	142.99	151.10	136.67	108.38	72.51	40.31	21.71	21.71	21.71	1.04	1.04	0.34	0.34	1,000+		
185-189	6.30	1.83	1.73	1.46	1.65	5.67	17.01	28.08	53.41	90.73	125.11	148.71	141.76	134.81	108.97	68.96	38.78	23.78	23.78	23.78	0.96	0.96	1.83	1.83	1,000-		
190-194	4.50	2.05	2.82	5.63	4.61	7.68	11.77	28.08	53.41	90.73	125.11	148.71	141.76	134.81	108.97	68.96	38.78	23.78	23.78	23.78	0.96	0.96	1.83	1.83	1,000+		
195-199	3.42	5.06	4.72	12.14	10.79	14.16	18.88	33.38	56.30	94.74	118.68	136.21	121.38	96.09	70.80	39.11	21.24	19.96	19.96	19.96	3.77	3.77	2.05	2.05	1,000+		
200-204	5.40	2.56	1.49	1.85	2.13	7.25	7.89	18.13	39.68	67.53	110.92	145.26	147.82	145.26	114.12	86.82	48.63	27.09	12.80	12.80	5.33	5.33	2.99	2.99	1,000-		
Total.	3.80	3.32	8.61	18.01	35.62	60.51	94.31	126.02	147.21	149.68	177.25	96.39	62.59	36.12	17.50	7.30	3.02	1.23	0.41	0.30	0.34	0.34	0.34	0.34	1,000-	

a The maximum or minimum value of the + or - did not exceed 0.03.

TABLE II.—Correlation between height and chest circumference (expiration), first million draft recruits.

SECTION A: ABSOLUTE NUMBERS DERIVED FROM SUMMATION OF SECTIONS.

Height, in inches.	Total.	Chest, in inches.										
		29	30	31	32	33	34	35	36	37	38	39
59	3,086	128	241	366	622	586	480	331	172	76	52	32
60	2,971	164	331	459	595	530	402	221	124	50	22	23
61	7,572	444	910	1,474	1,645	1,336	912	525	246	97	46	37
62	15,648	850	1,822	2,884	3,503	2,810	1,941	1,105	541	229	87	76
63	31,207	1,407	3,316	5,590	6,678	5,002	4,024	2,328	1,113	522	226	191
64	52,923	2,153	4,940	8,920	11,189	9,990	7,419	4,314	2,163	1,030	464	341
65	82,426	2,759	6,757	12,514	17,125	16,150	12,347	7,618	3,895	1,806	802	653
66	110,816	2,884	7,708	15,679	22,330	22,824	17,708	11,030	5,768	2,616	1,231	1,038
67	128,291	2,583	7,599	16,349	24,885	26,558	22,018	14,015	7,722	3,613	1,611	1,348
68	130,624	2,053	6,298	14,596	24,163	27,306	23,628	15,945	8,446	4,182	1,901	1,635
69	111,123	1,739	5,200	10,779	18,997	23,133	21,383	14,890	8,732	4,051	1,927	1,636
70	83,880	1,295	4,200	8,921	13,261	17,291	16,741	12,200	7,391	3,567	1,696	1,378
71	54,609	362	2,603	5,895	7,731	10,833	11,518	8,642	5,306	2,654	1,235	1,062
72	31,523	155	1,350	3,895	4,046	6,054	6,203	5,252	3,511	1,794	862	747
73	15,294	70	709	1,877	1,645	2,798	3,203	2,853	1,872	965	468	430
74	6,411	21	201	217	1,645	1,070	1,396	1,176	872	468	237	210
75	2,620	12	26	83	223	103	163	159	409	227	124	90
76	44	14	44	103	103	61	217	209	159	88	38	45
77	1,361	2	4	14	29	39	68	64	61	32	10	16
78	256	6	1	8	21	21	56	47	37	27	9	5
79	298	6	5	15	36	49	56	43	42	21	11	14
Total	873,159	18,063	49,090	103,294	159,379	175,858	152,663	103,414	59,015	28,175	13,151	11,027

Number of cases: 873,159. Height: Mean, 67.49 inches; standard deviation, 2.72±0.0014 inch. Chest circumference (expiration): Mean, 33.22 inches; standard deviation, 2.01±0.0010 inch. Correlation: 0.2304±0.0007.

SECTION B: RATIO PER 1,000 OF THE SEPARATE CHEST MEASUREMENTS TO EACH HEIGHT.

Height, in inches.	Proportion each chest per 1,000.	Chest, in inches.										Total. ^a
		29	30	31	32	33	34	35	36	37	3	39
59.....	3.53	41.46	78.09	118.60	201.56	189.89	155.53	107.26	55.74	24.63	10.85	10.37
60.....	3.35	50.15	113.32	137.12	203.70	181.44	137.02	75.70	42.45	17.12	7.53	7.87
61.....	3.67	58.64	120.18	194.68	204.05	176.44	122.47	69.34	32.49	12.81	6.08	4.89
62.....	18.15	53.04	114.96	181.98	221.04	177.31	120.45	66.73	34.14	14.45	5.49	4.80
63.....	33.74	43.09	106.26	176.24	214.00	188.12	128.94	81.52	35.67	16.73	7.24	6.12
64.....	60.61	40.08	93.34	168.55	211.43	183.77	126.19	74.32	40.87	19.46	8.77	6.44
65.....	94.40	33.47	81.98	151.81	207.77	185.94	149.80	92.42	47.25	21.91	9.73	7.92
66.....	20.13	20.03	66.95	141.49	201.51	203.97	159.80	96.54	52.05	23.61	11.11	9.37
67.....	146.92	20.13	58.15	127.43	193.99	209.04	180.88	104.25	67.19	28.16	12.56	10.51
68.....	146.90	15.72	48.14	111.66	183.00	208.19	192.51	122.06	67.72	32.01	15.24	12.52
69.....	127.27	1.65	38.61	97.00	170.96	200.14	196.00	134.00	78.98	36.45	20.24	16.42
70.....	98.97	8.81	31.03	82.31	158.10	200.14	196.00	134.00	78.98	36.45	20.24	16.42
71.....	58.94	6.63	24.72	56.37	128.34	198.39	200.14	134.00	78.98	36.45	20.24	16.42
72.....	37.50	4.82	19.35	46.39	108.09	183.07	200.14	134.00	78.98	36.45	20.24	16.42
73.....	3.34	3.34	12.32	31.88	108.09	183.07	200.14	134.00	78.98	36.45	20.24	16.42
74.....	3.34	4.58	12.32	31.88	108.09	183.07	200.14	134.00	78.98	36.45	20.24	16.42
75.....	1.94	1.94	12.32	31.88	108.09	183.07	200.14	134.00	78.98	36.45	20.24	16.42
76.....	1.41	1.41	12.32	31.88	108.09	183.07	200.14	134.00	78.98	36.45	20.24	16.42
77.....	1.41	1.41	12.32	31.88	108.09	183.07	200.14	134.00	78.98	36.45	20.24	16.42
78.....	1.41	1.41	12.32	31.88	108.09	183.07	200.14	134.00	78.98	36.45	20.24	16.42
79.....	1.41	1.41	12.32	31.88	108.09	183.07	200.14	134.00	78.98	36.45	20.24	16.42
Total.....	3.34	20.13	58.22	118.30	182.54	201.42	174.85	118.42	67.60	32.27	15.06	12.63

SECTION C: RATIO PER 1,000 OF THE SEPARATE HEIGHTS TO EACH CHEST MEASUREMENT.

Chest, in Inches.	Proportion each chest per 1,000.	Height, in inches.														Total. ^a							
		59	60	61	62	63	64	65	66	67	68	69	70	71	72		73	74	75	76	77	78	79
29.....	20.74	7.07	9.06	24.54	46.98	77.77	119.00	152.50	159.40	142.76	113.48	71.58	40.85	20.01	8.57	3.87	1.16	0.66	0.11	0.33	0.33	1,000+
30.....	56.22	4.91	6.74	18.54	37.12	67.55	100.63	137.64	157.02	154.60	128.12	87.39	53.03	27.50	12.45	4.09	1.61	0.59	0.26	0.08	0.33	1,000+
31.....	118.30	3.54	4.44	14.27	27.92	53.24	86.36	121.14	151.79	128.26	141.22	104.35	67.00	37.71	18.17	6.86	2.10	0.80	0.43	0.15	0.33	1,000+
32.....	182.54	3.90	3.73	9.69	21.96	41.90	70.20	107.45	140.10	156.14	151.60	119.19	83.20	48.50	25.38	10.36	4.05	1.40	0.65	0.18	0.33	1,000+
33.....	201.42	3.33	3.01	7.60	15.96	33.56	56.81	91.83	129.78	151.01	155.27	131.53	98.32	61.60	34.42	15.91	6.08	2.13	0.93	0.35	0.37	1,000+
34.....	174.85	3.14	2.63	5.97	12.71	26.36	48.60	80.87	115.98	144.21	154.75	140.13	109.66	75.45	43.32	20.98	9.08	3.48	1.42	0.45	0.37	1,000+
35.....	118.42	3.20	2.14	5.08	10.68	22.51	41.71	73.66	106.65	135.52	154.17	144.00	118.84	83.56	50.78	27.59	11.37	4.99	2.02	0.62	0.45	1,000+
36.....	67.60	2.91	2.10	4.17	8.13	18.86	36.56	66.00	97.74	130.84	149.90	147.96	125.25	89.91	56.49	32.28	14.78	6.93	2.99	1.03	0.63	1,000+
37.....	32.27	2.70	1.77	3.44	8.13	18.53	36.56	64.10	92.85	128.23	148.42	143.77	126.60	94.19	63.68	35.32	17.67	8.06	3.12	1.14	0.66	1,000+
38.....	15.06	2.95	1.67	3.50	6.62	17.18	35.26	60.99	93.53	122.51	151.40	146.53	126.12	93.91	63.55	35.59	18.02	9.43	2.89	1.76	0.84	1,000+
39.....	12.63	2.90	2.09	3.36	6.89	17.32	30.93	59.22	94.14	122.25	148.26	148.35	124.97	98.13	67.75	36.00	19.04	8.16	4.08	1.45	1.27	1,000+
Total.....	3.53	3.35	8.67	18.15	35.74	60.61	94.40	126.92	146.92	149.60	127.38	96.07	62.55	36.10	17.50	7.34	2.00	1.24	0.41	0.29	1,000+

^a The maximum or minimum value of the + or - did not exceed 0.03.

TABLE III.—*Correlation between weight and chest circumference (expiration), first million draft recruits.*
SECTION A: ABSOLUTE NUMBERS DERIVED FROM SUMMATION OF SECTIONS.

Chest, in inches.	Total.	Weight, in pounds.																					
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204
29	17,933	143	838	1,991	3,593	3,925	2,950	1,971	1,083	539	288	161	99	47	40	55	39	20	6	14	23	84	23
30	48,066	38	639	2,264	6,079	9,499	6,979	3,298	5,619	3,377	1,563	893	363	171	104	96	128	91	43	17	19	58	10
31	103,277	19	1,077	7,730	17,417	26,724	19,874	10,734	17,417	12,603	5,483	3,853	1,723	905	398	240	128	61	29	28	32	104	14
32	178,456	8	158	1,765	6,113	12,522	17,584	12,462	20,071	36,744	26,134	12,776	7,200	3,417	1,599	826	324	172	61	44	21	28	11
33	251,770	1	185	280	3,365	9,245	10,119	17,984	25,998	30,728	28,569	12,765	15,641	9,064	4,855	2,570	1,063	479	171	93	46	44	18
34	315,555	1	48	82	1,353	4,715	8,332	14,180	14,596	20,766	24,111	12,487	20,428	13,548	9,446	5,825	2,784	1,272	389	277	236	151	74
35	382,987	1	23	41	1,000	3,501	8,502	1,044	2,506	8,083	12,909	15,129	15,819	13,548	10,564	7,576	4,427	1,527	1,389	607	326	151	71
36	458,987	2	26	38	57	114	231	615	1,388	2,754	4,462	4,408	7,942	13,548	10,564	7,576	4,427	1,527	1,389	607	326	151	71
37	548,987	2	26	38	57	114	231	615	1,388	2,754	4,462	4,408	7,942	13,548	10,564	7,576	4,427	1,527	1,389	607	326	151	71
38	648,987	2	26	38	57	114	231	615	1,388	2,754	4,462	4,408	7,942	13,548	10,564	7,576	4,427	1,527	1,389	607	326	151	71
39	748,987	2	26	38	57	114	231	615	1,388	2,754	4,462	4,408	7,942	13,548	10,564	7,576	4,427	1,527	1,389	607	326	151	71
40	848,987	2	26	38	57	114	231	615	1,388	2,754	4,462	4,408	7,942	13,548	10,564	7,576	4,427	1,527	1,389	607	326	151	71
Total	872,419	213	2,313	7,391	21,382	41,665	63,866	85,072	100,715	107,129	101,040	88,316	72,618	53,688	39,998	29,141	19,052	12,692	8,310	5,596	3,853	2,967	5,432

Number of cases: 872,419. Weight: Mean, 141.59 pounds; standard deviation, 17.49 ± 0.0099 pound. Chest circumference (expiration): Mean, 33.23 inches; standard deviation, 2.03 ± 0.0010 inch. Correlation: 0.6607 ± 0.003 .

SECTION B: RATIO PER 1,000 OF THE SEPARATE WEIGHTS TO EACH CHEST MEASUREMENT.

Chest circumference, in inches.	Proportion each chest per 1,000.	Weight, in pounds.																						Total.
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204	
29	20.56	7.97	46.73	111.02	200.36	218.93	164.50	109.91	60.40	30.05	16.06	8.96	5.52	2.62	2.23	3.07	2.17	1.12	0.33	0.78	1.28	4.68	1.28	1,000+
30	56.23	7.77	13.03	46.77	123.92	193.43	203.42	160.16	114.54	68.84	32.47	14.15	7.40	3.49	2.12	1.96	1.18	.98	.27	.35	.39	1.18	.20	1,000+
31	118.38	1.05	3.94	16.75	58.19	121.28	173.06	190.84	168.64	122.03	72.45	37.31	16.68	7.79	3.85	2.32	1.24	.88	.46	.28	.22	.31	.14	1,000+
32	182.77	0.18	0.96	4.99	22.16	59.48	110.30	150.68	182.31	167.72	126.27	80.12	43.15	21.43	10.03	5.18	2.03	1.08	.38	.28	.13	.18	.06	1,000+
33	201.47	0.01	.48	1.59	7.77	24.15	57.57	102.31	147.33	173.09	162.53	129.51	88.98	51.57	27.62	14.62	6.05	2.73	.97	.26	.23	.06	.06	1,000+
34	174.86	0.01	.31	1.48	2.31	8.36	25.12	53.64	93.48	136.12	138.05	157.89	133.91	94.23	61.92	38.18	18.25	8.34	3.94	1.82	.95	.42	.18	1,000+
35	118.50	0.01	.22	.40	.97	3.89	10.10	24.45	49.98	87.78	124.87	134.91	135.29	132.98	102.19	73.28	42.82	24.44	13.44	6.06	3.15	1.46	.72	1,000+
36	67.48	0.03	.44	.96	1.21	2.45	3.92	10.45	23.58	46.78	76.31	108.89	134.02	135.29	107.92	84.22	54.22	34.87	24.44	13.44	6.06	3.15	1.46	1,000+
37	32.23	0.03	.71	.96	1.21	2.45	3.92	10.45	23.58	46.78	76.31	108.89	134.02	135.29	107.92	84.22	54.22	34.87	24.44	13.44	6.06	3.15	1.46	1,000+
38	14.96	0.03	.23	.67	1.21	2.45	3.92	10.45	23.58	46.78	76.31	108.89	134.02	135.29	107.92	84.22	54.22	34.87	24.44	13.44	6.06	3.15	1.46	1,000+
39	6.68	0.03	.23	.67	1.21	2.45	3.92	10.45	23.58	46.78	76.31	108.89	134.02	135.29	107.92	84.22	54.22	34.87	24.44	13.44	6.06	3.15	1.46	1,000+
40	5.86	0.03	.23	.67	1.21	2.45	3.92	10.45	23.58	46.78	76.31	108.89	134.02	135.29	107.92	84.22	54.22	34.87	24.44	13.44	6.06	3.15	1.46	1,000+
Total24	2.65	8.47	24.51	47.76	73.21	97.51	115.44	122.80	115.82	101.23	83.24	61.54	45.85	33.40	21.84	14.55	9.53	6.38	4.42	3.40	6.23	1,000+

SECTION C: RATIO PER 1,000 OF THE SEPARATE CHEST MEASUREMENTS TO EACH WEIGHT.

Weight, in pounds.		Proportion each weight per 1,000.	Chest, in inches.										Total. ^a	
			29	30	31	32	33	34	35	36	37	38		39
95-99	2.24	671.36	178.40	99.20	37.56	4.69	4.69	4.69	9.39	8.65	12.14	9.08	8.21	1,000-
100-104	2.45	362.36	276.26	175.96	68.31	36.75	20.75	9.94	11.24	8.65	8.65	5.01	5.68	1,000-
105-109	2.67	260.38	210.38	231.07	107.56	37.88	11.09	5.55	5.14	1.50	2.74	2.48	2.48	1,000-
110-114	2.81	188.04	168.04	285.99	185.26	63.84	18.51	8.40	2.67	1.06	2.74	2.18	1.56	1,000-
115-119	2.95	144.19	124.19	200.61	158.40	101.86	30.60	16.35	2.74	1.13	.69	.82	1.16	1,000-
120-124	3.10	104.29	84.29	146.25	118.44	68.40	40.00	20.46	3.62	1.13	.69	.82	1.16	1,000+
125-129	3.24	77.51	57.51	107.87	88.30	50.85	28.26	15.30	7.23	3.04	.83	.82	.60	1,000+
130-134	3.40	56.51	36.51	77.94	63.63	36.41	14.63	8.47	13.76	5.54	.78	.53	.60	1,000+
135-139	3.56	40.75	20.75	56.80	46.07	25.71	10.83	5.30	7.71	5.54	1.15	.53	.60	1,000+
140-144	3.72	29.03	9.03	40.82	32.77	18.01	7.74	3.77	4.46	10.97	2.24	.53	.60	1,000+
145-149	3.88	20.85	7.85	29.05	22.66	12.75	5.63	2.77	7.26	21.20	4.71	.88	.76	1,000+
150-154	4.04	15.82	5.82	21.63	16.66	9.77	4.33	2.11	10.37	35.65	8.95	1.60	1.45	1,000+
155-159	4.20	11.86	3.86	15.99	12.15	7.39	3.19	1.71	148.34	58.23	10.96	3.60	1.98	1,000+
160-164	4.36	8.88	2.88	11.99	9.08	5.33	2.16	1.18	165.71	88.20	31.83	7.10	4.15	1,000-
165-169	4.52	6.88	1.88	8.99	6.98	4.33	1.66	.88	118.01	124.70	49.17	14.14	8.19	1,000-
170-174	4.68	5.88	1.88	6.99	5.98	3.33	1.16	.60	218.01	162.61	70.15	26.72	19.15	1,000+
175-179	4.84	4.88	1.88	5.99	4.98	2.33	.66	.30	260.24	207.06	112.20	43.96	32.73	1,000+
180-184	5.00	3.88	1.88	4.99	3.98	1.33	.36	.00	250.47	206.23	154.27	71.06	65.22	1,000+
185-189	5.16	2.88	1.88	3.99	2.98	.33	.00	.00	212.54	224.82	191.82	136.52	100.44	1,000+
190-194	5.32	1.88	1.88	2.99	1.98	.33	.00	.00	172.33	223.86	210.23	198.53	133.13	1,000+
195-199	5.48	8.31	19.55	10.79	5.45	11.94	37.33	84.61	123.36	168.72	221.10	171.55	103.44	1,000+
200-204	5.64	4.23	1.84	2.58	1.66	2.03	5.15	13.62	49.60	116.72	177.84	187.96	437.78	1,000+
Total		20.56	56.23	118.38	182.77	201.47	174.86	118.50	67.48	32.23	14.98	6.68	5.86	1,000

^aThe maximum or minimum value of the + or - did not exceed 0.03.

TABLE IV.—Mean height, by groups and component sections, arranged in order of standing, with proportional weight and chest circumference (expiration) for each inch of height, also standard deviation for each height; first million draft recruits.

Group and section	Group and section No.	Description.	Number of men measured.	Mean height. Inches.	Standard deviation (height). Inches.	Mean weight. Pounds.	Mean chest. Mean height. Inches.
Average for the United States. Table I.							
Mountain whites	12		809,445	67.49	2.71	2,037	0.4920
Kentucky	1	0.7 per cent;	21,254	68.29	2.37	2,050	.4943
North Carolina	1	0.5 per cent;	4,033	68.21	2.51	2,081	.4940
South Carolina	1	0.5 per cent;	2,738	68.07	2.55	2,058	.4900
Tennessee	3	0.4 per cent;	1,864	68.19	2.83	2,060	.4940
Virginia	4	1.1 per cent;	5,900	68.43	2.61	2,060	.4810
West Virginia	1	0.9 per cent;	5,812	68.14	2.54	2,033	.4900
	1	3.7 per cent;	1,507	67.98	2.71	2,072	.4880
Agricultural, native white, South.	3		117,980	68.19	2.64	2,070	.4884
Alabama	3	Native percentage, white, 67.8 per cent; Negro, 31 per cent.	2,670	68.21	2.74	2,082	.4940
Arkansas	3	Native percentage, white, 66.9 per cent; Negro, 0.7 per cent.	1,550	68.64	2.60	2,080	.4840
Do.	3	Native percentage, white, 83.8 per cent; Negro, 12 per cent.	3,507	68.22	2.64	2,063	.4850
Kentucky	3	Native percentage, white, 76.4 per cent; Negro, 14.4 per cent.	11,466	67.85	2.52	2,060	.4940
Louisiana	3	Native percentage, white, 61 per cent; Negro, 34.4 per cent.	2,235	67.89	2.69	2,064	.4900
Maryland	3	Native percentage, white, 73.4 per cent; Negro, 14.3 per cent.	2,983	67.85	2.48	2,050	.4900
Mississippi	2	Native percentage, white, 61.3 per cent; Negro, 38.4 per cent.	13,638	68.44	2.48	2,094	.4880
Do.	3	Native percentage, white, 61.4 per cent; Negro, 37.5 per cent.	13,638	68.09	2.59	2,090	.4900
Do.	3	Native percentage, white, 70.7 per cent; Negro, 29.3 per cent.	1,329	68.03	2.51	2,090	.4880
North Carolina	3	Native percentage, white, 49.6 per cent; Negro, 51.7 per cent.	4,033	68.26	2.57	2,066	.4880
Do.	3	Native percentage, white, 57.1 per cent; Negro, 42.9 per cent.	2,653	68.21	2.72	2,071	.4850
Do.	3	Native percentage, white, 72.6 per cent; Negro, 27.4 per cent.	8,471	68.19	2.63	2,076	.4890
Oklahoma	3	Native percentage, white, 83.7 per cent; Negro, 16.3 per cent.	10,854	68.37	2.89	2,078	.4860
Tennessee	2	Native percentage, white, 74.1 per cent; Negro, 25.9 per cent.	6,308	68.39	2.60	2,040	.4840
Texas	2	Native percentage, white, 77.6 per cent; Negro, 22.4 per cent.	23,372	68.09	2.60	2,080	.4900
Do.	4	Native percentage, white, 62.3 per cent; Negro, 37.6 per cent.	2,723	68.09	2.70	2,090	.4870
Virginia	3	Native percentage, white, 64.8 per cent; Negro, 35.2 per cent.	5,812	68.12	2.73	2,090	.4880
West Virginia	2	Native percentage, white, 68.8 per cent; Negro, 31.2 per cent.	10,880	67.85	2.70	2,067	.4910
Mexican, sparsely settled.	14		11,064	68.16	2.69	2,080	.4870
Arizona	1	Indians, Chinese, and Japanese, 36.8 per cent; Mexicans, 63.2 per cent.	1,027	68.02	2.73	2,106	.4900
Do.	1	Indians, Chinese, and Japanese, 6.6 per cent; Mexicans, 93.4 per cent.	2,823	68.17	2.61	2,090	.4870
New Mexico	3	Mexicans, 14.3 per cent; native percentage, 85.6 per cent.	2,440	67.89	2.73	2,048	.4900
Texas	1	Mexicans, 17.1 per cent; native percentage, 82.9 per cent.	6,676	68.19	2.70	2,080	.4870

		10,000	68.12	2.61	2,000	4800
Indian, sparsely settled.	13					
Arizona.	1	1,077	68.02	2.73	2,106	4800
New Mexico.	1	293	67.26	2.90	2,068	4840
Oklahoma.	1	8,471	68.16	2.59	2,078	4850
South Dakota.	3	247	68.13	2.41	2,180	4860
German and Scandinavian, 10 per cent plus.	20					
Minnesota.	1	28,085	68.11	2.62	2,150	4851
Do.	2	6,461	68.44	2.54	2,170	4800
South Dakota.	1	7,601	68.14	2.63	2,170	4870
Wisconsin.	1	3,051	68.07	2.68	2,160	4820
Do.	2	3,297	68.13	2.66	2,130	4840
Do.	2	7,655	67.81	2.58	2,140	4850
Sparsely settled, not more than 3 per square mile	8					
California.	3	16,165	68.01	2.63	2,130	4829
Montana.	2	2,108	68.21	2.53	2,116	4800
Do.	2	6,531	68.17	2.57	2,150	4830
Nevada.	1	1,441	67.83	2.60	2,143	4870
New Mexico.	2	1,857	67.43	2.85	2,049	4830
Oregon.	2	1,077	68.13	2.52	2,140	4800
Utah.	1	1,224	68.16	2.64	2,114	4820
Wyoming.	1	1,927	67.79	2.63	2,130	4820
Native white of Scotch origin.	15					
Kentucky.	2	13,322	68.00	2.64	2,080	4844
North Carolina.	3	11,460	67.95	2.62	2,080	4840
Do.	2	2,053	68.24	2.72	2,074	4850
Scandinavians, 10 per cent	17					
Michigan.	1	31,009	67.96	2.63	2,150	4852
Minnesota.	1	2,344	67.10	2.61	2,160	5010
Do.	2	6,461	68.44	2.54	2,170	4860
Do.	2	7,601	68.14	2.63	2,170	4870
Do.	2	3,520	67.65	2.66	2,170	5020
North Dakota.	1	1,132	67.67	2.68	2,189	4880
Do.	2	3,307	68.03	2.48	2,159	4870
Do.	2	2,085	67.87	2.61	2,172	4890
South Dakota.	1	3,051	68.07	2.68	2,100	4820
Utah.	1	1,224	68.16	2.64	2,114	4820
Do.	2	2,781	67.73	2.56	2,108	4850
Washington.	2	6,501	67.87	2.70	2,140	4850
Wisconsin.	1	3,297	68.13	2.66	2,130	4840
Do.	2	7,655	67.81	2.58	2,140	4860
Desert.	9	6,121	67.86	2.72	2,080	4917
Arizona.	2	2,823	68.17	2.61	2,086	4870
Nevada.	1	1,441	67.83	2.60	2,143	4870
New Mexico.	2	1,857	67.43	2.85	2,049	4830

TABLE IV.—Mean height, by groups and component sections, arranged in order of standing, with proportional weight and chest circumference (expiration) for each inch of height; also standard deviation for each height; first million draft recruits.—Continued.

Group and section.	Group and section No.	Description.	Number of men measured.	Mean height. Inches.	Standard deviation (height). Inches.	Mean weight. Pounds.	Mean chest. Mean height. Inches.
Agricultural, Negroes, 45 per cent plus.	4		40,597	67.82	2.68	2,090	48.49
Alabama	2	Negro, 70.6 per cent; native parentage, white, 28.5 per cent	3,327	67.95	2.71	2,098	4890
Do	4	Negro, 72.8 per cent; native parentage, white, 26.9 per cent	669	68.16	2.61	2,115	4900
Arkansas	1	Negro, 55.3 per cent; native parentage, white, 44.7 per cent	4,945	68.05	2.63	2,083	4870
Georgia	2	Negro, 61 per cent; native parentage, white, 37.3 per cent	10,070	67.91	2.66	2,077	4920
Louisiana	1	Negro, 63 per cent; native parentage, white, 31.8 per cent	4,074	67.73	2.63	2,073	4910
Mississippi	1	Negro, 71.2 per cent; native parentage, white, 27.5 per cent	5,149	68.15	2.67	2,120	4880
North Carolina	4	Negro, 47.3 per cent; native parentage, white, 51.9 per cent	4,170	67.79	2.72	2,097	4900
South Carolina	2	Negro, 69.9 per cent; native parentage, white, 39.5 per cent	3,975	67.72	2.77	2,100	4910
Do	3	Negro, 62.2 per cent; native parentage, white, 37.7 per cent	3,804	68.07	2.69	2,080	4880
Tennessee	1	Negro, 44.2 per cent; native parentage, white, 55.8 per cent	2,218	68.46	2.65	2,110	4870
Texas	5	Negro, 51.1 per cent; native parentage, white, 37.3 per cent	1,416	68.46	2.65	2,110	4870
Virginia	2	Negro, 49.6 per cent; native parentage, white, 46.6 per cent	5,352	67.45	2.72	2,077	4900
Mountain	11		17,101	67.72	2.66	2,110	4921
Arkansas	2		1,559	68.64	2.60	2,050	4910
Massachusetts	1	foreign parentage, 30.7 per cent;	1,373	68.65	2.67	2,070	4920
Missouri	3	foreign parentage, 2.9 per cent;	1,139	68.63	2.61	2,080	4950
Montana	1	foreign parentage, 31.4 per cent;	5,117	67.82	2.66	2,150	4910
New Hampshire	1	foreign parentage, 21.6 per cent;	665	67.25	2.54	2,106	4910
New York	5	foreign parentage, 20 per cent;	786	67.16	2.69	2,074	4900
Do	8	foreign parentage, 24.7 per cent;	2,990	67.06	2.64	2,090	4970
Washington	3	foreign parentage, 20.6 per cent;	1,539	68.19	2.56	2,142	4900
Wyoming	1	foreign parentage, 22.3 per cent;	1,927	67.79	2.63	2,130	4920
Agricultural, mixed foreign and native white	2		97,340	67.62	2.66	2,110	4934
Colorado	4	69.5 per cent; foreign parentage, 16.2 per cent; foreign	1,227	68.06	2.70	2,087	4960
Illinois	8	54.1 per cent; foreign parentage, 31.5 per cent; foreign	2,461	67.77	2.63	2,110	4930
Indiana	2	76.2 per cent; foreign parentage, 16.8 per cent; foreign	937	68.12	2.49	2,120	4910
Iowa	1	60.7 per cent; foreign parentage, 34.2 per cent; foreign	12,136	68.09	2.56	2,139	4920
Kansas	2	72.9 per cent; foreign parentage, 16.1 per cent; foreign	5,504	68.18	2.54	2,108	4980

Michigan	2	foreign parentage, 29.4 per cent; foreign	12,567	67.63	2.55	2.10	.493
Nebraska	2	foreign parentage, 39.5 per cent; foreign	3,145	68.21	2.59	2.136	.490
New Jersey	2	foreign parentage, 21.7 per cent; foreign	8,966	66.83	2.70	2.073	.490
New York	7	foreign parentage, 17.9 per cent; foreign	6,465	67.45	2.64	2.036	.490
Ohio	2	; foreign parentage, 20 per cent; foreign	14,443	67.31	2.74	2.066	.4910
Pennsylvania	6	foreign parentage, 20.5 per cent; foreign	8,616	67.37	2.90	2.099	.4940
South Dakota	1	foreign parentage, 37.2 per cent; foreign	3,031	68.07	2.63	2.100	.4920
Vermont	(11)	foreign parentage, 33.7 per cent; foreign	2,077	67.12	2.52	2.061	.4960
Washington	1	foreign parentage, 22.9 per cent; foreign	5,176	68.01	2.60	2.139	.4920
Wisconsin	2	foreign parentage, 43.2 per cent; foreign	7,685	67.81	2.58	2.140	.4950
<hr/>							
Agricultural, native white, North, native white over 75 per cent, North.	1		66,865	67.60	2.63	2.09	.4900
<hr/>							
Illinois	3	33.2 per cent; foreign parentage, 10.3 per cent, per cent.	8,928	67.86	2.59	2.094	.4870
Indiana	3	; 32.5 per cent, foreign parentage, 11 per cent; per cent.	18,743	67.84	2.56	2.083	.4870
Iowa	2	73.1 per cent; foreign parentage, 17.7 per cent; per cent.	7,401	67.96	2.61	2.106	.4880
Ohio	3	78.7 per cent; foreign parentage, 13.7 per cent; per cent.	17,606	67.75	2.59	2.085	.4890
Pennsylvania	2	76.5 per cent; foreign parentage, 9.3 per cent, per cent.	14,218	68.73	2.62	2.085	.4970
<hr/>							
Mining	7		35,720	67.49	2.72	2.110	.4900
<hr/>							
Alabama	1	negro, 25.6 per cent	8,841	67.67	2.67	2.071	.4840
California	2	foreign parentage, 27.3 per cent;	943	67.69	2.64	2.154	.4960
Colorado	1	foreign parentage, 15.7 per cent;	1,066	68.06	2.79	2.061	.4890
Do.	3	foreign parentage, 27.1 per cent,	381	68.12	2.66	2.066	.4870
Do.	6	foreign parentage, 22.9 per cent;	1,223	67.65	2.71	2.090	.4840
Idaho	1	foreign parentage, 23.1 per cent,	4,031	68.10	2.57	2.163	.4950
Montana	1	foreign parentage, 31.4 per cent;	5,117	67.83	2.65	2.130	.4910
Nevada	1	foreign parentage, 25.8 per cent;	1,441	67.83	2.69	2.163	.4970
Pennsylvania	3	foreign parentage, 32.5 per cent;	7,306	68.65	2.57	2.105	.5000
Do.	4	foreign parentage, 18.1 per cent;	4,827	68.80	2.69	2.109	.4960
Utah	3	foreign parentage, 36.8 per cent;	563	67.65	2.78	2.127	.4940

TABLE IV.—Mean height, by groups and component sections, arranged in order of ascending, with proportional weight and chest circumference (expiration) for each inch of height; also standard deviation for each height; first million draft recruits—Continued.

Group and section	Group and section No.	Description.	Number of men measured.	Mean height. Inches.	Standard deviation (height). Inches.	Mean weight. Pounds.	Mean chest height. Inch.
Finland, 10 per cent	14		5,364	67.43	2.65	2,100	0.5016
Michigan	1	Large Finnish population; Scandinavians, 23.1 per cent.	2,344	67.10	2.61	2,100	.5010
Minnesota	3	Large Finnish population; Scandinavians, 31.1 per cent.	3,520	67.65	2.66	2,176	.5020
German and Austrian, 20 per cent plus	21		38,962	67.41	2.60	2,130	.4955
Illinois	1	Germans, 21.2 per cent; Austrians, 4.3 per cent.	6,303	67.43	2.67	2,123	.4950
Do.	4	Germans, 17.4 per cent; Austrians, 2.5 per cent.	4,236	67.59	2.64	2,115	.4940
Indiana	1	Germans, 17.2 per cent; Austrians, 4.1 per cent.	3,614	67.22	2.64	2,113	.4940
Minnesota	2	Germans, 22.3 per cent; Austrians, 2.8 per cent; Scandinavians, 16.8 per cent.	7,801	68.14	2.63	2,176	.4970
Ohio	1	Germans, 18.9 per cent; Austrians, 8.2 per cent.	17,204	67.66	2.67	2,111	.4960
Maritime	10		6,161	67.31	2.70	2,060	.4903
Maine	2	7.8 per cent;	328	67.59	2.39	2,091	.4970
Maryland	2	1.7 per cent.	1,068	67.37	2.60	2,060	.4900
Do.	4	1.8 per cent.	(a)	(a)	(a)	(a)	
Massachusetts	3	25.2 per cent;	1,177	68.90	2.70	2,070	.4910
North Carolina	5	0.9 per cent;	254	67.69	2.61	2,067	.4910
Virginia	1	3.6 per cent;	2,846	67.34	2.73	2,091	.4870
German and Austrian, 15 per cent plus.	22		126,994	67.27	2.72	2,120	.4954
Illinois	1	Germans, 15.2 per cent;	6,303	67.43	2.67	2,123	.4950
Do.	4	Germans, 11.4 per cent;	4,236	67.59	2.64	2,115	.4940
Indiana	1	Germans, 17.2 per cent;	3,614	67.22	2.64	2,113	.4940
Iowa	1	Germans, 17.2 per cent;	12,136	68.14	2.63	2,176	.4970
Minnesota	2	Germans, 22.3 per cent;	7,801	68.14	2.63	2,176	.4970
Nebraska	1	Germans, 18.9 per cent;	17,204	67.66	2.67	2,111	.4960
Do.	2	Germans, 18.9 per cent;	17,204	67.66	2.67	2,111	.4960
New Jersey	1	Germans, 18.9 per cent;	17,204	67.66	2.67	2,111	.4960
Ohio	1	Germans, 18.9 per cent;	17,204	67.66	2.67	2,111	.4960
Pennsylvania	3	Germans, 18.9 per cent;	17,204	67.66	2.67	2,111	.4960
Do.	7	Germans, 18.9 per cent;	17,204	67.66	2.67	2,111	.4960
Do.	1	Germans, 18.9 per cent;	17,204	67.66	2.67	2,111	.4960
Wisconsin	2	Germans, 18.9 per cent;	17,204	67.66	2.67	2,111	.4960
Do.	3	Germans, 18.9 per cent;	17,204	67.66	2.67	2,111	.4960
Do.	4	Germans, 18.9 per cent;	17,204	67.66	2.67	2,111	.4960

Russian, 10 per cent plus.		12, 076	67.11	2.68	2.120	.4978
16	Colorado.....	1,105	67.83	2.67	2.064	.4900
1	Kansas.....	1,087	68.30	2.57	2.122	.4860
3	North Dakota.....	2,005	67.87	2.61	2.172	.4900
3	Pennsylvania.....	7,305	66.55	2.57	2.105	.4900
2	South Dakota.....	791	67.92	2.53	2.170	.4950
6	Commuters.....	28,032	66.86	2.75	2.09	.4970
1	Illinois.....	6,303	67.43	2.67	2.123	.4960
1	New Jersey.....	17,795	66.72	2.74	2.078	.4970
1	New York.....	4,931	66.65	2.76	2.091	.4970
5	Eastern manufacturing	81,714	66.77	2.70	2.09	.4970
2	Connecticut.....	8,708	66.73	2.73	2.096	.4960
2	Massachusetts.....	18,447	66.67	2.67	2.07	.4970
2	New Hampshire.....	1,575	66.86	2.61	2.081	.4930
1	New Jersey.....	17,795	66.72	2.74	2.078	.4970
3	New York.....	3,150	66.87	2.66	2.092	.4960
1	Ohio.....	17,708	67.06	2.67	2.111	.4950
5	Pennsylvania.....	8,992	66.66	2.69	2.116	.4970
1	Rhode Island.....	3,928	66.40	2.61	2.06	.4940
19	French-Canadian, 10 per cent.....	25,892	66.67	2.65	2.07	.4966
3	Maine.....	1,267	67.07	2.58	2.06	.4950
2	Massachusetts.....	18,447	66.67	2.67	2.07	.4970
1	New Hampshire.....	665	67.25	2.54	2.106	.4910
2	Do.....	1,575	66.86	2.61	2.081	.4930
1	Rhode Island.....	3,928	66.40	2.61	2.06	.4940

(a) Not tabulated.

TABLE V.—Mean weight, by groups and component sections, arranged in order of standing, with proportional weight for each inch of height and chest circumference (expiration) for each pound of weight; also standard deviation for each weight; first million draft recruits.

Group and section.	Group and section No.	Description.	Number of men measured.	Mean weight.		Standard deviation (weight).	Mean weight.		Mean chest.
				Pounds.	Pounds.		Pounds.	Pounds.	
Average for the United States. Table I.									
Germans and Scandinavians, 10 per cent plus.									
20									
Minnesota.	1	Scandinavians, 37.4 per cent; Germans, 10.3 per cent.	898,445	141.54	17.42	2.097	146.06	17.00	2.200
Do.	2	Scandinavians, 16.8 per cent; Germans, 22.3 per cent.	28,063	146.06	17.00	2.150	146.06	17.00	2.200
South Dakota.	1	Scandinavians, 15.5 per cent; Germans, 10.7 per cent.	6,461	146.28	16.61	2.170	146.28	16.61	2.200
Do.	2	Scandinavians, 15.5 per cent; Germans, 10.7 per cent.	7,601	147.64	17.31	2.170	147.64	17.31	2.200
Wisconsin.	1	Scandinavians, 22.3 per cent; Germans, 13.6 per cent.	3,520	146.43	16.84	2.170	146.43	16.84	2.200
Do.	2	Scandinavians, 10.2 per cent; Germans, 26.3 per cent.	1,132	146.10	16.20	2.159	146.10	16.20	2.200
17									
Scandinavians, 10 per cent	17	Scandinavians, 10.3 per cent; Germans, 26.3 per cent.	3,005	146.93	16.23	2.172	146.93	16.23	2.200
Michigan.	1	Scandinavians, 23.1 per cent; large Finnish population.	3,005	147.48	16.83	2.160	147.48	16.83	2.200
Minnesota.	1	Scandinavians, 37.4 per cent; Germans, 10.3 per cent.	3,005	146.80	16.54	2.114	146.80	16.54	2.200
Do.	2	Scandinavians, 15.5 per cent; Germans, 10.7 per cent.	2,781	142.96	16.83	2.105	142.96	16.83	2.200
South Dakota.	1	Scandinavians, 13.9 per cent	3,005	147.48	16.83	2.172	147.48	16.83	2.200
Do.	2	Scandinavians, 15.5 per cent	3,005	146.93	16.23	2.172	146.93	16.23	2.200
Utah.	1	Scandinavians, 10.4 per cent	1,224	144.06	15.46	2.114	144.06	15.46	2.230
Do.	2	Scandinavians, 10.5 per cent	2,781	142.96	16.83	2.105	142.96	16.83	2.230
Washington.	1	Scandinavians, 13.5 per cent; Japanese, and Indians.	6,601	145.25	17.28	2.140	145.25	17.28	2.200
Do.	2	Scandinavians, 22.3 per cent; Germans, 13.6 per cent	3,297	145.13	16.93	2.130	145.13	16.93	2.230
Wisconsin.	1	Scandinavians, 22.3 per cent; Germans, 26.3 per cent.	7,685	144.94	17.13	2.140	144.94	17.13	2.230
Do.	2	Scandinavians, 10.2 per cent; Germans, 26.3 per cent.	5,804	145.76	16.86	2.150	145.76	16.86	2.200
Finns, 10 per cent									
Michigan.	1	Scandinavians, 23.1 per cent; large Finnish population.	2,344	144.74	16.83	2.160	144.74	16.83	2.230
Minnesota.	1	Scandinavians, 31.1 per cent; large Finnish population.	3,520	146.44	16.84	2.170	146.44	16.84	2.230
Sparsely settled, not more than 3 per square mile									
California.	1	Scandinavians, 23.1 per cent; large Finnish population.	16,165	144.94	16.93	2.130	144.94	16.93	2.210
Montana.	2	Sparsely settled; foreign born, white, 17.3 per cent.	2,108	144.30	17.53	2.116	144.30	17.53	2.210
Do.	3	Sparsely settled; foreign born, white, 18.1 per cent; Indians, Chinese, and Japanese, 6.1 per cent.	6,531	146.80	16.65	2.150	146.80	16.65	2.200
Nevada.	1	Sparsely settled; foreign born, white, 25.6 per cent; Japanese, 8.4 per cent	1,441	145.25	17.11	2.143	145.25	17.11	2.220
New Mexico.	1	Sparsely settled; foreign born, white, 9.2 per cent.	1,877	139.20	16.42	2.049	139.20	16.42	2.200
Oregon.	2	Sparsely settled; foreign born, white, 9.1 per cent.	1,077	145.82	16.64	2.140	145.82	16.64	2.200
Utah.	1	Sparsely settled; foreign born, white, 11.7 per cent.	1,224	144.06	15.46	2.114	144.06	15.46	2.230
Do.	2	Sparsely settled; foreign born, white, 11.7 per cent.	1,977	144.61	16.90	2.190	144.61	16.90	2.210
Wyoming.	1	Sparsely settled; foreign born, white, 18.6 per cent.	1,977	144.61	16.90	2.190	144.61	16.90	2.210
21									
Germans and Austrians, 20 per cent plus.	21	Sparsely settled; foreign born, white, 18.6 per cent.	38,902	143.27	18.05	2.130	143.27	18.05	2.230
Illinois.	1	Germans, 21.2 per cent; Austrians, 4.3 per cent.	6,303	143.19	17.96	2.123	143.19	17.96	2.230
Do.	4	Germans, 17.4 per cent; Austrians, 2.4 per cent.	4,266	143.02	17.92	2.115	143.02	17.92	2.230
Indiana.	1	Germans, 17.2 per cent; Austrians, 4.1 per cent.	3,614	142.07	18.15	2.113	142.07	18.15	2.240

Minnesota.....	2	Germans, 22.3 per cent; Austrians, 2.9 per cent; Scandinavians, 16.8 per cent.	7,001	147.64	17.31	2,179	.2300
Ohio.....	1	Germans, 18.9 per cent; Austrians, 8.5 per cent.	17,208	142.82	14.15	2,111	.2340
Mountain.....	11		17,103	142.97	16.76	2,110	.2330
Arkansas.....	2	98.9 per cent	1,559	140.78	14.90	2,050	.2300
Massachusetts.....	1	foreign percentage, 30.7 per cent; foreign	1,373	138.53	17.13	2,070	.2370
Missouri.....	3	foreign percentage, 3.9 per cent; foreign	1,139	142.49	15.68	2,090	.2340
Montana.....	1	foreign percentage, 31.4 per cent; foreign	5,117	145.70	16.65	2,150	.2300
New Hampshire.....	1	foreign percentage, 21.6 per cent; foreign	665	141.57	17.98	2,016	.2380
New York.....	5	foreign percentage, 20 per cent; foreign	795	139.30	16.74	2,074	.2390
Do.....	8	foreign percentage, 24.7 per cent; foreign	2,900	140.21	16.71	2,000	.2370
Washington.....	3	foreign percentage, 20.6 per cent; foreign	1,530	146.07	16.29	2,142	.2300
Wyoming.....	1	foreign percentage, 22.3 per cent; foreign	1,927	144.61	16.90	2,130	.2310
Agricultural, mixed foreign and native white.....	2		97,340	142.79	17.28	2,11	.2318
Colorado.....	4	foreign percentage, 18.2 per cent; foreign	1,227	142.05	16.20	2,097	.2330
Illinois.....	8	foreign percentage, 31.5 per cent; foreign	2,451	143.01	17.17	2,110	.2330
Indiana.....	2	foreign percentage, 16.4 per cent; foreign	837	144.45	17.34	2,120	.2310
Iowa.....	1	foreign percentage, 34.2 per cent; foreign	12,136	145.57	17.10	2,139	.2300
Kansas.....	2	foreign percentage, 16.1 per cent; foreign	8,504	143.50	17.21	2,105	.2310
Michigan.....	2	foreign percentage, 29.4 per cent; foreign	12,597	142.01	16.85	2,100	.2350
Nebraska.....	2	foreign percentage, 39.5 per cent; foreign	3,145	145.70	17.73	2,136	.2200
New Jersey.....	2	foreign percentage, 21.7 per cent; foreign	8,908	138.92	17.34	2,076	.2400
New York.....	7	foreign percentage, 17.9 per cent; foreign	6,465	141.53	17.62	2,068	.2300
Ohio.....	2	; foreign percentage, 20 per cent; foreign	14,443	141.10	17.31	2,008	.2340
Pennsylvania.....	6	foreign percentage, 30.5 per cent; foreign	8,616	141.40	16.93	2,090	.2300
South Dakota.....	1	foreign percentage, 37.2 per cent; foreign	3,051	146.49	18.54	2,160	.2200
Vermont.....	All.	foreign percentage, 33.7 per cent; foreign	2,077	140.33	16.43	2,001	.2380
Washington.....	1	foreign percentage, 22.9 per cent; foreign	5,176	145.50	17.10	2,139	.2300
Wisconsin.....	2	foreign percentage, 43.2 per cent; foreign	7,685	144.94	17.13	2,140	.2320

TABLE V.—Mean weight, by groups and component actions, arranged in order of standing, with proportional weight for each inch of height and chest circumference (expiration) for each pound of weight; also standard deviation for each weight, first million draft recruits—Continued.

Group and section.	Group and section No.	Description.	Number of men measured.	Mean height. Inches.	Standard deviation (height). Inches.	Mean weight. Pounds.	Mean chest. Inches.
German and Austrians, 15 per cent plus.	22		126,984	142.31	17.73	2,120	0, 2240
Illinois.....	1	German, 21.2 per cent; Austrians, 4.3 per cent.	6,303	143.19	17.88	2,123	2330
Do.....	4	German, 17.4 per cent; Austrians, 2.8 per cent.	4,236	143.02	17.92	2,115	2330
Indiana.....	1	German, 17.2 per cent; Austrians, 4.1 per cent.	3,614	142.07	18.15	2,113	2350
Iowa.....	1	German, 15.9 per cent; Austrians, 1.9 per cent.	12,136	145.67	17.10	2,139	2300
Minnesota.....	2	German, 22.3 per cent; Austrians, 2.9 per cent.	7,601	147.64	17.31	2,170	2300
Nebraska.....	1	German, 22.3 per cent; Austrians, 2.9 per cent.	7,629	144.37	17.48	2,120	2300
Do.....	2	German, 22.3 per cent; Austrians, 2.9 per cent.	3,145	145.70	17.73	2,136	2300
New Jersey.....	1	German, 17.85 per cent; Austrians, 1.9 per cent.	17,785	139.06	17.59	2,078	2300
Ohio.....	1	German, 17.205 per cent; Austrians, 1.8 per cent.	17,205	141.62	18.15	2,111	2340
Pennsylvania.....	3	German, 17.205 per cent; Austrians, 1.8 per cent.	7,205	140.10	17.17	2,105	2370
Do.....	5	German, 17.205 per cent; Austrians, 1.8 per cent.	8,892	141.09	17.02	2,116	2350
Do.....	7	German, 17.205 per cent; Austrians, 1.8 per cent.	17,243	139.55	17.56	2,063	2300
Wisconsin.....	1	German, 13.6 per cent; Austrians, 3.2 per cent.	3,297	145.13	16.93	2,130	2300
Do.....	2	German, 13.6 per cent; Austrians, 3.2 per cent.	7,665	144.94	17.13	2,140	2300
Do.....	4	German, 13.6 per cent; Austrians, 3.2 per cent.	5,895	144.35	17.48	2,140	2340
Russians, 10 per cent plus.	16		12,076	142.30	17.21	2,120	2340
Colorado.....	2	Russian, 8.3 per cent; Austrians, 4.3 per cent.	1,105	142.04	15.50	2,094	2340
Kansas.....	1	Russian, 13.1 per cent; Austrians, 4.3 per cent.	1,067	144.05	17.44	2,122	2300
North Dakota.....	3	Russian, 26.7 per cent; Austrians, 4.3 per cent.	2,065	147.46	16.83	2,172	2280
Pennsylvania.....	3	Russian, 21.1 per cent; Austrians, 4.3 per cent.	7,305	140.10	17.17	2,106	2270
South Dakota.....	2	Russian, 25.6 per cent; Austrians, 4.3 per cent.	7,664	147.22	16.15	2,170	2280
Mexican, sparsely settled.	14		10,779	142.14	17.36	2,090	2335
Arizona.....	1	Indian, Chinese, and Japanese, 26.6 per cent; Mexicans, 8.4 per cent.	1,027	143.29	16.93	2,106	2320
Do.....	2	Indian, Chinese, and Japanese, 6.6 per cent; Mexicans, 7.8 per cent.	2,823	142.93	17.34	2,096	2320
New Mexico.....	3	Mexican, 14.3 per cent; native percentage, 61.5 per cent.	8,640	139.01	17.36	2,048	2340
Texas.....	1	Mexican, 17.1 per cent; native percentage, 44.1 per cent.	6,676	141.85	17.40	2,050	2340
Mining.....	7		35,730	142.25	16.96	2,110	2330
Alabama.....	1	Native percentage, white, 71.5 per cent; Negro, 28.5 per cent.	8,941	140.31	16.41	2,071	2330
California.....	2	Native percentage, white, 71.5 per cent; Negro, 28.5 per cent.	943	145.64	16.86	2,154	2310
Colorado.....	1	Native percentage, white, 71.5 per cent; Negro, 28.5 per cent.	1,056	141.64	15.73	2,061	2360
Do.....	3	Native percentage, white, 71.5 per cent; Negro, 28.5 per cent.	381	142.13	15.50	2,066	2330
Do.....	6	Native percentage, white, 71.5 per cent; Negro, 28.5 per cent.	1,223	139.40	16.10	2,060	2360
Idaho.....	1	Native percentage, white, 71.5 per cent; Negro, 28.5 per cent.	4,031	145.31	15.20	2,133	2300
Montana.....	1	Native percentage, white, 71.5 per cent; Negro, 28.5 per cent.	5,117	143.70	16.63	2,150	2290
Nevada.....	1	Native percentage, white, 71.5 per cent; Negro, 28.5 per cent.	1,441	145.36	17.11	2,145	2300

Pennsylvania.	3	per cent; foreign percentage, 32.5 per cent; foreign cent.	7,306	140.10	17.17	2.103	2,370
Do.	4	per cent; foreign percentage, 18.1 per cent; foreign cent.	4,827	140.04	17.22	2.100	2,366
Utah.	3	per cent; foreign percentage, 36.9 per cent; foreign cent.	363	143.88	16.54	2.127	2,320
Desert.	9		6,121	142.08	17.23	2.080	2,340
Arizona.	2	cent; foreign percentage, 22.5 per cent; foreign cent.	2,823	142.05	17.34	2.086	2,330
Nevada.	1	cent; foreign percentage, 25.5 per cent; foreign cent.	1,441	145.33	17.11	2.143	2,320
New Mexico.	2	cent; foreign percentage, 6.2 per cent; foreign cent.	1,857	138.20	16.42	2.049	2,400
Indian, sparsely settled.	13		10,038	141.80	16.91	2.080	2,330
Arizona.	1	Indians, Chinese, and J Indians, 29.1 per cent; n	1,027	143.20	16.93	2.106	2,320
New Mexico.	1	Indians, 9.2 per cent, n	283	139.12	16.85	2.083	2,300
Oklahoma.	1	Per cent.	8,471	141.53	16.80	2.078	2,330
South Dakota.	3	Indians, 87.2 per cent; native percentage, white, 8.1 per cent.	247	143.20	16.77	2.180	2,290
Agricultural, Negroes, 45 per cent plus	4		49,303	141.61	16.64	2.090	2,244
Alabama.	2	Negro, 70.6 per cent; native percentage, white, 28.5 per cent.	3,227	142.57	16.77	2.066	2,230
Do.	2	Negro, 72.8 per cent; native percentage, white, 26.9 per cent.	3,669	144.21	16.81	2.115	2,200
Arkansas.	1	Negro, 55.3 per cent; native percentage, white, 41.7 per cent.	4,945	141.81	16.39	2.063	2,260
Georgia.	1	Negro, 61 per cent; native percentage, white, 37.3 per cent.	10,075	141.09	16.63	2.077	2,260
Louisiana.	1	Negro, 63 per cent; native percentage, white, 31.3 per cent.	4,074	140.47	16.45	2.120	2,260
Mississippi.	1	Negro, 71.2 per cent; native percentage, white, 27.8 per cent.	5,149	144.16	16.45	2.130	2,210
North Carolina.	4	Negro, 47.3 per cent; native percentage, white, 51.9 per cent.	3,575	142.13	17.01	2.067	2,330
South Carolina.	1	Negro, 59.9 per cent; native percentage, white, 39.5 per cent.	3,970	142.04	16.59	2.077	2,240
Do	2	Negro, 62.2 per cent; native percentage, white, 35.7 per cent.	3,804	138.90	15.70	2.000	2,330
Tennessee.	2	Negro, 44.2 per cent; native percentage, white, 54.8 per cent.	2,218	141.67	17.11	2.000	2,320
Texas.	1	Negro, 51.1 per cent; native percentage, white, 37.3 per cent.	1,346	144.66	13.23	2.110	2,310
Virginia.	2	Negro, 49.6 per cent; native percentage, white, 46.6 per cent.	5,352	140.10	16.43	2.077	2,290
Agricultural, native white, South.	3		117,548	141.44	16.33	2.070	2,240
Alabama.	3	Native percentage, white, 67.6 per cent; Negro, 31 per cent.	2,670	140.71	16.94	2.062	2,250
Arkansas.	2	Native percentage, white, 60.9 per cent; Negro, 0.7 per cent.	1,559	140.78	14.90	2.060	2,290
Do.	2	Native percentage, white, 63.9 per cent; Negro, 12 per cent.	3,007	140.77	16.13	2.063	2,250
Kentucky.	2	Native percentage, white, 76.4 per cent; Negro, 14.4 per cent.	11,460	140.03	16.76	2.064	2,240
Louisiana.	3	Native percentage, white, 61 per cent; Negro, 34.4 per cent.	5,235	140.13	16.23	2.004	2,290
Maryland.	2	Native percentage, white, 73.4 per cent; Negro, 14.8 per cent.	2,663	140.78	16.43	2.070	2,270
Mississippi.	2	Native percentage, white, 64.5 per cent; Negro, 33.4 per cent.	3,394	141.51	16.43	2.000	2,210
Missouri.	1	Native percentage, white, 94.4 per cent; Negro, 0.3 per cent.	13,588	141.67	17.03	2.080	2,240
Do.	3	Native percentage, white, 80.9 per cent; Negro, 38.1 per cent.	1,180	152.49	15.93	2.060	2,240
North Carolina.	2	Native percentage, white, 51.9 per cent; Negro, 47.3 per cent.	2,033	141.07	17.14	2.066	2,250
Do.	2	Native percentage, white, 57.1 per cent; Negro, 41.2 per cent.	3,403	141.63	16.35	2.074	2,250
Do.	6	Native percentage, white, 72.6 per cent; Negro, 13.7 per cent.	7,444	140.63	16.80	2.078	2,230
Oklahoma.	1	Native percentage, white, 82.7 per cent; Negro, 5 per cent.	8,471	141.63	16.93	2.040	2,270
Tennessee.	2	Native percentage, white, 74.1 per cent; Negro, 22 per cent.	10,958	142.62	16.38	2.060	2,210
Texas.	4	Native percentage, white, 77.6 per cent; Negro, 13.9 per cent.	6,808	139.40	17.29	2.060	2,230
Do	2	Native percentage, white, 53.3 per cent; Negro, 26.8 per cent.	2,722	142.31	17.06	2.060	2,270
Do	2	Native percentage, white, 64.8 per cent; Negro, 33.2 per cent.	2,865	140.77	16.28	2.065	2,270
Do	2	Native percentage, white, 66.8 per cent; Negro, 4.5 per cent.	10,960	141.62	16.96	2.067	2,260
West Virginia.	3						

TABLE V.—Mean weight, by groups and component sections, arranged in order of standing, with proportional weight for each inch of height and chest circumference (expiration) for each pound of weight; also standard deviation for each weight; first million draft recruits—(continued).

Group and section.	Group and section No.	Description.	Number of men measured.	Mean height. Inches.	Standard deviation (height). Inches.	Mean weight. Pounds.	Mean chest. Mean height. Inches.
Agricultural, native white, North, native white over 75 per cent, North.	1		66,865	141.22	17.45	2,000	2,240
Illinois.....	3	2 per cent; foreign parentage, 10.8 per cent, foreign cent.	8,928	142.13	17.25	2,004	2,233
Indiana.....	3	.5 per cent; foreign parentage, 11 per cent, foreign cent.	14,743	141.37	17.80	2,063	2,280
Iowa.....	2	1 per cent; foreign parentage, 17.7 per cent, foreign cent.	7,403	143.15	17.27	2,105	2,319
Ohio.....	3	7 per cent; foreign parentage, 13.7 per cent; foreign cent.	17,005	141.27	17.46	2,065	2,240
Pennsylvania.....	2	.5 per cent; foreign parentage, 9.8 per cent; foreign born, white, 7.9 per cent.	14,218	139.53	17.06	2,005	2,270
Maritime.....	10		6,161	141.38	16.86	2,000	2,250
Maine.....	2	7.8 per cent;	828	141.37	16.10	2,001	2,270
Maryland.....	2	1.7 per cent;	1,068	140.01	16.56	2,060	2,300
Do.....	4	1.3 per cent;	(*)	(*)	(*)	(*)	(*)
Massachusetts.....	3	25.2 per cent;	1,127	138.70	16.76	2,070	2,270
North Carolina.....	5	0.9 per cent;	254	141.27	15.86	2,067	2,260
Virginia.....	1	3.6 per cent;	2,886	140.82	17.25	2,001	2,300
Mountain whites.....	12		21,254	140.74	15.65	2,050	2,267
Kentucky.....	1	0.7 per cent;	4,033	139.92	15.26	2,051	2,270
North Carolina.....	1	0.5 per cent;	2,738	141.22	15.96	2,056	2,280
South Carolina.....	1	0.4 per cent;	1,664	140.42	16.72	2,080	2,260
Tennessee.....	3	1.1 per cent;	5,900	140.02	16.43	2,050	2,260
Virginia.....	4	0.9 per cent;	5,512	140.02	15.94	2,065	2,260
West Virginia.....	1	3.7 per cent,	1,507	140.55	16.45	2,072	2,280

TABLE VI.—Mean chest girth by groups and component sections, arranged on order of standing, with proportional chest circumference (expiration) for each inch of height and each pound of weight; also standard deviation for each chest measurement; first million draft recruits.

Group and section.	Group and section No.	Description.	Number of men measured.	Mean chest girth.	Standard deviation (chest).	Mean chest. Mean height.	Mean chest. Mean weight.
Average for the United States (Table II).							
Fin, 10 per cent.	18		873, 150	33.22	2.01	Inch. 0 4920	Pound. 0.2340
Michigan	1		5, 855	33.82	1.99	.5016	.2320
Minnesota	3	Large Finnish population; Scandinavians, 23.1 per cent.	2, 344	33.63	1.93	.5010	.2320
German and Scandinavian, 10 per cent plus.	20	Large Finnish population; Scandinavians, 31.1 per cent.	3, 320	33.65	1.98	.5020	.2320
Minnesota	1		26, 066	33.72	1.95	.4950	.2310
Minnesota	2	Scandinavians, 37.4 per cent; Germans, 10.2 per cent.	6, 461	33.86	1.86	.4850	.2280
Minnesota	2	Scandinavians, 16.8 per cent; Germans, 22.3 per cent.	7, 001	33.86	1.93	.4970	.2300
South Dakota	1	Scandinavians, 16.8 per cent; Germans, 10.7 per cent.	3, 051	33.51	1.96	.4920	.2280
Wisconsin	1	Scandinavians, 22.3 per cent; Germans, 13.5 per cent.	2, 297	33.64	1.89	.4940	.2300
Do.	2	Scandinavians, 10.2 per cent; Germans, 26.3 per cent.	7, 685	33.56	1.97	.4950	.2320
Scandinavian, 10 per cent	17		50, 953	33.65	1.95	.4950	.2310
Michigan	1		2, 344	33.63	1.96	.5010	.2320
Minnesota	1		6, 461	33.86	1.86	.4850	.2280
Do.	2		7, 001	33.86	1.93	.4970	.2300
Do.	3		3, 020	33.95	1.98	.5020	.2340
North Dakota	1		1, 122	33.72	1.91	.4980	.2300
Do.	2		3, 307	33.82	1.88	.4970	.2300
Do.	3		2, 005	33.70	1.86	.4900	.2280
South Dakota	1		3, 051	33.51	1.95	.4920	.2280
Utah	1		1, 224	33.59	1.82	.4920	.2300
Do.	2		2, 781	32.49	1.85	.4850	.2300
Washington	2		6, 001	33.41	1.96	.4920	.2300
Wisconsin	1		3, 297	33.68	1.89	.4940	.2320
Do.	2		7, 685	33.56	1.97	.4950	.2320
Sparsely settled, not more than 3 per square mile	8		16, 151	33.53	1.92	.4930	.2320
California	3		2, 108	33.40	1.97	.4900	.2310
Montana	3		6, 331	33.50	1.85	.4980	.2300
Nevada	1		1, 441	33.75	2.08	.4970	.2320
New Mexico	2		1, 857	33.26	1.84	.4900	.2300
Oregon	2		1, 077	33.40	1.91	.4900	.2300
Utah	1		1, 224	33.59	1.82	.4920	.2300
Wyoming	1		1, 927	33.38	1.89	.4920	.2310

German and Austrian, 20 per cent plus.	21	35,911	28.42	2.07	4900	2330
Illinois.....	1	6,808	23.38	2.02	4950	2330
Do.....	4	330	2.08	2.08	4950	2330
Indiana.....	1	9,614	33.43	2.12	4970	2330
Minnesota.....	1	7,601	23.38	1.98	4970	2330
Ohio.....	2	17,208	33.20	2.08	4950	2340
Russian, 10 per cent plus.	16	12,084	33.30	2.01	4976	2350
Colorado.....	2	1,108	33.30	1.75	4970	2340
Kansas.....	1	1,087	33.24	2.68	4980	2360
North Dakota.....	3	2,035	33.70	1.86	4980	2360
Pennsylvania.....	2	7,305	33.32	2.10	5000	2370
South Dakota.....	2	6,594	33.61	1.87	4980	2380
Agricultural, mixed foreign and native white.	2	97,319	33.36	2.00	4984	2340
Colorado.....	4	1,227	33.14	1.88	4980	2350
Illinois.....	8	2,451	33.42	1.98	4980	2360
Indiana.....	2	837	33.49	2.01	4910	2310
Iowa.....	1	12,136	33.54	1.98	4920	2300
Kansas.....	3	8,504	33.26	1.99	4980	2310
Michigan.....	3	12,567	33.42	1.98	4980	2350
Nebraska.....	2	3,145	33.41	1.95	4980	2360
New Jersey.....	3	8,968	33.40	2.12	4900	2400
New York.....	7	6,465	33.48	2.06	4960	2360
Ohio.....	2	14,443	33.06	1.99	4910	2340
Pennsylvania.....	6	8,616	33.29	1.96	4940	2360
South Dakota.....	1	3,031	33.51	1.95	4920	2360
Vermont.....	All.	2,077	33.43	1.90	4960	2360
Washington.....	1	5,176	33.47	2.02	4920	2300
Wisconsin.....	2	7,695	32.86	1.97	4950	2365
Desert.....	9	6,109	33.38	1.99	4920	2360
Arizona.....	2	1,027	33.28	1.91	4980	2360
Nevada.....	1	1,441	33.75	2.06	4970	2330
New Mexico.....	2	1,857	33.26	1.84	4980	2400

TABLE VI.—Mean chest girth by groups and component sections, arranged on order of standing, with proportional chest circumference (expiration) for each inch of height and each pound of weight; also standard deviation for each chest measurement; first million draft recruits—(continued).

Group and section.	Group and section No.	Description.	Number of men measured.	Mean height.		Standard deviation (height).	Mean weight.		Mean chest.	
				Inches.	Pounds.		Inches.	Pounds.	Inches.	Pounds.
German and Austrian, 15 per cent plus.	23		126,865	33.33	2.06		4954			
Illinois.	1	German, 21.2 per cent; Austrians, 4.3 per cent.	6,328	33.38	2.02		4950		2330	
Do.	4	German, 17.4 per cent; Austrians, 2.8 per cent.	4,226	33.45	2.08		4940		2330	
Indiana.	1	German, 17.4 per cent; Austrians, 4.1 per cent.	3,614	33.45	2.13		4970		2330	
Iowa.	1	German, 15.9 per cent; Austrians, 1.9 per cent.	12,136	33.64	1.95		4920		2300	
Minnesota.	2	German, 22.3 per cent; Austrians, 2.9 per cent.	7,601	33.66	1.93		4970		2300	
Nebraska.	1	German, 12.2 per cent; Austrians, 3.9 per cent.	7,020	33.41	1.95		4980		2300	
Do.	2	German, 13.5 per cent.	3,145	33.41	1.95		4980		2300	
New Jersey.	1	German, 18.9 per cent.	17,785	33.19	2.12		4970		2300	
Ohio.	1	German, 18.9 per cent.	17,306	33.20	2.06		4950		2340	
Pennsylvania.	3	German, 18.9 per cent.	7,306	33.22	2.10		4970		2370	
Do.	5	German, 18.9 per cent.	8,822	33.15	1.96		4950		2350	
Do.	7	German, 10.7 per cent; Austrians, 6.5 per cent.	8,616	33.01	2.08		4950		2360	
Wisconsin.	1	German, 13.6 per cent; Austrians, 3.2 per cent.	3,297	33.66	1.80		4940		2320	
Do.	2	German, 20.3 per cent; Austrians, 1.8 per cent.	7,685	33.56	1.97		4950		2320	
Do.	4	German, 27.1 per cent; Austrians, 4.3 per cent.	2,895	33.73	2.01		5000		2340	
Mountain.	11		17,811	33.33	1.96		4921		2330	
Arkansas.	2	foreign parentage, 30.7 per cent; foreign	1,559	33.29	1.90		4940		2360	
Massachusetts.	1	foreign parentage, 30.7 per cent; foreign	1,373	32.90	2.06		4920		2370	
Missouri.	3	foreign parentage, 3.9 per cent; foreign	1,130	33.30	1.76		4950		2340	
Montana.	1	foreign parentage, 31.4 per cent; foreign	5,117	33.31	1.93		4910		2300	
New Hampshire.	1	foreign parentage, 21.6 per cent; foreign	665	33.72	2.06		5010		2380	
New York.	5	foreign parentage, 20 per cent; foreign	796	33.17	2.01		4930		2390	
Do.	8	foreign parentage, 24.7 per cent; foreign	2,060	33.34	2.00		4970		2370	
Washington.	3	foreign parentage, 20.8 per cent; foreign	1,559	33.62	1.93		4980		2300	
Wyoming.	1	foreign parentage, 22.3 per cent; foreign	1,927	33.36	1.99		4920		2310	
Commuters.	6		28,960	33.33	2.06		4970		2390	
Illinois.	1	foreign parentage, 38.2 per cent; foreign	6,303	33.36	2.02		4950		2330	
New Jersey.	1	foreign parentage, 37.5 per cent; foreign	17,795	33.19	2.12		4970		2300	
New York.	1	foreign parentage, 27.6 per cent; foreign	4,934	33.16	2.06		4970		2340	

CHEST CIRCUMFERENCE—SECTIONS.

Mining.....	7	35,691	32.23	1.97	4929	234
Alabama.....	1	8,841	72.08	1.84	4840	2330
California.....	2	943	33.81	1.87	4980	2310
Colorado.....	1	1,056	33.32	1.77	4980	2350
Do.....	3	331	33.21	1.86	4870	2330
Do.....	6	1,223	32.70	1.89	4840	2350
Idaho.....	1	4,081	33.74	2.04	4680	2320
Montana.....	1	5,117	33.31	1.93	4910	2290
Nevada.....	1	1,441	33.75	2.06	4970	2330
Pennsylvania.....	3	7,306	33.32	2.10	5000	2370
Do.....	4	4,827	33.15	2.00	4940	2350
Utah.....	3	563	33.44	1.77	4940	2320
Mexican, sparsely settled.....	14	11,004	33.22	1.90	4874	235
Arizona.....	1	1,027	33.28	1.91	4880	2320
Do.....	2	2,823	33.26	1.90	4870	2320
New Mexico.....	3	82,63	33.63	1.83	4900	2340
Texas.....	1	6,676	33.34	1.98	4876	2340
Eastern manufacturing.....	5	81,568	33.20	2.08	4970	2380
Connecticut.....	2	8,708	33.34	2.20	4900	2390
Massachusetts.....	2	18,447	33.15	2.04	4970	2410
New Hampshire.....	2	1,575	32.96	2.00	4980	2370
New Jersey.....	1	17,796	33.19	2.12	4970	2380
New York.....	3	5,160	33.32	2.07	4980	2390
Ohio.....	1	17,208	33.20	2.08	4950	2340
Pennsylvania.....	5	8,802	33.15	1.96	4970	2350
Rhode Island.....	1	3,923	33.53	2.11	4940	2410

TABLE VI.—Mean chest girth by groups and component sections, arranged on order of standing, with proportional chest circumference (expiration) for each inch of height and each pound of weight; also standard deviation for each chest measurement; first million draft recruits—Continued.

Group and section.	Group and section No.	Description.	Number of men measured.	Mean height. Inches.	Standard deviation (height). Inches.	Mean weight. Pounds.	Standard deviation (weight). Pounds.	Mean chest. Inch.
Mountain white.	12		12, 154	33.20	1.87	150.2	1.87	20.27
Kentucky.	1	0.7 per cent; foreign	4, 083	33.16	1.80	150.0	1.80	20.70
North Carolina.	1	0.5 per cent; foreign	2, 738	33.64	1.62	150.0	1.62	20.80
South Carolina.	1	0.4 per cent; foreign	1, 664	32.97	1.63	149.0	1.63	20.50
Tennessee.	3	1.1 per cent; foreign	5, 900	32.93	1.65	149.0	1.65	20.50
Virginia.	4	per cent; foreign born.	3, 512	33.34	1.67	149.0	1.67	20.80
West Virginia.	1	3.7 per cent; foreign	1, 507	33.20	1.67	149.0	1.67	20.60
Agricultural, Negroes, 45 per cent plus.	1		49, 465	33.19	1.91	150.4	1.91	20.40
Alabama.	2	Negro, 70.8 per cent; white, 29.2 per cent.	3, 227	33.27	1.90	149.0	1.90	20.30
Do.	4	Negro, 72.4 per cent; white, 27.6 per cent.	6, 600	33.16	1.84	149.0	1.84	20.20
Arkansas.	1	Negro, 55.3 per cent; white, 44.7 per cent.	4, 945	33.18	1.95	149.0	1.95	20.30
Georgia.	2	Negro, 61 per cent; white, 39 per cent.	10, 070	33.33	1.91	149.0	1.91	20.60
Louisiana.	1	Negro, 63 per cent; white, 37 per cent.	4, 074	33.29	1.97	149.0	1.97	20.60
Mississippi.	1	Negro, 71.5 per cent; white, 28.5 per cent.	5, 149	33.24	1.88	149.0	1.88	20.10
North Carolina.	4	Negro, 47.3 per cent; white, 52.7 per cent.	4, 570	33.15	1.91	149.0	1.91	20.30
South Carolina.	2	Negro, 50.9 per cent; white, 49.1 per cent.	3, 975	33.20	1.85	149.0	1.85	20.40
Do.	3	Negro, 62.3 per cent; white, 37.7 per cent.	3, 804	33.05	1.85	149.0	1.85	20.40
Tennessee.	1	Negro, 44.2 per cent; white, 55.8 per cent.	2, 218	32.90	1.84	149.0	1.84	20.20
Texas.	5	Negro, 51.1 per cent; white, 48.9 per cent.	1, 346	33.35	2.05	149.0	2.05	20.10
Virginia.	2	Negro, 48.6 per cent; white, 51.4 per cent.	5, 352	33.07	1.89	149.0	1.89	20.60
Agricultural, native white, North; native white over 75 per cent North.	1		66, 838	33.13	1.90	150.0	1.90	20.40
Illinois.	3	83.2 per cent; foreign parentage, 16.8 per cent; per cent.	8, 928	33.07	1.94	149.0	1.94	20.20
Indiana.	3	82.5 per cent; foreign parentage, 17.5 per cent; per cent.	18, 743	33.06	2.00	149.0	2.00	20.30
Iowa.	2	73.1 per cent; foreign parentage, 26.9 per cent; per cent.	7, 401	33.20	1.92	149.0	1.92	20.10
Ohio.	3	78.7 per cent; foreign parentage, 21.3 per cent; per cent.	17, 606	33.13	2.00	149.0	2.00	20.40
Pennsylvania.	2	70.5 per cent; foreign parentage, 29.5 per cent; per cent.	14, 218	33.18	2.02	149.0	2.02	20.70
Indiana, sparsely settled.	13	foreign born, white, 7.9 per cent.	10, 038	33.13	1.89	149.0	1.89	20.40
Arizona.	1	Indians, Chinese, and Japanese, 86.6 per cent; Mexicans, 8.4 per cent.	1, 027	33.26	1.91	149.0	1.91	20.20
New Mexico.	1	Indians, 28.1 per cent; native parentage, white, 61.9 per cent.	283	33.25	1.81	149.0	1.81	20.60
South Dakota.	3	Indians, 87.2 per cent; native parentage, white, 8.1 per cent.	247	33.74	1.74	149.0	1.74	20.80
Oklahoma.	1	Indians, 9.2 per cent; native parentage, white, 75.8 per cent; Negro, 13.7 per cent.	8, 471	33.00	1.87	149.0	1.87	20.30

CHEST CIRCUMFERENCE—SECTIONS.

441

	19		25, 757	33.11	2.07	4906	2397
French-Canadians, 10 per cent.							
Maine.....	3	foreign	1,247	33.22	1.93	4950	2390
Massachusetts.....	2	foreign	18,447	32.15	2.04	4970	2410
New Hampshire.....	1	foreign	665	32.72	2.06	4910	2380
Do.....	2	foreign	1,575	32.96	2.00	4900	2370
Rhode Island.....	1	foreign	3,928	32.83	2.11	4940	2410
Agricultural, native white, South.	3		117,890	33.09	1.91	4954	2340
Alabama.....	3	white, 67.6 per cent; Negro, 31 per cent.	2,670	32.07	1.80	4910	2390
Arkansas.....	2	white, 93.9 per cent; Negro, 0.7 per cent.	1,559	33.29	1.90	4940	2380
Do.....	2	white, 83.9 per cent; Negro, 12 per cent.	3,007	33.10	1.78	4950	2350
Kentucky.....	2	white, 76.3 per cent; Negro, 14.4 per cent.	11,109	32.91	1.91	4940	2340
Do.....	2	white, 71 per cent; Negro, 34.4 per cent.	5,233	32.17	1.87	4900	2360
Louisiana.....	2	white, 71 per cent; Negro, 13.8 per cent.	2,393	32.39	1.99	4900	2370
Maryland.....	2	white, 73.4 per cent; Negro, 33.4 per cent.	2,394	32.65	1.96	4900	2310
Mississippi.....	2	white, 64.3 per cent; Negro, 32 per cent.	13,186	32.50	1.90	4900	2340
Missouri.....	1	white, 64.4 per cent; Negro, 32 per cent.	1,139	32.50	1.76	4950	2340
Do.....	2	white, 64.4 per cent; Negro, 32 per cent.	1,399	32.50	1.90	4950	2340
North Carolina.....	2	white, 60.8 per cent; Negro, 39.2 per cent.	2,083	32.16	1.82	4950	2340
Do.....	2	white, 57.1 per cent; Negro, 41.2 per cent.	2,744	32.09	1.85	4950	2340
Oklahoma.....	2	white, 72.6 per cent; Negro, 27.4 per cent.	8,471	32.09	1.87	4950	2330
Do.....	2	white, 62.1 per cent; Negro, 37.9 per cent.	10,958	32.22	1.95	4950	2330
Tennessee.....	2	white, 74.4 per cent; Negro, 25.6 per cent.	8,308	32.92	1.95	4940	2310
Texas.....	2	white, 77.8 per cent; Negro, 15.9 per cent.	22,372	32.90	1.96	4900	2310
Virginia.....	3	white, 62.8 per cent; Negro, 26.8 per cent.	3,772	32.16	1.89	4970	2370
Do.....	2	white, 64.3 per cent; Negro, 33.2 per cent.	3,866	32.36	1.94	4900	2370
West Virginia.....	2	white, 64.3 per cent; Negro, 33.2 per cent.	10,460	32.31	1.93	4910	2350
Maritime.....	10		6,157	33.09	2.04	4933	2350
Maine.....	2	7.8 per cent; foreign	828	33.64	1.95	4970	2370
Maryland.....	2	1.7 per cent; foreign	1,068	33.00	1.88	4900	2360
Do.....	4	1.3 per cent; foreign		(1)			
Massachusetts.....	3	25.2 per cent; foreign	1,127	32.88	2.12	4910	2370
North Carolina.....	5	0.9 per cent; foreign	264	33.24	1.94	4910	2350
Virginia.....	1	3.6 per cent; foreign	2,886	32.64	2.05	4970	2390
Native whites of Scotch origin.....	15		13,473	32.95	1.90	4944	2347
Kentucky.....	2	Native born, 6.9 per cent; foreign	11,469	32.90	1.91	4940	2340
North Carolina.....	3	Native born, 0.4 per cent; foreign	2,033	33.16	1.92	4950	2340

1 Not tabulated.

TABLE VIII.—Correlation between height and chest circumference (expiration): Group 1, agricultural, North, native white, 73 per cent plus, first million draft recruits.

Height, in inches.		Chest, in inches.									
Total.	29	30	31	32	33	34	35	36	37	38	39
59.....	247	6	24	58	56	40	27	13	6	5	2
60.....	176	10	23	50	35	19	14	9	1	1	3
61.....	420	21	33	85	72	35	28	19	8	10	3
62.....	880	60	138	213	165	102	49	28	17	17	14
63.....	2,961	105	215	431	316	229	142	60	41	22	25
64.....	5,981	172	334	751	630	473	257	126	65	32	41
65.....	8,998	243	538	1,279	1,188	778	502	253	105	74	58
66.....	8,478	248	658	1,719	1,819	1,047	742	387	157	108	88
67.....	10,107	298	878	2,118	2,819	1,647	996	481	248	135	133
68.....	10,437	370	1,058	2,044	2,945	1,793	1,173	641	289	106	145
69.....	8,823	114	353	2,041	2,945	1,793	1,173	641	289	106	145
70.....	6,688	61	220	1,381	1,869	1,097	689	373	192	77	108
71.....	4,280	24	131	620	1,381	880	517	267	102	65	40
72.....	2,455	12	56	337	490	460	443	276	102	31	15
73.....	1,404	5	21	123	182	232	214	143	68	14	15
74.....	470	2	24	45	96	108	82	53	25	5	4
75.....	152	3	4	23	32	40	21	6	5	3
76.....	79	5	7	15	11	9	17	1	1	3
77.....	22	3	2	3	4	4	1	1	3
78.....	15	4	4	5	1	1	1	1	1
79.....	27	2	2	4	9	1	4	4	1	1
Total.....	66,795	1,451	3,944	8,483	12,715	13,752	11,371	7,455	4,025	1,870	842

Number of cases: 66,795. Height: Mean, 67.59 inches; standard deviation, 2.63±0.003 inch. Chest circumference (expiration): Mean, 33.12 inches; standard deviation, 1.99±0.002 inch.

TABLE XV.—Correlation between weight and chest circumference (expiration): Group 3, agricultural, native white, South, first million draft recruits.

Chest circumference, In inches.	Total.	Weight, in pounds.																							
		95- 99	100- 104	105- 109	110- 114	115- 119	120- 124	125- 129	130- 134	135- 139	140- 144	145- 149	150- 154	155- 159	160- 164	165- 169	170- 174	175- 179	180- 184	185- 189	190- 194	195- 199	200- 204		
29.....	2,384	13	96	261	447	527	406	294	147	69	44	26	15	7	4	6	4	3	2	3	1	10	1		
30.....	6,744	4	71	257	748	1,211	1,387	1,226	781	543	253	111	63	30	22	16	8	5	2	1	3	5	3		
31.....	14,628	2	46	171	674	1,598	2,393	2,792	2,596	1,969	1,208	647	277	130	66	32	18	12	7	1	2	5	2		
32.....	22,423	2	22	78	396	1,094	2,215	3,444	4,215	3,846	2,935	1,968	1,161	528	273	180	64	25	12	7	4	2	2		
33.....	25,133	9	38	148	474	1,399	2,359	3,633	4,333	3,069	3,497	2,299	1,391	749	401	182	79	24	24	10	7	9		
34.....	20,839	5	7	36	163	435	1,018	1,926	2,863	3,301	3,327	2,767	1,964	1,368	839	422	213	94	53	22	9	4		
35.....	13,231	6	8	8	31	113	275	652	1,203	1,652	1,949	2,060	1,792	1,305	978	580	341	197	99	43	19	19		
36.....	6,901	3	5	12	17	29	66	156	306	525	723	901	942	908	729	588	405	235	171	96	40	34		
37.....	2,866	3	4	3	11	6	18	25	65	128	178	248	321	333	360	303	270	185	160	102	69	74		
38.....	1,305	5	5	4	5	7	8	14	18	17	27	54	78	117	127	137	129	129	106	91	93	134		
39.....	512	2	8	6	7	4	5	6	8	8	14	4	19	24	30	44	46	41	43	40	44	108		
40.....	483	1	14	5	5	6	2	6	5	5	8	5	4	4	9	12	23	15	26	27	32	265		
• Total.....	117,449	21	269	856	2,487	5,137	8,390	11,507	14,159	15,231	14,165	12,365	9,854	7,206	5,172	3,657	2,362	1,551	941	691	439	333	646		

Number of cases: 117,449. Weight: Mean, 141.44 pounds; standard deviation, 16.94±0.02 pound. Chest circumference (expiration): Mean, 33.09 inches; standard deviation, 1.93±0.0027 inch. Correlation: 0.5662±0.0011.

TABLE X.—Correlation between height and weight: Group 2, agricultural, mixed foreign and native white, North, first million draft recruits.

Height, in inches.	Weight, in pounds.																		
	93-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189
Total.	319	293	263	233	203	173	143	113	83	53	23	13	3	1	1	1	1	1	1
59.....	1	4	7	20	18	20	43	36	27	39	27	17	15	12	12	6	2	3	2
60.....	1	10	23	37	29	31	36	29	15	22	13	12	17	4	6	2	1	1	1
61.....	4	15	54	102	101	106	86	54	37	28	19	22	10	7	4	1	3	4	4
62.....	3	28	69	195	225	237	227	175	89	59	46	22	16	7	4	6	3	2	7
63.....	1	38	113	298	437	504	471	418	276	168	107	64	34	17	18	18	7	5	2
64.....	1	36	109	383	605	833	885	824	619	448	304	194	104	56	45	23	19	12	1
65.....	1	27	97	352	752	1,055	1,306	1,320	1,180	911	611	394	247	151	99	55	41	25	13
66.....	1	11	97	253	721	1,263	1,705	1,846	1,848	1,431	1,071	768	428	307	203	91	76	44	25
67.....	1	15	27	152	507	1,004	1,633	1,947	2,340	2,019	1,565	1,555	1,076	699	476	288	189	99	85
68.....	1	13	22	96	308	697	1,225	1,731	2,185	2,154	2,006	1,521	1,076	699	476	288	189	99	85
69.....	1	4	8	32	118	365	752	1,205	1,703	1,744	1,798	1,498	1,170	830	578	342	218	148	93
70.....	1	4	13	20	46	139	336	708	1,019	1,366	1,420	1,312	1,014	810	568	363	221	157	108
71.....	1	1	8	3	8	49	141	284	487	694	874	940	778	689	513	353	235	133	98
72.....	1	1	1	3	7	13	34	103	207	298	391	471	487	434	332	269	200	112	73
73.....	1	1	1	3	2	1	8	24	49	119	174	219	223	224	209	154	110	79	42
74.....	1	1	1	3	2	1	2	7	10	27	47	88	76	53	87	95	62	41	25
75.....	1	1	1	2	1	1	1	2	3	6	17	18	32	39	42	40	24	25	8
76.....	1	1	1	2	1	1	1	1	2	3	8	6	7	14	11	17	11	12	7
77.....	1	1	1	1	1	1	1	1	2	2	2	3	2	3	3	4	4	4	2
78.....	1	1	1	1	1	1	1	1	2	3	5	4	1	1	2	2	2	2	1
79.....	1	1	1	1	1	1	1	2	4	1	3	2	2	4	2	3	4	2	1
Total.....	97,340	12,207	650	1,951	3,889	6,384	8,893	10,718	12,104	11,542	10,508	8,728	6,455	4,916	3,548	2,332	1,556	982	655

Number of cases: 97,340. Height: Mean, 67.62 inches; standard deviation, 2.66±0.004 inch. Weight: Mean, 142.79 pounds; standard deviation, 17.28±0.026 pound.

TABLE XI.—Correlation between height and chest circumference (expiration): Group 2, agricultural, mixed foreign and native white, North, first million draft recruits.

Height in inches.	Chest, in inches.										Total.
	29	30	31	32	33	34	35	36	37	38	39
59	11	16	29	77	62	37	40	19	12	2	8
60	17	28	47	49	54	47	31	31	7	1	3
61	42	80	123	115	124	84	59	18	12	7	7
62	69	141	243	319	278	197	93	51	16	6	3
63	115	301	503	639	608	413	240	109	62	18	20
64	207	470	822	1,123	1,166	831	487	238	102	41	37
65	3,028	604	1,749	1,749	1,750	1,422	877	451	219	82	57
66	8,080	770	1,640	2,466	2,542	2,065	1,273	677	314	134	116
67	12,255	724	1,656	2,636	3,004	2,613	1,719	986	435	192	166
68	14,468	724	1,468	2,556	3,128	2,844	2,053	1,173	508	274	165
69	15,014	642	1,468	2,642	3,128	2,844	2,053	1,173	540	239	169
70	12,802	490	1,121	2,042	2,637	2,577	1,818	1,115	472	207	163
71	9,828	247	726	1,410	2,042	2,013	1,603	920	356	157	129
72	6,468	157	401	835	1,242	1,394	1,124	643	237	105	99
73	3,599	61	163	451	661	712	616	482	130	57	36
74	1,737	18	62	175	294	360	349	244	66	33	16
75	294	9	19	61	103	138	143	127	33	13	11
76	114	4	7	24	41	53	56	49	12	4	5
77	38	2	2	12	13	19	28	17	2	1	5
78	29	2	2	5	5	5	9	3	3	1	3
79	45	2	2	3	5	10	6	8	3	1	3
Total	1,693	4,006	10,247	16,749	19,807	17,829	12,632	7,363	3,541	1,576	1,205

Number of cases: 97,338. Height: Mean, 67.62 inches; standard deviation, 2.66±0.004 inch. Chest circumference (expiration): Mean, 33.37 inches; standard deviation, 2±0.003 inch.

TABLE XII.—Correlation between weight and chest circumference (expiration): Group 2, agricultural, mixed foreign and native white, North, first million draft recruits.

Chest circumference in inches.		Weight, in pounds.																								Total.
		95- 99	100- 104	105- 109	110- 114	115- 119	120- 124	125- 129	130- 134	135- 139	140- 144	145- 149	150- 154	155- 159	160- 164	165- 169	170- 174	175- 179	180- 184	185- 189	190- 194	195- 199	200- 204			
29	1,665	8	73	169	323	367	312	187	112	45	30	18	8	4	6	1	7	2	11	1			
30	4,697	4	54	201	539	878	908	586	586	370	181	65	34	10	7	9	8	2	2			
31	10,248	36	138	552	1,151	1,743	1,808	1,743	1,390	745	430	199	92	45	19	13	7	5	2			
32	16,750	16	77	300	901	1,702	1,861	1,743	1,390	745	430	199	92	45	19	13	7	3	2			
33	19,816	11	21	150	393	1,079	1,996	2,809	3,479	3,308	2,627	1,853	1,018	558	305	129	59	2	1			
34	17,827	1	6	13	35	127	404	915	1,637	2,387	2,908	2,654	2,453	1,725	1,064	658	245	140	11	1			
35	12,629	6	4	13	46	119	260	613	1,078	1,543	1,893	1,912	1,653	1,362	974	509	309	16	1			
36	7,366	7	6	8	10	34	67	153	325	543	817	1,003	907	999	809	620	434	36	21			
37	3,537	7	6	8	10	7	16	40	72	135	225	345	401	456	460	387	318	81	45			
38	1,574	9	7	5	2	4	8	6	17	30	45	88	92	154	182	199	171	109	58			
39	653	8	6	11	13	4	4	4	6	6	14	26	33	47	67	70	128	86	80			
40	537	8	9	12	10	15	5	9	3	6	6	8	15	12	8	23	40	59	93			
Total.....		97,319	13	237	657	1,951	3,898	6,337	8,979	10,753	12,044	11,533	10,504	8,740	6,449	4,917	3,545	2,342	1,500	978	659	441	352	330		

Number of cases: 97,319. Weight: Mean, 142.76 pounds; standard deviation, 17.27±0.03 pound. Chest circumference (expiration): Mean, 33.38 inches; standard deviation, 2.01±0.003 inch. Correlation: 0.6783±0.0012.

TABLE XIX.—Correlation between height and weight: Group 5, eastern manufacturing, first million draft recruits.

Height, in inches.	Total.	Weight, in pounds.																					
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204
59.....	395	2	13	18	25	30	31	53	46	39	35	15	31	15	11	15	4	4	1	2	3	2
60.....	425	...	16	44	52	78	64	44	38	22	20	15	9	9	5	8	2	2	3	3	...
61.....	1,161	4	40	86	182	209	188	142	104	75	38	26	23	14	6	3	6	4	2	1	2	7	1
62.....	2,548	4	45	190	352	418	420	347	252	198	120	64	48	38	21	11	5	4	2	2	3	5	...
63.....	4,541	1	56	193	463	683	744	685	561	409	268	182	90	76	49	31	13	11	4	4	2	8	...
64.....	6,967	3	52	169	482	896	1,034	1,009	967	793	538	362	258	142	81	68	33	28	16	12	5	7	...
65.....	9,681	...	34	148	480	959	1,263	1,355	1,351	1,067	938	643	435	264	208	136	63	57	37	25	18	15	...
66.....	11,791	...	25	75	366	832	1,238	1,608	1,885	1,590	1,318	1,019	672	429	324	204	142	84	61	41	31	21	...
67.....	12,171	...	13	52	189	559	939	1,447	1,685	1,721	1,558	1,235	919	650	382	275	206	105	85	49	38	30	...
68.....	11,047	...	12	30	86	334	676	1,447	1,685	1,721	1,558	1,235	919	650	382	275	206	105	85	49	38	30	...
69.....	8,613	...	3	10	33	127	323	605	1,070	1,368	1,441	1,292	964	705	513	377	244	169	107	67	57	41	...
70.....	5,779	...	7	9	13	51	120	289	492	1,063	1,135	1,067	923	704	535	377	244	169	107	67	57	41	...
71.....	3,452	...	4	13	34	95	197	299	443	455	409	392	408	330	224	152	92	73	53	39	...
72.....	1,797	...	2	11	11	33	59	132	181	194	231	201	196	138	171	121	84	62	33	37	...
73.....	336	...	1	2	2	10	21	39	78	92	83	88	83	93	59	45	31	29	19	9	...
74.....	2	2	4	6	17	19	27	40	30	30	47	26	23	24	12	9	11	...
75.....	118	1	1	2	1	2	6	1	14	20	10	12	8	10	7	4	2	3	...
76.....	51	5	5	6	6	6	6	1	3	3	1	...
77.....	13
78.....	15
79.....	19
Total.....	81,718	14	325	1,028	2,736	5,188	7,090	8,810	9,782	9,816	8,926	7,436	5,838	4,293	3,159	2,346	1,584	1,078	703	504	375	312	371

Number of cases: 81,718. Height: Mean, 66.77 inches; standard deviation, 2.70±0.005 inch. Weight: Mean, 139.48 pounds; standard deviation, 17.71±0.030 pound.

TABLE XIV.—Correlation between height and chest circumference (expiration): Group 3, agricultural, native white, South, first million draft recruits.

	Height, in inches.		Chest, in inches.									
	Total.	29	30	31	32	33	34	35	36	37	38	39
59.....	300	7	27	31	60	75	45	33	11	6	2	3
60.....	235	9	28	40	52	41	34	15	11	1	3	2
61.....	521	40	65	126	92	94	60	26	10	5	3	3
62.....	1,002	70	147	215	216	160	106	41	30	7	3	7
63.....	2,322	141	308	448	540	413	276	114	51	16	7	8
64.....	4,442	258	496	863	1,013	800	513	288	113	55	23	20
65.....	8,267	346	849	1,454	1,896	1,619	1,065	599	248	91	37	33
66.....	12,598	364	1,052	2,044	2,704	2,719	1,864	1,021	486	248	71	72
67.....	16,290	378	1,036	2,416	3,457	3,542	2,756	1,516	668	273	131	87
68.....	18,760	335	1,036	2,368	3,762	4,153	3,283	2,030	1,003	307	183	137
69.....	17,518	217	1,079	1,957	3,291	3,881	3,396	2,119	1,077	430	213	168
70.....	14,216	129	462	1,308	2,408	3,238	2,864	1,958	1,074	425	201	149
71.....	9,873	57	265	762	1,501	2,128	2,149	1,494	849	381	167	120
72.....	6,030	34	132	393	828	1,258	1,678	950	583	279	133	88
73.....	3,116	10	38	168	376	666	1,352	535	352	176	71	56
74.....	1,366	2	15	48	155	248	304	248	180	90	41	35
75.....	588	3	5	16	73	100	126	100	87	38	24	16
76.....	248	1	1	5	25	43	62	43	34	21	7	7
77.....	77	1	1	1	10	11	16	15	9	6	3	3
78.....	60	1	1	1	9	18	14	8	4	4	1	2
79.....	61	1	1	1	9	13	13	10	3	2	1	2
Total.....	117,990	2,421	6,776	14,084	22,478	26,217	21,003	13,163	6,924	2,864	1,324	1,013

Number of cases: 117,990. Height: Mean, 66.18 inches; standard deviation, 2.64±0.004 inch. Chest circumference (expiration): Mean, 33.09 inches; standard deviation, 1.91±0.003 inch.

TABLE XV.—Correlation between weight and chest circumference (expiration): Group 3, agricultural, native white, South, first million draft recruits.

Chest circumference, in inches.	Weight, in pounds.																		
	95- 99	100- 104	105- 109	110- 114	115- 119	120- 124	125- 129	130- 134	135- 139	140- 144	145- 149	150- 154	155- 159	160- 164	165- 169	170- 174	175- 179	180- 184	185- 189
Total.																			
28	2,394	13	96	447	527	405	294	147	69	44	26	15	7	4	6	4	3	2	3
29	6,744	4	71	748	1,211	1,387	1,226	781	543	253	111	63	30	22	16	8	5	2	1
30	14,626	2	46	674	1,598	2,383	2,792	2,596	1,969	1,208	647	277	130	65	32	18	12	7	2
31	22,423	2	22	396	1,094	2,215	3,444	4,215	3,846	2,935	1,968	1,161	528	273	130	64	25	12	4
32	25,133	...	9	38	474	1,390	2,359	3,533	4,333	4,080	3,497	2,299	1,391	749	401	182	79	24	7
33	20,839	...	5	7	163	113	1,018	1,926	2,866	3,301	3,327	2,767	1,904	1,368	839	422	213	94	53
34	13,231	...	6	8	31	113	275	653	1,203	1,652	1,849	2,060	1,792	1,305	978	580	341	197	99
35	6,901	...	3	5	17	29	66	156	306	525	733	901	942	908	729	588	406	235	171
36	2,896	...	3	4	3	11	6	25	65	128	178	248	321	333	360	303	270	185	102
37	1,305	...	5	4	5	7	8	14	18	17	27	54	78	117	127	137	129	129	105
38	512	...	2	6	7	4	5	7	8	8	14	8	19	24	30	44	46	41	43
39	1	5	9	6	2	6	5	5	8	5	4	4	9	12	23	15	28
40	483
* Total	117,449	21	269	856	2,457	5,137	8,390	11,507	14,189	15,281	12,385	9,854	7,206	5,172	3,657	2,362	1,551	941	691

Number of cases: 117,449. Weight: Mean, 141.44 pounds; standard deviation, 16.84±0.02 pound. Chest circumference (expiration): Mean, 33.09 inches; standard deviation, 1.93±0.027 inch. Correlation: 0.6663±0.0011.

TABLE XVI.—Correlation between height and weight: Group 4, agricultural, Negro, 45 per cent plus, first million draft recruits.

Height, in inches.	Total.	Weight, in pounds.																					
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204
59	166	1	1	2	9	16	11	14	21	17	12	17	22	9	4	5	3	1	1				1
60	132	3	16	18	20	27	10	14	10	10	9	6	6	7	4	2	2	1					
61	284	3	16	37	45	48	30	22	26	19	21	6	6	4	5	2	2	1					
62	632	2	7	37	81	114	93	67	47	26	26	14	11	8	5	2	2	3	2	2	7	1	4
63	1,311		27	42	129	207	229	206	175	282	166	122	159	37	26	17	6	7	4	4	2	1	4
64	2,440		13	61	206	380	426	337	562	740	968	795	554	343	204	103	64	35	10	10	1	2	4
65	4,133		64	213	396	531	656	610	866	897	968	740	515	358	205	109	67	31	15	13	7	6	11
66	5,951	1	19	45	196	385	608	863	866	1,020	1,066	968	737	517	338	209	105	49	34	28	21	9	31
67	7,093		30	102	309	571	855	1,020	1,066	1,066	1,120	967	737	517	338	209	105	49	34	28	21	9	31
68	7,726		44	101	463	776	1,013	967	1,066	1,066	1,120	967	737	517	338	209	105	49	34	28	21	9	31
69	6,840		4	10	40	127	274	456	748	863	1,013	968	795	560	390	244	161	88	45	34	22	6	21
70	5,226		4	10	40	127	274	456	748	863	1,013	968	795	560	390	244	161	88	45	34	22	6	21
71	3,573		1	3	15	38	114	197	456	587	706	719	723	483	414	284	183	108	76	34	21	19	34
72	2,106		1	2	12	30	90	145	206	253	298	298	285	225	150	136	87	67	40	24	13	24	24
73	1,103		1	1	3	16	22	58	80	106	118	118	131	161	136	87	67	40	24	13	14	10	10
74	440		1	1	5	2	9	14	29	38	49	46	59	22	23	10	21	13	11	4	2	11	4
75	193				2	1	4	3	10	12	16	20	23	8	10	6	7	1	1	3	2	1	4
76	69				4			1	2	4	4	4	1	1	3	5	1	4	6	2	1	2	2
77	31				1	1	1	1	1	2	2	2	1	1	1	1	1	1	1	1	1	2	2
78	21																						
79	33																						
Total.....	49,503	13	125	303	1,119	2,144	3,377	4,833	5,618	6,109	6,077	5,287	4,444	3,128	2,386	1,670	1,023	647	436	251	145	106	183

Number of cases: 49,503. Height: Mean, 67.82 inches; standard deviation, 2.08±0.006 inch. Weight: Mean, 141.61 pounds; standard deviation 16.64±0.006 pound.

TABLE XVII.—Correlation between height and chest circumference (expiration): Group 4, agricultural, Negro, 45 per cent plus, first million draft recruits.

	Total.	Chest, in inches.																
		29	30	31	32	33	34	35	36	37	38	39						
59.	167	5	9	26	30	30	32	30	32	20	8	4	2	1				
60.	131	4	23	26	30	12	17	17	17	9	6	2	2	2				
61.	282	17	44	47	67	53	25	25	25	17	9	2	2	1				
62.	631	27	80	122	159	109	78	78	78	34	12	8	2	3				
63.	1,301	52	171	262	314	241	143	143	143	60	27	17	11	8				
64.	2,421	106	286	442	599	446	323	323	323	156	74	27	7	3				
65.	4,101	107	282	442	599	446	323	323	323	171	74	27	33	8				
66.	5,900	137	412	655	893	859	652	652	652	381	171	70	22	18				
67.	7,064	141	395	866	1,205	1,311	978	978	978	516	277	97	52	35				
68.	7,997	102	370	881	1,424	1,516	1,368	1,368	1,368	734	394	133	79	37				
69.	6,814	74	259	620	1,149	1,492	1,492	1,492	1,492	929	520	190	79	55				
70.	5,227	30	165	400	827	1,147	1,075	1,075	1,075	791	469	224	107	40				
71.	3,606	23	80	242	528	424	462	462	462	604	336	161	84	51				
72.	2,140	14	37	116	242	195	250	250	250	364	254	123	61	43				
73.	1,139	7	21	48	104	65	108	108	108	89	50	77	38	22				
74.	446	4	5	23	38	28	39	39	39	35	30	16	13	10				
75.	194	2	1	3	18	12	16	16	16	9	9	4	7	4				
76.	73	1	1	1	1	8	7	7	7	6	5	3	1	2				
77.	32	1	1	1	1	1	5	5	5	4	2	2	2	2				
78.	18	1	1	1	1	1	4	4	4	3	2	2	2	2				
79.	33	1	1	2	5	3	3	3	3	11	2	3	3	2				
Total.....	49,447	863	2,591	5,673	9,162	10,378	9,227	5,917	3,287	1,381	638	340						

Number of cases: 49,447. Height: Mean, 67.84 inches; standard deviation, 2.69±0.008 inch. Chest circumference (expiration): Mean, 33.20; standard deviation, 1.91±0.004 inch.

TABLE XVIII.—Correlation between weight and chest circumference (expiration): Group 4, agricultural, Negro, 45 per cent plus, first million draft recruits.

Chest circumference, in inches.	Total.	Weight, in pounds.																					
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204
29.....	863	4	34	89	159	177	136	111	59	32	27	8	6	3	2	5	2	1	1	2	5
30.....	2,647	3	39	116	325	457	487	445	317	207	105	59	35	20	10	5	4	1	1	1	2	10
31.....	5,721	1	25	82	338	637	913	1,078	974	661	498	259	132	47	34	16	12	7	5	5	1	1
32.....	9,212	1	18	42	219	519	968	1,464	1,527	1,556	1,988	778	461	240	117	55	20	15	14	7	3	3
33.....	10,384	5	11	60	231	589	1,069	1,487	1,775	1,729	1,336	853	544	299	162	80	30	30	40	22	12
34.....	9,185	5	4	16	77	210	492	860	1,177	1,515	1,422	1,260	948	592	380	167	83	40	22	12	2
35.....	5,853	1	3	3	18	51	141	258	508	669	903	969	744	614	455	243	129	70	42	21	7
36.....	3,266	4	6	4	10	14	31	74	165	267	365	465	450	435	329	251	185	97	49	16	17
37.....	1,360	3	1	1	11	17	25	70	103	123	158	196	184	139	98	88	64	35	18
38.....	180	7	3	3	1	1	3	6	3	7	25	29	42	73	63	85	70	62	60	34	26
39.....	4	3	1	2	2	2	1	2	1	7	3	4	12	15	13	14	16	15	11	14
40.....	164	4	5	6	2	2	2	4	2	2	2	5	3	1	7	7	12	10	9	9	12
Total.....	49,465	9	149	364	1,135	2,136	3,374	4,849	5,594	6,113	6,088	5,287	4,441	3,103	2,385	1,676	1,023	645	403	280	145	115	161

Number of cases: 49,465. Weight: Mean, 131.58 pounds; standard deviation, 16.64±0.04 pound. Chest circumference (expiration): Mean, 33.19 inches; standard deviation, 1.92±0.0041 inch. Correlation: 0.639±0.0018.

TABLE XIX.—Correlation between height and weight: Group 5, eastern manufacturing, first million draft recruits.

Total.	Weight, in pounds.																							Total.
	95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204		
395	2	13	18	25	30	31	53	46	39	35	15	31	15	11	15	4	4	1		2	3	2		
400		16	44	52	78	64	44	38	22	20	15	9	5	5	8	2	2				3			
425	4	40	86	182	209	188	142	104	75	86	26	23	14	6	3	6	4				7	1		
1,161	1	45	190	352	418	420	347	252	198	120	64	48	38	21	11	5	4	2	1	1	3	5		
2,548	4	56	193	463	633	744	665	561	409	298	182	90	76	41	31	13	11	4	4	4	5	8		
5,541	3	32	109	482	886	1,034	1,009	793	558	363	208	258	142	81	68	33	28	16	12	12	18	25		
6,967		34	148	440	969	1,253	1,365	1,351	1,254	838	643	435	254	208	126	63	57	37	25	18	15	28		
9,681		25	75	366	832	1,238	1,368	1,368	1,590	1,318	1,019	672	429	324	204	142	106	86	60	37	30	33		
11,791		13	52	186	529	939	1,447	1,685	1,721	1,558	1,235	919	650	382	275	244	165	90	66	40	30	39		
12,171		12	30	86	324	676	1,070	1,368	1,460	1,441	1,292	964	705	513	375	244	165	90	66	40	37	39		
14,048		3	10	33	127	323	605	943	1,083	1,135	1,077	682	521	408	330	244	169	107	67	57	41	50		
8,613		7	9	13	51	130	289	492	667	764	737	652	521	408	330	244	169	107	67	57	41	50		
5,779		3		6	13	34	95	197	299	443	455	409	392	286	235	171	121	94	62	33	39	56		
5,452		2		6	6	11	33	59	132	181	194	231	292	286	235	171	121	94	62	33	39	56		
1,797		1	2		1	2	10	21	33	78	92	83	88	83	83	59	45	31	29	19	11	8		
798		1	2		1	2	7	17	19	27	7	14	20	10	12	8	10	7	4	2	3	10		
74				1			2	6	2	1	1	5	5	6	6	6	6	0	1	3	3	1		
336					1			1	2	1	1	1	2	2	2	2	2	2	1	1	1	2		
118										3	2		3	1	1	1	1	1	1	1	1	2		
75										2	1		1	2	2	2	2	2	1	1	1	2		
51										2	2		1	1	1	1	1	1	1	1	1	2		
76										2	2		1	1	1	1	1	1	1	1	1	2		
13										2	2		1	1	1	1	1	1	1	1	1	2		
77										2	2		1	1	1	1	1	1	1	1	1	2		
78										2	2		1	1	1	1	1	1	1	1	1	2		
79										2	2		1	1	1	1	1	1	1	1	1	2		
81,718	14	325	1,028	2,735	5,188	7,090	8,810	9,782	9,816	8,926	7,436	5,838	4,293	3,156	2,346	1,584	1,078	703	508	375	312	371		

Number of cases: 81,718. Height: Mean, 66.77 inches; standard deviation, 2.70±0.005 inch. Weight: Mean, 139.48 pounds; standard deviation, 17.71±0.030 pound.

TABLE XX.—Correlation between height and chest circumference (expiration): Group 5, Eastern manufacturing, first million draft recruits.

Height, in inches.	Total.	Chest, in inches.										
		29	30	31	32	33	34	35	36	37	38	39
59	383	16	37	37	82	67	54	42	27	10	7	4
60	425	23	51	53	98	82	65	19	16	6	6	1
61	1,161	54	130	231	250	215	133	84	36	14	7	5
62	2,539	129	272	458	521	469	332	203	90	38	17	17
63	4,533	178	439	703	904	833	642	395	193	66	43	29
64	6,951	267	540	1,033	1,458	1,315	1,070	590	249	156	68	45
65	9,696	359	758	1,396	1,943	1,773	1,463	843	331	270	134	96
66	11,773	349	800	1,406	2,282	2,317	1,812	1,244	694	336	163	156
67	12,156	273	723	1,406	2,226	2,432	2,116	1,356	781	439	205	149
68	14,021	193	723	1,243	2,028	2,207	1,894	1,353	813	415	215	191
69	8,605	118	332	894	1,438	1,714	1,538	1,157	715	341	167	147
70	6,752	59	220	481	910	1,130	1,120	774	509	272	130	114
71	3,469	34	95	288	483	643	670	525	335	178	104	64
72	1,791	9	39	223	386	372	372	257	194	103	61	25
73	797	1	14	108	146	162	162	130	96	53	31	11
74	386	1	4	46	58	65	65	54	40	31	9	8
75	118	1	2	10	24	28	28	19	11	8	3	2
76	51	2	2	6	7	10	9	9	7	3	2	2
77	12	1	2	2	2	2	2	2	2	2	2	2
78	14	1	2	2	3	3	3	3	2	2	2	2
79	16	1	2	2	6	6	1	5	2	2	2	2
Total	81,569	2,047	5,018	10,234	14,920	15,799	13,552	9,209	5,438	2,725	1,371	1,255

Number of cases: 81,569. Height: Mean, 66.77 inches; standard deviation, 2.69 ± 0.004 inch. Chest circumference (expiration): Mean, 33.18 inches; standard deviation 2.08 ± 0.003 inch.

TABLE XXI.—Correlation between weight and chest circumference (expiration): Group 5, eastern manufacturing, first million draft recruits.

Chest circumference, in inches.	Total.	Weight, in pounds.																		195-199	200-204
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194
29	2,031	13	113	251	452	486	317	180	85	48	21	15	11	8	6	4	4	3	1	1	12
30	5,011	2	75	312	748	1,069	992	1,800	480	269	130	50	31	9	7	12	8	7	2	3	12
31	10,228	2	50	263	775	1,538	1,896	1,929	1,582	1,013	605	285	127	58	34	18	12	11	6	4	2
32	14,922	1	21	108	449	1,227	1,991	2,624	2,530	2,208	1,641	857	483	227	104	63	34	10	2	7	3
33	15,812	1	14	40	197	593	1,174	1,921	2,506	2,769	2,419	1,771	1,145	650	253	190	72	41	12	5	3
34	13,549	1	11	11	44	169	500	933	1,540	2,023	2,211	2,015	1,669	1,035	719	401	194	87	44	22	12
35	9,211	1	5	6	14	56	143	278	606	938	1,259	1,398	1,315	1,121	814	586	301	191	96	44	22
36	6,436	1	4	4	6	7	23	81	175	332	504	653	749	710	688	503	429	229	153	90	46
37	2,726	1	1	3	6	5	9	13	34	71	126	210	240	313	351	327	297	236	143	115	76
38	1,368	1	1	3	3	13	7	9	12	22	25	49	85	119	132	161	144	149	131	111	80
39	705	1	1	2	2	13	14	9	8	8	10	17	4	23	28	45	62	75	74	63	57
40	599	1	1	2	13	11	14	9	8	8	10	17	4	4	10	17	22	38	28	41	61
Total	81,598	18	293	1,006	2,714	5,173	7,069	8,784	9,868	9,713	8,962	7,368	5,833	4,277	3,176	2,327	1,579	1,077	693	516	373
																					305
																					474

Number of cases: 81,598. Weight: Mean, 139.37 pounds; standard deviation, 17.81±0.03 pound. Chest circumference (expiration): Mean, 33.20 inches; standard deviation, 2.11±0.003 inch. Correlation: 0.6977±0.0012.

TABLE XXIII.—Correlation between height and chest circumference (expiration): Group 6, commuter, first million draft recruits.

	Height, in inches.	Chest, in inches.										Total.
		29	30	31	32	33	34	35	36	37	38	39
59	159	8	17	18	27	28	26	15	10	4	4	2
60	164	11	24	26	35	31	30	8	3	3	2	1
61	164	15	46	71	89	80	62	32	14	9	2	2
62	164	40	96	145	164	156	126	79	41	9	6	6
63	167	70	167	237	338	296	218	127	70	21	17	9
64	170	80	172	384	452	413	370	206	125	76	21	14
65	172	109	279	466	608	657	508	358	186	84	45	33
66	177	116	278	565	780	778	626	448	260	115	62	55
67	183	121	278	503	729	841	714	507	276	146	71	43
68	183	121	228	503	716	784	690	505	318	174	80	69
69	183	121	228	503	716	784	690	505	318	174	80	69
70	183	121	228	503	716	784	690	505	318	174	80	69
71	183	121	228	503	716	784	690	505	318	174	80	69
72	183	121	228	503	716	784	690	505	318	174	80	69
73	183	121	228	503	716	784	690	505	318	174	80	69
74	183	121	228	503	716	784	690	505	318	174	80	69
75	183	121	228	503	716	784	690	505	318	174	80	69
76	183	121	228	503	716	784	690	505	318	174	80	69
77	183	121	228	503	716	784	690	505	318	174	80	69
78	183	121	228	503	716	784	690	505	318	174	80	69
79	183	121	228	503	716	784	690	505	318	174	80	69
Total	28,994	737	1,771	3,542	5,072	5,594	4,839	3,425	2,059	1,026	493	436

Number of cases: 28,994. Height: Mean, 66.87 inches; standard deviation, 2.74±0.008 inch. Chest circumference (expiration): Mean, 33.23 inches; standard deviation, 2.09±0.006 inch.

TABLE XXIV.—Correlation between weight and chest circumference (expiration): Group 6, commuter, first million draft recruits.

Chest circumference, in inches.	Total.	Weight, in pounds.																							
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204		
29	693	7	37	78	170	156	104	72	35	10	5	3	5	1	2	4	1	2							
30	1,772	1	31	111	254	379	352	275	188	97	33	17	11	7	4	4	1	1	1	1	2	3			
31	3,528	1	16	73	274	529	643	690	526	387	200	98	53	27	10	27	7	2	2	3					
32	5,073		12	51	149	418	714	825	952	780	540	309	150	93	38	57	26	9	2	1					
33	5,563		4	18	70	190	385	661	917	942	874	687	426	243	99	57	26	9	2	1					
34	4,840		2	9	17	59	179	341	551	780	806	686	515	350	258	160	63	35	12	13	10	4			
35	3,422			1	5	15	51	106	216	354	462	557	482	409	321	213	105	58	39	35	15	7			
36	2,056				2	2	8	33	65	117	185	244	308	261	262	211	163	89	35	34	15	12			
37	1,025				3	4	1	1	4	25	56	77	102	121	117	133	108	90	58	45	31	19			
38	492				3	3	3	1	2	8	30	30	22	51	52	53	48	54	54	30	31	17			
39	241				3	8	10	5	2	5	5	3	10	7	10	17	8	23	23	37	24	18			
40	245					11	10	10	5	5	5	11	2	2	7	8	8	18	15	18	20	20			
Total	28,980	9	102	354	958	1,773	2,450	3,005	3,475	3,579	3,176	2,702	2,086	1,572	1,180	894	553	363	242	196	137	106	128		

Number of cases: 28,980. Weight: Mean, 139.80 pounds; standard deviation, 17.69±0.050 pound. Chest circumference (expiration): Mean, 33.26 inches; standard deviation, 2.12±0.006 inch.

TABLE XXV.—Correlation between height and weight: Group 7, mining, first million draft recruits.

Height, in inches.	Total.	Weight, in pounds.																					
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204
59	123	1	4	3	3	8	9	13	5	11	11	19	13	12	2	4	2	2	1				1
60	121	4	4	5	17	13	20	8	10	15	7	6	2	5	4	3	1						
61	319	7	25	46	56	56	37	44	30	21	17	8	2	4	4	3	2	2	2				
62	615	1	15	23	82	84	101	94	85	51	29	13	5	5	2	2	2	1	1	1	2	2	
63	1,299	2	18	43	117	193	189	179	162	132	89	62	30	12	11	8	7	1	2	1	4	2	2
64	2,148	8	39	121	225	304	336	315	283	253	176	113	74	51	30	19	11	11	5	11	6	12	5
65	3,394	1	4	22	144	268	394	468	547	672	736	812	545	377	242	202	98	61	38	24	16	12	9
66	4,733	1	5	25	64	191	384	510	738	803	893	954	545	377	242	202	98	61	38	24	16	12	9
67	5,205	1	8	25	64	191	384	510	738	803	893	954	545	377	242	202	98	61	38	24	16	12	9
68	5,205	1	8	25	64	191	384	510	738	803	893	954	545	377	242	202	98	61	38	24	16	12	9
69	4,504	3	1	19	38	155	289	157	238	386	440	473	455	344	288	224	127	63	38	21	14	9	8
70	3,409	1	1	6	22	71	157	157	238	386	440	473	455	344	288	224	127	63	38	21	14	9	8
71	2,307	1	1	1	1	2	14	48	122	183	255	325	329	302	215	161	107	81	69	27	18	16	18
72	1,620	1	1	1	1	2	2	14	50	66	122	157	182	153	150	125	82	52	43	28	11	6	11
73	620	1	1	1	1	2	2	11	18	20	50	66	80	86	72	64	59	44	25	8	6	12	12
74	250	1	1	1	1	1	1	2	3	10	20	24	24	25	35	29	27	24	17	10	7	3	12
75	120	1	1	1	1	1	1	3	1	3	7	7	17	12	16	14	13	8	6	4	1	3	12
76	49	1	1	1	1	1	1	1	1	1	1	1	5	6	10	2	3	7	6	1	3	1	3
77	21	1	1	1	1	1	1	1	1	1	1	1	2	3	3	3	2	3	2	4	1	3	1
78	14	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
79	13	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Total.	35,730	5	88	265	762	1,466	2,367	3,229	4,034	4,369	4,380	3,824	3,226	2,398	1,722	1,334	768	528	371	193	131	89	130

Number of cases: 35,730. Height: Mean, 67.49 inches; standard deviation, 2.72±0.007 inch. Weight: Mean, 142.25 pounds; standard deviation, 16.86±0.045 pound.

TABLE XXVI.—Correlation between height and chest circumference (expiration): Group 7, mining, first million draft recruits.

Height, in inches.		Chest, in inches.										
Total.		29	30	31	32	33	34	35	36	37	38	39
50	130	4	9	17	20	23	27	16	8	4	1	1
51	130	8	13	16	22	25	30	18	12	5	2	1
52	319	22	45	64	56	64	38	20	13	4	2	2
53	614	30	57	115	138	112	80	48	25	17	7	4
54	1,256	71	108	212	274	241	167	108	47	42	17	10
55	2,143	82	191	357	451	400	314	191	88	74	31	19
56	3,375	104	269	517	661	664	531	326	176	99	36	41
57	4,722	143	305	633	899	984	776	517	276	150	61	49
58	5,259	194	390	697	1,035	1,097	892	611	361	180	81	52
59	5,196	238	419	764	1,096	1,085	956	669	389	164	94	59
60	4,502	43	157	260	542	669	712	524	342	144	62	45
61	3,403	33	100	269	312	467	516	308	238	72	40	26
62	2,303	22	51	148	150	254	284	202	129	36	12	8
63	1,253	7	17	32	66	113	138	123	72	21	8	2
64	620	5	7	25	35	51	45	39	24	8	2
65	248	6	6	12	14	20	32	20	9	3	2
66	120	3	3	5	8	9	4	4	1
67	48	1	1	1	2	4	2	2	1
68	20
69	14
70	11
Total	35,686	731	1,835	4,089	6,404	7,221	6,452	4,395	2,522	1,136	504	397

Number of cases: 35,686. Height, mean, 67.49 inches; standard deviation, 2.72±0.007 inch. Chest circumference (expiration): Mean, 33.26 inches; standard deviation, 1.96±0.006 inch.

TABLE XXVII.—(Correlation between weight and chest circumference (expiration): Group 7, mining, first million draft recruits.

Chest circumference, in inches.		Weight, in pounds.																				Total.	
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194		195-199
29		5	36	82	138	148	124	86	47	31	8	5	3	1	1	5	1					3	1
30		1	22	73	229	317	370	342	227	123	54	29	16	8	5	4	2				1		2
31		1	15	59	234	434	694	782	725	534	314	160	79	50	16	10	4	3			3		2
32				3	26	105	655	931	1,185	1,111	911	590	329	194	61	51	9	2	1				2
33				5	7	40	374	663	1,185	1,277	1,090	894	718	468	215	155	47	24	7			3	
34				1	4	8	411	708	1,024	1,412	1,048	1,048	663	533	429	278	136	70	35	7		2	
35				1	1	19	39	76	201	360	542	676	663	533	429	278	136	70	35	25		5	
36				1	1	3	12	17	47	113	203	299	315	354	324	253	186	120	59	24	14	20	
37				2	2		2	2	11	22	36	72	102	100	158	153	116	101	86	49	29	24	
38				2	2		2	2	4	2	3	15	30	35	46	52	54	47	31	28	20	26	
39				2	2		2	2	3	2	3	3	2	9	13	17	18	15	25	20	8	24	
40					2		1		3	2	6	3	3		4	4	3	7	11	17	11	12	
Total		35,691	6	85	255	782	1,481	2,394	3,225	4,026	4,349	4,370	3,824	3,223	2,418	1,700	1,330	798	527	366	198	130	91

Number of cases: 35,691. Weight: Mean, 142.28 pounds; standard deviation, 16.90±0.05 pound. Chest circumference (expiration): Mean, 33.23 inches; standard deviation, 1.97±0.01 inch. Correlation: 0.6754±0.0019.

Number of cases: 35,691. Weight: Mean, 142.28 pounds; standard deviation, 16.90±0.05 pound. Chest circumference (expiration): Mean, 33.23 inches; standard deviation, 1.97±0.01 inch. Correlation: 0.6764±0.0019.

TABLE XXIX.—Correlation between height and chest circumference (expiration): Group 8, sparsely settled, not more than 3 per square mile, first million draft recruits.

Height, in inches.	Total.	Chest, in inches.											
		29	30	31	32	33	34	35	36	37	38	39	
59	34	2	4	5	12	3	7	1	
60	25	3	3	10	2	4	6	1	1	
61	67	3	10	10	17	11	10	2	4	
62	185	10	19	33	53	38	10	14	4	
63	332	13	31	83	83	66	59	33	8	
64	719	14	62	106	161	141	127	81	34	13	1	
65	1,278	23	103	259	329	294	230	121	67	30	6	
66	2,183	24	117	317	417	338	250	121	67	30	13	2	
67	3,310	20	95	266	440	502	436	275	170	96	16	
68	4,558	26	73	223	473	563	516	377	217	103	31	
69	5,768	9	57	169	351	471	471	398	237	103	21	
70	7,845	9	35	108	244	360	398	343	194	92	36	
71	1,318	3	20	64	161	261	292	251	138	62	25	
72	1,733	2	27	184	116	184	156	104	45	18	
73	333	2	6	38	60	79	68	52	25	13	
74	163	2	11	23	40	38	23	20	3	
75	82	4	10	20	17	15	7	4	
76	25	1	3	2	6	1	2	2	
77	10	4	3	
78	7	1	
79	7	
Total	16,143	157	606	1,456	2,704	3,348	3,228	2,325	1,316	579	249	175	

Number of cases: 16,143. Height: Mean, 68.01 inches; standard deviation, 2.63 \pm 0.010 inch. Chest circumference (expiration): Mean, 33.52 inches; standard deviation, 1.90 \pm 0.007 inch.

TABLE XXX.—Correlation between weight and chest circumference (expiration): Group 8, sparsely settled, not more than 3 per square mile, first million draft recruits.

Chest circumference, in inches.	Weight, in pounds.																						Total.
	95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204	
29	1	5	13	28	39	27	20	8	5	1	13	1	3	2	2	1	1			1			
30	1	7	27	48	106	132	117	72	38	30	70	21	13	7	4	1	4			2			
31	1	2	17	57	128	241	277	283	197	122	278	140	59	38	14	3	7	1		2	1		
32	1	4	11	43	112	237	368	471	489	411	278	337	203	114	63	22	7	2		1			
33	1	4	11	43	112	237	368	464	604	546	491	463	351	209	161	67	38	15	3	2			
34	1	4	11	43	112	237	368	464	604	546	491	463	351	209	161	67	38	15	3	2			
35	1	4	11	43	112	237	368	464	604	546	491	463	351	209	161	67	38	15	3	2			
36	1	4	11	43	112	237	368	464	604	546	491	463	351	209	161	67	38	15	3	2			
37	1	4	11	43	112	237	368	464	604	546	491	463	351	209	161	67	38	15	3	2			
38	1	4	11	43	112	237	368	464	604	546	491	463	351	209	161	67	38	15	3	2			
39	1	4	11	43	112	237	368	464	604	546	491	463	351	209	161	67	38	15	3	2			
40	1	4	11	43	112	237	368	464	604	546	491	463	351	209	161	67	38	15	3	2			
41	1	4	11	43	112	237	368	464	604	546	491	463	351	209	161	67	38	15	3	2			
42	1	4	11	43	112	237	368	464	604	546	491	463	351	209	161	67	38	15	3	2			
43 and over	1	4	11	43	112	237	368	464	604	546	491	463	351	209	161	67	38	15	3	2			
Total	16, 151	2	18	72	204	463	848	1,268	1,708	1,934	2,006	1,868	1,587	1,268	904	725	436	310	189	103	80	59	101

Number of cases: 16,151. Weight: Mean, 144.88 pounds; standard deviation, 16.94±0.064 pound. Chest circumference (expiration): Mean, 33.53 inches; standard deviation, 1.92±0.007 inch.

TABLE XXXI.—Correlation between height and weight: Group 9, desert, first million draft recruits.

Height, in inches.	Total.	Weight, in pounds.																					
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204
59	22		1	1	5			2	6	1	1	2			2		1				1		1
60	12			2				1	1		2				1								
61	35		2	6	6	2		5	3														
62	91		7	14	13	7	12	10	6		2												
63	150		4	12	21	33	23	17	15		7												
64	302		2	5	27	38	54	38	45		4		1		1								
65	505		4	6	21	46	73	81	75	68	47	28	19	11	9	5	4	2	3	1	1		
66	716			9	17	36	90	109	115	111	74	58	36	26	20	16	8	7	5	2	1		
67	888			3	18	41	72	108	162	141	95	102	75	63	38	35	17	10	9	5	2	2	4
68	894			1	5	18	54	92	128	129	144	92	75	63	38	38	18	17	11	6	5	3	6
69	881					7	14	30	44	86	110	134	121	108	81	43	41	13	8	3	2	2	7
70	648					3	11	26	60	78	98	91	82	72	47	31	20	16	8	5	3	2	6
71	449					1	3	8	22	39	57	73	63	43	38	38	18	17	11	6	5	3	4
72	288						3	2	9	11	36	20	40	38	48	23	19	12	11	5	6	3	4
73	141								1	2	7	4	5	27	17	12	12	7	4	3	4	3	1
74	66									1				8	10	10	7	4	2	3	1	4	4
75	25													3	6	2	2	2	1	1	1	2	2
76	12													2	3	3	1	1					
77	2													1	1	2	2	1					
78	3															3	1	1					
79	2																1	1					
Total.....	6,121	19	56	133	238	423	569	722	764	738	638	520	406	282	216	126	92	63	34	27	13		42

Number of cases: 6,121. Height: Mean, 67.87 inches; standard deviation, 2.72 ± 0.017 inch. Weight: Mean, 142.08 pounds; standard deviation, 17.23 ± 0.106 pound.

TABLE XXXII.—Correlation between height and chest circumference (expiration): Group 9, desert, first million draft recruits.

Height, in inches.	Chest circumference, in inches.										
	29	30	31	32	33	34	35	36	37	38	39
59	22	1	2	4	10	3	3	1	1	1	
60	12	1	1	2	2	3	1	1	1		
61	34	1	9	8	1	1	1	2	2		
62	91	7	18	27	15	6	10	1	1		
63	159	11	21	37	40	23	15	4	3		2
64	302	20	51	72	87	51	26	7	7		
65	505	35	72	90	109	83	42	32	18		3
66	713	38	104	149	164	123	75	31	16		5
67	888	49	107	180	207	161	99	66	22		7
68	883	34	73	177	208	182	113	68	28		6
69	860	15	75	143	170	141	135	70	31		4
70	646	8	57	92	123	105	60	49	29		13
71	449	2	30	64	96	90	54	55	24		12
72	258	4	15	31	37	60	29	44	26		4
73	140	6	6	14	27	30	8	16	6		7
74	66	1	1	5	9	6	3	7	8		1
75	25	1	2	1	6	6	3	3	1		2
76	12	1	1	1	2	1	1	1	1		1
77	2	1	1	1	1	1	1	1	1		1
78	2	1	1	1	1	1	1	1	1		1
79	2	1	1	1	1	1	1	1	1		1
Total	6,110	265	644	1,097	1,279	1,098	784	449	221	94	80

Number of cases: 6,110. Height: Mean, 67.87 inches; standard deviation, 2.72±0.017 inch. Chest circumference (expiration): Mean, 33.37 inches; standard deviation, 1.98±0.012 inch.

TABLE XXXIII.—Correlation between weight and chest circumference (expiration): Group 9, desert, first million draft recruits.

Chest circumference, in inches.		Weight, in pounds.																						
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204	
Total.		99	1	4	10	17	18	19	14	10	4	14	22	1	1	1	1	1	1	1	1	1	1	
29		262	1	6	15	32	37	52	39	37	15	44	8	3	4	2	3	1	1	1	1	1	1	
30		645	5	18	40	65	125	118	112	64	49	22	15	4	4	2	1	1	1	1	1	1	1	
31		1,098	3	8	28	65	112	169	148	177	154	100	54	21	9	3	1	2	1	1	1	1	1	
32		1,278	1	4	13	37	72	142	206	244	181	132	106	56	27	26	7	10	3	1	1	1	1	
33		1,098	1	1	12	14	33	55	115	153	179	169	138	97	69	62	28	19	12	8	1	1	1	
34		796	1	1	1	8	24	40	78	109	117	108	103	66	62	33	25	24	18	4	2	2	2	
35		450	1	1	1	1	5	11	23	40	46	60	86	67	36	24	18	24	12	4	7	2	8	
36		221	1	1	1	1	1	2	2	9	14	22	34	31	34	18	24	22	12	4	7	2	8	
37		94	1	1	1	1	1	1	1	1	5	9	4	7	14	11	12	14	6	3	4	3	10	
38		43	1	1	1	1	1	1	1	1	1	1	1	4	7	14	11	3	7	4	4	3	6	
39		35	1	1	1	1	1	1	1	1	2	1	1	1	1	1	5	3	1	1	2	6	17	
40																								
Total.....		6,109	2	18	56	133	238	423	566	721	761	735	636	519	408	282	216	123	95	68	28	25	15	43

Number of cases: 6,109. Weight: Mean, 142.08 pounds; standard deviation, 17.26±0.105 pound. Chest circumference (expiration): Mean, 33.38 inches; standard deviation, 1.99±0.012 inch.

TABLE XXXIV.—Correlation between height and weight: Group 10, maritime, first million draft recruits.

Height, in inches.	Total.	Weight, in pounds.																					
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204
59.....	33			2	3	5	1	1	4	2	4	2	2	2	1	2	2	1	1		1		
60.....	28			4	2	5	1	1	1	1	6	1	2	1	2	1	2						
61.....	81	1	4	6	9	11	8	12	4	2	4	5	3	3	4	2	1				1		
62.....	63			4	3	13	19	24	36	39	19	10	13	9	8	12	5	1	1				
63.....	114	2	4	3	13	24	35	49	60	56	48	36	29	24	10	12	10	6	9	1			
64.....	243	1	3	13	24	41	53	60	64	103	98	92	82	51	29	16	10	6	9	1			
65.....	369		3	6	24	41	53	60	64	103	98	92	82	51	29	16	10	6	9	1			
66.....	621		2	6	30	64	103	98	92	82	51	29	16	10	6	9	1	1	2	1			
67.....	827			3	21	55	99	114	139	127	100	96	69	53	32	28	24	12	6	6			
68.....	940		1	6	12	39	77	106	143	157	117	119	85	62	38	26	24	12	6	6			
69.....	904			7	17	22	53	86	116	135	115	109	77	67	43	30	23	13	9	4			
70.....	594				1	10	25	60	73	91	124	109	77	67	43	30	23	13	9	4			
71.....	325					2	9	25	45	76	103	73	55	41	23	15	10	12	5	4			
72.....	205						1	8	17	33	44	49	44	28	30	15	10	12	5	4			
73.....	78							4	9	10	18	23	25	28	9	14	8	4	2	2			
74.....	38							1	2	2	7	8	13	9	6	4	3	3	3	1			
75.....	17																						
76.....	5																						
77.....	2																						
78.....	3																						
Total.....	6,161	4	17	58	147	307	508	617	755	797	747	607	454	387	251	173	115	90	57	32	21	14	23

Number of cases: 6,161. Height: Mean, 67.31 inches; standard deviation, 2.70±0.016 inch. Weight: Mean, 140.38 pounds; standard deviation, 16.86±0.103 pound.

TABLE XXXV.—Correlation between height and chest circumference (expiration): Group 10, maritime, first million draft recruits.

Height, in inches.	Total.	Chest circumference, in inches.											
		29	30	31	32	33	34	35	36	37	38	39	
54 and under.....	33	2	2	4	4	9	2	6	1	2		1	
59.....	30	3	3	6	5	5	3	3	2				
60.....	6	9	12	9	5	3	3	3	1			1	
61.....	13	15	22	21	16	14	12	13	7	1			
62.....	113	20	47	54	51	46	40	25	4				
63.....	244	27	62	82	82	70	76	40	5	1			
64.....	367	22	43	59	107	117	111	40	11	4			
65.....	620	28	62	107	145	185	145	61	27	9	3	2	
66.....	824	26	57	124	189	200	145	111	39	13	5	3	
67.....	941	26	47	116	198	209	139	111	47	23	6	4	
68.....	904	25	47	116	157	209	162	98	68	26	10	9	
69.....	752	18	34	69	132	139	123	87	59	21	12	19	
70.....	594	8	23	49	89	135	123	68	31	14	9	14	
71.....	324	2	10	24	52	73	70	39	31	20	3	3	
72.....	206	2	3	15	24	42	41	24	27	20	2	5	
73.....	78	1	1	2	15	6	4	12	15	7	2	4	
74.....	38	1	1	1	9	2	5	7	6	1			
75.....	17	1	1	2	2	2	1	2	1	1		1	
76.....	5	1	1	1	1	1	1	1	1	1			
77.....	2	1	1	1	1	1	1	1	1	1			
78.....	3	1	1	1	1	1	1	1	1	1			
Total.....	6,157	214	395	797	1,186	1,270	984	607	394	176	62	72	

Number of cases: 6,157. Height: Mean, 67.31 inches; standard deviation, 2.70±0.016 inch. Chest circumference (expiration): Mean, 33 inches; standard deviation, 2.03±0.012 inch.

TABLE XXXVI.—Correlation between weight and chest circumference (expiration): Group 10, maritime, first million draft recruits.

Chest circumference, in inches.	T total.	Weight, in pounds.																					
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204
29	213	3	10	25	38	43	32	24	16	7	8	5	2	2	2	2	1	2
30	396	1	5	10	41	66	100	74	43	27	13	35	6	7	4	4	1	
31	798	1	2	8	33	90	128	146	137	113	69	17	25	14	14	6	3	1	2	
32	1,185	2	19	56	154	168	210	189	104	58	71	31	16	10	6	
33	1,270	4	9	43	67	131	195	235	194	146	108	71	36	25	13	
34	965	2	1	2	25	54	95	138	147	157	130	92	56	39	25	
35	905	2	1	2	13	39	66	88	94	76	71	65	44	28	19	
36	395	2	2	2	2	14	19	44	51	60	47	46	44	28	17	
37	177	
38	61	
39	39	
40	32	
Total.....	6,156	5	17	57	145	303	507	616	755	799	744	607	473	344	257	170	116	90	58	33	20	14	26

Number of cases: 6,156. Weight: Mean, 140.43 pounds; standard deviation, 18.90±0.103 pound. Chest circumference (expiration): Mean, 33 inches; standard deviation, 2.04±0.012 inch.

TABLE XXXVIII.—Correlation between height and chest circumference (expiration): Group 11, mountain, first million draft recruits.

Height, in inches.	Total.	Chest circumference, in inches.										
		29	30	31	32	33	34	35	36	37	38	39
59.....	57	2	4	10	14	11	8	5	1	2		1
60.....	47	1	9	8	10	6	3	5	3	1		
61.....	120	6	16	23	19	26	18	6	5	1		
62.....	260	11	31	51	60	44	31	18	8	4	1	
63.....	476	13	50	87	108	87	73	24	24	4		
64.....	857	17	80	152	193	180	124	49	34	15	10	3
65.....	1,539	40	108	224	318	331	260	142	63	33	9	11
66.....	2,071	49	124	257	403	451	381	227	115	40	15	9
67.....	2,577	40	122	328	509	534	450	322	170	65	16	21
68.....	2,565	31	105	249	468	590	484	331	162	95	26	24
69.....	2,267	23	71	200	391	483	472	294	183	78	41	31
70.....	1,762	15	41	128	260	353	354	201	178	63	40	19
71.....	1,223	6	24	83	164	222	278	192	150	45	23	18
72.....	708	4	8	33	90	136	158	116	77	25	12	16
73.....	336	2	2	11	37	58	80	59	40	11	4	3
74.....	135	3	2	2	9	18	38	28	19	4		
75.....	59	1	1	1	4	8	11	12	11	2	3	3
76.....	19	1	1	1	2	2	3	4	2	2		2
77.....	10				1		4		3	1		1
78.....	8						2		1	2		
79.....	7					1	2		2	1		
Total.....	17,101	259	800	1,849	3,060	3,541	3,284	2,127	1,251	575	226	179

Number of cases: 17,101. Height: Mean, 67.72 inches; standard deviation, 2.89±0.010 inch. Chest circumference (expiration): Mean, 33.32 inches; standard deviation, 1.94±0.007 inch.

TABLE XXXIX.—*Correlation between weight and chest circumference (expiration): Group 11, mountain, first million draft recruits.*

Chest circumference, in inches.	Total.	Weight, in pounds.																					
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204
29	251	1	10	10	49	57	50	32	19	14	4	3	1	2	1	1	1	1	1	1	1	1	1
30	800	5	17	17	84	152	183	145	111	44	41	8	5	18	8	4	4	1	1	1	3	3	3
31	1,850	2	14	14	69	191	304	356	325	264	163	84	41	167	73	32	21	4	3	1	3	3	3
32	3,061	2	10	10	51	154	268	455	539	558	420	263	167	232	103	61	19	9	3	3	3	3	3
33	3,542	1	4	4	20	85	178	340	503	611	532	474	365	232	103	119	65	39	16	8	4	2	5
34	3,235	1	1	2	1	27	59	172	307	408	503	499	458	284	217	178	115	57	20	13	4	2	5
35	2,127	1	1	1	1	6	18	37	98	168	269	311	332	284	214	178	115	57	20	13	4	2	5
36	1,249	1	1	1	1	1	2	13	27	61	94	144	150	160	152	147	104	55	37	13	12	11	10
37	1,578	1	1	1	1	1	2	6	3	10	25	39	56	67	76	84	62	61	26	16	15	10	20
38	227	1	1	1	1	1	2	2	1	2	3	6	8	16	19	33	27	22	23	11	7	9	18
39	99	1	1	1	1	1	2	2	1	2	2	1	3	3	3	3	8	9	11	7	3	5	39
40	86	1	1	1	1	1	2	2	1	1	2	1	1	3	3	1	1	3	7	7	3	5	39
Total.....	17,103	1	23	00	282	674	1,067	1,563	1,937	2,141	2,055	1,852	1,583	1,221	825	657	407	263	180	89	64	45	94

Number of cases: 17,103. Weight: Mean, 142.97 pounds; standard deviation, 16.78±0.061 pound. Chest circumference (expiration): Mean, 33.33 inches; standard deviation, 1.96±0.007 inch.

TABLE XL.I.—Correlation between height and chest circumference (expiration): Group 12, mountain whites, first million draft recruits.

Height, in inches.		Total.	Chest circumference, in inches.										
			29	30	31	32	33	34	35	36	37	38	39
50	55	4	4	7	11	5	14	2	2	2	2	1	3
51	34	4	2	6	16	5	4	5					
52	60	1	6	10	13	13	8	5					
53	136	8	14	35	29	24	12	9	3				
54	241	18	30	74	65	52	34	14	4				
55	377	41	57	166	171	132	97	44	18	4			
56	1,389	95	128	228	328	288	187	112	46	18	2		
57	2,064	174	247	468	466	461	324	153	78	26	11	1	7
58	2,068	211	309	549	546	534	350	147	75	55	22	22	28
59	2,475	251	394	715	715	694	430	167	102	77	27	33	33
60	2,161	298	460	853	853	815	542	202	136	75	33	48	16
61	2,688	42	70	138	440	608	549	405	233	80	48	16	20
62	7,908	8	14	39	285	399	415	323	179	88	19	20	25
63	1,113	5	11	48	148	257	232	172	128	45	26	15	20
64	577	2	4	14	70	123	129	101	64	45	10	9	9
65	258	1	2	4	25	42	56	46	29	18	4	4	4
66	108		2	2	5	9	24	25	23	10	4		
67	44				7	13	6	6	11	1			
68	16					4	3	4	2	1			
69	5					2	2	1					
70	7						1	2	1	2			
Total	21,233	328	1,001	2,368	4,062	4,661	3,926	2,671	1,353	546	215	162	

Number of cases: 21,233. Height: Mean 68.29 inches; standard deviation, 2.57±0.008 inch. Chest circumference (expiration): Mean, 33.20 inches; standard deviation, 1.86±0.006 inch.

TABLE XLII.—Correlation between weight and chest circumference (expiration): Group 12, mountain whites, first million draft recruits.

Chest circumference, in inches.	Weight, in pounds.																			
	95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194
Total.	325	325	325	325	325	325	325	325	325	325	325	325	325	325	325	325	325	325	325	325
29	5	8	37	70	72	48	31	26	6	4	8	4	2	2	1	2	1	1	1	1
30	1	13	43	107	206	229	163	119	56	38	13	9	9	9	10	7	1	2	1	3
31	1	13	37	134	211	439	489	402	275	150	77	44	23	28	14	7	1	4	1	2
32	1	13	37	134	211	439	489	402	275	150	77	44	23	28	14	7	1	4	1	2
33	1	13	37	134	211	439	489	402	275	150	77	44	23	28	14	7	1	4	1	2
34	1	13	37	134	211	439	489	402	275	150	77	44	23	28	14	7	1	4	1	2
35	1	13	37	134	211	439	489	402	275	150	77	44	23	28	14	7	1	4	1	2
36	1	13	37	134	211	439	489	402	275	150	77	44	23	28	14	7	1	4	1	2
37	1	13	37	134	211	439	489	402	275	150	77	44	23	28	14	7	1	4	1	2
38	1	13	37	134	211	439	489	402	275	150	77	44	23	28	14	7	1	4	1	2
39	1	13	37	134	211	439	489	402	275	150	77	44	23	28	14	7	1	4	1	2
40	1	13	37	134	211	439	489	402	275	150	77	44	23	28	14	7	1	4	1	2
Total	21,254	21,254	21,254	21,254	21,254	21,254	21,254	21,254	21,254	21,254	21,254	21,254	21,254	21,254	21,254	21,254	21,254	21,254	21,254	21,254

Number of cases: 21,254. Weight: Mean, 140.28 pounds; standard deviation, 16.10±0.05 pound. Chest circumference (expiration): Mean, 33.20 inches; standard deviation, 1.87±0.01 in. Correlation: 0.6532±0.0077.

TABLE XIII.—Correlation between height and weight: Group 18, Indian, sparsely settled, first million draft recruits.

Total.	Height, in inches.	Weight, in pounds.																						
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204	
29	59		2								3	1	1	1	1	2	2							
40	60					1						2					1							
81	61			2	6	7	6	4				5	2				1				1	2		
82	62			4	14	21	9	13	12			5	3	3	3	1	1							
210	63	1	2	7	32	40	28	47	48	20	22	14	5	10	11	4	3		1					
396	64		5	7	28	60	82	50	47	46	42	20	10	20	17	6	5							
712	65		8	11	41	83	99	115	114	75	59	71	33	43	42	12	11							
1,103	66		7	7	31	71	146	167	169	167	113	83	49	67	67	36	14	6	2					
1,427	67		2	8	17	71	144	192	217	209	231	133	83	87	87	53	33	8	7					
1,629	68		4	4	11	46	100	158	260	255	227	192	112	112	112	56	32	12	5					
1,411	69		4	1	13	55	109	153	260	222	227	194	149	143	122	71	57	17	9					
1,225	70		4			7	27	46	119	160	189	184	154	122	77	58	33	17	9					
845	71		1			1	10	23	61	87	100	113	109	95	79	58	39	23	11					
482	72		2			1	4	14	4	34	60	67	67	59	56	39	33	14	10					
239	73									14	22	27	22	35	31	30	18	15	8					
104	74									2	5	7	17	15	16	10	6	5	5					
52	75									3	2	1	5	8	7	5	3	3	2					
16	76														2	4	2	2	2					
3	77																							
7	78																							
79	79																							
10,038	Total.	1	43	82	186	405	727	913	1,206	1,307	1,268	1,065	799	632	475	335	238	121	77	52	39	37	60	

Number of cases: 10,038. Height: Mean, 68.12 inches; standard deviation, 2.61 \pm 0.012 inch. Weight: Mean, 141.89 pounds; standard deviation, 16.91 \pm 0.080 pound.

TABLE XLIV.—Correlation between height and chest circumference (expiration): Group 13, Indian, sparsely settled, first million draft recruits.

Height, in inches.	Chest circumference, in inches.											Total.
	29	30	31	32	33	34	35	36	37	38	39	
59	1	2	3	8	11	4	1	1	1	1	1	29
60	1	1	1	4	2	1	1	1	1	1	1	11
61	5	10	18	9	5	1	1	1	1	1	1	40
62	3	9	16	22	13	16	10	13	1	1	1	92
63	9	18	36	63	39	28	28	27	1	1	1	210
64	37	77	77	95	66	89	45	42	3	3	3	396
65	50	153	153	156	151	173	10	13	8	4	1	711
66	24	50	254	254	244	246	86	60	19	7	3	1,103
67	22	80	155	306	317	306	133	82	24	16	10	1,427
68	28	73	207	334	357	306	198	94	21	18	13	1,620
69	35	73	206	269	318	273	178	97	26	21	11	1,411
70	11	53	157	207	259	264	174	81	35	17	8	1,224
71	7	38	126	134	196	158	129	81	18	11	10	845
72	3	18	56	69	89	111	82	51	6	5	4	482
73	3	8	36	26	43	27	41	28	7	3	1	239
74	1	4	13	10	19	14	12	17	2	1	1	104
75	1	1	2	6	8	5	8	9	2	1	1	51
76	1	1	1	2	4	1	1	1	1	1	1	16
77	1	1	1	2	2	1	1	1	1	1	1	3
78	1	1	1	2	1	1	1	1	1	1	1	7
79	1	1	1	2	1	1	1	1	1	1	1	5
Total	178	468	1,260	1,979	2,144	1,829	1,151	616	201	129	80	10,035

Number of cases: 10,035. Height: Mean, 68.12 inches; standard deviation, 2.61 ± 0.012 inch. Chest circumference (expiration): Mean, 33.13 inches; standard deviation, 1.87 ± 0.009 inch.

TABLE XLV.—(Correlation between weight and chest circumference (expiration): Group 13, Indian, sparsely settled, first million draft recruits.

Chest circumference, in inches.		Weight, in pounds.																							
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204		
29	178		12	19	37	36	28	16	13	9	1	1	4	1	1	1					1	2			
30	469		4	9	51	81	105	77	56	55	11	11	4	2	2	6	2								
31	1,260		11	10	55	129	207	243	198	177	97	69	23	1	5	6	2	2			1	1			
32	1,982	1	16	7	27	87	203	312	335	337	268	157	95	32	11	10	2	1	1						
33	2,146			5	14	51	141	161	324	377	365	297	189	103	59	36	2								
34	1,828					1	9	29	176	253	317	278	239	174	131	79	46	27	8	1					
35	1,150					1	1	15	43	88	157	178	157	173	132	88	60	17	13	6					
36	615							3	10	22	43	60	78	97	88	67	55	34	23	11	8	6	5		
37	201								1	2	6	12	11	23	24	21	21	11	16	10	6	7	12		
38	130									2	2	2	2	6	15	18	11	15	15	11	12	7			
39	37									2	2	2	2	5	4	3	4	2	4	5	1	5			
40	42			2						2	1				2	2	1	2	1	1	5	24			
Total	10,038	1	43	52	186	405	726	915	1,207	1,306	1,268	1,065	798	632	475	335	238	121	74	55	39	37	60		

Number of cases: 10,038. Weight: Mean, 141.89 pounds; standard deviation, 16.91±0.08 pound. Chest circumference (expiration): Mean, 33.13 inches; standard deviation, 1.80±0.01 inch. Correlation: 0.6775±0.0036.

TABLE XLVI.—Correlation between height and weight: Group 14, Mexican, sparsely settled, first million draft recruits.

Height, in inches.	Total.	Weight, in pounds.																			
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	154-159	155-160	165-169	170-174	175-179	184-189	185-189	190-194	195-204
59	30			2	4	2	1	4	3	3	3	1	3	2	1			1			
60	17			2		5		1													
61	48		4	5	6	5	7	7	5	2	2	3	1					1			
62	110		6	10	14	18	9	11	13	8	6	7	1								
63	241		7	17	28	39	43	29	20	23	3	4	4					1			
64	456		6	14	40	63	78	62	60	50	29	21	5								
65	799		3	15	71	78	111	124	103	93	70	40	30	17	18	1		1			
66	1,128		1	6	28	83	134	164	168	156	125	94	57	41	21	5	4	1			
67	1,473		4	13	32	81	144	183	235	225	165	154	85	57	32	14	11	10	1	1	
68	1,647			8	16	38	97	164	255	215	245	195	132	95	67	38	26	17	8	2	
69	1,402			1	5	21	48	84	131	185	208	196	172	109	93	54	40	27	9	5	
70	1,413			2	6	22	47	88	155	189	209	181	171	121	101	51	37	28	12	6	
71	889	1		3	5	11	6	21	53	89	107	110	122	94	101	53	29	24	12	7	
72	619			2	4			7	28	36	73	71	90	79	65	32	16	12	8	4	
73	283						1	1	5	12	29	21	52	33	37	41	24	14	7	5	
74	119				1			1		5	10	12	11	18	21	8	9	4	4	1	
75	61									2	3		6	5	10	7	3	4	5	2	
76	22									2		1		2	2	4	2	4	6	1	
77	7														1		1				
78	3															2					
Total	10,779	33	97	257	464	738	978	1,245	1,296	1,288	1,087	943	885	553	375	245	167	119	65	53	30

Number of cases: 10,779. Height: Mean, 68.16 inches; standard deviation, 2.89±0.012 inch. Weight: Mean, 142.14 pounds; standard deviation, 17.30±0.080 pound.

TABLE XLVII.—Correlation between height and chest circumference (expiration): Group 14, Mexican, sparsely settled, first million draft recruits.

Height, in inches.		Chest circumference, in inches.											
Total.		29	30	31	32	33	34	35	36	37	38	39	
59	28	1	3	6	9	3	3	1	1	1	
60	17	6	4	6	2	3	2	
61	48	10	5	5	14	12	1	1	2	1	1	
62	112	14	14	17	35	16	8	8	3	
63	246	11	25	44	59	54	28	11	14	5	5	
64	468	19	65	93	89	88	58	27	60	7	7	
65	816	33	61	144	189	159	107	100	40	14	13	
66	1,154	20	85	174	241	257	186	164	45	25	25	
67	1,511	46	107	193	309	327	227	223	81	32	32	
68	1,995	77	160	331	531	374	281	272	99	44	44	
69	1,568	23	59	150	267	304	240	212	109	42	42	
70	1,326	17	46	128	211	203	254	187	93	51	51	
71	819	8	17	70	132	149	188	125	59	44	44	
72	637	3	16	37	86	99	139	111	70	60	60	
73	385	2	5	23	55	55	84	52	37	12	11	
74	181	2	9	20	20	34	21	16	12	12	
75	61	1	1	2	3	16	13	1	1	1	
76	22	2	4	
77	7	
78	3	
79	
Total	11,061	221	586	1,271	2,029	2,373	1,011	1,327	719	347	159	122	

Number of cases: 11,061. Height: Mean, 68.16 inches; standard deviation, 2.63 ± 0.12 inch. Chest circumference (expiration): Mean, 33.22 inches; standard deviation, 1.97 ± 0.040 inch.

TABLE XLVIII.—Correlation between weight and chest circumference (expiration): Group 14, Mexican, sparsely settled, first million draft recruits.

Chest circumference, in inches.	Weight, in pounds.																						
	95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204	
Total																							
29																							
30																							
31																							
32																							
33																							
34																							
35																							
36																							
37																							
38																							
39																							
40																							
Total	10,785	37	93	246	461	740	979	1,247	1,293	1,286	1,098	946	680	556	372	245	166	120	64	53	30	73	

Number of cases: 10,785. Weight: Mean, 142.28 pounds; standard deviation, 17.54±0.08 pound. Chest circumference (expiration): Mean, 33.22 inches; standard deviation, 1.99±0.01 inch. Correlation: 0.6859±0.0084.

TABLE L.—Correlation between height and chest circumference (expiration): Group 15, native whites of Scotch origin, first million draft recruits.

	Height, in inches.										Chest circumference, in inches.									
	Total.																			
	29	30	31	32	33	34	35	36	37	38	39									
59	27	2	4	7	4	5	1	2												
60	42	3	4	11	7	7	1	1												
61	69	7	11	6	18	8	3	3												
62	140	13	15	27	28	14	5	5												
63	294	25	44	60	47	32	10	7												
64	584	37	73	153	93	53	32	12												
65	1,016	50	112	238	198	116	60	24												
66	1,485	35	116	362	337	201	104	48												
67	1,943	35	144	446	409	306	154	82												
68	2,144	36	122	491	453	379	196	111												
69	1,975	19	94	406	442	345	220	122												
70	1,550	18	43	253	379	310	195	116												
71	1,042	5	28	172	229	200	156	93												
72	624	3	11	109	141	130	88	50												
73	298		2	38	74	66	49	34												
74	136		3	21	22	30	22	17												
75	52		5	6	11	10	10	7												
76	27		1	2	1	7	4	3												
77	8				1	2	2	1												
78	1																			
79	12			3	4	2	2	1												
Total	13,469	303	826	1,855	2,811	2,904	2,213	1,315	730	270	129	113								

Number of cases; 13,469. Height: Mean, 68.01 inches; standard deviation, 2.63±0.011 inch. Chest circumference (expiration): Mean, 32.94 inches; standard deviation, 1.89±0.008 inch.

TABLE II.—Correlation between weight and chest circumference (expiration): Group 15, native whites of Scotch origin, first million draft recruits.

Chest circumference, in inches.	Weight, in pounds.																			
	95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194
29																				
30	298	3	13	45	56	66	48	11	6	24	16	1	3	1	1	1	1	1	2	
31	827	5	33	103	137	177	160	84	60	24	16	4	3	1	1	1	1	1		
32	1,855	1	4	18	79	231	349	301	286	129	74	34	14	5	3	1	1	1	1	
33	2,813			10	58	165	263	576	487	336	217	125	64	29	10	5	1	1	1	
34	2,909			4	14	49	159	299	440	467	423	251	156	70	34	11	9	3	1	
35	2,216			1	5	12	38	212	305	343	369	306	219	152	85	49	16	9	4	
36	1,730			1	2	3	8	56	113	158	185	196	168	146	110	67	40	23	10	
37	1,310			1	1	3	2	18	22	55	82	82	92	106	95	69	41	29	23	
38	270			1	1	2	2	2	3	8	18	21	23	36	25	37	32	20	20	
39	130			1	1	1	2	1	1	2	1	1	2	4	1	1	1	15	18	
40	53			1	1	1	1	2	1	1	1	1	2	1	1	1	1	7	2	
41	57																	2	2	
Total	13,473	4	24	115	319	687	1,078	1,436	1,748	1,520	1,389	1,024	750	548	381	232	158	108	80	39

Number of cases: 13,473. Weight: Mean, 140.36 pounds; standard deviation, 16.96±0.07 pound. Chest circumference (expiration): Mean, 32.95 inches; standard deviation, 1.90±0.01 inch. Correlation: 0.7099±0.0029.

TABLE LII.—Correlation between height and weight: Group 16, Russian, 10 per cent plus, first million draft recruits.

Height, In Inches.	Total.	Weight, in pounds.																				
		100- 99	105- 109	110- 114	115- 119	120- 124	125- 129	130- 134	135- 139	140- 144	145- 149	150- 154	155- 159	160- 164	165- 169	170- 174	175- 179	180- 184	185- 189	190- 194	195- 199	200- 204
50	51	3	1	3	5	2	8	2	4	6	4	5	2	2	3							
51	55	3	1	9	14	2	3	4	7	3	3	1	2					1				
52	136	2	13	11	32	8	23	9	13	3	3	1					1					
53	253	9	12	33	33	40	49	34	18	7	5	1										
54	497	10	18	43	74	78	78	71	52	38	12	8					1					
55	863	8	18	41	87	110	133	135	99	88	36	27	39	13	10		4					
56	1,313	3	17	56	94	146	138	220	192	142	106	66	39	26	18		7					
57	1,769	1	35	136	166	154	212	262	239	232	172	127	84	52	38		10					
58	1,813	7	13	54	118	189	189	212	273	274	214	153	128	78	53		26					
59	1,784	3	7	1	31	62	124	212	259	256	224	201	145	98	84		17					
60	1,311	1	1	2	8	43	77	122	147	183	175	141	133	98	65		9					
61	986	70	33	4	7	33	55	33	99	120	139	141	109	92	70		10					
62	317	71	632		7	10	35	33	58	91	92	85	68	40	36		17					
63	157	73	137		8	31	29	48	42	46	43	42	46	23	30		12					
64	63	74	63		1	5	1	13	13	13	13	12	14	26	20		8					
65	25	75	15		5	3	1	3	1	3	2	2	15	9	7		7					
66	6	76	6										2	2	2		3					
67	8	77	7										1	2	4		1					
68	1	78	8														2					
69	1	79	1														2					
70	6	80	1														1					
Total	12,076	39	94	271	538	789	1,091	1,399	1,423	1,425	1,265	1,057	829	622	436	274	183	130	69	65	34	43

Number of cases: 12,076. Height: Mean, 67.11 inches; standard deviation, 2.68±0.012 inch. Weight: Mean, 142.30 pounds; standard deviation, 17.21±0.075 pound.

TABLE LIII. Correlation between height and chest circumference (expiration): Group 16, Russian, 10 per cent plus, first million draft recruits.

Height, in inches.	Chest circumference, in inches.											
	Total.	29	30	31	32	33	34	35	36	37	38	39
59	58	4	4	8	10	9	11	6	2	2	1	1
60	54	5	7	7	10	11	7	6	1	1	1	1
61	134	8	16	23	30	23	11	12	8	1	1	1
62	252	10	22	44	64	52	36	14	9	1	1	1
63	525	27	43	73	107	105	76	50	28	10	4	2
64	857	28	61	119	168	154	152	95	47	22	5	6
65	1,310	41	86	192	231	264	228	139	79	24	15	11
66	1,768	48	91	214	324	360	304	214	119	49	21	14
67	1,812	24	75	209	324	371	309	245	144	60	29	22
68	1,780	14	59	186	318	364	313	243	154	71	34	24
69	1,310	11	33	108	200	260	281	193	124	53	30	17
70	985	2	15	68	137	200	225	143	105	53	25	12
71	680	4	4	36	89	116	143	85	69	43	16	17
72	316	1	2	11	36	67	67	62	40	14	13	7
73	157	1	1	6	10	22	31	38	28	6	4	6
74	63	1	1	1	3	4	4	11	9	4	3	1
75	25	1	1	1	3	1	1	7	4	2	2	1
76	8	1	1	1	1	1	1	4	1	1	1	1
77	1	1	1	1	1	1	1	1	1	1	1	1
78	1	1	1	1	1	1	1	1	1	1	1	1
79	6	1	1	1	1	1	1	3	1	1	1	1
Total	12,057	228	527	1,365	2,075	2,390	2,215	1,570	973	421	205	148

Number of cases: 12,057. Height: Mean, 67.11 inches; standard deviation, 2.69 ± 0.012 inch. Chest circumference (expiration): Mean, 33.39 inches; standard deviation, 2 ± 0.009 inch.

TABLE LVI.—Correlation between height and chest circumference (expiration): Group 17, Scandinavian, 10 per cent, first million draft recruits.

Height, in inches.	Chest circumference, in inches.											
	Total.	29	30	31	32	33	34	35	36	37	38	39
59.....	172	3	8	14	30	33	28	23	17	10	5	1
60.....	94	3	8	17	17	15	13	12	5	1	1	2
61.....	286	15	24	35	46	45	36	15	10	6	3	1
62.....	524	18	42	97	116	98	92	40	14	5	1	1
63.....	1,161	26	93	192	244	249	182	88	24	18	10	5
64.....	2,249	57	155	325	467	482	380	223	98	87	34	11
65.....	3,880	86	216	486	809	856	668	418	202	171	118	28
66.....	5,937	93	271	685	1,155	1,296	1,051	723	379	275	144	39
67.....	7,430	72	295	713	1,288	1,537	1,516	1,043	528	347	98	65
68.....	8,161	53	215	610	1,314	1,747	1,718	1,208	715	375	144	90
69.....	7,274	34	184	469	1,011	1,472	1,542	1,194	783	348	143	94
70.....	5,744	27	104	303	706	1,094	1,274	1,016	645	317	159	99
71.....	3,800	7	45	193	404	710	840	711	453	244	114	79
72.....	2,254	3	28	85	230	366	463	440	311	182	97	49
73.....	1,140	1	7	34	91	183	244	225	166	111	41	27
74.....	525	2	1	9	38	67	110	98	94	54	34	18
75.....	215	1	1	5	16	21	39	37	52	18	15	10
76.....	74	1	1	2	2	4	15	19	13	14	4	4
77.....	30	1	1	1	1	6	5	6	7	4	1	1
78.....	19	1	1	1	1	3	3	3	4	5	1	1
79.....	22	1	1	1	2	2	2	3	4	2	3	2
Total.....	50,961	501	1,699	4,286	7,987	10,296	10,221	7,548	4,544	2,258	984	627

Number of cases: 50,961. Height: Mean, 67.96 inches; standard deviation, 2.63 ± 0.006 inch. Chest circumference (expiration): Mean, 33.63 inches; standard deviation, 1.94 ± 0.004 inch.

TABLE I.VII. Correlation between weight and chest circumference (expiration): Group 17, Scandinavians, 10 per cent, first million draft recruits.

Chest circumference, in inches.	Total.	Weight, in pounds.																							
		95- 99	100- 104	105- 109	110- 114	115- 119	120- 124	125- 129	130- 134	135- 139	140- 144	145- 149	150- 154	155- 159	160- 164	165- 169	170- 174	175- 179	180- 184	185- 189	190- 194	195- 199	200- 204		
29	498	2	15	52	87	96	93	61	38	20	15	6	2	2	2	3	1	2	1	1	1	1	3		
30	1,699	1	9	40	159	288	352	317	230	150	65	42	12	7	25	3	6	4	3	3	1	1	3		
31	4,285	1	6	37	174	352	663	821	803	616	355	213	108	55	25	15	5	4	3	3	1	1	1		
32	7,967	1	2	16	72	302	716	1,099	1,395	1,500	1,224	775	464	223	95	63	18	12	13	2	6	4	1		
33	10,281	1	6	4	19	126	367	802	1,373	1,768	1,885	1,483	1,081	665	386	189	71	33	51	13	6	4	1		
34	10,232	1	2	3	10	41	131	382	735	1,253	1,538	1,789	1,553	1,116	760	509	238	97	61	21	6	4	1		
35	7,559	1	1	1	1	13	39	106	274	553	802	1,062	1,235	1,083	924	642	385	214	115	51	21	13	5		
36	4,545	1	2	2	2	3	10	30	75	139	267	449	570	627	651	590	452	278	179	116	49	37	17		
37	2,253	1	1	3	3	7	3	1	9	39	70	129	216	261	322	301	256	213	156	101	75	47	43		
38	966	1	1	1	4	2	3	1	2	9	9	25	44	50	89	108	134	101	118	99	62	41	79		
39	379	1	1	1	1	1	1	1	1	1	4	6	5	15	23	26	9	33	34	40	42	41	70		
40	247	1	1	1	3	2	6	1	3	1	5	2	1	11	4	9	9	13	13	12	20	20	119		
Total	50,953	4	45	166	534	1,261	2,381	3,622	4,946	6,027	6,239	6,001	5,291	4,125	3,294	2,460	1,618	1,001	689	439	275	210	335		

Number of cases: 30,953. Weight: Mean, 146.15 pounds; standard deviation, 17.01 \pm 0.036 pound. Chest circumference (expiration): Mean, 33.65 inches; standard deviation, 1.96 \pm 0.004 inch.

TABLE LX.--(Correlation between height and chest circumference (expiration): Group 18, Finn, 10 per cent, first million draft recruits.

Height, in inches.	Total.	Chest circumference, in inches.										
		29	30	31	32	33	34	35	36	37	38	39
59	15											
60	22	1	2	1	4	2	1	4	1	2		
61	46	1	3	5	3	3	4	1	2	1		
62	102	5	3	7	12	8	9	2	1	3		
63	216	3	17	13	26	18	23	7	4	1		
64	263	9	3	22	36	53	38	22	12	9		
65	565	9	16	40	76	78	80	34	19	5		
66	759	9	20	51	107	119	98	70	57	21		
67	873	8	26	72	138	155	134	110	67	31		
68	886	5	30	66	152	185	170	125	88	47		
69	767	2	18	50	136	185	170	115	89	65		
70	547	2	15	39	85	165	152	126	102	44		
71	361	1	7	28	70	98	93	71	82	27		
72	180	1	2	13	35	76	71	40	40	33		
73	41	1	3	9	16	26	32	39	23	14		
74	83		1	3	8	9	17	21	9	12		
75	23		1	1	1	5	4	8	12	3		
76	1						1	3	6	2		
77	1							1	1	1		
78	1											
79	1											
Total	5,855	49	163	421	915	1,145	1,123	889	614	322	140	85

Number of cases: 5,855. Height: Mean, 67.43 inches; standard deviation, 2.65 ± 0.017 inch. Chest circumference (expiration): Mean, 33.82 inches; standard deviation, 1.98 ± 0.012 inch.

TABLE LX.—Correlation between weight and chest circumference (expiration): Group 18, Finn, 10 per cent, first million draft recruits.

Chest circumference, in inches.	Total.	Weight, in pounds.																					
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204
29	49			7	9	6	7	12	5	1			1					1				1	
30	163		1	5	9	30	35	26	24	17	5	5	2	1		1							
31	421		1	8	19	40	64	97	74	58	27	14	9	3	1		1	1					
32	915			3	9	45	85	141	164	161	150	73	46	21	8	4	4						
33	1,145				2	15	53	88	166	235	191	153	114	62	40	12	8	1			1		
34	1,125				1	7	24	60	87	145	183	196	155	117	67	39	24	19	6	7	1		
35	1,967			1		3	13	21	41	71	78	124	139	121	100	82	39	35	23	14	9	2	
36	614						6		13	25	46	80	78	87	77	71	49	25	15	6	6	6	1
37	322									8	18	34	28	42	51	49	28	16	17	13	10	2	6
38	149				1				2	4	5	6	11	9	13	19	15	4	2	4	6	7	11
39	58			1							3		1		6	3	5	4	3	2	1	1	11
40	27										1			2		3							
41																							
42																							
Total.....	5,855	2	25	50	146	281	451	577	725	707	685	584	400	363	287	169	125	73	48	34	19	35	

Number of cases: 5,855. Weight: Mean, 145.80 pounds, standard deviation, 16.88±0.11 pound. Chest circumference (expiration): Mean, 33.82 inches; standard deviation, 1.99±0.01 inch. Correlation: 0.6727±0.0048.

TABLE LXI.—Correlation between height and weight: Group 19, French-Canadian, 10 per cent, first million draft recruits.

Total.		Weight, in pounds.																					
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204
59	101	1	2	3	6	5	7	13	15	11	9	3	9	7	5	1	1	1	1	1	1	1	1
60	112	1	4	10	15	23	17	26	13	11	7	10	2	2	3	1	1	1	1	1	1	1	1
61	376	1	18	30	71	72	82	80	81	29	8	10	9	2	1	1	1	1	1	1	1	1	1
62	845	3	17	69	131	151	142	110	112	49	41	18	10	7	6	3	4	3	2	2	1	1	1
63	1,532	1	23	77	199	267	242	212	170	124	93	51	22	20	23	10	10	6	7	6	1	1	1
64	3,210	2	23	56	196	355	362	363	306	240	126	100	64	41	19	19	25	12	11	10	5	3	3
65	5,640	3	13	44	176	358	478	510	451	367	264	208	131	65	60	36	26	16	10	12	9	7	7
66	3,840	1	13	22	143	315	476	537	574	484	373	303	192	135	98	54	30	19	14	12	8	5	5
67	3,833	1	11	15	63	187	342	467	568	593	444	341	271	185	137	112	59	53	28	23	16	11	13
68	3,478	1	6	13	24	120	256	381	451	482	440	393	286	221	131	97	67	58	28	18	16	11	13
69	2,581	1	3	2	11	32	90	222	277	354	340	324	269	221	131	98	89	52	47	31	17	14	9
70	1,732	1	3	5	5	15	43	82	145	215	248	233	214	150	98	82	72	50	41	31	13	9	14
71	1,098	1	1	1	6	16	30	76	89	146	132	116	64	57	39	25	27	15	11	9	6	1	7
72	76	2	2	1	1	3	3	2	2	11	19	31	27	27	20	20	8	5	5	6	1	1	1
73	31	1	1	1	1	1	1	1	1	1	3	2	3	3	1	1	3	2	2	2	2	3	3
74	16	1	1	1	1	1	1	1	1	1	1	1	3	5	1	1	3	1	2	1	1	1	1
75	4	1	1	1	1	1	1	1	1	1	2	1	1	1	1	1	1	1	1	1	1	1	1
76	6	1	1	1	1	1	1	1	1	1	2	1	1	1	1	1	1	1	1	1	1	1	1
77	7	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
78	7	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
79	7	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Total	25,862	5	133	349	1,044	1,868	2,528	3,004	3,181	3,065	2,618	2,215	1,724	1,269	852	652	427	323	201	131	109	87	67

Number of cases: 25,862. Height: Mean, 66.67 inches; standard deviation, 2.65 ± 0.008 inch. Weight: Mean, 137.88 pounds; standard deviation, 17.38 ± 0.052 po und.

TABLE LXII.—Correlation between height and chest circumference (expiration): Group 19, French-Canadian, 10 per cent, first million draft recruits.

Height, in inches.	Chest circumference, in inches.											
	29	30	31	32	33	34	35	36	37	38	39	
59.....	103	4	7	8	21	18	19	10	7	5	2	
60.....	111	5	14	16	22	25	15	4	6	1	1	
61.....	375	22	40	78	83	65	42	25	9	1	1	
62.....	839	40	95	159	190	162	102	50	25	7	6	
63.....	1,526	63	166	268	312	311	198	117	61	17	13	
64.....	2,303	93	185	388	525	443	321	184	95	36	20	
65.....	3,203	111	235	474	714	611	458	264	157	77	42	
66.....	3,280	101	274	531	790	753	595	362	206	112	60	
67.....	3,424	79	214	450	763	759	686	413	238	115	47	
68.....	3,475	65	177	396	613	724	578	443	263	108	61	
69.....	3,575	31	74	273	440	506	301	342	215	96	40	
70.....	1,720	15	65	162	284	357	302	226	152	71	45	
71.....	1,037	11	36	77	157	175	188	170	108	57	34	
72.....	506	4	11	41	61	92	90	82	56	28	13	
73.....	76	2	2	14	22	40	47	30	21	15	10	
74.....	30	1	2	8	13	11	15	14	6	9	5	
75.....	17	1	1	3	9	9	6	4	4	1	2	
76.....	3	1	1	2	1	4	1	1	1	1	1	
77.....	6	1	1	1	1	1	1	1	1	1	1	
78.....	8	1	1	1	1	1	1	1	1	1	1	
79.....	8	1	1	1	1	1	1	1	1	1	1	
Total.....	25,772	645	1,597	3,340	5,016	5,065	4,189	2,786	1,605	753	395	381

Number of cases, 25,772. Height: Mean, 66.67 inches; standard deviation, 2.65±0.008 inch. Chest circumference (expiration): Mean, 33.11 inches; standard deviation, 2.05±0.006 inch.

TABLE LXIII.—Correlation between weight and chest circumference (expiration): Group 19, French-Canadian, 10 per cent, first million draft recruits.

Chest circumfer- ence, in inches.	Total.	Weight, in pounds.																					
		95- 99	100- 104	105- 109	110- 114	115- 119	120- 124	125- 129	130- 134	135- 139	140- 144	145- 149	150- 154	155- 159	160- 164	165- 169	170- 174	175- 179	180- 184	185- 189	190- 194	195- 199	200- 204
29.....	666	2	41	78	158	161	106	54	24	19	5	5	5	1	2	1	1	2	2	2	3	2	2
30.....	1,549	1	30	99	282	353	324	238	125	78	32	5	5	3	10	2	4	2	1	1	2	3	2
31.....	3,343		16	96	291	534	648	636	511	298	176	68	27	14	2	5	6	4	2	3	1	3	2
32.....	5,013		7	41	193	491	741	947	976	632	516	240	120	62	25	12	9	4	3	1	2	1	1
33.....	5,072		1	10	79	246	446	718	831	888	701	502	321	165	74	43	21	18	3	1	2	1	1
34.....	4,183		4	2	10	61	182	307	531	653	664	640	476	300	173	99	47	21	8	4	2	4	1
35.....	2,789		2	2	4	25	48	73	201	286	371	432	460	325	217	169	80	55	20	8	4	10	1
36.....	1,612				3	4	3	30	57	100	157	199	190	218	102	103	118	70	40	29	24	12	10
37.....	754			1	3	2	5	2	9	17	25	56	75	92	43	53	44	52	32	31	20	15	19
38.....	368				2	1	1	2	6	4	7	7	25	43	8	14	16	18	32	13	17	12	35
39.....	197		1	1	2	1	2	2	3	3	2	2	5	2	1	6	5	9	8	13	24	19	88
40.....	186				1	1	2	2	2	2	3	1	2	2	1	1	6	9	5	13	24	19	
Total.....	25,787	3	103	330	1,025	1,889	2,506	3,009	3,274	2,980	2,639	2,157	1,711	1,224	899	631	423	322	199	131	110	85	157

Number of cases: 25,787. Weight: Mean, 138.15 pounds; standard deviation, 17.99±0.06 pound. Chest circumference (expiration): Mean, 33.11 inches; standard deviation, 2.07±0.01 inch. Correlation: 0.7169±0.0026.

TABLE LXIV.—Correlation between height and weight: Group 20, German and Scandinavian, 10 per cent plus, first million draft recruits.

Height, in inches.	Weight, in pounds.																						
	95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204	
59.....	1	1	2	1	1	1	1	1	1	1	1	1
60.....	103	1	4	1	1	1	1	1	1	1	1	1
61.....	53	1	7	1	1	1	1	1	1	1	1	1
62.....	108	1	4	1	1	1	1	1	1	1	1	1
63.....	214	1	16	1	1	1	1	1	1	1	1	1
64.....	549	2	8	2	2	2	2	2	2	2	2	2
65.....	1,104	2	22	2	2	2	2	2	2	2	2	2
66.....	1,987	2	57	2	2	2	2	2	2	2	2	2
67.....	3,141	1	9	1	1	1	1	1	1	1	1	1
68.....	4,034	1	2	1	1	1	1	1	1	1	1	1
69.....	4,590	4	5	4	4	4	4	4	4	4	4	4
70.....	4,116	1	1	1	1	1	1	1	1	1	1	1
71.....	3,311	
72.....	2,215	
73.....	1,372	
74.....	674	
75.....	318	
76.....	126	
77.....	46	
78.....	21	
79.....	14	
80.....	19	
Total.....	28,005	1	24	80	286	652	1,226	1,856	2,662	3,270	3,448	3,424	2,934	2,287	1,876	1,417	948	565	395	263	144	122	195
Total.....																							

Number of cases: 28,005. Height: Mean, 68.11 inches; standard deviation, 2.62±0.007 inch. Weight: Mean, 146.66 pounds; standard deviation, 17.00±0.048 pound.

TABLE LXV.—*Correlation between height and chest circumference (expiration): Group 20, German and Scandinavian, 10 per cent plus, first million draft recruits.*

Height, in inches.	Total.	Chest circumference, in inches.										
		29	30	31	32	33	34	35	36	37	38	39
59.....	104	1	4	9	17	19	16	15	12	6	4	1
60.....	83	1	3	10	9	8	8	9	2	1	1	1
61.....	108	8	11	13	15	27	18	9	3	2	2	
62.....	214	6	16	40	41	42	44	20	4	2		
63.....	536	12	35	87	125	114	86	37	22	9		
64.....	1,103	22	78	157	221	244	169	122	57	17	5	4
65.....	1,987	40	108	248	415	442	345	211	84	42	17	9
66.....	3,138	46	126	373	601	704	568	379	191	89	36	15
67.....	4,024	33	145	399	699	867	821	599	275	144	55	25
68.....	4,491	31	109	330	704	993	990	711	395	196	83	49
69.....	4,114	19	85	246	593	794	904	698	459	207	77	62
70.....	3,300	16	56	165	387	636	746	581	376	217	84	56
71.....	2,208	3	33	113	225	399	499	410	273	143	69	51
72.....	1,372	1	13	49	144	213	298	255	194	115	61	31
73.....	675	2	4	16	50	109	142	149	101	62	29	11
74.....	317	1	1	7	19	33	69	66	55	33	22	10
75.....	124	1	1	3	11	13	24	22	28	11	8	3
76.....	40			2	1	2	10	12	10	9		2
77.....	21					1	4	4	6	2		1
78.....	14						2	4		1		1
79.....	19					2	1	2	4	1	3	2
Total.....	29,031	241	829	2,240	4,220	5,666	5,732	4,305	2,551	1,313	563	371

Number of cases: 29,031. Height: Mean, 68.11 inches; standard deviation, 2.62 ± 0.007 inch. Chest circumference (expiration): Mean, 33.72 inches; standard deviation, 1.93 ± 0.003 inch.

TABLE LXVI.—Correlation between weight and chest circumference (expiration) Group 20, German and Scandinavian, 10 per cent plus, first million draft recruits.

Chest circumference, in inches.	Total.	Weight, in pounds.																					
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204
28	240	1	8	24	48	48	42	24	14	12	10	4	1	1	1		1						
29	829		3	23	82	144	170	147	108	75	30	23	9	3	3	2	4						
30	2,239		4	17	93	201	336	406	431	319	194	121	59	29	17	6	2	1					
31	4,222		2	8	33	164	382	560	736	782	611	455	240	128	60	36	11	8	5	1	1		
32	5,696		4	1	12	64	192	441	774	987	1,064	826	569	355	208	119	40	18	6	2	3	1	
33	5,757		1	3	6	16	72	202	408	733	863	1,022	883	603	432	289	138	43	29	7	4	3	
34	4,306																						
35	2,552			1	2	2	6	15	59	159	288	468	646	704	603	432	367	222	122	73	23	9	10
36	1,310			1	1	3	2	6	35	68	145	239	310	367	378	344	266	162	94	61	20	25	10
37	1,565			1	1	1	1	2	5	1	37	65	133	154	184	172	159	126	89	66	45	24	23
38	218										1	4	2	9	11	16	18	20	22	26	31	22	49
39	30																						
40	152																						
Total.....	28,056	1	27	82	282	652	1,222	1,856	2,080	3,262	3,428	3,421	2,931	2,288	1,875	1,416	953	565	365	258	146	121	186

Number of cases: 28,056. Weight: Mean, 146.67 pounds; standard deviation, 17.01 \pm 0.048 pound. Chest circumference (expiration): Mean, 33.72 inches; standard deviation, 1.95 \pm 0.006 inch.

TABLE LXVII.—Correlation between height and weight: Group 21, German and Austrian, 20 per cent plus, first million draft recruits.

Height, in inches.	Total.	Weight, in pounds.																190-199	200-204					
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174			175-179	180-184	185-189		
59	197	1	4	3	8	18	10	20	20	15	26	12	17	5	8	16	7	1	1	1	1	1	4	
60	110	2	2	8	15	15	12	13	6	6	8	6	3	3	4	4	1	1	1	2	2	4	3	
61	332	1	11	18	48	46	34	38	31	24	17	18	14	11	3	1	1	1	1	1	1	1	4	
62	695	2	17	47	86	120	112	84	69	64	29	17	14	14	11	5	1	1	1	1	1	1	4	
63	1,416	1	17	68	119	213	235	203	187	171	80	62	35	25	14	15	1	1	1	1	1	1	4	
64	2,358	1	16	68	120	247	309	356	346	271	232	132	77	60	45	23	9	9	3	8	6	3	7	
65	3,905	1	18	62	120	247	337	465	497	575	513	387	288	180	91	50	26	18	20	15	9	4	8	
66	5,138	1	14	44	156	337	465	497	575	713	770	653	499	339	209	121	107	63	44	53	23	10	31	
67	5,954	1	20	120	266	441	665	635	778	911	782	683	457	335	230	141	97	54	53	35	29	19	42	
68	5,865	1	20	120	266	441	665	635	778	911	782	683	457	335	230	141	97	54	53	35	29	19	42	
69	4,949	1	5	14	36	97	265	446	651	763	789	755	625	445	328	238	146	83	54	35	29	19	42	
70	3,525	1	1	4	13	40	150	240	454	583	664	679	587	386	354	278	184	125	71	35	26	27	46	
71	2,200	1	2	4	4	29	47	128	249	365	445	460	439	355	289	212	165	94	62	45	31	28	46	
72	1,509	1	4	2	2	2	5	15	41	86	151	233	286	298	273	179	189	124	71	34	18	18	39	
73	261	1	1	1	1	2	4	18	4	15	116	144	176	163	114	116	99	71	42	28	28	17	39	
74	104	1	1	1	1	1	1	4	4	9	3	23	19	33	33	32	23	35	29	25	9	7	22	
75	38	1	1	1	1	1	1	1	1	1	1	5	11	11	12	15	10	6	4	10	11	4	15	
76	16	1	1	1	1	1	1	1	1	1	1	2	3	2	6	3	6	3	7	11	3	3	8	
77	11	1	1	1	1	1	1	1	1	1	1	1	1	1	2	1	1	1	2	3	2	1	1	
78	19	1	1	1	1	1	1	1	1	1	1	1	1	1	1	2	2	2	2	2	1	1	3	
79	38,962	6	108	312	800	1,674	2,489	3,387	4,231	4,678	4,515	4,141	3,354	2,589	2,011	1,520	997	655	467	296	211	171	350	
Total.....																								

Number of cases: 38,962. Height: Mean, 67.41 inches; standard deviation, 2.69±0.007 inch. Weight: Mean, 143.37 pounds; standard deviation, 18.05±0.044 pound.

TABLE LXVIII.—Correlation between height and chest circumference (expiration): Group 21, German and Austrian, 20 per cent plus, first million draft recruits.

Height, in inches.	Total.	Chest circumference, in inches.										
		29	30	31	32	33	34	35	36	37	38	39
59.....	197	9	8	18	41	35	35	20	19	4	6	2
60.....	110	1	13	18	26	15	19	5	8	2	2	1
61.....	332	20	34	66	60	68	34	26	14	6	3	2
62.....	697	37	81	128	133	124	87	66	25	11	13	2
63.....	1,401	55	137	232	312	263	185	108	66	22	22	8
64.....	2,357	84	184	372	481	437	370	207	122	53	51	19
65.....	3,805	114	291	532	754	747	579	380	209	107	144	41
66.....	5,139	117	281	643	952	1,117	905	547	307	144	109	83
67.....	5,954	109	334	651	991	1,070	1,109	692	362	208	85	63
68.....	5,857	79	224	548	991	1,144	1,120	837	476	230	109	99
69.....	4,951	42	163	442	760	957	980	719	475	213	97	104
70.....	3,525	31	98	219	495	736	743	514	355	188	77	69
71.....	2,200	16	48	159	256	416	487	367	250	123	72	56
72.....	1,360	5	29	77	153	217	307	240	145	97	49	41
73.....	609	3	7	25	66	102	124	105	81	51	24	21
74.....	261	1	3	10	20	38	47	49	51	19	14	9
75.....	104	3	8	15	14	24	17	10	6	7
76.....	38	1	3	5	10	10	5	4
77.....	16	2	1	4	3	3
78.....	11	1	2	3	2	1
79.....	19	2	4	1	4	1
Total.....	38,943	723	1,934	4,147	6,582	7,662	7,112	4,924	3,026	1,469	703	631

Number of cases: 38,943. Height: Mean, 67.41 inches; standard deviation, 2.69±0.007 inch. Chest circumference (expiration): Mean, 33.40 inches; standard deviation, 2.05±0.005 inch.

TABLE LXIX.—Correlation between weight and chest circumference (expiration): Group 21, German and Austrian, 20 per cent plus, first million draft recruits.

Chest circumference, in inches.	Total.	Weight, in pounds.																					
		95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200 and over.
29.....	702	4	26	83	141	152	121	64	43	21	16	6	4	2	1	1	1	1	1	4
30.....	1,822	27	114	239	377	381	307	221	146	65	37	15	10	3	4	1	1	1	4
31.....	4,144	1	20	70	233	519	705	768	716	480	301	162	91	33	18	13	4	2	5	2	2	2	2
32.....	6,579	1	6	29	135	397	705	1,051	1,208	1,113	782	554	300	160	71	40	14	6	3	1	1	1	1
33.....	7,657	11	7	46	160	385	1,737	1,089	1,380	1,348	1,002	673	415	205	118	33	18	9	4	5	4	3
34.....	7,113	3	3	13	34	141	331	1,089	1,956	1,127	1,185	979	675	400	263	148	53	22	11	9	17	23
35.....	4,923	3	1	5	13	30	83	217	411	568	744	713	674	531	306	233	136	96	21	17	13	22
36.....	3,025	2	2	1	3	9	28	72	127	217	316	395	398	427	379	242	153	103	77	35	32	41
37.....	1,496	1	1	1	2	1	5	14	27	61	97	133	163	166	191	184	151	99	60	58	32	41
38.....	1,707	1	7	2	1	4	4	3	5	21	35	46	66	69	95	82	77	58	30	32	56
39.....	345	1	2	1	2	2	3	9	7	7	13	20	30	34	37	32	20	35	78
40.....	288	1	2	5	5	4	2	5	4	2	5	8	5	12	13	17	18	17	20	138
Total.....	38,911	6	108	310	794	1,675	2,485	3,384	4,223	4,674	4,510	4,137	3,349	2,588	2,009	1,518	996	655	467	293	212	170	348

Number of cases: 38,911.
Correlation: 0.7079 ± 0.0017 .

Weight: Mean, 143.27 pounds; standard deviation, 18.04 \pm 0.04 pound.

Chest circumference (expiration): Mean, 33.42 inches; standard deviation 2.07 \pm 0.01 inch.

TABLE LXX.—Correlation between height and weight: Group 22, German and Austrian, 15 per cent plus, first million draft recruits.

Total.	Weight, in pounds.																							Total.....
	95-99	100-104	105-109	110-114	115-119	120-124	125-129	130-134	135-139	140-144	145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200 and over.		
533	5	14	25	29	45	43	69	57	43	57	41	38	22	19	22	9	3	2	1	2	2	5		
531	5	17	51	70	83	67	53	41	30	23	26	16	14	9	15	6	4	2	2	2	1	5		
1,306	9	36	98	139	228	178	174	108	98	53	34	35	24	13	9	14	6	4	2	2	1	10		
2,649	11	60	175	323	417	421	353	279	217	140	73	55	34	24	14	14	6	5	4	1	12	16		
5,143	6	50	208	445	710	814	800	707	496	306	229	198	90	47	40	25	11	11	14	9	8	17		
8,493	14	54	192	503	944	1,212	1,253	1,064	754	464	302	235	116	63	55	34	11	11	14	19	18	26		
12,958	1	38	140	340	606	1,006	1,563	2,437	2,585	2,344	1,563	1,351	886	639	502	323	172	118	48	34	51	66		
16,913	1	1	113	255	455	795	1,295	2,447	2,585	2,344	1,563	1,351	886	639	502	323	172	118	48	34	51	66		
19,021	7	80	204	286	1,205	7,498	2,437	2,585	2,344	1,563	1,351	886	639	502	323	172	118	48	34	51	66		
18,657	10	37	114	178	300	403	2,383	2,585	2,344	1,563	1,351	886	639	502	323	172	118	48	34	51	66		
15,275	4	12	32	66	182	403	1,737	1,767	2,063	2,084	1,842	1,370	909	727	488	332	202	119	88	70	117		
11,184	5	4	21	51	136	283	1,144	1,441	1,521	1,401	1,011	674	436	303	204	133	82	74	56	44	66		
7,045	2	5	12	18	56	143	278	468	754	916	935	596	361	249	169	124	91	64	44	30	57		
4,858	1	2	5	8	23	53	123	223	335	490	537	439	291	206	119	82	64	44	30	57			
1,821	2	2	5	13	42	60	100	174	192	241	206	119	82	64	44	30	57	44	66		
314	1	1	1	7	9	20	24	58	84	92	106	111	82	64	44	30	57	44	66		
104	1	5	2	7	15	27	37	44	41	30	26	17	12	8	1	1		
52	3	4	2	4	4	5	7	11	19	7	4	3	2	1	1		
34	1	4	2	2	2	2	2	2	2	2	2	2	2	1	1		
8		
42		
126,994	42	336	1,138	2,912	5,876	8,782	11,668	14,114	15,295	14,707	13,139	10,734	8,245	6,141	4,729	2,992	2,007	1,315	900	625	465	832		

Number of cases: 126,994. Height: Mean, 67.27 inches; standard deviation, 2.72±0.004 inch. Weight: Mean, 142.31 pounds; standard deviation, 17.73±0.024 pound.

TABLE LXXI.—Correlation between height and chest circumference (expiration): Group 22, German and Austrian, 15 per cent plus, first million draft recruits.

Height, in inches.	Total.	Chest circumference, in inches.										
		29	30	31	32	33	34	35	36	37	38	39
59.....	548	27	47	58	111	102	83	59	36	9	13	3
60.....	530	20	65	78	117	96	69	42	17	8	5	2
61.....	1,307	60	149	268	253	242	151	96	46	19	8	6
62.....	2,655	144	281	482	534	498	358	216	88	28	14	12
63.....	5,125	268	510	828	1,073	986	729	412	220	64	47	28
64.....	8,486	328	692	1,286	1,753	1,646	1,209	764	420	180	69	49
65.....	12,048	411	1,028	1,858	2,558	2,511	2,041	1,318	644	328	142	109
66.....	16,912	442	1,090	2,286	3,316	3,433	2,775	1,828	969	438	214	181
67.....	19,010	378	1,033	2,211	3,449	3,800	3,378	2,292	1,259	624	284	212
68.....	18,620	281	1,061	1,861	3,191	3,766	3,532	2,421	1,450	720	339	260
69.....	15,273	158	799	1,351	2,326	3,021	2,997	2,241	1,405	658	300	274
70.....	11,159	99	346	1,357	1,586	2,203	2,203	1,729	1,085	598	250	213
71.....	7,059	44	178	462	1,863	1,360	1,427	1,156	765	396	203	175
72.....	4,034	10	70	230	475	1,724	1,860	1,672	496	271	132	94
73.....	1,859	5	18	91	199	263	384	360	240	140	71	58
74.....	819	3	12	35	71	119	172	151	133	65	32	26
75.....	312	2	3	7	21	47	57	64	49	30	18	14
76.....	105	1	2	5	8	12	24	27	13	9	2	3
77.....	54	1	1	7	12	12	6	6	1	5
78.....	32	2	7	4	4	9	4
79.....	40	2	6	9	5	7	3
Total.....	126,887	2,639	6,839	14,168	21,909	24,971	22,624	15,871	9,377	4,616	2,147	1,726

Number of cases: 126,887. Height: Mean, 67.27 inches; standard deviation, 2.72±0.004 inch. Chest circumference (expiration): Mean, 33.32 inches; standard deviation, 2.04±0.008 inch.

TABLE LXXIV.—Correlation between stature and weight in white and colored troops, demobilization.

Height, in centimeters.	Total.	Weight, in pounds.										Mean weight.	
		100-109	110-119	120-129	130-139	140-149	150-159	160-169	170-179	180-189	190-199		200 and over.
148-149	18		1	4	5	2	4	4	2				140.06
150-151	42	6	11	20	6	2	4	2	1				126.11
152-153	120	10	48	29	21	6	4	5	1				123.33
154-155	324	21	88	111	66	27	6	3	2				125.55
156-157	622	33	162	194	153	61	18	6	5				126.59
158-159	1,210	42	285	436	269	115	39	13	6				127.49
160-161	2,300	60	414	728	634	317	102	37	8				129.98
162-163	3,569	71	491	1,077	1,055	556	199	68	35				132.02
164-165	5,408	66	465	1,445	1,730	967	459	152	57				134.76
166-167	7,056	35	457	1,564	2,100	1,732	747	260	123				137.52
168-169	8,978	29	342	1,662	2,743	2,284	1,797	488	158				139.81
170-171	9,567	16	253	1,276	2,609	2,609	1,649	801	224				143.10
172-173	162	4	152	960	2,327	2,758	1,918	904	366				145.42
174-175	9,636	7	93	643	1,322	2,478	2,083	1,137	503				148.21
176-177	5,802	3	56	334	1,211	2,068	1,986	1,176	534				151.26
178-179	4,031	4	22	157	739	1,468	1,560	1,068	877				154.13
180-181	2,637	4	23	98	361	841	1,068	617	463				156.92
182-183	1,596	4	5	47	168	539	694	417	351				159.14
184-185	1,029	4	10	18	60	220	376	417	267				162.99
186-187	555	3	8	14	34	92	214	222	182				166.53
188-189	221	1	4	10	27	66	109	135	98				168.07
190-191	111	1	2	5	8	23	35	25	41				168.74
192-193	48	1	4	4	6	18	10	4	13				169.21
194-195	31	1	4	4	4	7	10	4	10				169.52
196-197	17	1	1	1	7	2	3	3	4				169.80
198-199	4	1	1	1	1	1	1	1	1				169.82
200-201	2	1	1	1	1	1	1	1	1				169.80
202-203													169.80
204-205													169.80
Total	81,558	424	3,393	10,815	18,151	19,243	14,488	8,487	3,915	1,679	591	372	
Mean height		163.33	164.61	167.31	169.78	172.24	174.30	176.14	177.38	179.16	178.73	179.93	

Weight: Mean, 144.92 pounds; standard deviation, 17.09±0.0285 pound. Height: Mean, 171.93 centimeters; standard deviation, 6.70±0.0112 centimeter. Correlation: 0.5198±0.0017.

TABLE LXXVI.—Correlation between leg length and knee height, white troops, demobilization.

Knee height, in centimeters.		Total.	Leg length, in centimeters.																		Mean leg length.			
			56-57	58-59	60-61	62-63	64-65	66-67	68-69	70-71	72-73	74-75	76-77	78-79	80-81	82-83	84-85	86-87	88-89	90-91		92 and over.		
38-39		839	3	18	29	61	115	139	143	137	98	41	38	2	1					1				On
40-41		2,680	12	26	139	271	418	496	491	423	285	175	104	16	9	7	2	3					88.17	
42-43		7,721	16	42	175	427	963	1,360	1,451	1,319	899	525	304	58	18	18	4	4					88.92	
44-45		14,515	25	85	123	389	1,018	2,018	2,910	3,010	2,240	1,374	808	127	45	45	9	4	3	4			70.16	
46-47		17,743	11	17	92	269	580	1,384	2,807	3,921	3,631	2,436	1,542	276	79	26	11	6	5				71.53	
48-49		14,711	4	18	71	210	523	896	1,530	2,447	2,908	2,040	1,565	408	145	40	13	9	2	11			72.59	
50-51		8,994	5	18	19	73	222	596	875	1,213	1,330	1,496	1,373	362	197	82	33	16	3	3			73.59	
52-53		5,007	1	3	4	13	43	155	449	866	860	736	977	268	221	155	60	37	10	3			75.69	
54-55		2,453	1	1	3	5	5	22	90	299	421	419	383	120	111	69	48	22	14	3			77.12	
56-57		914	1	1			2	7	12	65	113	143	163	44	41	22	17	12	8	1			77.88	
58-59		273					1	3	8	17	13	30	49	8	8	2	3	4	1				76.96	
60-61		47						1	1		2	4	2			2	2	2	1				78.64	
62-63		14																						
Number measured		76,141	79	177	655	1,722	3,903	7,052	10,767	13,657	12,837	10,040	7,213	2,119	985	379	189	95	31	48				
Not measured		21,051																						
Total		97,192																						
Mean knee height.....cm.		44.50	44.20	43.83	44.34	44.71	45.33	45.99	46.71	47.32	47.96	48.58	49.26	50.21	51.02	51.85	52.21	52.02	51.79	48.54				

Leg length: Mean, 71.69 centimeters; standard deviation, 4.71±0.0081 centimeter. Knee height: Mean, 47.08 centimeter; standard deviation, 3.62±0.006 centimeter. Correlation: 0.4178±0.0020.

TABLE LXXVII.—*Correlation between chest circumference (rest) and weight, white troops, demobilization.*

Weight, in pounds.	Total.	Chest circumference, in centimeters.										Mean chest circumference.
		68-77	78-81	82-85	86-89	90-93	94-97	98-101	102-105	106-109	110 and over.	
100-109.....	418	84	166	118	24	10	8	3	4	1	2	81.32
110-119.....	3,365	211	1,078	1,409	546	79	25	9	4	2	2	82.74
120-129.....	10,697	176	1,570	4,730	3,388	707	88	31	2	2	3	84.74
130-139.....	17,959	80	856	5,546	7,887	3,050	472	61	3	1	3	86.77
140-149.....	18,892	53	311	2,715	8,111	5,988	1,512	192	9	3	1	88.90
150-159.....	14,040	47	149	808	4,110	5,883	2,613	444	32	3	1	90.72
160-169.....	8,207	29	45	241	1,152	3,229	2,636	777	87	10	1	93.27
170-179.....	3,747	9	28	80	261	955	1,458	760	172	23	1	94.76
180-189.....	1,488	2	9	17	66	178	1,512	496	166	40	2	96.86
190-199.....	545	3	3	10	13	53	113	183	106	50	11	98.82
200 and over.....	348	2	2	3	7	26	95	97	69	47	103.03
Number measured.....	79,706	696	4,215	15,676	25,561	20,049	9,463	3,051	662	201	72	
Not measured.....	16,168	
Total.....	95,874	
Mean weight..... lbs.	126.77	126.66	132.99	141.04	150.19	159.58	169.36	180.61	189.32	191.17	

Weight: Mean, 144.67 pounds; standard deviation, 16.92±0.0283 pound. Chest circumference (rest): Mean, 88.91 centimeters; standard deviation, 5.19±0.009 centimeter. Correlation: 0.6598±0.0013.

TABLE LXXVIII.—Correlation between chest circumference (rest) and neck circumference, white troops, demobilization.

	Neck circumference, in centimeters		Chest circumference (rest), in centimeters										Mean chest circumference	
	Total		66-71	71-77	77-81	82-85	86-89	90-93	94-97	98-101	102-105	106-109		110 and over
26 and under	131	131	1	4	2	53	69	17	2	3	88.36
29	65	65	2	5	7	40	4	2	25	7	93.61
30	219	219	7	31	77	180	182	49	24	9	87.94
31	314	314	13	63	123	418	183	33	17	9	87.95
32	1,123	1,123	21	123	340	1,796	1,162	40	19	15	87.14
33	2,282	2,282	33	123	380	4,407	3,897	247	205	40	10	87.49
34	1,133	1,133	23	124	1,354	4,407	3,010	1,182	683	166	17	87.10
35	20,094	20,094	26	116	1,344	5,404	8,010	4,739	2,178	408	36	88.74
36	20,026	20,026	26	48	1,670	5,877	8,648	5,300	2,218	769	153	14	90.77
37	18,047	18,047	21	26	269	1,827	5,468	5,300	2,539	906	181	24	91.94
38	10,031	10,031	11	15	112	1,611	2,175	3,491	7,322	694	191	44	92.58
39	4,428	4,428	4	3	44	150	2,650	1,384	1,490	402	160	46	95.62
40	1,716	1,716	1	1	11	33	186	384	150	120	75	24	98.14
41	1,492	1,492	1	1	1	4	25	61	19	37	37	8	101.97
42	147	147	1	1	1	2	4	14	5	11	11	5	99.12
43	52	52	2	1	2	1	2	4	1	1	1	1	94.26
44	44	44	1	1	1	1	6	3	1	2	1	1	95.66
45	23	23	1	1	1	1	7	1	2	2	1	1	92.44
46	17	17	1	1	1	1	5	2	2	3	1	1	92.60
47	10	10	1	1	1	1	1	2	2	2	1	1	90.70
48	5	5	1	1	1	1	1	1	2	1	94.83
49	6	6	1	1	1	1	1	1	2	1	87.72
50	18	18	1	1	1	6	4	1	2	1	87.72
Number measured	93,271	93,271	165	599			30,615	24,120	11,163	3,400	903	999	61	
Not measured	93,603	93,603												
Total	93,874	93,874												
Mean neck circumference			34.90	33.86	34.48	35.02	35.72	36.45	37.19	37.88	38			

Chest circumference (rest): Mean, 88.79 centimeters, standard deviation, 5.19±0.0080 centimeters. Neck circumference: Mean, 35.94 centimeters, standard deviation, 1.80±0.003 centimeter. Correlation, 0.5001±0.0016.

TABLE LXXIX.—Correlation between chest circumference (rest) and transverse pelvic diameter, white troops, demobilization.

Transverse pelvic diameter, in centimeters.	Chest circumference (rest), in centimeters.										Mean chest circumference (rest).
	Total.	68-77	78-81	82-85	86-89	90-93	94-97	98-101	102-105	106-109	110
19 and under.....	3			1	1						1
20.....	80	8	2	19	18		9	1			88.83
21.....	216	31	55	62	47		15	4			87.70
22.....	335	53	83	98	52		16	7			86.47
23.....	734	32	100	256	203		37	8			86.19
24.....	1,480	29	198	495	515		181	10		1	86.14
25.....	3,235	58	409	1,113	1,066		461	18			85.73
26.....	6,112	117	653	1,858	1,807		1,067	55			85.82
27.....	9,588	136	826	2,717	3,397		2,477	97			86.34
28.....	14,419	137	898	3,551	5,123		3,387	211			86.88
29.....	15,817	99	678	2,977	5,474		1,769	402			87.04
30.....	13,885	80	513	2,150	4,442		1,975	551			88.94
31.....	10,014	45	299	1,369	2,910		1,687	607			89.42
32.....	7,243	30	175	931	2,024		1,269	496			90.04
33.....	5,069	89	380	567	3,398		1,923	401			90.58
34.....	2,956	11	42	245	703		614	300			91.97
35.....	1,676	9	34	97	352		403	191			92.66
36.....	828	4	11	76	177		183	118			92.90
37.....	448	1	15	55	108		66	58			91.95
38.....	365	3	13	69	127		71	24			90.81
39.....	364	3	13	62	137		47	16			90.62
40.....	202	1	10	44	64		21	3			88.96
41.....	79	1	1	10	28		12	4			91.11
42 and over.....	331		17	41	100		54	24			90.50
Number measured.....	95,479	838	5,073	18,868	30,641	24,160	11,183	3,606	805	223	82
Not measured.....	365										
Total.....	95,874										
Mean transverse pelvic diameter.....	cm.	27.96	27.99	28.50	28.22	29.35	30.62	31.31	32.41	33.41	34.06

Chest circumference (rest): Mean, 88.78 centimeters; standard deviation, 5.17±0.0090 centimeter. Transverse pelvic diameter: Mean, 29.45 centimeters; standard deviation, 2.90±0.005 centimeter. Correlation: 0.3073±0.0021.

TABLE LXXX.—Correlation between chest transverse and chest antero-posterior, while troops, demobilization.

Chest, antero-posterior, in centimeters.	Total.	Chest, transverse, in centimeters.															Mean chest trans- verse. Cm.	
		18-19	20-21	22-23	24-25	26-27	28-29	30-31	32-33	34-35	36-37	38-39	40-41	42-43	44-45	46-47		48-49
14-15.....	19				2	8	5	2	1	2		2			1			28.50
16-17.....	377		3	12	56	149	128	25	211	32	23	13	3	1		2	2	27.08
18-19.....	8,606	6	43	71	716	2,882	3,309	1,201	2,855	331	132	118	20	11	1	3	6	27.87
20-21.....	40,585	7	187	228	1,605	9,486	16,664	9,520	3,891	713	303	111	30	12	4	4	8	28.61
22-23.....	36,105		159	189	673	5,099	13,490	11,511	3,891	483	106	31	10	2	2	1	2	29.37
24-25.....	8,795		26	26	87	673	2,459	3,192	1,694	483	106	7	7	1	3	1		30.25
26-27.....	1,291	1	8	9	20	70	273	381	315	156	46	13	1		2			30.92
28-29.....	432		5	1	10	58	116	119	56	33	17	13	7		2		1	30.32
30-31.....	182		3	4	5	18	35	62	21	16	8	7	3		1		1	30.53
32-33.....	142		1	5	2	12	18	44	36	13	5	4						31.01
34-35.....	21						3	11	3	3								30.60
36-37.....	5							2			1							32.50
38-39.....	4											1						31.00
40-41.....	9							6										29.61
Total.....	96,583	15	426	546	3,176	18,468	36,502	26,106	8,475	1,793	541	307	86	26	15	11	20	
Mean chest antero-posterior...cm.....		20.37	21.57	21.37	20.59	20.92	21.40	21.94	22.52	23.23	23.00	22.44	22.31	21.65	24.10	21.77	22.5	

Chest, transverse: Mean, 29.02 centimeters; standard deviation, 2.40±0.0037 centimeter. Chest, antero-posterior: Mean, 21.58 centimeters; standard deviation, 1.87±0.0029 centimeter. Correlation: 0.271±0.0024.

TABLE LXXXI.—Correlation between waist circumference and transverse pelvic diameter, while troops, demobilization.

Transverse pelvic diameter, in centimeters.	Total.	Waist circumference, in centimeters.											Mean waist circumference.	
		50-63	64-67	68-71	72-75	76-79	80-83	84-87	88-91	92-95	96-99	100-103		104 and over.
19 and under.....	25	2	3	20	2	1	1	7	2				1	Ow.
20.....	80	10	17	9	22	18	17	16	4					69.18
21.....	216	7	42	27	62	43	36	22	8					70.50
22.....	333	7	87	94	81	55	45	22	10			1		70.50
23.....	730	7	228	215	228	111	19	22	10			1		73.74
24.....	11,473	10	418	755	534	302	69	25	11			1		73.11
25.....	3,250	30	96	755	1,245	775	210	50	11			2		73.42
26.....	6,123	45	183	1,150	2,245	1,692	577	124	5					73.63
27.....	9,608	47	235	1,496	3,841	4,898	2,472	271	6					74.65
28.....	14,460	73	222	1,567	4,250	5,179	3,572	728	4					73.36
29.....	15,828	55	158	1,778	5,719	6,623	4,530	1,812	10					76.58
30.....	13,955	55	104	1,933	3,719	5,179	3,572	1,368	32					77.75
31.....	10,060	36	75	560	2,622	4,139	3,520	1,523	3					78.92
32.....	7,274	23	43	458	1,698	2,623	2,534	1,229	6					80.17
33.....	5,096	24	21	230	1,796	2,623	2,534	1,109	8					80.64
34.....	2,971	11	8	230	1,372	1,725	1,173	715	19					81.64
35.....	1,683	10	3	53	409	388	376	476	25					82.92
36.....	823	5	5	29	95	188	188	320	20					82.92
37.....	453	3	3	33	73	108	108	140	16					81.80
38.....	363	4	4	33	81	108	72	55	23					82.92
39.....	362	1	1	23	98	97	66	36	8					78.77
40 and over.....	492	4	7	26	93	142	106	61	23					79.01
Number measured.....	95,658	452	1,495	9,492	23,138	27,698	18,682	8,938	3,678	1,218	533	193	133	79.71
Not measured.....	499													
Total.....	96,157													
Mean transverse pelvis.....	cm.	28.59	27.58	28.04	28.63	29.35	30.06	30.76	31.41	32.06	32.47	33.51	33.96	

Waist circumference: Mean, 77.57 centimeters; standard deviation, 6.00±0.0063 centimeter. Transverse pelvis: Mean, 29.13 centimeters; standard deviation, 2.85±0.0044 centimeter. Correlation: 0.3510±0.0019.

TABLE LXXXII.—Correlation between length of arm and forearm, white troops, demobilization.

Forearm, in centimeters.	Total.	Arm length, in centimeters.																		Mean arm length.		
		60-61	62-63	64-65	66-67	68-69	70-71	72-73	74-75	76-77	78-79	80-81	82-83	84-85	86-87	88-89	90-91	92-93	94-95		96-97	98-99
20	85																					
21	117																					
22	259	2	3	4	5	3	12	8	17	13												
23	1,096	3	7	8	23	38	26	13	10	12	16	23	13	9	9	5	1	3	1		1	
24	3,745	1	8	24	60	151	222	35	26	19	67	33	12	15	24	5	5	3		1		
25	10,404	1	6	12	49	288	676	211	860	97	276	111	42	42	11	4	1	2				
26	18,146	2	1	12	49	320	1,035	2,036	2,513	2,081	3,379	550	251	98	48	16	16	1				
27	20,374	2	3	5	17	126	717	2,049	3,714	4,216	3,469	2,047	1,045	447	180	69	16	5				
28	15,194	1	3	5	11	16	261	915	2,393	4,133	4,871	3,610	2,188	1,158	510	228	36	16				
29	8,037				2	16	67	197	393	1,747	3,554	3,681	2,681	1,591	817	386	102	27				
30	3,301				2	4	53	161	361	1,404	2,946	1,965	1,300	717	565	352	128	50				
31	1,011				1	4	80	184	400	894	1,436	88	131	190	215	182	74	54				
32	240				2	4	3	4	3	11	25	88	29	10	50	59	38	22				
33	83				1	3	2	2	2	3	5	10	10	16	20	20	10	13				
34-35	368				5	3	24	5	50	56	72		49		42	31	12	18				
Number measured	82,492	10	30	99	254	997	3,108	6,382	10,703	13,373	14,532	12,200	8,934	5,630	3,315	1,949	566	274	126	70	40	
Not measured	13,832																					
Total	96,324																					
Mean forearm...cm.		24.20	24.33	24.22	24.24	24.59	25.12	25.53	26.00	26.46	26.91	27.41	27.83	28.25	28.58	28.93	29.34	29.77	29.79	28.94	28.50	

Arm length: Mean, 78.57 centimeters; standard deviation, 4.69 ± 0.0078 centimeter. Forearm length: Mean, 26.91 centimeters; standard deviation, 1.73 ± 0.0098 centimeter. Correlation: 0.587 ± 0.0018 .

TABLE LXXXIII.—Correlation between stature and sitting height, white troops, demobilization.

[illegible]

Height: Mean, 171.99 centimeters; standard deviation, 6.63 \pm 0.0102 centimeter. Sitting height: Mean, 90.39 centimeters; standard deviation, 3.51 \pm 0.0054 centimeter. Correlation: 0.6628 \pm 0.0012

TABLE LXXXIV.—*Correlation between stature*

Height: Mean, 171.99 centimeters; standard deviation, 6.66 ± 0.0102 centimeter. Span: Mean, 175.35 centimeters; standard deviation, 7.95 ± 0.0122 centimeter. Correlation: 0.7944 ± 0.0008 .

TABLE LXXXV.—*Correlation between stature*

Sternal notch, in centimeters.	Total.	Height, in centimeters.												
		148-149	150-151	152-153	154-155	156-157	158-159	160-161	162-163	164-165	166-167	168-169	170-171	172-173
120-121	31	2	6	2	1	1	5	2	3			4	2	2
122-123	102		12	23	16	4	6	4	11	8		1	6	4
124-125	228		5	35	82	52	7	6	3	11	3	3	1	4
126-127	545		4	33	109	183	120	27	8	9	13	17	5	2
128-129	1,198	1	3	14	57	217	444	296	77	7	19	19	14	5
130-131	2,627	1	9	5	28	130	494	931	652	194	57	51	28	20
132-133	4,377	2	2	5	13	25	189	763	1,602	1,158	370	96	55	22
134-135	7,018		1	6	18	22	52	298	1,188	2,456	1,894	680	150	50
136-137	9,401	1	1	4	11	29	20	115	385	1,642	3,217	2,728	947	189
138-139	11,767	3	1	1	8	34	37	93	125	511	1,769	3,841	3,623	1,348
140-141	13,234		2	1	2	6	12	47	52	145	453	1,754	4,230	4,009
142-143	12,721	2	1	3	6	4	10	22	51	68	134	426	1,781	3,898
144-145	11,563	4			4	4	5	14	35	72	89	166	563	1,805
146-147	8,466	1			7	6	9	7	1	31	76	105	195	391
148-149	5,779	4	1	1	4	10	5	11	21	5	27	59	122	146
150-151	3,534			1			4	7	5	9	11	9	38	29
152-153	2,017					1		2	3	8	6	3	5	8
154-155	1,023				1				5	13	6	4	4	12
156-157	545							1	1	5	10	7	6	5
158-159	245									2	6	11	7	4
160 and over	219						1	6	4	3	2	10	10	5
Number measured	96,439	21	47	136	373	710	1,422	2,663	4,222	6,267	8,161	9,994	11,691	11,547
Not measured	753													
Total	97,192													
Mean sternal notch		139.07	127.65	127.66	128.81	129.70	130.64	132.37	133.43	135.31	136.84	138.37	140.16	141.71
cm.														

Height: Mean, 171.99 centimeters; standard deviation, 6.66 ± 0.0102 centimeter. Sternal notch: Mean, 141.13 centimeters; standard deviation, 5.97 ± 0.0091 centimeter. Correlation: 0.8567 ± 0.0006 .

and span, white troops, demobilization.

Height, in centimeters.																Mean height.
174-175	176-177	178-179	180-181	182-183	184-185	186-187	188-189	190-191	192-193	194-195	196-197	198-199	200-201	202-203	204-210	Cm.
1		2	1	2	1		1									170.10
	1		1	1	1											158.39
3	2	2		1			1									158.57
4	1	2	1				1									159.10
3	2	1		1												159.79
4		2				1	1									160.78
15	6	4		1		2	2				1					161.94
34	13	7	4	1	1	4	4			1						163.31
66	38	12	6	1	2			1		1						164.36
180	55	33	4	6	7	6	2	1	2	1						165.85
420	167	96	16	4	7	7	5	2	4		2				1	167.19
835	383	153	30	12	7	10	7	7	3	1	1				1	168.65
1,406	685	282	81	18	9	7	7	1	8	4	1				1	169.98
1,982	1,189	530	157	57	11	10	11	1	6	1	1					171.23
1,983	1,687	963	365	98	33	6	9	7	3	2	3					172.61
1,547	1,624	1,314	629	215	103	41	19	8	4	4	3					173.96
1,059	1,334	1,243	931	415	151	64	29	6	5		3	2				175.47
715	928	908	825	551	228	101	39	11	3	3	1	3				176.90
339	554	710	728	574	343	145	58	14	3	3	1	2	1			177.95
156	295	434	514	455	317	189	73	17	5			1				179.18
52	123	233	291	328	271	174	112	36	10	1	3					180.40
13	43	98	137	198	184	139	87	45	14	3	6					182.00
11	26	39	55	95	113	100	72	42	23	7	2	2			1	183.09
	8	12	23	45	64	65	54	28	18	2	1	1				184.25
4	4	12	8	23	26	33	28	23	10	9	1	3			1	185.12
			4	4	7	13	10	7	12	4	2	2	1			185.09
				1		9	3	2	1	3						188.35
1								2	6	1	4				2	188.61
10,833	9,169	7,176	4,814	3,106	1,887	1,128	639	261	140	49	37	18	4	2	5	190.00
178.07	180.03	181.86	184.03	186.11	187.62	188.78	189.04	190.97	190.34	190.29	187.04	189.50	195.50	204.50	182.10	

and height of sternal notch, white troops, demobilization.

Height, in centimeters.																Mean height.
174-175	176-177	178-179	180-181	182-183	184-185	186-187	188-189	190-191	192-193	194-195	196-197	198-199	200-201	202-203	204-210	Cm.
1																159.72
3	5	1	1				1									159.24
3	3	6	2			1										157.14
1	6	4	6	1			3	1								159.90
13	5	3	4	4	4	2	1	1								159.58
28	13	2	6	6	10	7	2			1						161.36
63	67	34	5	6	4	9	7	1	1	3					1	163.34
66	61	30	14	7	7	11	11	1		2	1					165.40
292	109	91	33	16	5	11	10	4	3	5	2					167.24
1,734	416	186	81	41	16	16	8	6	13	1	1				2	169.28
3,943	1,760	436	74	39	24	18	8	9	8	4	1	1				173.42
3,318	3,555	1,709	342	73	39	28	12	5	8	4	8	1				175.32
1,035	2,355	2,613	1,216	256	71	34	39	7	4		1	3				177.42
241	657	1,540	1,726	885	208	59	33	10	7	1	1	3	1			179.08
46	115	377	913	1,062	490	153	45	8	8	2	1	2				181.32
12	27	72	343	497	627	310	77	11		1	2	1	1			183.14
8	8	20	16	145	281	307	148	35	5	2	2					184.58
5	1	4	11	32	78	129	163	70	13	3		1				185.78
1	2	3		15	11	20	61	61	37	2	2					186.32
2	7	16	6	8	12	9	13	30	35	20	11	5	2	1	1	184.81
10,818	9,172	7,148	4,799	3,095	1,887	1,126	642	260	142	47	36	19	6	2	6	
143.12	144.61	146.14	148.01	149.65	151.18	151.82	152.63	154.50	153.46	152.24	149.83	150.60	150.83	147.50	141.83	

TABLE LXXXVI.—Correlation between stature and height of pubic arch, white troops, demobilization.

Pubic arch, in centimeters.		Height, in centimeters.																				Mean height
		145-149	150-154	155-159	160-164	165-169	170-174	175-179	180-184	185-189	190-194	195-199	200-204	205-209	210-214	215-219	220-224	225-229	230-234	235-239	240-244	
70-74	285	2	11	8	20	10	31	33	36	41	44	55	57	61	67	74	83	94	107	124	144	167.08
75-79	480	1	12	18	27	36	44	55	67	80	94	107	124	144	167	194	224	259	304	359	424	165.58
80-84	770	1	4	24	75	94	100	107	114	124	136	149	167	194	224	259	304	359	424	499	584	163.98
85-89	1,581	2	4	26	98	155	238	348	504	714	994	1,364	1,844	2,444	3,194	4,044	5,044	6,244	7,644	9,244	11,044	129.08
90-94	3,021	2	4	26	102	187	307	474	714	1,044	1,494	2,044	2,744	3,544	4,444	5,444	6,544	7,744	9,044	10,444	11,944	135.67
95-99	6,276	3	5	14	29	56	94	144	214	304	414	544	694	864	1,044	1,244	1,444	1,644	1,844	2,044	2,244	137.42
100-104	12,631	3	3	16	34	70	124	194	284	394	514	644	794	944	1,114	1,294	1,474	1,644	1,814	1,984	2,144	139.46
105-109	14,969	3	3	16	34	70	124	194	284	394	514	644	794	944	1,114	1,294	1,474	1,644	1,814	1,984	2,144	141.50
110-114	14,228	3	3	16	34	70	124	194	284	394	514	644	794	944	1,114	1,294	1,474	1,644	1,814	1,984	2,144	143.54
115-119	14,060	3	3	16	34	70	124	194	284	394	514	644	794	944	1,114	1,294	1,474	1,644	1,814	1,984	2,144	145.58
120-124	11,051	2	1	4	9	14	20	31	44	59	74	89	104	119	134	149	164	179	194	209	224	147.62
125-129	11,217	2	1	4	9	14	20	31	44	59	74	89	104	119	134	149	164	179	194	209	224	149.66
130-134	4,273	1	1	3	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	151.70
135-139	2,218	1	1	3	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	153.74
140-144	1,203	1	1	3	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	155.78
145-149	85	1	1	3	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	157.82
150-154	40	1	1	3	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	159.86
Number measured	91,395	21	45	137	1,390	3,941	7,700	9,453	11,054	10,940	10,253	9,687	8,842	8,014	7,248	6,511	5,764	5,017	4,270	3,523	2,776	2,029
Not measured	5,827																					
Total	97,192																					
Mean cubic arch, ... cm.	85.2077	16.76.80																				

Height: Mean, 172.02 centimeters; standard deviation, 5.77 ± 0.011 centimeter. Pubic height: Mean, 86.83 centimeters; standard deviation, 5.06 ± 0.006 centimeter. Correlation: 0.6330 ± 0.0012.

TABLE LXXXV 11.—*Correlation between stature and sitting height, various groups, measurements.*

Sitting height, in centimeters.	Height, in centimeters.																			
	148- 149	150- 151	152- 153	154- 155	156- 157	158- 159	160- 161	162- 163	164- 165	166- 167	168- 169	170- 171	172- 173	174- 175	176- 177	178- 179	180- 181	182- 183	184- 185	186- 187
Total.	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
75-77	1	2	2	1	1	1	1	1	2	1	1	1	1	1	1	1	1	1	1	1
78-79	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
80-81	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
82-83	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
84-85	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
86-87	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
88-89	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
90-91	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
92-93	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
94-95	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
96-97	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
98-99	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
100-01	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
102-03	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
104-05	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
106-07	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Number measured...	6,433	2	9	13	23	55	87	159	314	462	562	706	743	745	586	470	313	207	138	74
Not measured...	60																			
Total	6,493																			

Height: Mean, 171.99 centimeters; standard deviation .690±0.041 centimeter. Sitting height: Mean, 87.35 centimeter; standard deviation, 3.48±0.021 centimeter. Correlation: 0.6088±0.0053.

TABLE LXXXIX.—Correlation between stature and height of sternal notch, colored troops, demobilization.

Height of sternal notch, in centimeters.	Stature, in centimeters.																			
	148-149	150-151	152-153	154-155	156-157	158-159	160-161	162-163	164-165	166-167	168-169	170-171	172-173	174-175	176-177	178-179	180-181	182-183	184-185	186-187
120-121.....	2																			
122-123.....	4																			
124-125.....	10	1	2	6	1	4			1	1				1						
126-127.....	27	1	5	7	8	4			1	2	2	1	1	1		1	1	1		
128-129.....	59	1	1	5	15	16	7	12	33	12	8	3	2	3	5	2	2	2		
130-131.....	95		2	5	28	21	28	77	147	52	11	17	2	2	5	1	1	1		
132-133.....	232	2			5	11	41	115	174	82	82	94	23	6	13	6	3	2	1	
134-135.....	393		1		2	1	17	72	152	78	203	278	233	175	46	13	4	2	2	1
136-137.....	533				1	2	1	24	78	203	278	233	233	302	160	44	10	5	7	1
138-139.....	669		1		1	1	4	5	29	70	210	278	233	302	160	44	10	5	7	1
140-141.....	810	1			1	1	1	4	5	29	70	210	278	302	160	44	10	5	7	1
142-143.....	877		1		1	1	1	2	4	5	29	70	210	302	160	44	10	5	7	1
144-145.....	962				1	1	1	1	3	7	4	27	74	153	208	134	34	4	2	1
146-147.....	648				2	1	2	1	1	4	4	24	69	153	208	134	34	4	2	1
148-149.....	496				2	2	1	1	1	4	2	6	16	44	120	166	99	26	4	4
150-151.....	293								1	1		1	3	12	18	73	91	66	16	5
152-153.....	193					1	1		1	1			2	1	2	19	54	58	40	12
154-155.....	133								1	1				1	2	3	16	31	41	27
156-157.....	67															3	17	19	12	8
158-159.....	29															1	1	3	5	2
160 and over.....	22															1	1	2	2	1
Number measured.....	6,454	2	9	13	23	56	88	161	318	466	565	661	704	749	532	469	315	207	138	75
Not measured.....	39																			
Total.....	6,493																			

Height: Mean, 171.97 centimeters; standard deviation, 6.91±0.0410 centimeter. Sternal notch: Mean, 142.39 centimeters; standard deviation, 6.05±0.0359 centimeter. Correlation: 0.8382±0.0022.

TABLE XC.—Correlation between stature and height of pubic arch, colored troops, demobilization.

Public height, in centimeters.	Total.	Stature, in centimeters.																									
		148-149	150-151	152-153	154-155	156-157	158-159	160-161	162-163	164-165	166-167	168-169	170-171	172-173	174-175	176-177	178-179	180-181	182-183	184-185	186-187	188-189	190-191	192-193	194-195	196-197	198-199
70-71	6						1	1	1		2	1							1								
72-73	18						2	5			2	1		2													
74-75	27	1					3	4			3	6															
76-77	47						6				3	7	6														
78-79	107						7				12	13	14	9	2	6	5										
80-81	229						12				27	36	29	21	14	8	4										
82-83	370						22				59	39	26	21	11	14	12	4	3	1			3		1		
84-85	558						12				67	59	42	25	21	14	10	8	5	2							
86-87	782						6				117	102	80	42	25	21	11	10	8	5	2						
88-89	939						1				131	145	120	88	46	33	11	10	8	5	2						
90-91	959						3				148	177	164	140	125	107	99	53	24	8	3		1	1			
92-93	800						1				164	177	164	140	125	107	99	53	24	8	3		1	1			
94-95	606										157	177	164	140	125	107	99	53	24	8	3		1	1			
96-97	406										140	125	107	99	53	24	8	3	1	4	1		2	1			
98-99	241										115	135	114	72	25	22	15	6	1	2	1		2	1			
100-101	92										34	59	75	78	50	37	47	36	22	14	17		7	5	3		
102-103	20										7	16	7	12	19	14	17	7	5	3	3		2	1			
104-105	13										2	2	3	2	2	3	3	3	3	3	3		3	1			
Number measured	6,220	2	7	12	21	53	85	157	308	443	545	633	684	725	717	561	455	302	202	134	74	44	24	17	10	3	2
Not measured	273																										
Total	6,493																										

Height: Mean, 171.98 centimeters; standard deviation, 6.88±0.142 centimeter. Height of pubic arch: Mean, 89.42 centimeters; standard deviation, 5.27±0.032 centimeter. Correlation: 0.6948±0.0044.

..... centimeters; mean knee height, 69.0±0.0435 centimeter. Correlation:

Knee height, in centimeters.	Height, in centimeters.																				Total.
	149-150-151-152-153-154-155-156-157-158-159-160-161-162-163-164-165-166-167-168-169-170-171-172-173-174-175-176-177-178-179-180-181-182-183-184-185-186-187-188-189-190-191-192-193-194-195-196-197-198-199																				
35 and under.....	16																				16
36-37.....	44																				44
38-39.....	74																				74
40-41.....	178																				178
42-43.....	403																				403
44-45.....	934																				934
46-47.....	1,404																				1,404
48-49.....	1,249																				1,249
50-51.....	835																				835
52-53.....	388																				388
54-55.....	151																				151
56-57.....	52																				52
58-59.....	21																				21
60-61.....	2																				2
62 and over.....																					
Number measured.....	5,725																				5,725
Not measured.....	768																				768
Total.....	6,493																				6,493

Height: Mean, 172.05 centimeters; standard deviation, 6.90±0.0435 centimeter. Knee height: Mean, 47.26 centimeters; standard deviation, 3.64±0.0229 centimeter. Correlation: 0.4763±0.0069.

TABLE XCII.—Correlation between leg length and knee height, colored troops, demobilization.

Knee height, in centimeters.	Leg length, in centimeters.												
	60-61	62-63	64-65	66-67	68-69	70-71	72-73	74-75	76-77	78-79	80-81	82-83	84-85
Total.	74	5	4	6	8	14	13	13	8	2	4	1	1
38-39	1	2	8	6	15	37	24	25	24	13	11	1	1
40-41	2	8	19	16	69	78	61	41	36	32	22	3	1
42-43	3	10	32	49	114	174	139	177	126	52	22	2	1
44-45	3	6	22	35	111	220	259	253	221	136	68	8	1
46-47	8	8	14	47	111	220	259	253	221	173	92	13	4
48-49	2	2	6	19	48	188	226	236	193	139	101	42	6
50-51	3	3	3	8	14	52	79	146	58	70	72	17	2
52-53	3	3	2	3	8	26	25	58	16	19	18	27	9
54-55	1	1	2	1	1	12	11	10	10	7	13	11	3
56-57	49	1	1	1	5	5	4	10	6	5	2	7	2
Number measured.	5,595	16	39	89	207	388	776	955	917	631	380	192	78
Not measured.	6,498												35
Total.	6,493												20

Leg length: Mean, 74.38 centimeters; standard deviation, 4.59±0.029 centimeter. Knee height: Mean, 47.32 centimeters; standard deviation, 3.37±0.0229 centimeter. Correlation: 0.4305±0.0073.

TABLE XCIII.—Correlation between chest circumference (rest) and weight, colored troops, demobilization.

Weight, in pounds.	Chest circumference, in centimeters.												
	68-73	74-77	78-81	82-85	86-89	90-93	94-97	98-101	102-109				
Total.	10	3	3	40	6			2	1				
100-109	79	1	0	25	55	11		1					
110-119	289	3	8	144	260	44		2					
120-129	602	2	2	283	586	153		1					
130-139	757	4	21	171	386	267		6					
140-149	735	1	14	75	321	204		10					
150-159	434	1	4	15	127	73		18					
160-169	228	1	2	7	28	91		16					
170-179	110	1	1	1	10	31		10					
180-189	52	1	1	1	1	7		5					
190-199	23					1		10					
200 and over.								80					
Number measured.	3,319	13	22	184	707	1,184	809	304	16				
Not measured.	3,036												
Total.	6,355												

Weight: Mean, 149.53 pounds; standard deviation, 17.53±0.045 pound. Chest circumference (rest): Mean, 88.14 centimeters; standard deviation, 4.79±0.040 centimeter. Correlation: 0.4519±0.0067.

TABLE XCIV.—Correlation between chest circumference (rest) and neck circumference, colored troops, demobilization.

Neck circumference, in centimeters.		Chest circumference, in centimeters.							
Total.	73 and under.	74-77	78-81	82-85	86-89	90-93	94-97	98-101	102-107
29.....	1		2	2	1	7	1		
30.....	16	1	3	6	3	4			
31.....	14	6	17	24	12	2			
32.....	61	15	42	69	38	10			
33.....	177	38	103	218	137	45	5	1	
34.....	519	8	87	400	403	146	17	3	
35.....	1,074	12	91	380	624	265	65	7	
36.....	1,474	9	32	194	589	423	132	22	
37.....	1,415	2	11	59	280	338	163	29	6
38.....	920	3	4	23	80	131	120	26	12
39.....	398	1		6	19	46	52	26	12
40.....	161			1	2	10	11	11	1
41.....	36					1	3	5	2
42.....	11								
43.....	2								
44.....	1								
Number measured.....	16	57	392	1,412	2,192	1,456	590	132	33
Not measured.....	75								
Total.....	6,355								

Chest circumference (rest): Mean, 87.97 centimeters; standard deviation, 4.84 ± 0.029 centimeter. Neck circumference: Mean, 36.37 centimeters; standard deviation, 1.72 ± 0.010 centimeter. Correlation: 0.5172 ± 0.0082 .

TABLE XCV.—Correlation between chest circumference (rest) and transverse pelvis, colored troops, demobilization.

Transverse pelvis, in centimeters.		Chest circumference, in centimeters.									
		68-73	74-77	78-81	82-85	86-89	90-93	94-97	98-101	102-109	
Total.											
20.	8										
21.	22			1	1	3	2	1			
22.	14			3	3	7	3		1		
23.	35			1	5	6					
24.	126	1	2	15	14	10	11	6			
25.	331	3	10	50	50	30	39	19			
26.	744	8	7	66	114	100	198	45	1		
27.	1,003	6	12	59	308	369	268	19	3		
28.	1,128	2	7	59	245	462	275	84	7		
29.	999	2	6	37	185	342	213	133	11		
30.	708	1	3	27	101	241	161	96	22		
31.	530	1	1	12	70	174	115	83	22		
32.	342			6	42	96	46	58	19		
33.	158			5	15	41	23	29	14		
34.	67			2	6	11	11	8	3		
35.	32			1	1	5	5	5			
36.	19				2	7	4	2			
37.	8				2	2	1	2			
38.	22			1	6	9	3	2			
39.	17				3	7	5	1			
40.	7				2	2	1	2			
41.	4					3		1			
42 and over.	22			1	3	7	6	2			
Number measured.	6,345	20	57	395	1,429	2,209	1,474	592	136	33	
Not measured.	10										
Total.	6,355										

Chest circumference (rest): Mean, 87.9 centimeters; standard deviation, 4.86 \pm 0.029 centimeter. Transverse pelvis: Mean, 28.54 centimeters; standard deviation, 2.64 \pm 0.016 centimeter. Correlation: 0.3297 \pm 0.0075.

TABLE X(VI).—Correlation between chest transverse and chest antero-posterior, colored troops, demobilization.

Chest, antero-posterior, in centimeters.	Chest, transverse, in centimeters.														
	18-19	20-21	22-23	24-25	26-27	28-29	30-31	32-33	34-35	36-37	38-39	40-41	42-43	44-45	
Total.															
14-15	3						1								
16-17	23				2	13	3								
18-19	777	1	6	4	8	330	150	36	2	1	1				
20-21	3,116		24	44	202	842	211	22	50	3	5	1		1	
22-23	2,090		9	98	575	1,318	704	237	50	9	9				
24-25	2,357			13	244	766	132	73	19				1	1	
26-27	50			5	26	100	17	9	8	1					
28-29	16			1	1	13	4	3							
30-31	13				1	3	1	3	1						
32-33	2			1			1	1							
34-35	3					1	1					1			
Numbered measured.	6,450	1	39	35	199	2,548	1,862	573	102	14	15	2	1	2	
Not measured.	43														
Total.	6,493														

Chest, transverse: Mean, 29.05 centimeters standard deviation, 2.26 ± 0.013 centimeter. Chest, antero-posterior: Mean, 21.21 centimeters; standard deviation, 1.74 ± 0.010 centimeter. Correlation: 0.2267 ± 0.0080 .

TABLE XCVII.—Correlation between waist circumference and transverse diameter of pelvis, colored troops, demobilization.

	Transverse pelvis, in centimeters.	Total.	Waist circumference, in centimeters.											
			63 and under.	64-67	68-71	72-75	76-79	80-83	84-87	88-91	92-95	96-99	100-109	
20.	5			1	2	2							
21.	18			4	2	4							
22.	91	2		1	5	9	1						
23.	14												
24.	35		5	13	8	7	2						
25.	126	3		33	43	33	4	1					
26.	337	2	10	94	129	83	14	1					
27.	752	3	19	137	259	235	62	2					
28.	1,017	3	11	104	342	374	147	1					
29.	1,146	5	7	82	267	449	244	2					
30.	1,014	5	6	43	169	339	252	16					
31.	720	3	2	26	116	238	182	6					
32.	339	4	1	18	62	151	160	5					
33.	343	1	1	12	39	87	86	9					
34.	158	1	1	3	15	32	47	12					
35.	68	1			3	14	15	10					
36.	32			2	5	5	2	3					
37.	20					1	1	1					
38.	6							3					
39 and over.	1					1		1	2	1			
Number measured.	6,354	33	71	576	1,499	2,099	1,247	533	205	59	36	26	
Not measured.	91												
Total.	6,445												

Waist circumference: Mean, 77.82 centimeters; standard deviation, 5.71±0.034 centimeter. Transverse pelvis: Mean, 28.42 centimeters; standard deviation, 2.35±0.014 centimeter. Correlation: 0.4456±0.0098.

TABLE XCVIII.—Correlation between arm length and forearm, colored troops, demobilization.

Forearm, in centimeters.	Arm length, in centimeters.															
	68-69	70-71	72-73	74-75	76-77	78-79	80-81	82-83	84-85	86-87	88-89	90-91	92-93	94-95	96-97	98-99
20.	13				1	1	1	2	3	1	1					
21.	12			3	5	1	1	1	1	2			3			
22.	5			1		1										
23.	17	2	2	3	1	1	2	1		1	1	2				
24.	54	4	13	10	9	6	12	1	1		1					
25.	235	6	54	51	42	40	59	30	6	8	3	1	1			
26.	613	3	72	116	151	138	176	120	52	22	9	4				
27.	1,071	2	26	141	206	267	301	214	123	62	30	4	1			
28.	1,208	4	15	57	133	265	311	245	163	121	82	17	8			
29.	1,060	3	6	12	54	136	211	138	159	117	63	24	19	2	2	1
30.	1,662	2	1	3	9	35	117	138	26	75	23	9	10	9	2	1
31.	317	1		1	2	2	28	43	58	22	19	7	8	4	1	
32.	115						3	12	4	6	6	3	1			
33.	55						1	2		1		4				
34.	18			1												
35.	8		2													
36.	8					3	2	1		1	2					
37.	9			3		3	1	3		1			1			
38.	15			1		1	2	1	6		2	2				
39.	11				1					1						
Number measured.	5,514	16	88	399	614	909	921	822	632	441	246	119	52	22	8	5
Not measured.	979															
Total.	6,493															

Arm length: Mean, 80.79 centimeters; standard deviation, 4.76±0.0306 centimeter. Forearm: Mean, 28.20 centimeter; standard deviation, 2.03±0.013 centimeter. Correlation: 0.5782±0.0060.

	42	75	585	1,603	4,947	10,800	18,745	23,286	18,666	10,704	4,441	1,528	277	40	11	8	90.42
Number measured.....	96,387																
Not measured.....	1,280																
Total.....	97,156																
Mean chest circumference..... cm.....	88.98	86.61	87.26	86.37	86.80	87.14	87.83	88.80	89.03	90.39	91.16	91.16	91.80	94.40	91.77	90.00	

Sitting height: Mean, 90.41 centimeters; standard deviation, 3.45 \pm 0.005 centimeter. Chest circumference (rest): 88.79 centimeters; standard deviation, 5.09 \pm 0.0078 centimeter. Correlation: 0.2415 \pm 0.0021 centimeter.

TABLE C.—Association between blouse groups and weight, white troops, demobilization.

Chest circumference (rest), in centimeters.																						
Weight, in pounds.																						
68-73	74-77	78-81 s.	78-81 m.	79-81 l.	82-85 s.	82-85 m.	82-85 l.	86-89 s.	86-89 m.	86-89 l.	90-93 s.	90-93 m.	90-93 l.	94-97 s.	94-97 m.	94-97 l.	98-101 s.	98-101 m.	98-101 l.	102-105	106-109	110 and over.
16	68	71	90	5	4	51	63	16	18	13	3	5	2	5	7	1	1	3	3	4	1	2
28	146	268	714	66	65	898	456	323	210	118	42	30	2	6	18	2	1	5	5	4	2	2
36.5	28	183	526	44	43	592	456	1,462	1,908	576	962	1,913	175	168	293	11	11	15	3	4	2	3
10.97	28	146	188	248	3,433	3,689	705	1,462	1,908	576	962	1,913	175	168	293	11	23	25	3	3	1	3
17.569	27	53	54	531	1,490	1,586	531	1,986	5,947	1,350	935	4,272	81	323	1,145	44	81	108	3	9	1	3
18.592	27	53	130	129	1,344	1,404	1,254	2,777	2,585	1,247	462	3,964	1,387	298	2,184	131	157	260	27	32	3	1
14.00	19	28	9	60	342	434	32	277	548	149	528	1,252	1,252	298	2,184	208	181	525	71	87	10	1
8.207	10	19	28	80	32	92	144	54	550	117	32	420	503	50	108	300	105	541	114	172	23	2
3.747	5	4	2	22	5	37	8	19	125	117	12	56	110	18	341	153	56	338	102	166	40	2
1.468	1	1	2	4	3	9	8	33	32	4	21	28	28	56	56	17	15	101	67	106	50	11
80-189	1	2	3	4	3	7	1	6	4	3	4	2	2	12	12	14	4	48	43	97	69	47
90-199	1	1	2	3	1	1	1	3	6	4	4	2	2	5	5	5	4	48	43	97	69	47
200 and over	348	1	1	2	1	1	1	3	6	4	4	2	2	5	5	5	4	48	43	97	69	47
79,706	160	536	641	835	4,034	9,863	1,749	5,055	16,496	4,010	2,936	12,879	4,274	1,053	7,399	1,011	644	1,969	438	632	201	72
16,168																						
Not measured																						
Total	95,874																					
Mean weight, lbs.	126.24	119.94	125.77	134.71	125.36	131.65	139.56	133.66	141.24	140.52	141.62	149.56	157.97	150.07	156.50	170.09	161.57	166.57	178.52	180.61	186.33	191.17
Blouse group designation	1	2 s.	2 m.	2 l.	3 s.	3 m.	3 l.	4 s.	4 m.	4 l.	5 s.	5 m.	5 l.	6 s.	6 m.	6 l.	7 s.	7 m.	7 l.	8	9	10

Weight: Mean, 144.67 pounds.

TABLE CII.—Association between blouse groups and chest transverse diameter, white troops, demobilization.

Chest, transverse, in centimeters.		Total.	Chest circumference (rest), in centimeters.																							
			68-73	74-77	78-81 s.	78-81 m.	78-81 l.	82-85 s.	82-85 m.	82-85 l.	86-89 s.	86-89 m.	86-89 l.	90-93 s.	90-93 m.	90-93 l.	94-97 s.	94-97 m.	94-97 l.	98-101 s.	98-101 m.	98-101 l.	102- 105	106- 109	110 and over.	
19.....	19	5	2	2	1	1	2	3	9	2	1	1	1	2	2	1	15	1	1	1	1	1	1	1	1	
20.....	154	1	2	2	7	31	7	3	19	7	31	0	1	20	11	7	15	1	1	8	3	5	3	3	3	
21.....	305	2	4	2	16	64	15	8	42	13	64	13	8	47	13	7	24	4	1	8	3	5	3	3	3	
22.....	251	1	4	2	10	47	12	28	17	13	47	13	8	43	18	3	24	4	3	5	3	5	3	3	3	
23.....	311	1	14	10	32	111	17	51	17	20	46	20	10	36	6	3	21	1	1	6	2	1	1	1	1	
24.....	582	0	58	40	118	24	32	131	17	18	75	18	9	37	11	4	30	3	4	4	4	2	1	1	1	
25.....	682	28	128	107	385	82	166	665	174	361	361	71	32	64	25	5	20	7	9	15	4	2	1	1	1	
26.....	631	24	134	179	733	169	342	1,070	552	1,190	206	206	114	173	74	17	92	21	17	31	8	6	1	1	1	
27.....	1,915	19	101	179	782	222	510	2,699	923	4,831	550	550	201	1,143	267	50	242	78	48	110	19	8	3	1	1	
28.....	1,469	28	75	131	590	181	474	2,786	1,152	4,527	977	977	656	2,527	730	144	762	177	89	251	56	30	5	1	1	
29.....	18,618	30	45	64	328	135	256	2,012	918	1,281	3,241	1,281	739	3,611	1,172	275	1,954	254	135	431	81	32	24	5	1	
30.....	15,596	24	38	24	101	36	153	1,052	603	826	3,241	862	739	3,611	1,172	275	1,954	254	135	431	81	32	24	5	1	
31.....	10,232	12	22	8	23	22	60	1,454	236	390	1,701	479	409	2,238	1,140	458	123	176	140	234	68	24	6	1	1	
32.....	5,705	13	4	2	16	3	23	174	120	142	709	242	238	1,136	207	73	609	115	78	234	68	24	6	1	1	
33.....	2,668	1	3	2	16	3	23	174	120	142	709	242	238	1,136	207	73	609	115	78	234	68	24	6	1	1	
34.....	1,204	1	3	2	16	3	23	174	120	142	709	242	238	1,136	207	73	609	115	78	234	68	24	6	1	1	
35.....	566	1	3	2	16	3	23	174	120	142	709	242	238	1,136	207	73	609	115	78	234	68	24	6	1	1	
36.....	320	1	3	2	16	3	23	174	120	142	709	242	238	1,136	207	73	609	115	78	234	68	24	6	1	1	
37.....	206	1	3	2	16	3	23	174	120	142	709	242	238	1,136	207	73	609	115	78	234	68	24	6	1	1	
38.....	160	1	3	2	16	3	23	174	120	142	709	242	238	1,136	207	73	609	115	78	234	68	24	6	1	1	
39.....	147	1	3	2	16	3	23	174	120	142	709	242	238	1,136	207	73	609	115	78	234	68	24	6	1	1	
40.....	55	1	3	2	16	3	23	174	120	142	709	242	238	1,136	207	73	609	115	78	234	68	24	6	1	1	
41.....	29	1	3	2	16	3	23	174	120	142	709	242	238	1,136	207	73	609	115	78	234	68	24	6	1	1	
42 and over.....	115	1	3	2	16	3	23	174	120	142	709	242	238	1,136	207	73	609	115	78	234	68	24	6	1	1	
Number measured.....	95,590	195	647	758	3,331	992	2,059	11,961	4,367	5,956	19,924	4,797	3,406	15,523	5,154	1,217	8,765	1,215	774	2,324	514	905	223	82	82	
Not measured.....	284																									
Total.....	95,874																									
Mean chest transverse, cm.....		29.96	26.77	27.14	27.62	27.52	27.84	28.20	28.46	28.76	28.05	28.35	29.35	29.53	29.83	30.00	30.35	30.80	30.87	31.18	31.25	32.17	33.17	34.32	34.32	
Blouse group designa- tion.....		1	2 s.	2 m.	2 l.	3 s.	3 m.	3 l.	4 s.	4 m.	4 l.	5 s.	5 m.	5 l.	6 s.	6 m.	6 l.	7 s.	7 m.	7 l.	8	9	10	10	10	

Chest, transverse: Mean, 26.01 centimeters.

TABLE 'CIII.- Association between blouse groups and chest diameter, antero-posterior, white troops, demobilization.

Chest antero-posterior. In centimeters.		Chest circumference (rest), in centimeters.																Total.	110 and over.			
		66-73	74-77	78-81	82-85	82-85	82-85	86-89	86-89	90-93	90-93	94-97	94-97	98-101	98-101	102-105	106-109					
14.....	6						1	2										1	1			
15.....	16	2	12	1	2	1	1	3														
16.....	57	6	40	18	58	20	11	63	30	14	32	6	9	1	1	1	1					
17.....	317	19	89	73	287	65	101	457	141	84	243	44	44	19	39	18	1	2	2			
18.....	1,707	39	177	150	1,016	192	345	1,201	453	291	1,336	291	291	86	361	113	6	8	4			
19.....	6,829	29	183	225	1,047	262	635	3,542	1,382	1,396	4,063	865	865	422	1,568	473	18	52	4			
20.....	16,628	29	210	235	1,047	262	635	3,542	1,382	1,396	4,063	865	865	422	1,568	473	18	52	4			
21.....	22,573	42	276	335	1,354	3,372	541	3,372	1,354	1,945	6,168	1,364	1,364	903	3,996	1,135	45	145	37	24	3	
22.....	22,187	22	45	61	285	1,077	298	1,876	850	1,397	4,979	1,268	1,268	1,071	4,722	1,569	309	346	74	61	6	
23.....	13,511	21	27	15	94	69	84	578	302	367	2,125	621	621	646	3,113	1,087	352	396	141	117	11	
24.....	6,244	5	10	3	32	20	16	161	106	149	625	222	222	249	1,342	488	169	145	118	190	25	2
25.....	2,461	3	4	3	42	31	43	152	62	62	163	62	62	62	163	163	169	145	118	190	25	2
26.....	396	2		1	13	6	3	25	12	14	64	16	16	19	84	44	11	150	23	44	11	10
27.....	220	2		3	6	3	6	21	6	8	34	10	11	34	44	44	11	150	23	44	11	10
28.....	202	1		1	6	1	3	9	10	8	30	11	11	30	44	44	11	150	23	44	11	10
29.....	119			2	6	2	6	26	7	9	28	8	8	18	12	12	6	23	3	6	13	4
30.....	57			2	2	2	2	14	3	7	20	5	5	7	8	8	1	19	3	5	11	5
31.....	85			1	1	1	1	8	1	3	13	7	7	3	8	8	1	19	3	5	11	5
32.....	58				3	3	3	1	5	6	14	7	7	3	9	9	5	13	3	6	13	4
33.....	16				3	3	3	10	2	2	10	2	2	2	2	2	2	14	3	6	13	4
34.....	29				3	3	3	10	2	2	10	2	2	2	2	2	2	14	3	6	13	4
35 and over.....	29				3	3	3	10	2	2	10	2	2	2	2	2	2	14	3	6	13	4
Number measured.....	95,590	183	640	756	3,323	996	2,063	11,937	4,896	5,963	10,924	4,803	3,511	15,528	5,157	1,217	774	2,319	515	806	223	82
Not measured.....	95,284																					
Total.....	95,874																					
Mean chest antero-posterior diameter, cm.....	19.95	19.93	20.08	20.50	20.54	20.65	20.80	20.82	21.30	21.46	21.88	21.97	22.13	22.56	22.73	22.94	23.45	23.46	23.67	24.48	25.34	26.43
Blouse group designation.....	1	2 s.	2 m.	2 l.	3 s.	3 m.	3 l.	4 s.	4 m.	4 l.	5 s.	5 m.	5 l.	6 s.	6 m.	6 l.	7 s.	7 m.	7 l.	8	9	10

Chest antero-posterior diameter: Mean, 21.57 centimeters.

Chest antero-posterior diameter: Mean, 21.57 centimeters.

TABLE CIV.—Association between blouse groups and transverse pelvic diameter, white troops, demobilization.

Chest circumference (rest), in centimeters.																							
Transverse pelvic diameter, in centimeters.	68-73	74-77	78-81 s.	78-81 m.	78-81 l.	82-85 s.	82-85 m.	82-85 l.	86-89 s.	86-89 m.	86-89 l.	90-93 s.	90-93 m.	90-93 l.	94-97 s.	94-97 m.	94-97 l.	98-101 s.	98-101 m.	98-101 l.	102-105	106-109	110 and over.
Total.	30	3	2	1	1	4	21	1	1	13	2	5	11	2	1	7	1	1	3	1	1	1	1
19 and under.	80	5	2	1	1	4	21	1	1	13	2	5	11	2	1	7	1	1	3	1	1	1	1
20.	216	3	6	3	3	6	42	6	8	46	8	15	30	2	2	12	1	1	5	2	2	1	1
21.	335	2	6	19	14	13	63	22	9	64	11	36	36	2	2	12	1	1	5	2	2	1	1
22.	335	2	6	19	14	13	63	22	9	64	11	36	36	2	2	12	1	1	5	2	2	1	1
23.	724	6	26	24	40	167	40	167	40	124	28	14	52	31	4	26	4	7	4	3	4	1	1
24.	1,480	5	24	45	30	240	77	714	240	328	62	26	124	31	4	40	10	12	4	1	3	1	1
25.	3,235	8	50	73	45	175	106	106	272	632	122	96	269	66	10	86	9	10	12	4	1	1	1
26.	6,112	19	98	124	83	363	363	363	559	1,310	238	219	704	144	43	186	9	10	12	4	1	1	1
27.	9,388	31	201	402	97	517	517	517	881	2,166	368	378	2,222	260	78	419	39	31	55	25	21	4	2
28.	15,419	51	321	609	154	841	841	841	1,200	3,068	545	621	2,215	240	144	868	78	40	126	25	21	4	2
29.	21,819	81	452	823	207	1,071	1,071	1,071	1,369	3,531	785	662	2,907	765	232	1,041	136	103	245	54	48	4	2
30.	28,885	13	65	77	149	257	257	257	369	3,725	750	532	2,654	893	210	1,557	327	133	377	69	90	15	6
31.	37,243	28	112	112	161	321	321	321	435	4,351	901	574	3,250	567	754	165	327	162	377	67	147	20	7
32.	47,093	48	127	177	200	384	384	384	511	5,111	1,011	683	4,350	567	754	165	327	162	377	67	147	20	7
33.	57,093	6	23	106	106	313	313	313	429	4,299	849	1,011	5,111	567	754	165	327	162	377	67	147	20	7
34.	67,093	4	23	118	118	324	324	324	429	4,299	849	1,011	5,111	567	754	165	327	162	377	67	147	20	7
35.	77,093	3	6	4	4	42	42	42	58	5,589	1,118	1,118	6,702	567	754	165	327	162	377	67	147	20	7
36.	87,093	3	6	4	4	42	42	42	58	5,589	1,118	1,118	6,702	567	754	165	327	162	377	67	147	20	7
37.	97,093	1	1	11	11	44	44	44	60	6,003	1,200	1,200	7,203	567	754	165	327	162	377	67	147	20	7
38.	107,093	1	1	11	11	44	44	44	60	6,003	1,200	1,200	7,203	567	754	165	327	162	377	67	147	20	7
39.	117,093	1	1	11	11	44	44	44	60	6,003	1,200	1,200	7,203	567	754	165	327	162	377	67	147	20	7
40.	127,093	1	1	11	11	44	44	44	60	6,003	1,200	1,200	7,203	567	754	165	327	162	377	67	147	20	7
41.	137,093	1	1	11	11	44	44	44	60	6,003	1,200	1,200	7,203	567	754	165	327	162	377	67	147	20	7
42 and over.	331	1	2	13	2	8	27	6	23	66	11	16	55	11	5	44	5	4	17	3	6	2	2
Number measured.	95,479	643	757	3,325	991	2,061	11,948	4,859	5,953	19,902	4,796	3,508	15,503	5,149	1,217	8,756	1,210	772	2,320	514	805	223	82
Not measured.																							
Total.	95,479	643	757	3,325	991	2,061	11,948	4,859	5,953	19,902	4,796	3,508	15,503	5,149	1,217	8,756	1,210	772	2,320	514	805	223	82
Mean transverse pelvic diameter, cm.	27.95	27.44	27.92	28.66	27.96	28.39	28.07	28.64	28.23	29.22	28.24	29.78	30.47	30.14	30.59	31.27	30.98	31.51	31.67	31.95	33.41	34.68	
Blouse group designation.	1	2 a.	2 m.	2 l.	3 a.	3 m.	3 l.	4 a.	4 m.	4 l.	5 a.	5 m.	5 l.	6 a.	6 m.	6 l.	7 a.	7 m.	7 l.	8	9	10	

Transverse pelvic diameter: Mean, 29.45 centimeters.

TABLE CV.—Association between blouse groups and neck circumference, white troops, demobilization.

Neck circumference, in centimeters.	Total.	Chest circumference (rest), in centimeters.																				110 and over.		
		68-73	74-77	78-81 s.	78-81 m.	78-81 l.	82-85 s.	82-85 m.	82-85 l.	86-89 s.	86-89 m.	86-89 l.	90-93 s.	90-93 m.	90-93 l.	94-97 s.	94-97 m.	94-97 l.	98-101 s.	98-101 m.	98-101 l.		102-105	106-109
26 and under.	151	1	4			2																		
29	55	2	5			6																		
30	219	7	31	1	20	17	2	16	15	1	1	1	4	12	17	1	23	6	1	4	2	3	1	
31	314	13	93	16	40	52	9	52	19	35	38	33	3	33	13	5	17	6	6	6	2	2	1	
32	1,133	21	133	59	230	17	73	278	67	41	41	6	6	22	6	3	13	3	2	2	2	2	2	
33	4,286	43	383	145	566	126	225	1,207	364	298	718	134	7	36	6	3	13	3	2	2	2	2	2	
34	11,353	29	124	230	894	230	512	2,899	996	2,581	2,581	146	58	152	37	6	38	6	3	9	3	10	2	
35	20,094	28	116	177	827	240	593	3,451	1,447	5,179	2,581	474	221	867	721	27	168	10	7	29	4	10	2	
36	22,628	26	48	89	436	145	371	2,390	1,116	5,057	3,524	1,203	691	2,732	1,418	109	681	63	47	88	21	17	2	
37	18,047	21	26	26	170	103	180	1,085	553	3,466	1,433	398	444	2,000	1,450	252	1,719	205	112	251	45	36	4	
38	10,051	11	15	8	64	40	57	1,370	184	3,466	1,433	398	444	2,000	1,450	252	1,719	205	112	251	45	36	4	
39	4,426	4	3	1	31	12	2	118	73	124	401	126	150	803	310	113	1,051	158	149	149	104	104	14	
40	1,716	1	1	2	4	5	2	21	10	30	100	38	48	230	106	43	381	75	84	254	64	100	46	
41	492							4		7	10	8	9	111	2	3	14	20	28	72	20	75	24	
42	147							2			3	1	1	4	4	1	3	2	1	6	4	8	5	
43	52							1			2	3	1	1	1	1	1	2	2	2	1	4	1	
44	23							1			1	3	1	1	1	1	1	1	1	1	1	1	1	
45	22							1			1	3	1	1	1	1	1	1	1	1	1	1	1	
46	17							1			1	3	1	1	1	1	1	1	1	1	1	1	1	
47	16							1			1	3	1	1	1	1	1	1	1	1	1	1	1	
48	5							1			1	3	1	1	1	1	1	1	1	1	1	1	1	
49	6							1			1	3	1	1	1	1	1	1	1	1	1	1	1	
50	18							1			1	3	1	1	1	1	1	1	1	1	1	1	1	
Number measured.	95,271	166	599	754	3,311	981	2,053	11,946	4,858	5,927	19,890	4,708	3,494	15,480	5,146	1,211	8,737	1,215	772	2,315	513	801	223	81
Not measured.	603																							
Total.	95,874																							
Mean neck circumference, cm.		34.36		34.22	34.45	34.80	34.85	34.98	35.20	35.60	35.72	35.89	36.30	36.44	36.61	37.08	37.19	37.35	37.81	37.89	37.97	38.82	39.71	40.77
Blouse group designation.		1		2 s.	2 m.	2 l.	3 s.	3 m.	3 l.	4 s.	4 m.	4 l.	5 s.	5 m.	5 l.	6 s.	6 m.	6 l.	7 s.	7 m.	7 l.	8	9	10

Neck circumference: Mean, 35.96 centimeters.

TABLE CVI.—Association between blouse groups and total arm length, white troops, demobilization.

Arm length, in centimeters.	Chest circumference (rest), in centimeters.																Total.
	68-73	74-77	78-81	78-81	82-85	82-85	86-89	86-89	90-93	90-93	94-97	94-97	98-101	98-101	102-105	106-109	110 and over.
68-65.....	8	9	26	31	34	36	67	37	12	13	2	7	2	3	1	4	1
66-67.....	310	8	64	107	263	125	180	169	66	63	11	24	48	21	2	7	1
68-69.....	1,210	25	113	287	774	271	501	1,592	206	349	132	318	132	51	4	25	4
70-71.....	3,700	12	130	453	1,516	374	845	2,970	409	832	239	737	239	83	15	47	14
72-73.....	7,020	47	152	660	2,218	591	1,115	3,670	578	1,800	331	1,199	331	129	28	174	29
74-75.....	12,707	22	116	500	2,324	822	1,027	3,823	656	2,646	559	1,634	559	156	373	26	10
76-77.....	15,710	32	105	500	2,324	822	1,027	3,823	656	2,646	559	1,634	559	156	373	26	10
78-79.....	16,910	31	82	69	1,931	577	949	3,932	823	2,619	679	1,634	679	116	414	41	15
80-81.....	14,112	17	37	17	1,328	555	330	1,973	250	1,116	538	1,116	538	92	401	127	12
82-83.....	9,985	14	37	17	791	313	155	1,121	124	1,116	538	1,116	538	92	401	127	15
84-85.....	6,184	8	6	84	588	179	151	1,121	124	1,116	538	1,116	538	92	401	127	15
86-87.....	3,521	7	3	12	179	151	151	1,121	124	1,116	538	1,116	538	92	401	127	7
88-89.....	1,896	7	3	12	179	151	151	1,121	124	1,116	538	1,116	538	92	401	127	7
90-91.....	242	1	1	7	68	33	33	251	33	325	34	34	34	34	42	13	2
92-93.....	103	1	1	7	12	15	11	251	33	325	34	34	34	34	42	13	2
94-95.....	113	1	1	7	12	15	11	251	33	325	34	34	34	34	42	13	2
96 and over.....	113	1	1	7	12	15	11	251	33	325	34	34	34	34	42	13	2
Number measured..	94,940	617	741	3,264	11,858	4,839	5,896	19,831	4,783	15,440	5,134	8,727	1,208	771	2,303	510	80
Not measured.....	834																
Total.....	95,874																
Mean arm length.....																	
Blouse group designation.....	76.78	74.19	76.09	78.14	76.67	78.65	76.19	78.17	80.13	77.13	77.85	76.91	82.72	78.94	82.77	81.64	81.65
	1	2 s.	2 m.	2 l.	3 m.	3 l.	4 s.	4 m.	4 l.	5 s.	6 m.	6 l.	7 s.	7 m.	7 l.	8	10

Arm length: Mean, 76.42 centimeters; standard deviation, 4.58±0.0013 centimeter.

TABLE CVII.—*Correlation between chest circumference (rest) and sitting height, colored troops, demobilization.*

[Basis of construction of blouse groups shown by heavy lines; circle symbols are the "blouse" group designations. For relative frequency of "groups" see Table 121.]

		Sitting height, in centimeters.													Total.
		70-77	78-79	80-81	82-83	84-85	86-87	88-89	90-91	92-93	94-95	96-97	98-99		
68-69	2	①			2										
70-71	7					2	3		1						
72-73	11			1			3	3		2					
74-75	16	②		3	2	4	2	3	2						
76-77	41			6	7	12	10	6	1						
78-79	110		②	1	6	14	31	31	18	7	1				
80-81	286		4	23	42	73	70	37	22	8	6				
82-83	504	③	1	7	69	111	131	84	44	20	6	2	1		
84-85	927		2	8	35	118	193	229	187	96	37	17	4	1	
86-87	1,113		③	1	8	44	97	232	226	148	56	18	5	3	
88-89	1,088		3	7	30	69	266	249	172	85	17	8	2		
90-91	836	④		4	10	22	118	199	208	81	20	11	3		
92-93	643		1	1	6	26	76	153	168	123	55	23	9	3	
94-95	388														
96-97	206	⑤		1	7	11	32	74	86	58	25	11	1		
98-99	98					5	24	30	46	52	28	15	2	3	
100-01	44		⑤		1	2	6	17	25	14	13	9	4	2	
102-03	16				1	4	7	6	11	8	5	2			
104-05	8	⑥				2		2	3	5	2	2			
106-07	3							1	2	1	1	1			
108-09	4					1				2					
Number measured	6,345	9	42	198	513	1,108	1,500	1,354	924	480	164	60	20		
Not measured	90														
Total	6,445														

Chest circumference (rest): Mean, 87.99 centimeters; standard deviation, 4.76±0.0283 centimeter. Sitting height: Mean, 87.35 centimeters; standard deviation, 3.43±0.0203 centimeter. Correlation: 0.3012±0.0077.

TABLE CVIII.—Association between blouse groups and weight, colored troops, demobilization.

Chest circumference (rest), in centimeters.																							
Weight, in pounds.		68-73	74-77	78-81 s.	78-81 m.	78-81 l.	82-85 s.	82-85 m.	82-85 l.	86-89 s.	86-89 m.	86-89 l.	90-93 s.	90-93 m.	90-93 l.	94-97 s.	94-97 m.	94-97 l.	98-101 s.	98-101 m.	98-101 l.	102-105	106-109
100-109	10	1	3	2	1	1	16	23	1	1	3	2	2	10	1	2	1	2	1	1	1	1	1
110-119	79	1	6	8	16	8	45	88	11	15	39	1	13	10	1	2	1	1	1	1	1	1	1
120-129	289	3	8	13	43	9	120	198	28	45	194	11	25	114	2	2	5	4	1	1	1	1	1
130-139	602	2	2	7	34	6	130	338	48	64	332	22	25	114	14	10	10	3	1	1	1	1	1
140-149	757	4	4	4	15	4	140	449	116	47	32	51	33	197	37	15	32	3	2	4	1	1	1
150-159	735	1	1	1	6	7	5	5	41	29	13	257	51	33	197	37	15	32	3	2	4	1	1
160-169	434	1	1	1	3	1	1	3	12	6	107	14	14	159	31	12	46	15	2	8	1	1	1
170-179	228	1	1	1	3	2	1	3	3	2	17	9	2	64	25	13	42	26	6	10	2	2	1
180-189	110	1	1	1	1	1	1	1	1	1	5	5	2	15	14	2	28	19	11	14	3	5	4
190-199	52	1	1	1	1	1	1	1	1	1	1	1	1	6	1	1	2	2	7	3	4	3	3
200 and over	23	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Number measured	3,319	13	22	35	119	30	113	461	133	114	954	116	89	594	126	53	170	81	11	57	12	12	4
Not measured	3,036																						
Total	6,355																						
Mean weight, lbs.	135.27	128.77	128.21	131.47	140.50	129.63	136.34	145.85	140.38	147.24	154.16	151.47	157.09	163.31	164.31	167.50	175.61	164.50	181.87	176.50	183.25	183.25	183.25
Blouse group designation	1	2	3 s.	3 m.	3 l.	4 s.	4 m.	4 l.	5 s.	5 m.	5 l.	6 s.	6 m.	6 l.	7 s.	7 m.	7 l.	8 s.	8 m.	8 l.	9	9	9

Weight: Mean, 149.50 pounds.

TABLE 'IX.—Association between blouse groups and shoulder width, colored troops, demobilization.

Chest circumference (rest), in centimeters.																						
Shoulder width, in centimeters.	Chest circumference (rest), in centimeters.																					
	66-73	74-77	78-81 s.	78-81 m.	78-81 l.	82-85 s.	82-85 m.	82-85 l.	86-89 s.	86-89 m.	86-89 l.	90-93 s.	90-93 m.	90-93 l.	94-97 s.	94-97 m.	94-97 l.	98-101 s.	98-101 m.	98-101 l.	102-105	106-109
34.....	3	2	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
35.....	7	4	1	2	2	1	4	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
36.....	11	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
37.....	23	3	11	21	3	6	14	1	4	9	1	2	4	1	3	6	2	1	1	1	1	1
38.....	74	2	10	33	5	28	49	9	11	37	7	6	28	4	3	8	3	1	1	1	1	1
39.....	201	4	14	41	7	44	114	23	30	100	19	8	52	16	1	18	3	2	3	3	3	3
40.....	455	1	10	17	8	66	207	47	45	245	32	27	130	15	16	18	3	2	3	3	3	3
41.....	813	3	15	58	8	66	207	46	61	375	51	42	263	45	15	12	12	7	10	7	1	1
42.....	1,121	1	3	10	25	36	160	47	44	397	40	39	250	46	21	74	24	2	21	3	3	3
43.....	1,211	1	3	10	25	36	160	47	44	397	40	39	250	46	21	74	24	2	21	3	3	3
44.....	1,046	2	2	19	2	14	87	28	29	350	17	27	178	33	15	68	24	2	18	5	10	1
45.....	632	4	3	2	2	5	39	14	21	154	13	18	100	25	14	41	21	2	18	5	10	1
46.....	383	1	1	2	1	2	12	8	7	49	6	3	40	14	14	10	7	1	9	4	5	2
47.....	104	1	1	1	1	1	1	1	2	15	3	1	12	3	5	5	4	1	6	1	3	1
48.....	170	1	1	1	1	1	1	1	1	2	1	1	1	2	3	1	1	1	1	1	4	4
49.....	32	1	1	1	1	1	1	1	1	2	1	1	1	2	5	1	1	1	1	1	1	1
50 and over.....	13	1	1	1	1	1	1	1	1	1	1	1	1	2	1	1	1	1	1	1	1	1
Number measured.....	0, 299	56	88	257	45	264	930	225	257	1,742	191	172	1,089	204	107	334	141	20	94	22	25	8
Not measured.....	66	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Total.....	6, 355	66	88	257	45	264	930	225	257	1,742	191	172	1,089	204	107	334	141	20	94	22	25	8
Mean shoulder width.....	41.72	40.41	40.44	40.95	41.29	41.26	41.82	42.30	42.17	42.73	43.30	43.52	43.72	44.01	44.26	44.74	45.35	44.25	45.67	46.27	47.55	47.55
Blouse group designation.....	1	2	3 s.	3 m.	3 l.	4 s.	4 m.	4 l.	5 s.	5 m.	5 l.	6 s.	6 m.	6 l.	7 s.	7 m.	7 l.	8 s.	8 m.	8 l.	9	9

Shoulder width: Mean, 42.99 centimeters; standard deviation, 2.15 ± 0.030 centimeter.

TABLE CX.—Association between blouse groups and transverse diameter of chest, colored troops, demobilization.

Transverse chest, in centimeters.	Chest circumference (rest), in centimeters.																				
	68-73	74-77	78-81 s. m.	78-81 l.	82-85 s. m.	82-85 l.	86-89 s. m.	86-89 l.	90-93 s. m.	90-93 l.	94-97 s. m.	94-97 l.	98-101 s. m.	98-101 l.	102-105	106-109					
Total.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1					
19.....	15	2	1	1	2	3	4	10	1	4	2	1	3	1	1	1					
20.....	24	2	1	1	3	1	3	7	1	4	1	3	1	1	1	1					
21.....	16	2	1	1	1	1	7	3	1	1	2	1	1	1	1	1					
22.....	21	2	2	2	1	1	9	1	1	1	1	2	1	1	1	1					
23.....	24	2	2	2	1	1	2	2	1	1	1	1	1	1	1	1					
24.....	64	2	5	9	4	1	2	2	1	1	1	1	1	1	1	1					
25.....	134	4	12	19	13	41	3	20	2	23	2	1	1	1	1	1					
26.....	359	3	12	19	88	17	75	5	6	50	2	8	2	1	1	1					
27.....	688	3	11	18	64	178	34	390	37	133	10	35	2	1	1	1					
28.....	1,211	5	7	23	57	248	66	416	48	260	35	16	3	10	2	2					
29.....	1,295	5	7	10	29	43	54	416	48	260	35	16	3	10	2	2					
30.....	1,109	1	3	2	17	4	348	41	34	256	51	25	33	1	15	2					
31.....	1,708	2	8	4	30	99	31	172	14	202	41	25	72	4	2	4					
32.....	404	1	3	1	15	8	16	78	13	95	25	15	60	4	4	4					
33.....	164	1	3	4	2	5	35	4	6	28	10	7	18	4	7	1					
34.....	77	1	3	4	8	1	11	3	1	15	3	9	6	3	5	2					
35.....	19	1	3	4	1	1	2	1	1	4	1	2	5	1	2	1					
36.....	10	1	3	4	1	1	1	1	1	1	1	1	1	1	1	1					
37.....	6	1	2	2	2	1	2	2	1	1	1	1	1	1	1	1					
38.....	9	1	2	2	1	1	3	3	1	1	1	1	1	1	1	1					
39.....	9	1	2	2	1	1	3	3	1	1	1	1	1	1	1	1					
40 and over.....	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1					
Number measured.....	6,339	20	57	91	258	46	267	934	228	1,757	193	338	143	20	93	22					
Not measured.....	16														24	8					
Total.....	6,355																				
Mean chest circum- ference.....cm.	27.70	26.44	26.88	27.31	27.54	27.84	28.13	28.54	28.48	29.08	29.67	30.44	30.74	31.60	31.49	31.91					
Blouse group design- ation.....	1	2	3 s.	3 m.	3 l.	4 s.	4 m.	4 l.	5 s.	5 m.	6 s.	7 s.	7 m.	8 s.	8 m.	8 l.					

Chest circumference: Mean, 26.01 centimeters.

TABLE CXII.—Association between blouse groups and transverse pelvic diameter, colored troops, demobilization.

Transverse pelvis, in centimeters.		Chest circumference (rest), in centimeters.																				
		68-73	74-77	78-81 s.	78-81 m.	79-81 l.	82-85 s.	82-85 m.	82-85 l.	86-89 s.	86-89 m.	86-89 l.	90-93 s.	90-93 m.	90-93 l.	94-97 s.	94-97 m.	94-97 l.	98-101 s.	98-101 m.	98-101 l.	102-105 106-109
20	8				1	1	1	1	1	2	2		2				1					
21	21		2	1	1	1	2	3	3	4	1		3									
22	22		3	2	1	2	3	3	3	3			2						1			
23	34	1	10	3	3	2	10	1	5	10	2		1				4	2				
24	123	3	5	21	8	21	23	8	11	33	6		26				2	7				
25	331	3	7	21	41	39	66	9	11	83	10		30				3	1				
26	774	5	12	38	71	59	178	21	41	218	15		17				0	2				
27	1,073	12	15	33	9	52	209	44	49	342	18		21				28	11				
28	1,003	2	7	13	21	30	103	32	52	368	32		27				55	12				
29	799	2	5	8	24	14	72	15	25	190	26		24				77	35				
30	530	1	3	6	17	9	43	18	22	132	20		19				48	3				
31	342		2	3	6	22	11	11	10	72	14		13				16	2				
32	135		1	1	3	4	11	4	1	30	10		2				12	1				
33	67				1	1	4	1	1	7	3		4				3	3				
34	37				1	1	2	1	1	2	1		1				1	2				
35	16												3				3	4				
36	19												1				2	2				
37	8												1				1	1				
38	22												1				2	1				
39	17												1				1	1				
40	4												1				2	1				
41	4												1				2	1				
42 and over	22												5				2	2				
Number measured	6,345	20	57	91	258	46	268	934	227	1,756	194	173	1,097	204	110	339	143	20	94	22	25	8
Not measured	10																					
Total	6,355																					
Mean transverse pelvic diameter	27.10	26.67	26.81	27.08	27.98	27.09	27.09	27.69	28.44	28.03	29.18	28.66	29.15	29.61	29.63	29.83	30.08	30.35	30.78	31.14		32.85
Blouse group designation	1	2	3 s.	3 m.	3 l.	4 s.	4 m.	4 l.	5 s.	5 m.	5 l.	6 s.	6 m.	6 l.	7 s.	7 m.	7 l.	8 s.	8 m.	8 l.		0

Transverse pelvic diameter: Mean, 28.54 centimeters.

TABLE CXIII.—Association between blouse groups and neck circumference, colored troops, demobilization.

Neck circumfer- ences, in centimeters.		Chest circumference (rest), in centimeters.																					
Total.		73 and under.	74-77	78-81 s.	78-81 m.	78-81 l.	82-85 s.	82-85 m.	82-85 l.	86-89 s.	86-89 m.	86-89 l.	90-93 s.	90-93 m.	90-93 l.	94-97 s.	94-97 m.	94-97 l.	98-101 s.	98-101 m.	98-101 l.	102-105	106-109
29	1			1	1	1	1	1	1	1	1	1	1	1	1		1						
30	16		1	1	3			6	3		3	1		6									
31	14		6	5	1	1	5	16	9	5	12	2		2									
32	61		15	15	22	1	14	46	9	5	29	4		5									
33	177	2	8	25	64	14	54	134	30	26	103	8		36									
34	519	2	15	64	14	54	134	30	26	103	8			114	13	4	12	1	1	1			
35	1,074	6	12	17	67	3	71	273	56	53	316	34	19	223	36	18	36	11	2	1	6		
36	1,474	3	9	19	59	13	75	235	70	77	500	47	36	223	67	27	85	40	4	13	5	6	
37	1,415	1	2	7	17	8	32	131	70	61	468	60	49	307	55	33	89	41	6	18	5	6	
38	920	1	3	10	10	1	12	59	18	30	221	29	7	18	96	17	16	73	31	3	20	3	10
39	398	1	1	3	3	1	1	7	4	4	69	2	2	35	6	10	28	14	2	19	5	8	4
40	161									2	15	2		8	1	8	3		8	3	1		
41	36																3			5			
42	11																						
43	2																						
44	1																						
Number measured	6,280	16	57	89	257	46	267	921	224	259	1,741	192	172	1,083	201	110	338	142	17	93	22	25	8
Not measured	75																						
Total	6,355																						
Mean neck circum- ference.....cm.	35.37	34.40	34.53	34.91	35.20	35.32	35.50	35.70	36.06	36.28	36.42	36.90	36.95	37.14	37.56	37.86	37.99	37.82	38.57	38.68	38.42	38.42	38.42
Blouse group des- ignation.....	1	2	3 s.	3 m.	3 l.	4 s.	4 m.	4 l.	5 s.	5 m.	5 l.	6 s.	6 m.	6 l.	7 s.	7 m.	7 l.	8 s.	8 m.	8 l.	9	9	9

Neck circumference: Mean 36.37 centimeters.

TABLE CXIV.—Association between blouse groups and total arm length, colored troops, demobilization.

Arm length, in centimeters.		Chest circumference (rest), in centimeters.																						
		68-73	74-77	78-81 s.	78-81 m.	78-81 l.	82-85 s.	82-85 m.	82-85 l.	86-89 s.	86-89 m.	86-89 l.	90-93 s.	90-93 m.	90-93 l.	94-97 s.	94-97 m.	94-97 l.	98-101 s.	98-101 m.	98-101 l.	102-105	106-109	
Total.																								
60-65	5																							
66-67	2																							
68-69	19																							
70-71	103	1	2	3	1	1	1	1	6	11	22	5	4	2	1	2	1	1	1	1	1	1	1	1
72-73	249	1	4	16	10	12	17	17	6	23	68	8	14	17	4	2	7	9	3	3	3	3	3	3
74-75	464	2	7	23	15	36	39	93	14	28	127	14	26	82	7	9	13	30	14	5	4	4	4	4
76-77	723	2	10	47	23	55	55	157	31	54	325	23	34	153	21	13	20	64	18	2	2	2	2	2
78-79	1,042	5	17	12	59	10	40	177	36	41	411	29	24	202	32	16	18	70	18	5	5	5	5	5
80-81	1,057	5	10	6	40	8	36	177	45	41	294	28	32	176	37	13	13	43	33	2	2	2	2	2
82-83	1,881	1	3	4	11	4	18	108	30	28	265	35	11	164	47	10	10	42	21	1	1	1	1	1
84-85	684																							
86-87	460																							
88-89	244																							
90-91	115																							
92-93	52																							
94-95	21																							
96 and over	14																							
Number measured.	6,135	19	51	86	252	44	259	915	221	248	1,705	190	165	1,064	202	102	318	134	18	93	19	24	6	
Not measured	220																							
Total	6,355																							
Mean arm length		79.03	76.26	76.06	77.87	79.18	77.45	79.05	80.33	78.62	80.41	82.15	80.22	82.01	83.36	81.70	83.20	84.14	81.17	84.46	86.18	83.50		
Blouse group designation.		1	2	3 s.	3 m.	3 l.	4 s.	4 m.	4 l.	5 s.	5 m.	5 l.	6 s.	6 m.	6 l.	7 s.	7 m.	7 l.	8 s.	8 m.	8 l.	9		

Arm length: Mean, 80.56 centimeters; standard deviation, 4.76±0.0213 centimeter.

TABLE CXV.—Correlation between waist circumference and leg length, while troops, demobilization.

(Basis of construction of breeches groups shown by heavy lines: circle symbols are the "breeches" group designations. For relative frequency of "groups" see Table 211.)

Leg length, in centimeters.	Total.	Waist circumference (rest), in centimeters.										Mean waist circumference (rest).		
		63 and under.	64-67	68-71	72-75	76-79	80-83	84-87	88-91	92-95	96-99		100-103	104-107
50-51.....	24		2	1	4	7	3	4	2	1				79.02
52-53.....	39		1		10	14	3	3	4					77.50
54-55.....	41		1	5	12	13	5	3	1		1			76.90
56-57.....	103		2	12	39	22	19	7		2				76.42
58-59.....	265	①	17	38	70	76	37	15	6	4	1	⑪	1	76.14
60-61.....	923		35	169	297	211	122	44	18	16		1	2	75.66
62-63.....	2,357		82	374	733	611	314	128	66	18	12		2	75.90
64-65.....	4,943		169	791	1,519	1,101	811	298	130	62	17	10	4	76.26
66-67.....	9,790		231	1,300	2,615	3,002	1,524	645	264	97	52	14	12	76.82
68-69.....	14,201		242	1,692	3,869	4,191	2,448	1,035	411	146	72	29	19	77.18
70-71.....	17,304		260	1,764	4,305	5,007	3,374	1,559	628	186	90	32	26	77.74
72-73.....	16,103		203	1,397	3,766	4,760	3,336	1,583	602	217	90	36	16	78.14
74-75.....	12,410		119	921	2,708	3,661	2,719	1,353	567	171	102	24	21	78.62
76-77.....	8,513		72	566	1,694	2,556	1,925	1,003	449	121	51	23	16	78.94
78-79.....	4,861		32	293	936	1,419	1,062	664	280	81	33	8	11	79.26
80-81.....	2,390		13	118	402	679	571	357	162	44	21	9	3	79.86
82-83.....	1,063		4	53	175	261	272	159	85	35	3	6		80.18
84-85.....	1,405		3	13	58	115	107	63	30	14	5			80.34
86-87.....	202		2	2	23	63	58	26	13	6	1	2	2	81.14
88-89.....	96		1	7	14	21	24	21	8	1	1			80.26
90-91.....	36			2	9	10	7	7	2	1				79.38
92-93.....	22		2	2	4	3	3	3	1		2			77.96
94-95.....	22				4	8	3	1	1					78.58
96-97.....	10			2	1	5	4	2						76.26
98-99.....	14		1		1	2	6	3						80.94
100-101.....	3				2			1		1				77.50
Number measured.....	96,157	454	1,500	9,523	23,265	27,848	18,780	8,987	3,705	1,223	555	194	133	77.96
Not measured.....	1,006													
Total.....	97,163													
Mean leg length.....	cm.	70.90	69.34	70.06	70.76	71.52	72.08	72.63	72.78	72.34	72.26	72.46	71.98	

Leg length: Mean, 71.44 centimeters; standard deviation, 4.72 centimeters. Waist circumference: Mean, 77.87 centimeters; standard deviation, 6.08 centimeters. (Correlation: 0.169 ± 0.0021 .)

TABLE CXVIII.—Association between breeches groups and thigh circumference, white troops, demobilization.

Thigh circumference, in centimeters.	Breeches groups (waist circumference, in centimeters).																Total.	104 and over.							
	50-53	54-57	58-71	58-71	69-71	72-75	72-75	76-79	80-83	80-83	90-93	84-87	84-87	88-91	88-91	92-95			98-99	100-103.					
42	15	18	2	19	7	21	5	11	2	5	5	2	1	2	1	1	1	1							
43	256	11	45	9	10	52	19	24	2	8	8	2	1	1	1	1	1	1							
44	434	19	65	26	14	10	92	31	3	3	3	2	2	2	2	2	2	2							
45	914	27	145	48	32	32	39	46	6	6	6	3	3	3	3	3	3	3							
46	1,568	42	217	79	372	72	62	122	11	25	25	11	8	8	8	8	8	8							
47	2,357	43	216	147	690	103	130	291	16	64	64	27	20	20	20	20	20	20							
48	4,566	31	256	190	1,004	134	206	553	26	134	134	38	30	30	30	30	30	30							
49	6,638	24	196	208	1,084	167	330	1,065	55	285	285	100	80	80	80	80	80	80							
50	9,226	18	127	234	1,080	158	413	1,792	143	695	695	155	119	119	119	119	119	119							
51	10,078	21	83	157	812	119	440	2,748	208	1,494	1,494	252	168	168	168	168	168	168							
52	11,147	16	44	109	640	76	361	2,710	396	1,945	1,945	368	242	242	242	242	242	242							
53	10,922	15	36	71	360	67	256	2,710	396	1,945	1,945	368	242	242	242	242	242	242							
54	9,441	9	18	29	208	33	161	2,104	558	1,454	1,454	447	304	304	304	304	304	304							
55	7,903	6	12	21	101	19	118	1,454	558	1,454	1,454	447	304	304	304	304	304	304							
56	5,730	8	3	2	54	8	27	1,084	25	84	84	280	191	191	191	191	191	191							
57	4,290	8	3	2	3	3	3	1,084	25	84	84	280	191	191	191	191	191	191							
58	2,959	3	4	2	2	2	2	96	23	55	55	183	124	124	124	124	124	124							
59	1,128	3	2	1	1	1	1	31	6	18	18	69	44	44	44	44	44	44							
60	799	1	1	1	2	2	2	14	2	12	12	39	24	24	24	24	24	24							
61	554	1	1	1	2	2	2	9	4	9	9	30	18	18	18	18	18	18							
62	352	1	1	1	2	2	2	8	4	8	8	21	12	12	12	12	12	12							
63	243	1	1	1	2	2	2	10	2	10	10	5	5	5	5	5	5	5							
64	181	1	1	1	2	2	2	5	2	5	5	3	3	3	3	3	3	3							
65	95	1	1	1	2	2	2	3	1	3	3	1	1	1	1	1	1	1							
66 and over	595	9	6	7	18	8	49	55	7	55	55	8	66	66	66	66	66	66							
Number measured	95,183	309	1,461	1,367	6,984	1,045	2,662	17,106	3,296	5,005	17,506	5,090	2,821	13,713	2,131	1,137	6,906	880	465	2,636	577	1,213	553	193	132
Not measured	939																								
Total	96,157																								
Mean thigh circumference	49.17	47.95	49.51	49.58	49.73	50.90	50.99	51.13	52.39	52.49	52.51	54.23	54.19	54.06	55.63	55.73	55.40	56.80	57.04	57.04	58.61	58.61	59.24	61.10	61.76
Breeches group designation	1	2	3 s.	3 m.	3 l.	4 s.	4 m.	4 l.	5 s.	5 m.	5 l.	6 s.	6 m.	6 l.	7 s.	7 m.	7 l.	8 s.	8 m.	8 l.	9	10	11	12	

Thigh circumference: Mean, 52.71 centimeters.

TABLE CXIX.—Association between breeches groups and suprapatella, white troops, demobilization.

Suprapatella, in centimeters.		Waist circumference, in centimeters.																								101 and over.					
		63 and under.	64-67 s.	68-71 s.	68-71 m.	68-71 l.	72-75 s.	72-75 m.	72-75 l.	76-79 s.	76-79 m.	76-79 l.	80-83 s.	80-83 m.	80-83 l.	84-87 s.	84-87 m.	84-87 l.	88-91 s.	88-91 m.	88-91 l.	92-95 s.	92-95 m.	92-95 l.	96-99 s.		96-99 m.	96-99 l.	100-103 s.	100-103 m.	100-103 l.
32.....	1,815	29	191	89	387	61	62	56	173	58	35	91	11	4	51	8	1	27	5	9	11	14	5	4	4	6	6	6	6	6	6
33.....	3,473	41	266	168	756	116	151	140	355	118	24	135	22	8	52	6	4	10	14	11	20	6	5	5	5	5	5	5	5	5	5
34.....	6,637	49	307	251	1,099	156	303	350	959	246	87	304	56	17	106	13	8	32	6	18	20	10	13	6	4	4	4	4	4	4	4
35.....	10,814	66	367	296	1,394	207	485	636	2,026	540	179	755	136	55	222	25	14	60	24	29	18	24	17	13	2	2	2	2	2	2	2
36.....	14,058	65	192	229	1,272	165	530	879	3,038	946	317	1,616	219	72	402	78	20	89	41	20	41	28	17	2	2	2	2	2	2	2	2
37.....	15,508	64	85	162	927	148	471	606	3,435	999	451	2,298	315	139	1,128	89	41	207	79	88	31	42	31	4	4	4	4	4	4	4	4
38.....	14,014	42	64	73	491	88	308	814	3,013	805	560	2,550	402	184	1,128	146	67	321	103	133	133	68	31	4	4	4	4	4	4	4	4
39.....	10,722	32	34	70	287	51	175	549	2,102	648	453	2,266	372	196	1,181	138	51	384	77	100	133	68	31	4	4	4	4	4	4	4	4
40.....	7,363	29	22	23	134	25	83	370	1,290	346	314	2,018	285	155	1,094	142	83	428	103	133	133	68	31	4	4	4	4	4	4	4	4
41.....	5,023	23	55	29	146	19	59	189	625	184	204	1,023	142	113	820	101	68	383	83	197	190	68	31	4	4	4	4	4	4	4	4
42.....	6,510	14	17	11	110	23	47	168	638	189	214	1,156	180	174	1,072	139	113	708	139	487	487	252	252	252	252	252	252	252	252	252	252
Total.....	96,157	454	1,500	1,391	7,073	1,059	2,664	5,057	17,644	5,147	2,838	13,802	2,140	1,147	6,955	885	470	2,653	592	1,223	555	555	555	555	555	555	555	555	555	555	555
Mean suprapatella circumference, cm.....	36.34	34.99	35.41	35.74	35.72	36.25	36.38	37.11	37.25	37.33	38.09	38.22	38.25	38.92	39.96	38.94	39.47	39.63	39.41	40.22	40.22	40.22	40.29	40.29	40.65	40.65	40.65	40.65	40.65	40.65	40.65
Breeches group designation.....	1	2	3 s.	3 m.	3 l.	4 s.	4 m.	5 s.	5 m.	5 l.	6 s.	6 m.	6 l.	7 s.	7 m.	7 l.	8 s.	8 m.	8 l.	9	9	9	10	10	11	11	12	12	12	12	12

Suprapatella: Mean, 37.34 centimeters; standard deviation, 2.45 ± 0.0066 centimeter.

TABLE CXX.—Association between breeches groups and patella circumference, white troops, demobilization.

Patella circumference, in centimeters.		Total.	Waist circumference, in centimeters.																104 and over.						
			63 and under.	64-67	68-71	68-71 m.	68-71 l.	72-75	72-75 m.	72-75 l.	76-79	76-79 m.	76-79 l.	80-83	80-83 m.	80-83 l.	84-87	84-87 m.		84-87 l.	88-91	88-91 m.	88-91 l.	92-95	96-99
31 and under.....	976	26	54	40	99	10	31	135	27	49	90	26	20	109	22	10	84	24	1	47	18	29	19	6	6
32.....	1,745	45	134	122	337	28	114	391	45	88	180	47	30	87	22	9	35	6	2	11	3	10	6	3	3
33.....	4,821	58	326	286	884	79	332	1,200	148	333	597	123	85	220	25	14	69	15	7	16	2	9	9	4	2
34.....	10,459	95	431	376	1,587	192	615	2,790	348	769	1,656	315	218	618	67	66	183	57	19	56	10	18	14	4	2
35.....	16,926	65	299	310	1,520	274	698	4,065	645	1,198	2,364	731	508	1,903	168	164	507	54	49	141	20	42	20	3	4
36.....	19,637	61	170	158	1,328	220	519	4,065	820	1,176	2,305	1,093	628	2,529	311	209	1,077	87	67	288	46	96	42	4	7
37.....	17,627	34	45	60	648	143	245	2,710	872	1,533	3,831	2,269	667	3,260	476	265	1,470	167	88	54	155	54	12	15	8
38.....	12,025	26	20	24	262	67	78	2,136	394	405	2,329	933	401	2,563	481	186	1,083	182	99	547	125	214	90	37	14
39.....	6,724	20	11	6	70	32	35	1,375	156	157	883	447	205	1,320	331	135	1,083	165	76	516	83	255	92	23	7
40.....	3,338	17	5	4	24	9	13	124	53	50	335	166	70	680	178	63	624	93	37	340	92	189	107	71	67
41.....	1,879	6	5	5	14	5	4	33	15	9	104	57	13	263	69	26	362	75	25	263	75	206	102	37	67
Total.....	98,157	454	1,500	1,391	7,073	1,059	2,684	17,248	3,323	5,057	17,644	5,147	2,838	13,802	2,140	1,147	6,955	885	470	2,653	582	1,223	555	194	133
Mean patella circumference.....cm.	34.90	34.15	34.25	35.88	35.41	34.90	35.48	35.03	35.61	36.20	36.64	36.92	36.92	37.38	36.87	37.48	37.69	37.42	37.95	38.10	38.33	38.27	39.08	39.65	12
Breeches group designation.....	1	2	3 s.	3 m.	3 l.	4 s.	4 m.	4 l.	5 s.	5 m.	5 l.	6 s.	6 m.	6 l.	7 s.	7 m.	7 l.	8 s.	8 m.	8 l.	9	9	10	11	12

Patella circumference: Mean, 36.21 centimeters; standard deviation, 1.98 centimeters.

TABLE CXXI.—Association between breeches groups and calf circumference, while troops, demobilization.

Calf circumference, in centimeters.	Waist circumference, in centimeters.															
	Total.	63 and under.	64-67	68-71	68-71	72-75	72-75	72-75	72-75	76-79	76-79	76-79	76-79	80-83	80-83	84-87
29.....	557	13	67	141	16	26	119	23	16	43	10	10	10	3	29	4
30.....	2,472	35	239	124	126	600	600	81	106	256	44	8	8	14	79	8
31.....	6,021	47	348	246	1,204	1,651	1,651	292	315	718	175	18	18	35	247	14
32.....	12,115	72	381	246	1,787	2,255	2,255	590	682	2,076	564	50	50	102	783	22
33.....	17,024	60	238	289	1,478	2,466	2,466	743	1,172	3,618	950	114	114	194	1,733	35
34.....	18,651	62	111	182	1,017	1,666	1,666	709	1,108	4,141	1,183	194	194	412	2,651	58
35.....	16,485	66	49	73	505	92	3,586	500	860	3,502	1,036	276	276	466	3,286	92
36.....	11,346	51	9	38	188	112	1,010	236	500	2,031	668	362	362	421	2,490	104
37.....	6,303	20	17	9	70	30	327	84	205	868	352	157	157	250	1,371	158
38.....	3,039	10	20	6	37	16	104	32	54	253	123	103	103	98	601	88
39 and over.....	2,044	17	41	80	11	16	112	29	36	138	40	26	26	65	223	49
Number measured.....	96,087	453	1,500	7,087	1,059	2,680	17,240	3,319	5,055	17,644	5,145	2,835	13,763	2,136	1,147	6,980
Not measured.....	96,157															
Total.....																
Mean calf circumference.....																
Breeches group designation.....																
1.....	33.57	32.07	32.49	32.60	32.80	33.02	33.27	33.48	33.78	34.04	34.25	34.56	34.83	34.99	35.16	35.47
2.....																
3.....																
4.....																
5.....																
6.....																
7.....																
8.....																
9.....																
10.....																
11.....																
12.....																
13.....																
14.....																
15.....																
16.....																
17.....																
18.....																
19.....																
20.....																
21.....																
22.....																
23.....																
24.....																
25.....																
26.....																
27.....																
28.....																
29.....																
30.....																
31.....																
32.....																
33.....																
34.....																
35.....																
36.....																
37.....																
38.....																
39 and over.....																
Number measured.....	96,087	453	1,500	7,087	1,059	2,680	17,240	3,319	5,055	17,644	5,145	2,835	13,763	2,136	1,147	6,980
Not measured.....	96,157															
Total.....																
Mean calf circumference.....																
Breeches group designation.....																
1.....	33.57	32.07	32.49	32.60	32.80	33.02	33.27	33.48	33.78	34.04	34.25	34.56	34.83	34.99	35.16	35.47
2.....																
3.....																
4.....																
5.....																
6.....																
7.....																
8.....																
9.....																
10.....																
11.....																
12.....																
13.....																
14.....																
15.....																
16.....																
17.....																
18.....																
19.....																
20.....																
21.....																
22.....																
23.....																
24.....																
25.....																
26.....																
27.....																
28.....																
29.....																
30.....																
31.....																
32.....																
33.....																
34.....																
35.....																
36.....																
37.....																
38.....																
39 and over.....																

Calf circumference: Mean, 34.09 centimeters; standard deviation, 2.02 ± 0.0045 centimeters.

TABLE CXXII.—*Correlation between waist circumference and leg length, colored troops, demobilization.*

[Basis of construction of breeches groups shown by heavy lines; circle symbols are the "breeches" group designations. For relative frequency of "groups," see Table 121.]

Leg length, in centimeters.	Waist circumference, in centimeters.											
	63 and under.	64-67	68-71	72-75	76-79	80-83	84-87	88-91	92-95	96-99	100-103	104 and over.
Total.												
54-55.....	2	③	1		④	1						
56-57.....	1				1							
58-59.....	4	②	1	⑤	2	⑥	1					
60-61.....	22	1	3	6	10	2						
62-63.....	49	①	1	17	8	6	⑦	⑧	⑨	⑩	⑪	1
64-65.....	109	3	10	25	35	11	8	3	3	1	1	3
66-67.....	241	10	41	68	78	28	8	3				
68-69.....	464	2	71	133	131	67	35	7	⑬	1	1	1
70-71.....	890	7	99	240	277	164	58	30	8	3	1	1
72-73.....	1,028	2	10	282	348	180	73	26	6	4	1	2
74-75.....	1,083	8	94	261	371	215	77	29	7	7	1	3
76-77.....	1,031	6	66	237	334	216	104	38	11	10	1	1
78-79.....	712	8	42	131	223	174	84	24	7	4	1	1
80-81.....	430	1	5	72	133	110	44	22	13	2	1	1
82-83.....	214	2	13	24	72	61	26	12	1	2	1	
84-85.....	87	1	4	13	29	24	6	7	1	1	1	
86-87.....	46		2	4	17	9	8	4	1			1
88 and over.....	32		1	4	12	8	4	1	2			
Number measured.....	6,445	34	71	588	1,519	2,080	1,276	586	209	60	38	11
Not measured.....	75											15
Total.....	6,520											

Waist circumference: Mean, 77.83 centimeters; standard deviation, 5.76 ± 0.034 centimeter.

TABLE CXXIII.—Association between breeches groups and transverse diameter of pelvis, colored troops, demobilization.

Transverse pelvis, in centimeters.	Total.	Waist circumference, in centimeters.													
		63 and under.	64-67	68-71	72-75	76-79	80-83	84-87	88-91	92-95	96-99	100-103	104-109		
20.	5														
21.	18			1		1									
22.	14		1	3		5									
23.	33		2	1		2									
24.	125		5	8		10									
25.	325	2	10	20	2	12									
26.	752	2	19	47	2	16									
27.	1,017	2	12	103	28	30	4								
28.	1,014	2	17	14	55	42	14								
29.	1,014	2	6	53	171	194	22								
30.	1,720	3	2	30	26	27	22								
31.	520	4	1	20	12	7	13								
32.	343	1	1	11	18	4	31								
33.	158	1	1	2	12	3	20								
34.	68			2	1	1	11								
35.	22			2	2	1	2								
36.	20				1	1	1								
37.	5														
38.	4														
39 and over.	1														
Number measured	6,354	33	71	416	88	247	1,004	248	1,314	493	114	925	208	51	394
Not measured.	91														
Total.	6,445														
Mean transverse pelvic diameter.....cm.	27.97	26.39	25.93	26.92	27.36	27.02	27.52	28.07	27.43	28.26	28.75	28.36	29.26	29.75	29.98
Breeches group designation		1	2	3 s.	4 s.	5 s.	6 m.	7 s.	8 m.	9	10	11	12	13	14

Transverse pelvic diameter: Mean, 28.42 centimeters.

TABLE CXXIV.—Association between breeches groups and knee height, colored troops, demobilization.

[illegible]

Knee height: Mean, 47.30 centimeters.

TABLE V. Association between blouse groups and suprapatella circumference, colored troops, demobilization.

		Waist circumference, in centimeters																					
Suprapatella circumfer- ence, in centimeters.	Total and un- der.	64-67		68-71		72-75		76-79		80-83		84-87		88-91		92-95		96-99		100-103		104-109	
		s.	m.	s.	m.	s.	m.	s.	m.	s.	m.	s.	m.	s.	m.	s.	m.	s.	m.	s.	m.	s.	m.
32	85	3	17	4	15	5	3	5	3	2	4	2	1	1	2								
33	188	16	50	9	44	11	9	16	8	4	5	6	9	4	2								
34	378	12	12	19	25	10	36	3	12	34	1	9	3	2	1								
35	650	13	14	38	41	29	101	43	9	34	16	3	24	1	2								
36	846	10	11	47	53	44	209	64	7	75	34	6	35	1	2								
37	1,042	3	10	53	203	44	53	270	92	10	134	4	35	6	1								
38	1,010	5	3	33	37	41	241	94	32	200	41	6	39	17	2	15	2	6	4	2	1		
39	763	3	1	2	20	62	16	33	173	12	176	41	12	73	9	2	18	6	6	2	2		
40	590	1	1	4	9	45	9	23	128	45	153	20	8	64	16	2	21	5	8	4	1		
41	369	3	11	4	21	5	9	60	17	180	20	6	64	16	3	23	9	3	6	3	6		
42 and over	522	1	6	2	4	21	4	11	70	19	89	12	81	17	5	63	17	33	17	8	12		
Number measured.	6,443	34	70	425	89	251	1,018	265	1,330	495	949	211	52	306	88	16	147	46	80	37	11	15	
Not measured.	2																						
Total	6,445																						
	38.65	34.80	35.50	35.75	36.05	36.45	36.43	37.32	37.66	37.48	38.66	38.62	38.45	39.12	39.46	39.52	39.38	40.15	40.43	40.43		41.12	
	1	2	3 s.	3 m.	3 l.	4 s.	4 m.	4 l.	5 s.	5 m.	5 l.	6 m.	6 l.	7 s.	7 m.	7 l.	8 s.	8 m.	9	10		11	

38536°-21-36

Suprapatella circumference: Mean, 37.61 centimeters; standard deviation, 2.43 ± 0.0214 centimeter.

TABLE CXXVII.—Association between breeches groups and circumference at patella, colored troops, demobilization.

Patella circumference, in centimeters.	Total.	Waist circumference, in centimeters.																							
		63 and under.	64-67 s.	68-71 m.	69-71 l.	72-75 s.	72-75 m.	72-75 l.	76-79 s.	76-79 m.	76-79 l.	80-83 s.	80-83 m.	80-83 l.	84-87 s.	84-87 m.	84-87 l.	88-91 s.	88-91 m.	88-91 l.	92-95	96-99	100-103	104-109	
31 and under.....	53	1	3	4	4	3	3	2	2	8	4	1	1	3	1	8	3	1	1	3	1	1	1	1	
32.....	83	1	13	4	15	4	14	5	4	6	1	1	1	5	2	1	1	2	1	1	2	1	1	1	
33.....	249	4	18	16	53	22	52	9	19	29	7	7	7	2	2	9	1	2	2	2	2	2	2	2	
34.....	575	4	12	20	108	50	144	23	30	97	25	15	15	8	5	28	2	3	6	6	5	5	5	5	
35.....	1,030	4	18	15	110	65	270	46	63	215	61	77	77	28	6	52	6	3	9	3	7	7	7	7	
36.....	1,207	4	10	10	63	55	242	64	59	312	94	145	145	44	6	71	18	4	17	3	10	10	10	10	
37.....	1,270	8	6	5	49	36	164	40	45	209	93	20	230	43	14	73	21	3	36	13	11	11	11	11	
38.....	941	3	1	1	16	8	80	31	26	98	56	10	117	47	9	79	14	4	32	11	6	6	6	6	
39.....	574	4	1	1	6	3	29	17	14	37	19	5	75	29	4	50	15	6	28	6	12	12	12	12	
40.....	308	1	1	1	1	2	9	4	2	8	6	1	19	6	3	27	7	11	22	10	11	11	11	11	
41 and over.....	154	1	1	2	1	1	2	1	2	8	6	1	19	6	3	27	7	11	22	10	11	11	11	11	
Number measured.....	6,444	34	70	74	425	88	251	1,018	249	1,330	496	115	949	212	52	396	88	16	147	46	60	37	11	15	
Not measured.....	1																								
Total.....	6,445																								
Mean patella circumfer- ence, cm.....		36.21	34.19	34.28	34.95	35.27	35.67	36.18	35.81	36.49	36.87	36.70	37.34	37.78	37.35	37.77	37.97	37.06	38.67	38.63	38.58	38.46	39.35	39.35	
Breeches group designa- tion.....		1	2	3 s.	3 m.	4 s.	4 m.	4 l.	5 s.	5 m.	5 l.	6 s.	6 m.	6 l.	7 s.	7 m.	7 l.	8 s.	8 m.	8 l.	9	10	11	11	

Patella circumference: Mean, 36.52 centimeters; standard deviation, 1.99±0.0175 centimeter.

TABLE XXVIII. -Association between breeches groups and circumference of cals, colored troops, demobilization.

[illegible]

Calf circumference: Mean, 34.71 centimeters; standard deviation, 2.10 ± 0.0125 centimeter.

TABLE CXXIX.—Comparative frequency distribution of "blouse" groups, by States of nativity, white troops (absolute numbers).

State	Total	Chest circumference (rest), in centimeters													110 and over
		68-77	78-81	78-81	82-85	82-85	86-89	86-89	89-92	92-101	92-101	102-105	106-109		
		s.	m.	m.	s.	m.	s.	m.	s.	m.	s.	m.	s.		
Alabama.....	1,532	17	14	19	81	189	106	79	344	104	7	11	3		
Alaska.....	12														
Arizona.....	127	1	3	3	1	6	12	10	17	7	1	1	1		
Arkansas.....	2,257	26	101	37	40	304	187	125	625	185	5	7	1		
California.....	446	6	4	4	6	52	19	34	96	21	5	7	1		
Colorado.....	216	7	2	2	2	23	17	12	43	13	1	1	1		
Connecticut.....	977	12	10	8	32	133	33	85	197	39	11	1	1		
Delaware.....	275	2	2	2	4	26	12	23	63	18	1	1	1		
District of Columbia.....	190	2	2	1	4	26	13	14	37	13	1	1	1		
Florida.....	746	11	7	12	16	113	43	43	151	44	8	3	3		
Georgia.....	2,333	16	15	100	37	248	152	108	567	166	8	4	4		
Idaho.....	1,187	1	1	2	13	5	6	31	31	6	1	1	1		
Illinois.....	6,490	65	41	162	120	721	314	354	1,223	266	66	76	11		
Indiana.....	3,816	40	21	172	59	375	251	190	874	230	12	34	3		
Iowa.....	1,570	5	4	46	24	170	58	85	263	102	9	21	2		
Kansas.....	961	10	2	33	14	87	45	80	211	64	11	12	1		
Kentucky.....	2,718	24	23	125	45	377	192	140	633	177	11	17	2		
Louisiana.....	1,313	5	7	12	19	101	60	80	297	36	8	8	2		
Maine.....	604	8	21	3	13	94	34	38	134	34	14	7	2		
Maryland.....	1,008	21	15	63	16	143	65	59	246	28	11	10	6		
Massachusetts.....	4,750	27	33	212	49	126	217	239	960	137	46	24	6		
Michigan.....	3,660	27	25	120	36	335	155	266	728	135	40	24	6		
Minnesota.....	1,513	17	7	29	68	171	77	88	308	43	15	23	2		
Mississippi.....	1,355	12	3	33	33	160	90	76	333	43	12	13	1		
Missouri.....	2,565	23	21	120	43	346	216	131	649	186	10	21	4		
Montana.....	263	1	1	2	12	12	37	13	174	16	1	1	1		
Nebraska.....	660	5	4	15	17	63	37	33	147	39	10	10	1		
Nevada.....	114	1	1												
New Hampshire.....	410	4	3	2	11	40	21	19	97	14	3	10	3		
New Jersey.....	3,102	26	46	133	33	474	118	243	630	108	24	31	10		
New Mexico.....	226	1	1	2	11	15	15	6	43	15	1	1	1		
New York.....	8,001	101	119	351	312	1,393	383	790	1,771	270	99	76	27		
North Carolina.....	1,188	15	8	11	12	53	54	54	234	66	10	27	1		
North Dakota.....	363	1	1	2	12	12	16	11	59	14	7	9	2		
Ohio.....	4,927	71	47	226	73	165	934	462	1,967	342	49	30	9		
Oklahoma.....	2,196	19	11	23	165	241	196	98	516	215	7	13	3		
Oregon.....	1,081	4	16	23	30	70	36	49	233	60	9	8	1		
Pennsylvania.....	10,630	78	114	396	303	1,404	365	975	2,194	378	99	41	26		
Rhode Island.....	397	1	7	2	6	55	19	28	40	10	7	2	2		
South Carolina.....	630	3	5	3	11	83	40	31	137	24	4	3	1		
South Dakota.....	409	1	1	2	4	41	14	115	530	164	1	16	3		
Tennessee.....	2,533	18	20	95	46	336	146	113	580	164	7	14	16		
Texas.....	2,809	21	13	69	50	407	260	142	789	316	22	36	15		
Utah.....	102														
Vermont.....	441	3	3	2	6	62	24	24	56	20	3	1	1		
Virginia.....	1,405	11	14	28	11	37	139	70	292	82	11	4	4		
Washington.....	1,984	16	7	46	15	81	173	88	108	202	11	19	5		
West Virginia.....	1,640	9	12	35	16	29	150	81	82	332	16	21	5		
Wisconsin.....	2,640	11	5	52	12	219	91	130	432	119	53	16	3		
Wyoming.....	74	2	1	1	1	10	1	4	10	4	2	2	1		
Others.....	1,043	9	10	36	10	117	35	76	213	47	10	11	5		
Grand total.....	95,467	440	756	3,342	1,077	12,016	4,834	5,946	10,968	4,919	775	2,392	806		
									5,175	1,210	8,601	1,221	228		

TABLE (XXXI).—Comparative frequency distribution of "breches" groups, by States of nativity, white troops (absolute numbers).

State.	Waist circumference, in centimeters.										104 and over
	Total.	10-13	14-17	18-21	22-25	26-29	30-33	34-37	38-41	42-45	
Alabama.....	1,540	3	10	9	84	20	16	229	68	1	3
Alaska.....	12	1	1	1	1	1	1	1	1	1	1
Arizona.....	125	1	1	1	1	1	1	1	1	1	1
Arkansas.....	2,280	4	21	111	186	37	15	361	101	52	1
California.....	4,460	1	2	1	39	7	7	88	19	10	1
Colorado.....	217	1	1	1	21	6	4	40	19	18	1
Connecticut.....	883	2	10	32	73	9	44	179	20	91	1
District of Columbia.....	775	3	5	7	17	2	6	48	15	15	1
Florida.....	752	2	7	7	48	10	15	130	33	142	1
Georgia.....	2,391	7	21	162	35	37	27	407	104	71	3
Idaho.....	2,161	1	3	11	182	37	14	1,067	317	260	1
Illinois.....	6,549	36	126	65	465	111	141	1,067	1,067	1,067	4
Indiana.....	3,923	17	81	52	398	50	108	1,067	1,067	1,067	3
Iowa.....	1,579	4	14	11	85	5	58	273	40	117	1
Kansas.....	1,964	1	14	11	40	9	19	173	28	55	1
Kentucky.....	2,747	15	62	30	322	49	54	605	123	86	2
Louisiana.....	1,355	2	6	8	10	8	18	126	26	64	1
Maine.....	1,887	5	2	28	45	2	34	126	23	58	1
Maryland.....	1,014	6	38	25	97	10	28	192	110	405	1
Massachusetts.....	4,748	33	114	150	401	28	279	992	221	818	6
Michigan.....	3,640	19	58	70	279	33	126	702	100	465	4
Minnesota.....	1,925	8	21	21	100	15	57	252	53	117	2
Mississippi.....	1,610	5	45	30	52	5	17	218	47	37	1
Missouri.....	2,074	15	45	30	271	12	5	556	101	114	1
Montana.....	263	1	2	18	41	2	23	106	23	70	1
Nebraska.....	804	6	9	18	41	2	4	141	24	141	1
Nevada.....	17	1	1	1	1	1	1	1	1	1	1
New Hampshire.....	411	4	10	17	44	6	22	67	5	20	1
New Jersey.....	3,119	17	80	105	256	26	135	627	75	216	3
New Mexico.....	225	2	3	3	14	1	4	556	8	9	1
New York.....	9,060	46	206	210	788	93	420	1,704	264	645	20
North Carolina.....	1,194	5	14	4	76	13	8	190	53	1,540	1
North Dakota.....	1,349	2	1	2	17	3	11	58	6	37	1
Ohio.....	6,772	42	118	164	567	53	230	1,358	194	401	5
Oklahoma.....	2,199	11	16	7	148	33	14	173	28	45	1
Oregon.....	1,052	5	8	48	148	18	5	126	96	8	1
Pennsylvania.....	10,607	41	158	172	788	73	400	1,981	231	705	12
Rhode Island.....	384	1	1	1	40	3	19	87	34	34	1
South Carolina.....	630	5	5	9	34	7	8	108	33	16	1
South Dakota.....	407	2	2	3	18	2	7	62	5	28	1
Tennessee.....	2,560	9	26	7	128	22	23	413	92	96	6
Texas.....	3,845	13	32	14	223	55	34	612	103	102	6
Utah.....	104	1	1	6	23	4	4	23	8	4	1
Vermont.....	442	6	5	1	28	1	21	57	58	38	1
Virginia.....	1,423	4	21	13	94	15	23	237	68	37	2
Washington.....	197	11	22	9	96	55	6	235	100	26	1
West Virginia.....	1,639	10	26	10	85	17	24	201	35	90	2
Wisconsin.....	2,643	13	20	22	144	42	53	375	115	106	2
Wyoming.....	79	1	1	1	8	1	11	429	19	11	1
Others.....	1,124	5	5	26	84	7	43	296	34	82	36
Grand total.....	96,157	454	1,500	1,391	7,074	1,058	2,484	17,248	3,323	5,067	193

BREECHES GROUPS—STATE DISTRIBUTION.

565

State.	Total.	Waist circumference, in centimeters																100-103 s.	104 and over.
		63 and under.	64-67 s.	68-71 m.	72-75 s.	76-79 m.	80-83 s.	84-91 m.	92-95 s.	96-99 m.	100-103 s.	104 and over.							
Alabama.....	393	7	21	7	0	53	21	17	76	30	11	1	2	2	3				
Alaska.....	4																		
Arizona.....	4																		
Arkansas.....	306	1	3	2	21	5	11	36	17	5	26	7	3	1	2		1		
California.....	22				3	2		1	3										
Colorado.....	6																		
Connecticut.....	10																		
Delaware.....	25																		
District of Columbia.....	40																		
Florida.....	270	3	5	26	2	11	49	6	9	55	27	5	1	6	8	1	1		
Georgia.....	1,034	13	10	57	21	51	173	53	41	191	98	13	4	24	6	8	1		
Idaho.....	118	1	3	3	6	4	6	13	3	8	27	6	1	2	2	3			
Illinois.....	66	1	1	4	5	1	2	4	6	4	4	2							
Indiana.....	17																		
Iowa.....	34																		
Kansas.....	187																		
Kentucky.....	13																		
Louisiana.....	704	8	7	29	10	24	96	16	25	186	46	18	3	17	6	5	1		
Maine.....	135																		
Maryland.....	21																		
Massachusetts.....	44																		
Michigan.....	13																		
Minnesota.....	493																		
Mississippi.....	120																		
Missouri.....	13																		
Montana.....	46																		
Nebraska.....	1																		
Nevada.....	1																		
New Hampshire.....	46																		
New Jersey.....	1																		
New Mexico.....	25																		
New York.....	113																		
North Carolina.....	618																		
North Dakota.....	6																		
Ohio.....	120																		
Oklahoma.....	119																		
Oregon.....	6																		
Pennsylvania.....	144																		
Rhode Island.....	2																		
South Carolina.....	198																		
South Dakota.....	13																		
Tennessee.....	231																		
Texas.....	553																		
Utah.....	2																		
Vermont.....	1																		
Virginia.....	532																		
Washington.....	10																		
West Virginia.....	42																		
Wisconsin.....	13																		
Wyoming.....	70																		
Others.....																			
Grand total	46,944	77	53	390	123	282	1,090	272	297	1,480	523	150	1	49	71	44	13	17	

* Includes 499 mixed races which were not previously counted: 6,004—699—6,446.

2,025	...	1	2	11	39	59	92	127	201	194	264	231	259	208	109	93	67	32	14	13	4	2	...	173.21								
7,688	...	4	9	12	25	60	92	126	165	179	200	207	181	146	87	73	49	30	22	8	4	2	...	173.21								
2,675	3	9	10	13	34	69	108	156	290	347	312	269	214	136	89	52	38	14	6	3	1	1	...	173.18								
81	...	1	4	1	2	7	8	10	10	11	6	6	1	2	...	2	173.13							
102,304	23	66	150	395	7631	494	2,793	4,499	6,462	8,669	10,569	12,348	12,226	11,503	9,696	7,594	5,060	3,283	2,019	1,209	698	288	156	60	37	21	4	2	3	172.00		
Not measured
102,577
Total

TABLE 'XXXIV.—Comparative frequency distribution of statures, by Q. M. C. distribution zones, demobilization—Continued.

SECTION C: PROPORTIONAL NUMBER OF EACH 1,000 STATURES IN THE VARIOUS DISTRIBUTION ZONES.

Stature, in centimeters.	Zone 1.	Zone 2.	Zone 4.	Zone 5.	Zone 7.	Zone 8.	Zone 9.	Zone 10.	Zone 11.	Zone 12.	Zone 13.	Total.
148-149.....	43.48	347.83	86.96	280.87	86.96	86.96	86.96	1,000
150-151.....	145.45	363.63	36.36	90.91	290.91	36.36	18.18	18.18	1,000
152-153.....	146.67	406.67	33.33	46.67	240.00	33.33	26.67	20.00	20.00	6.67	20.00	1,000
154-155.....	138.54	433.25	30.23	25.19	287.15	45.34	7.56	15.11	5.04	12.59	1,000
156-157.....	121.05	428.95	38.16	38.16	268.42	22.37	30.26	13.16	7.89	6.58	25.00	1,000
158-159.....	127.70	420.95	41.89	44.59	268.24	33.11	16.89	14.86	14.19	1.35	16.22	1,000
160-161.....	109.87	398.87	31.96	63.55	282.59	42.73	23.70	15.44	14.00	2.87	24.42	1,000
162-163.....	83.54	356.54	31.50	77.38	295.82	60.72	29.67	21.91	10.50	4.57	27.85	1,000
164-165.....	104.53	323.22	32.43	72.99	303.08	63.07	31.39	25.61	14.36	2.52	26.80	1,000
166-167.....	83.95	288.73	33.19	81.98	328.17	67.18	35.38	29.83	17.34	3.01	31.22	1,000
168-169.....	72.63	267.75	31.76	97.29	327.30	76.62	36.03	34.80	16.97	3.22	35.65	1,000
170-171.....	71.77	237.22	33.13	110.98	329.68	85.40	38.41	37.43	18.75	2.27	34.99	1,000
172-173.....	61.11	223.15	34.57	119.69	315.80	90.94	43.91	45.55	19.42	3.52	42.35	1,000
174-175.....	56.43	204.50	32.88	124.71	328.95	92.09	46.23	45.26	22.50	4.88	41.60	1,000
176-177.....	46.84	180.73	34.33	135.75	327.75	97.91	47.15	53.97	22.64	3.72	49.21	1,000
178-179.....	41.56	170.97	32.45	155.54	310.42	107.52	44.59	59.23	22.69	4.49	50.53	1,000
180-181.....	44.28	153.91	31.88	163.55	300.93	113.17	50.19	66.91	24.99	2.76	47.43	1,000
182-183.....	32.04	153.19	30.82	164.48	300.27	112.60	51.87	68.66	23.80	4.88	57.37	1,000
184-185.....	25.79	126.49	35.22	179.56	305.06	116.07	44.15	78.87	27.28	3.97	57.54	1,000
186-187.....	34.02	135.27	24.90	178.42	285.47	122.82	62.24	65.56	34.85	7.47	48.96	1,000
188-189.....	30.61	123.91	26.24	164.72	330.90	113.70	72.89	56.85	27.70	52.48	1,000
190-191.....	13.94	128.92	27.87	223.00	268.29	94.07	76.65	62.72	41.81	3.48	56.23	1,000
192-193.....	25.64	115.38	19.23	243.59	288.46	96.15	83.33	57.69	25.64	6.41	38.46	1,000
194-195.....	17.24	155.17	17.24	155.17	327.59	103.45	103.45	51.72	68.97	1,000
196-197.....	51.28	102.56	25.64	205.13	282.05	128.21	76.92	25.64	51.28	51.28	1,000
198-199.....	100.00	200.00	300.00	200.00	50.00	50.00	100.00	1,000
200-201.....	200.00	800.00	1,000
202-203.....	1,000.00	1,000
204-205.....	1,000.00	1,000
206-207.....	1,000.00	1,000
208-209.....	500.00	500.00	1,000
Average proportion for each zone.....	66.01	237.63	32.90	115.62	316.15	85.58	40.85	42.73	19.62	3.52	39.40	1,000

State.	Total.	Weight, in pounds.										Mean weight.	
		100-109	110-119	120-129	130-139	140-149	150-159	160-169	170-179	180-189	190-199	200 and over.	
Alabama	383	4	19	48	81	81	81	35	26	8			Lbs. 144.79
Alaska	12				1	2	2	3	2	2			162.00
Arizona	125		2	14	25	27	28	16	5	8			148.34
Arkansas	2,538	8	91	267	510	617	514	295	152	49	25	10	146.83
California	414	2	13	55	86	113	70	40	18	12	3	2	145.37
Colorado	208		4	18	49	55	39	23	13	4	1	2	147.34
Connecticut	550	9	37	97	124	128	77	48	21	7	1	1	141.05
Delaware	189	1	9	33	42	57	30	17	4	2	2		142.22
District of Columbia	184	3	6	33	42	57	28	17	6	1		1	140.80
Florida	140		7	24	38	54	19	11	5	1			141.50
Georgia	446	2	26	60	107	106	63	45	23	8	2	4	143.84
Idaho	153		6	13	19	32	37	23	16	5	1	1	150.97
Illinois	6,462	32	247	791	1,405	1,608	1,164	674	331	133	50	27	145.42
Indiana	3,804	12	140	522	889	880	696	369	186	60	26	24	144.78
Iowa	1,543	6	35	111	277	350	331	254	101	55	15	8	150.05
Kansas	978	2	16	80	187	236	199	121	72	36	20	9	150.14
Kentucky	2,753	7	95	398	653	623	497	282	125	46	19	8	144.50
Louisiana	1,726	6	63	191	367	413	335	198	89	36	18	10	146.41
Maine	209		12	39	43	50	34	15	9	4		3	142.97
Maryland	983	10	57	156	240	245	150	67	31	18	4	5	141.81
Massachusetts	1,320	14	96	263	322	302	174	93	33	12	9	2	139.74
Michigan	3,618	15	123	467	834	875	629	405	171	63	17	19	145.07
Minnesota	1,882	5	34	137	309	404	428	298	154	68	31	14	151.37
Mississippi	1,566	6	60	139	311	382	319	196	91	35	17	10	147.54
Missouri	2,752	10	93	335	635	653	484	313	127	62	24	16	145.70
Montana	245		2	20	46	46	63	36	18	9	1	4	151.11
Nebraska	791	2	29	44	129	175	167	129	66	34	11	5	151.23
Nevada	16				1	3	5	2	2				149.50
New Hampshire	94	2	12	27	10	23	7	12				1	136.95
New Jersey	3,103	23	218	579	800	708	416	211	80	36	17	15	140.29
New Mexico	221	1	11	29	46	60	44	14	11	2	1	2	144.00
New York	8,965	82	598	1,628	2,319	2,067	1,201	629	268	105	41	27	140.43
North Carolina	570	2	18	64	127	147	96	65	30	13	4	4	146.17
North Dakota	335	1	4	24	56	76	70	62	29	12	1		150.89
Ohio	6,900	35	264	928	1,571	1,649	1,291	714	277	102	47	22	144.4

TABLE CXXXVI.—Comparative frequency distribution of chest circumference (rest), by Q. M. C. distribution zones, white and colored troops, at demobilization.

SECTION A: ABSOLUTE NUMBERS.

Chest circumference, in centimeters.	Total.	Zone 1.	Zone 2.	Zone 4.	Zone 5.	Zone 7.	Zone 8.	Zone 9.	Zone 10.	Zone 11.	Zone 12.	Zone 13.
60-64.....	41	3	21	1	3	11	1	1	1	1	1	1
65-69.....	54	6	15	7	30	18	3	2	12	2	1	10
70-74.....	272	18	62	7	30	88	31	11	56	19	6	56
75-79.....	2,088	165	553	101	271	592	197	72	658	667	247	60
80-84.....	17,385	1,265	4,499	577	2,226	5,045	1,692	658	667	247	60	449
85-89.....	39,796	2,556	9,549	1,281	4,822	12,209	3,742	1,684	1,725	616	127	1,455
90-94.....	29,821	1,909	6,749	993	3,294	9,723	2,268	1,291	1,339	723	125	1,407
95-99.....	10,190	641	2,241	314	965	3,685	628	378	456	309	34	529
100-104.....	1,831	130	437	62	124	705	110	58	74	46	3	82
Number measured.....	101,478	6,693	24,126	3,336	11,740	32,086	8,672	4,154	4,333	1,993	356	3,989
Not measured.....	853											
Total.....	102,333											
Mean chest circumference.....	88.62	88.41	88.37	88.50	88.15	88.96	87.96	88.66	88.90	89.97	88.78	89.65
Standard deviation.....	5.12	5.25	5.18	5.18	4.92	5.20	4.89	4.88	4.97	5.06	4.96	4.96

SECTION B: PROPORTIONAL NUMBER OF THE VARIOUS CHEST CIRCUMFERENCES (REST) TO EACH 1,000 FOR A ZONE.

60-64.....	0.40	0.45	0.87	0.30	0.26	0.34	0.12	0.18	0.92	0.50	0.25	0.25
65-69.....	53	90	62	2.10	2.56	2.74	3.57	2.65	2.77	1.00	2.81	2.51
70-74.....	2.68	2.69	2.57	2.10	2.56	2.74	3.57	2.65	2.77	1.00	2.81	2.51
75-79.....	20.58	24.65	22.92	30.28	23.08	18.45	22.72	17.33	12.92	9.53	16.85	14.04
80-84.....	171.32	189.01	186.48	172.96	189.60	157.24	195.11	158.40	153.93	123.93	168.54	112.56
85-89.....	392.16	381.89	395.80	383.99	410.73	390.51	431.50	405.39	398.11	324.13	356.74	361.75
90-94.....	293.88	285.22	279.73	297.66	280.57	303.03	261.53	310.78	309.02	362.77	351.12	352.72
95-99.....	100.42	95.77	92.89	94.13	82.18	115.16	72.42	90.99	105.24	155.05	95.51	132.62
100-104.....	18.05	19.42	18.11	18.58	10.56	21.97	12.68	13.96	17.08	23.08	8.43	20.56
Total.....	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00

SECTION C: PROPORTIONAL NUMBER OF EACH 1,000 CHEST CIRCUMFERENCES (REST) IN THE VARIOUS DISTRIBUTION ZONES.

Chest circumference, in centimeters.	Zone 1.	Zone 2.	Zone 4.	Zone 5.	Zone 7.	Zone 8.	Zone 9.	Zone 10.	Zone 11.	Zone 12.	Zone 13.	Total
60-64.....	73.17	512.20	24.39	73.17	268.29	24.39	24.39	1,000
65-69.....	111.11	277.78	92.59	333.33	55.56	37.04	74.07	18.52	1,000
70-74.....	66.18	227.94	25.74	110.29	323.53	113.97	40.44	44.12	7.35	3.68	36.76	1,000
75-79.....	79.02	264.85	48.37	129.79	283.52	94.35	34.48	26.82	9.10	2.87	26.82	1,000
80-84.....	72.76	258.79	33.19	128.04	290.19	97.33	37.85	38.37	14.21	3.45	25.83	1,000
85-89.....	64.23	239.95	32.19	121.17	306.79	94.03	42.32	43.35	16.23	3.19	36.56	1,000
90-94.....	64.02	226.32	33.30	110.46	326.05	76.05	43.29	44.90	24.24	4.19	47.18	1,000
95-99.....	62.90	219.92	30.81	94.70	362.61	61.63	37.10	44.75	30.32	3.34	51.91	1,000
100-104.....	71.00	238.67	33.86	67.72	385.04	60.08	31.68	40.42	25.12	1.64	44.78	1,000
Average chest for each zone.....	65.96	237.75	32.87	115.09	316.19	85.46	40.91	42.70	19.64	3.51	39.31	1,000

TABLE CXXXVII.—Comparative frequency distribution of waist circumference, by Q. M. C. distribution zones, white and colored troops, at demobilization.

SECTION A: ABSOLUTE NUMBERS.

Waist circumference, in centimeters.	Total.	Zone 1.	Zone 2.	Zone 4.	Zone 5.	Zone 7.	Zone 8.	Zone 9.	Zone 10.	Zone 11.	Zone 12.	Zone 13.
60-64.....	350	30	108	12	23	116	28	3	9	7	3	11
65-69.....	4,373	404	1,192	163	345	1,541	348	63	121	53	8	135
70-74.....	24,442	1,895	6,372	765	2,496	7,985	2,072	674	898	387	77	821
75-79.....	36,986	2,464	8,758	1,242	4,355	11,441	3,180	1,565	1,715	705	122	1,439
80-84.....	22,916	1,237	5,042	784	2,915	7,153	1,988	1,166	1,069	538	101	923
85-89.....	8,971	446	1,880	264	1,200	2,744	796	512	391	229	29	480
90-94.....	2,412	144	520	82	277	788	185	125	95	51	9	136
95-99.....	806	44	205	28	97	252	63	35	27	16	39
100-104.....	240	23	64	13	24	71	10	11	11	2	3	8
105-109.....	74	2	22	4	10	14	7	2	7	2	1	3
110.....	6	1	1	2	1	1
Number measured.....	101,576	6,689	24,164	3,357	11,743	32,107	8,678	4,157	4,343	1,990	353	3,995
Not measured.....	757
Total.....	102,333
Mean waist circumference.....	77.92	77.05	77.53	77.94	78.48	77.79	77.94	79.37	78.35	78.81	78.49	78.73
Standard deviation.....	5.96	5.94	6.00	6.04	5.80	5.99	5.83	5.64	5.69	5.83	6.03	6.12

SECTION B: COMPARATIVE NUMBER OF THE VARIOUS WAIST CIRCUMFERENCES TO EACH 1,000 FOR A ZONE.

60-64.....	3.45	4.49	4.47	3.57	1.96	3.61	3.23	.72	2.07	3.52	8.50	2.75
65-69.....	43.06	60.40	49.33	45.56	29.38	48.01	40.10	15.15	27.86	26.63	22.66	33.79
70-74.....	240.64	253.30	263.70	227.88	212.55	248.71	238.76	162.12	206.77	194.49	218.13	205.51
75-79.....	364.13	368.37	362.43	369.97	370.85	356.34	366.44	376.47	394.99	354.27	345.61	360.20
80-84.....	225.60	184.93	206.66	233.54	248.23	222.79	229.08	280.48	246.14	270.34	286.12	231.04
85-89.....	88.32	66.68	77.80	78.64	102.19	85.48	91.72	123.16	90.03	115.07	82.15	120.15
90-94.....	23.75	21.53	21.52	24.43	23.59	24.55	21.32	30.07	21.87	25.62	25.50	34.04
95-99.....	7.94	6.58	8.48	8.34	8.26	7.85	7.26	8.42	6.22	8.04	9.76
100-104.....	2.36	3.44	2.65	3.67	2.04	2.21	1.15	2.65	2.53	1.00	8.50	2.00
105-109.....	.73	.30	.91	1.19	.85	.44	.81	.48	1.61	1.00	2.83	.75
110.....	.060409	.06	.12	.24
Total.....	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00

SECTION C: PROPORTIONAL NUMBER OF EACH 1,000 WAIST CIRCUMFERENCES IN THE VARIOUS DISTRIBUTION ZONES.

Waist circumference, in centimeters.	Zone 1.	Zone 2.	Zone 4.	Zone 5.	Zone 7.	Zone 8.	Zone 9.	Zone 10.	Zone 11.	Zone 12.	Zone 13.	Total.
60-64.....	85.71	308.57	34.29	65.71	331.43	80.00	8.57	25.71	20.00	8.57	31.43	1,000
65-69.....	92.39	272.58	37.27	78.89	352.39	79.58	14.41	27.67	12.12	1.83	30.87	1,000
70-74.....	77.53	260.70	31.30	102.12	326.69	84.77	27.58	36.74	15.83	3.15	33.59	1,000
75-79.....	66.62	236.79	33.58	117.75	309.33	85.98	42.31	46.37	19.06	3.30	38.91	1,000
80-84.....	53.96	220.02	34.21	127.20	312.14	96.75	50.88	46.65	23.48	4.41	40.28	1,000
85-89.....	49.72	209.56	29.43	133.76	306.87	88.73	57.07	43.58	25.53	3.23	53.51	1,000
90-94.....	59.70	215.59	34.00	114.84	326.70	76.70	51.82	39.39	21.14	3.73	56.38	1,000
95-99.....	54.59	254.34	34.74	120.35	312.66	78.16	43.42	33.50	19.85	48.39	1,000
100-104.....	95.83	266.67	54.17	100.00	295.83	41.67	45.83	45.83	8.33	12.50	33.33	1,000
105-109.....	27.03	297.30	54.05	135.14	189.19	94.59	27.03	94.59	27.03	13.51	40.54	1,000
110.....	166.67	166.67	333.33	166.67	166.67	1,000
Average proportion for each zone.....	65.85	237.89	33.05	115.61	316.00	85.43	40.93	42.76	19.59	3.48	39.33	1,000

TABLE CXXXVIII.—Comparative frequency of eye color in the various States of nativity of demobilized men.

State.	Total.	Clear blue.	Blue with brown spots.	Light brown.	Dark brown.	No color.
Alabama.....	1,932	246	881	274	515	16
Alaska.....	13	7	2	1	3	
Arizona.....	130	43	32	24	30	1
Arkansas.....	2,582	1,064	423	381	701	13
California.....	483	189	108	84	97	5
Colorado.....	227	93	46	47	41	
Connecticut.....	997	464	138	192	198	5
Delaware.....	300	127	71	29	72	1
District of Columbia.....	231	87	34	31	79	
Florida.....	1,024	97	443	146	328	10
Georgia.....	3,403	330	1,433	460	1,138	42
Idaho.....	164	77	23	46	17	1
Illinois.....	6,708	3,112	1,363	1,221	995	17
Indiana.....	3,955	1,265	1,616	450	598	26
Iowa.....	1,610	679	451	251	220	9
Kansas.....	1,015	433	248	167	157	10
Kentucky.....	2,934	565	1,510	280	545	34
Louisiana.....	2,079	362	315	387	1,006	9
Maine.....	694	365	90	162	60	17
Maryland.....	1,142	387	222	191	331	11
Massachusetts.....	4,795	2,365	521	1,043	739	127
Michigan.....	3,728	1,821	626	728	538	15
Minnesota.....	1,951	969	485	280	211	6
Mississippi.....	2,102	582	435	375	604	16
Missouri.....	2,847	651	1,420	266	471	39
Montana.....	266	122	41	64	38	1
Nebraska.....	823	353	218	124	126	2
Nevada.....	18	2	5	7	4	
New Hampshire.....	414	201	59	101	47	6
New Jersey.....	3,188	1,374	477	665	653	19
New Mexico.....	230	69	38	54	68	1
New York.....	9,240	3,845	1,247	1,716	2,384	48
North Carolina.....	1,815	479	366	210	734	26
North Dakota.....	358	158	101	58	41	
Ohio.....	7,094	3,027	1,336	1,387	1,297	47
Oklahoma.....	2,316	1,008	408	404	486	10
Oregon.....	1,070	529	164	184	190	3
Pennsylvania.....	10,901	4,381	1,795	2,409	2,257	59
Rhode Island.....	403	186	41	89	84	3
South Carolina.....	829	128	296	128	257	20
South Dakota.....	416	177	114	63	60	2
Tennessee.....	2,815	426	1,463	442	462	22
Texas.....	4,374	1,511	904	787	1,145	27
Utah.....	105	51	18	24	12	
Vermont.....	447	229	49	99	41	29
Virginia.....	1,930	614	339	243	723	11
Washington.....	2,025	996	332	361	334	12
West Virginia.....	1,697	726	335	317	311	8
Wisconsin.....	2,677	1,441	474	453	273	6
Wyoming.....	80	31	15	20	13	1
Total.....	102,577	38,354	23,571	17,955	21,824	793

TABLE CXXXIX.—Comparative frequency of hair color in various States of nativity of demobilized men.

State.	Total.	No color.	Flaxen.	Light brown.	Medium brown.	Dark brown.	Red.	Red and black.
Alabama.....	1,932	19	16	287	782	776	20	32
Alaska.....	13		1	5	7			
Arizona.....	130	1	2	25	20	72		10
Arkansas.....	2,582	12	131	490	471	1,116	31	301
California.....	483	4	20	126	87	213	5	28
Colorado.....	227		14	49	44	100	3	17
Connecticut.....	997	4	60	160	187	518	16	52
Delaware.....	300	2	3	55	54	168	2	16
District of Columbia.....	231	1	8	40	23	143	1	15
Florida.....	1,024	7	8	152	351	477	17	12
Georgia.....	3,403	39	29	449	1,183	1,619	43	41
Idaho.....	164	1	11	45	25	78	2	2
Illinois.....	6,708	20	329	2,000	1,649	2,238	109	363
Indiana.....	3,955	20	237	995	1,241	1,343	49	70
Iowa.....	1,610	4	122	422	376	622	15	49
Kansas.....	1,015	6	62	261	205	425	14	42
Kentucky.....	2,934	28	63	563	1,083	1,081	39	77
Louisiana.....	2,079	10	29	270	237	1,360	13	160
Maine.....	694	17	38	118	147	360	3	11
Maryland.....	1,142	7	53	201	186	604	19	72
Massachusetts.....	4,795	126	281	804	849	2,498	67	170
Michigan.....	3,728	8	280	1,190	646	1,484	40	80
Minnesota.....	1,951	4	195	587	430	629	40	66
Mississippi.....	2,102	13	43	411	381	1,159	15	80
Missouri.....	2,847	24	97	646	1,003	962	34	81
Montana.....	266	23	62	45	116	6	14	
Nebraska.....	823	3	51	219	162	350	11	27
Nevada.....	18			6	1	11		
New Hampshire.....	414	7	25	67	85	205	8	17
New Jersey.....	3,188	17	152	723	431	1,667	50	148
New Mexico.....	230	3	11	28	31	144		13
New York.....	9,240	39	347	1,765	1,224	5,212	138	515
North Carolina.....	1,815	25	38	228	278	1,207	14	25
North Dakota.....	358	2	25	102	78	131	6	11
Ohio.....	7,094	30	472	2,183	1,176	3,007	67	159
Oklahoma.....	2,316	7	103	465	441	1,057	28	215
Oregon.....	1,070	63	302	234	414	21	32	4
Pennsylvania.....	10,901	57	529	2,329	1,588	5,703	136	559
Rhode Island.....	403	2	19	52	84	219	6	21
South Carolina.....	829	19	9	110	261	400	6	24
South Dakota.....	416	6	40	107	92	154	4	13
Tennessee.....	2,815	21	40	488	1,255	930	39	42
Texas.....	4,374	19	310	742	712	2,044	64	479
Utah.....	106		15	29	18	39	2	2
Vermont.....	447	28	26	90	73	214	5	11
Virginia.....	1,930	13	67	309	275	1,143	20	103
Washington.....	2,025	11	107	550	450	811	33	63
West Virginia.....	1,697	6	84	392	244	871	23	83
Wisconsin.....	2,677	8	157	880	646	799	20	127
Wyoming.....	80	1	5	22	19	23	2	8
Total.....	102,577	747	5,132	22,506	21,656	46,446	1,329	4,516

INDEX.

	Page.
Admiral Islands, shoulder breadth of inhabitants of.....	204
Africa, Central, average weight of adult males of tribes of.....	120
African Negroes:	
relative arm length in	219
thoracic index for	207, 209
African tribes:	
relative leg length of	221
relative pubic height of	200
relatively small pelvis of	215
smallest thigh circumference found in	227
Age:	
and stature of Civil War volunteers	35
distribution of Civil War volunteers and World War troops	65
mean stature of each, 18 to 25 years, United States Army recruits	73
of recruits	64
Ages of soldiers (officers and men) serving in Civil and World Wars	64
Agricultural groups:	
foreign and native white—	
weight distribution of draft recruits of	133, 134
mean chest circumference of draft recruits of	150
mean height of	108
mean weight of draft recruits of	132
native white, South—	
height distribution of	109, 110
mean chest circumference of draft recruits of	150
mean height of	108
mean weight of draft recruits of	132
weight distribution of draft recruits of	133, 134
Negroes, 45 per cent plus—	
height distribution of	109, 110
mean chest circumference of draft recruits of	150
mean height of	108
mean weight of draft recruits of	132
weight distribution of draft recruits of	133, 134
North, native white over 73 per cent—	
height distribution of	109, 110
mean chest circumference of draft recruits of	150
mean weight of draft recruits of	132
weight distribution of draft recruits of	133, 134
of sections—	
mixed foreign and white, index of build for draft recruits of	173
native, North, index of build of draft recruits of	173
Negro, index of draft recruits of	173
Negro, relative chest circumference of draft recruits of	150
northern, native white, relative chest circumference of draft recruits of	150
relative chest circumference of draft recruits of	150
southern, native white, index of build for draft recruits of	173
southern white, relative chest circumference of	150
Ainos, relative pubic height of	200
sitting height of	190
Alabama:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919	282
with clear blue eyes, demobilization, 1919	281
with dark brown eyes, demobilization, 1919	283
with dark brown hair, demobilization, 1919	291
with flaxen hair, demobilization, 1919	288
with light brown eyes, demobilization, 1919	283
with light brown hair, demobilization, 1919	290
with medium brown hair, demobilization, 1919	290
with red hair, demobilization, 1919	289

Alabama—Continued.	Page.
average weight of draft recruits from, at mobilization, 1917-1918, and demobilization, 1919	122
comparative view of mean height and mean weight of men from	124
increase in stature of soldiers at demobilization over stature of recruits, 1917-1919	76
index of build for, demobilization, 1919	167
index of build for recruits from, mobilization, 1917-1918	166
mean chest circumference (expiration), of draft recruits from	142
mean stature of draft recruits from	75
mean stature of soldiers from, demobilization, 1919	76
relative chest circumference of recruits from	144
Alaska:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919	282
with clear blue eyes, demobilization, 1919	281
with dark brown eyes, demobilization, 1919	283
with flaxen hair, demobilization, 1919	288
with light brown eyes, demobilization, 1919	283
with light brown hair, demobilization, 1919	290
with medium brown hair, demobilization, 1919	290
average weight of draft recruits from, at mobilization, 1917-1918, and demobilization, 1919	122
comparative view of mean height and mean weight of men from	124
difference of weight of draft recruits from, at mobilization, 1917-1918, and demobilization, 1919	123
increase in stature of soldiers at demobilization over stature of recruits, 1917-1919	76
index of build for, demobilization, 1919	167
index of build for recruits from, at mobilization, 1917-1918	166
mean chest circumference (expiration) of draft recruits from	142
mean stature of draft recruits from	75
mean stature of soldiers from, at demobilization, 1919	76
relative chest circumference of recruits from	144
American Indians, relative leg length of	221
Anorchism, cryptorchidism, hypospadias, and monorchism:	
chest circumference of drafted men with	382
correlation between height and chest circumference (expiration), in recruits with	381
correlation between height and weight in recruits with	380
robustness of drafted men with	382
stature of drafted men with	379
weight of drafted men with	382
Anthropoids (see also Apes, anthropoid):	
transverse diameter of pelvis in	215
Anthropological work in connection with the Army, 1917-1919, history of the	49
Anthropologists:	
Army, daily reports from	55
list of, who supervised taking measurements in camps	56
Anthropology, importance of, in the Army	34
Anthropometric work in connection with draft recruits	49
Anthropometry, importance of, in the Army	45
Apparatus used in taking measurements	57
Apes, anthropoid, shoulder breadth of	204
Arizona:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919	282
with clear blue eyes, demobilization, 1919	281
with dark brown eyes, demobilization, 1919	283
with dark brown hair, demobilization, 1919	291
with flaxen hair, demobilization, 1919	288
with light brown eyes, demobilization, 1919	283
with light brown hair, demobilization, 1919	290
with medium brown hair, demobilization, 1919	290
average weight of draft recruits, at mobilization, 1917-1918, and demobilization, 1919	122
comparative view of mean height and mean weight of men from	124
difference of weight of draft recruits from, at mobilization, 1917-1918, and demobilization, 1919	123
increase in stature of soldiers at demobilization over recruits, 1917-1919	76
index of build for, at demobilization, 1919	167
index of build for recruits from, at mobilization, 1917-1918	166
mean chest circumference (expiration), of draft recruits from	142
mean stature of draft recruits from	75
mean stature of soldiers from, at demobilization, 1919	76
relative chest circumference of recruits from	144

Arkansas:

absolute and relative numbers of veterans from—	Page.
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits from, at mobilization, 1917-1918, and demobilization, 1919.....	122
comparative view of mean height and mean weight of men from.....	124
difference of weight of draft recruits from, at mobilization, 1917-1918, and demobilization, 1919.....	123
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build for, at demobilization, 1919.....	167
index of build for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, demobilization, 1919.....	96
relative chest circumference of recruits from.....	144

Arm and forearm:

correlation of, in white troops.....	42
length of, correlation between, colored troops, demobilization.....	531
length of, correlation between, white troops, demobilization.....	516

Arm length:

and forearm, correlation between, Negro troops.....	267
and forearm, correlation between, white troops.....	262
average, in color races, demobilization, 1919.....	235
comparison of, in white and Negro troops.....	40, 233
general discussion.....	219
mean, in white troops.....	219
relative, in African Negroes.....	219
in Bavarians.....	219
in French.....	219
in Mawambu pygmies.....	219
in Ba-Binga (African race).....	220
in Bugu (African race).....	220
in Lobi (African race).....	220
standard deviation of.....	220
summary of, of approximately 6,000 colored troops, demobilization.....	234
summary of, of approximately 100,000 white troops, demobilization.....	234
total, and blouse groups, association between, colored troops, demobilization.....	547
total, and blouse groups, association between, white troops, demobilization.....	539

Armenians, relative leg length of.....

221

Army, history of anthropological work in connection with, 1917-1919.....

49

Asiatic peoples, ratio of pubic height to total stature of.....

199

Asthma:

and chest circumference.....	43
chest circumference of drafted men with.....	357
correlation between height and chest circumference (expiration), in draft recruits with.....	359
correlation between height and weight in recruits with.....	358
Pignet's index of build for drafted men with.....	360
robustness in drafted men with.....	360
stature of drafted men with.....	356
weight of drafted men with.....	356

Astigmatism, chest circumference in drafted men with.....

322

correlation between height and chest circumference in recruits with.....

322

correlation between height and weight in recruits with.....

321

in drafted men.....

319

Pignet's index of build in drafted men with.....

322

robustness of drafted men with.....

322

stature of drafted men with.....

319

weight of drafted men with.....

320

Athletes, weights of.....

118, 119

Australians, calf circumference of.....

230

Austria, standards of stature in.....

45

Austrian and German groups:	Page.
height distribution of	109, 110
mean chest circumference of draft recruits of	150
mean height of	108
mean weight of draft recruits of	132
of sections, relative chest circumference of draft recruits of	150
weight distribution of draft recruits of	133, 134
Ba-Binga (African race):	
relative arm length of	220
relative thigh circumference for	227
Ba-Tua tribe (African):	
pelvic diameter of	215
thigh circumference in	227
Bavarians:	
approximate average stature of	47
chest circumference of	139
relative arm length of	219
relative forearm length of	220
relative shoulder breadth of	204
transverse diameter of pelvis of	215
Upper, average weight of adult male	120
Belgians:	
approximate average stature of	47
average weight of adult male	120
ratio of pubic height to total stature of	199
relative chest circumference of	139
relative shoulder breadth of	204
sitting height of	190
Belgium, standards of stature in	45
Bertillon's proportions, identification by, belongs to anthropology	34
Black (one-quarter) race, comparative frequency distribution of measurements of, demobilization	236-240
Blouse groups	271
and antero-posterior diameter of chest, association between, colored troops, demobilization	544
and chest diameter, antero-posterior, association between, white troops, demobilization	536
and chest transverse diameter, association between, white troops, demobilization	535
and neck circumference, association between, colored troops, demobilization	546
and neck circumference, association between, white troops, demobilization	538
and shoulder width, association between, colored troops, demobilization	542
and shoulder width, association between, white troops, demobilization	534
and suprapatella circumference, association between, colored troops, demobilization	559
and total arm length, association between, colored troops, demobilization	547
and total arm length, association between, white troops, demobilization	539
and transverse diameter of chest, colored troops, demobilization	543
and transverse pelvic diameter, association between, colored troops, demobilization	545
and transverse pelvic diameter, association between, white troops, demobilization	537
and weight, association between, colored troops, demobilization	541
and weight, association between, white troops, demobilization	533
comparative frequency distribution of, by States of nativity, colored troops	563
comparative frequency distribution of, by States of nativity, white troops	562
dimensions associated with, Negro troops, demobilization	273
dimensions associated with, white troops, demobilization	273
Blouses, measurements for	271
Body, size and proportions of, essential to proper cutting of uniforms	34
Bosnian-Herzegovinians (soldiers), average stature of	68
Breeches groups:	
and calf circumference, association between, white troops, demobilization	554
and circumference at patella, association between, colored troops, demobilization	560
and circumference of calf, association between, colored troops, demobilization	561
and knee height, association between, colored troops, demobilization	557
and knee height, association between, white troops, demobilization	550
and patella circumference, association between, white troops, demobilization	553
and suprapatella, association between, white troops, demobilization	552
and thigh circumference, association between, colored troops, demobilization	558
and thigh circumference, association between, white troops, demobilization	551
and transverse diameter of pelvis, association between, colored troops, demobilization	556
and transverse pelvic diameter, association between, white troops, demobilization	549
comparative frequency distribution of, by States of nativity, colored troops	565
comparative frequency distribution of, by States of nativity, white troops	564

Breeches groups—Continued.	Page.
dimensions associated with, Negro troops, demobilization.....	275
dimensions associated with, white troops, demobilization.....	274
Breeches, measurements for.....	274
British army, minimum stature in.....	45
British, inhabitants of United Kingdom of Great Britain and Ireland, average stature of...	68
Bugu (African race):	
relative arm length of.....	220
thoracic index for.....	207, 209
Build (<i>see also</i> Robustness).....	37
average index for men at demobilization, 1919.....	164
average index of, for recruits, World War.....	164
comparison of index of, in recruits of 1917-1918 and in veterans of 1919 and 1864-1865...	38, 168
distribution index of, by States.....	165
importance of the index of.....	162
index of.....	37
and Pignet's index of robustness of recruits found with specified diseases and defects.	408
at mobilization, by States, 1917-1918.....	166
average, in color races, demobilization, 1919.....	235
by groups of sections, 1917-1918.....	173
by sections.....	170, 172
by States, demobilization, 1919.....	167
calculated by three methods, first million draft recruits.....	164
comparison of, of men at demobilization in 1865 and 1919.....	170
for mean stature and weight.....	164
in eight European races, demobilization, 1919.....	243, 244
increase in, at demobilization, 1919, over mobilization, 1917-1918.....	168
mean, of color races.....	174
mean, of eight European races of men at demobilization.....	173
of Civil War veterans (white troops) and World War veterans (white and Negro troops).....	164, 165
of recruits, by sections, 1917-1918.....	172
Pignet's, for men of various heights with average chest and weight for certain per cents of men of each height.....	188
method of determining.....	163
variability of, by States.....	37
Bulgarians:	
chest circumference of.....	139
of western Bulgaria, average stature of.....	68
Bushmen:	
pelvic diameter of.....	215
relative pubic height of.....	200
Calf circumference.....	39
absolute and relative, in eight European races, demobilization, 1919.....	231
and breeches groups, association between, colored troops, demobilization.....	561
and breeches groups, association between, white troops, demobilization.....	554
average, in color races, demobilization, 1919.....	235
comparative frequency distribution of, in eight European races, demobilization.....	231
comparison of, in eight European races.....	230, 231
general discussion of.....	230
in eight European races at demobilization, 1919.....	243
in European groups.....	230
in Malay groups.....	230
in Mongoloid groups.....	230
in Negro troops.....	230
in white troops.....	230
maximum, directions for measuring.....	59
standard deviation of.....	230
summary of, of approximately 6,000 colored troops, demobilization.....	234
summary of, of approximately 100,000 white troops, demobilization.....	234
California:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289

California—Continued.	Page.
average weight of draft recruits, at mobilization, 1917-1918, and demobilization, 1919..	122
comparative view of mean height and mean weight of men from.....	124
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build for—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144
Calipers, wooden sliding, used in taking measurements.....	57
Camps, number of men measured at.....	56
Army anthropology an aid in detecting.....	34
Cardiac disorders (<i>see also</i> , Heart).	
Cardiac hypertrophy:	
chest circumference in drafted men with.....	334
correlation between height and chest circumference in recruits with.....	333
correlation between height and weight in recruits with.....	332
in drafted men.....	331
Pignet's index of build in drafted men with.....	334
robustness in drafted men with.....	334
stature of drafted men with.....	331
weight of drafted men with.....	331
Cards, statistical perforated.....	63
Cavalry, stature of, in Great Britain.....	45
Characteristics, racial, and classification of troops.....	34
Centimeters, table for converting into inches.....	410
Chest:	
antero-posterior diameter of—	
and thoracic index.....	206
and blouse groups, association between, colored troops, demobilization.....	544
average, in color races, demobilization, 1919.....	235
directions for measuring.....	58
in eight European races, demobilization, 1919.....	243
summary of, of approximately 6,000 colored troops, demobilization.....	234
summary of, of approximately 100,000 white troops, demobilization.....	234
comparative frequency distribution of antero-posterior diameter of, in each of eight European races, demobilization, 1919.....	212
comparative frequency distribution of transverse diameter of, in each of eight European races, demobilization, 1919.....	211
Chest circumference (<i>see also</i> , Chest girth).....	37
and asthma.....	43
and diseases of the heart.....	43
and neck circumference, Negro troops, correlation between.....	266
and neck circumference, correlation between, white troops.....	260
and sitting height, Negro troops, correlation between.....	266
and sitting height, correlation between, in white troops.....	42
and sitting height, correlation between, white troops, demobilization.....	532, 533
and stature, correlation between.....	256
and stature, correlation between, Civil War recruits.....	256
and stature, correlation between, recruits, 1917-1918.....	256
is, Negro troops, correlation between.....	267
is between crista, correlation between, white troops.....	258
, correlation between, white troops.....	261
.....	43
.....	43
n.....	258
lation between.....	265
n, white troops.....	42, 260
recruits found with special diseases or defects.....	410
.....	139, 140
lization, 1919.....	235
Civil War data.....	138
s.....	139
arious classes of.....	140
tribution of, mobilization, 1917-1918, and demobiliza-.....	139
tates.....	140, 142, 144
.....	136
relation to various diseases and defects.....	296

Chest circumference—Continued.	Page.
important in relation to general size.....	137
in drafted men—	
mean, by groups of sections.....	150, 151, 152
with asthma.....	357
with astigmatism.....	322
with cardiac hypertrophy.....	334
with cryptorchidism, hypospadias, anorchism, and monorchism.....	382
with defective physical development.....	375
with defective and deficient teeth.....	363
with enlarged inguinal rings.....	369
with exophthalmic goiter.....	309
with flat-foot.....	370
with hernia.....	364
with hyperopia.....	318
with hypertrophic tonsillitis.....	326
with mitral insufficiency.....	338
with mitral stenosis.....	339
with myopia.....	315
with overweight and obesity.....	379
with pulmonary tuberculosis.....	301
with simple goiter.....	305
with simple tachycardia.....	330
with underweight.....	376
with valvular diseases of the heart.....	342
with varicocele.....	347
with varicose veins.....	347
in eight European races at demobilization, 1919.....	152, 154, 155, 243
in relation to blouse groups.....	271
in relation to stature.....	48
increase in mean, of veterans.....	37
level of nipples, directions for measuring.....	58
mean, at expiration.....	138
mean, by groups of sections.....	149, 150, 151, 153
mean, by sections.....	144, 145, 146, 147
mean, of first million draft recruits.....	37
medico-military importance of.....	136
methods of measurement.....	137
occupational significance of.....	137
of men of color races.....	156
of recruits, comparison of, with other countries.....	139
racial significance of.....	137
relative, by States, first million draft recruits.....	144
requirements of Army Regulations regarding.....	298
standard deviation of, by sections.....	147, 148
variability according to race.....	37
Chest circumference (expiration):	
and height classes.....	158
and height, correlation between—	
first million draft recruits.....	420, 421
group 1, agricultural, North, native white, 73 per cent plus.....	443
group 2, agricultural, mixed foreign and native white, North, first million draft recruits.....	446
group 3, agricultural, native white, South, first million draft recruits.....	449
group 4, agricultural, Negro, 45 per cent plus, first million draft recruits.....	452
group 5, eastern manufacturing, first million draft recruits.....	455
group 6, commuter, first million draft recruits.....	458
group 7, mining, first million draft recruits.....	461
group 8, sparsely settled, first million draft recruits.....	464
group 9, desert, first million draft recruits.....	467
group 10, maritime, first million draft recruits.....	470
group 11, mountain, first million draft recruits.....	473
group 12, mountain whites, first million draft recruits.....	476
group 13, Indian, first million draft recruits.....	479
group 14, Mexican, first million draft recruits.....	482
group 15, native whites of Scotch origin, first million draft recruits.....	485
group 16, Russian, first million draft recruits.....	489
group 17, Scandinavian, first million draft recruits.....	491
group 18, Finn, first million draft recruits.....	494
group 19, French-Canadian, first million draft recruits.....	497
group 20, German and Scandinavian, first million draft recruits.....	500

Chest circumference—Continued.

and height, correlation between—Continued.	Page.
group 21, German and Austrian, first million draft recruits.....	503
group 22, German and Austrian, first million draft recruits.....	506
and weight classes, first million draft recruits.....	162
and weight, correlation between—	
first million draft recruits.....	422, 423
group 1, agricultural, North, native white, first million draft recruits.....	444
group 2, agricultural, mixed foreign and white, North, first million draft recruits.....	447
group 3, agricultural, native white, South, first million draft recruits.....	450
group 4, agricultural, Negro, 45 per cent plus, first million draft recruits.....	453
group 5, eastern manufacturing, first million draft recruits.....	456
group 6, commuter, first million draft recruits.....	459
group 7, mining, first million draft recruits.....	462
group 8, sparsely settled, first million draft recruits.....	465
group 9, desert, first million draft recruits.....	468
group 10, maritime, first million draft recruits.....	471
group 11, mountain, first million draft recruits.....	474
group 12, mountain whites, first million draft recruits.....	477
group 13, Indian, first million draft recruits.....	480
group 14, Mexican, sparsely settled, first million draft recruits.....	483
group 15, native whites of Scotch origin, first million draft recruits.....	486
group 16, Russian, first million draft recruits.....	489
group 17, Scandinavian, first million draft recruits.....	492
group 18, Finn, first million draft recruits.....	495
group 19, French-Canadian, first million draft recruits.....	498
group 20, German and Scandinavian, first million draft recruits.....	501
group 21, German and Austrian, first million draft recruits.....	504
group 22, German and Austrian, first million draft recruits.....	507
distribution, by groups of sections.....	153
distribution by height, first million draft recruits.....	161
distribution by special diseases or defects.....	392, 393, 402, 403
heights, and weights, various, shown for United States, first million draft recruits....	157
mean, by sections, first million draft recruits.....	145
mean, by States, first million draft recruits.....	142
of native American white draft recruits of Civil War.....	142
standard deviation of, by sections, first million draft recruits.....	148
weight, and height, measurements showing proportionate measurements of two of them to the total of the third, first million draft recruits.....	177
Chest circumference (rest):	
absolute and relative, in eight European races, demobilization.....	154
and neck circumference, correlation between, colored troops, demobilization.....	527
and neck circumference, correlation between, white troops, demobilization.....	512
and sitting height, association between, colored troops.....	540
and transverse pelvis, correlation between, colored troops, demobilization.....	528
and transverse pelvic diameter, correlation between, white troops, demobilization....	513
and weight, correlation between, white troops, demobilization.....	511
comparative frequency distribution of, in eight European races, demobilization.....	155
comparative frequency distribution of, by Quartermaster Corps distribution zones. white and colored troops, at demobilization.....	571
summary of, of approximately 6,000 colored troops, demobilization.....	234
summary of, of approximately 100,000 white troops, demobilization.....	234
Chest diameter.....	39
absolute and relative, demobilization, 1919.....	210
antero-posterior—	
and blouse groups, association between, white troops, demobilization.....	536
and transverse, correlation between, colored troops, demobilization.....	529
and transverse, correlation between, white troops.....	261
and transverse, correlation between, white troops, demobilization.....	514
and transverse, of Negro troops, correlation between.....	266
comparison of, in eight European races.....	209
mean.....	209
standard deviation of.....	209
mean transverse.....	208
transverse (<i>see also</i> , Thoracic index)—	
and antero-posterior, in white troops.....	42
and blouse groups, association between, colored troops, demobilization.....	543
and blouse groups, association between, white troops, demobilization.....	535
at level of nipples, directions for measuring.....	58
average, in color races, demobilization, 1919.....	215
comparison of, in eight European races.....	208, 210

Chest diameter—Continued.	
transverse—Continued.	Page.
general discussion.....	207
in eight European races at demobilization, 1919.....	243
in Navajo Indians.....	207
of the English.....	208
of the French.....	208
of Germans.....	208
of Hebrews.....	208
of Irish.....	208
of Italians.....	208
of Poles.....	208
of Scotch.....	208
standard deviation of.....	208
summary of, of approximately 6,000 colored troops, demobilization.....	234
summary of, of approximately 100,000 white troops, demobilization.....	234
Chest girth (<i>see also</i> , Chest circumference):	
and neck girth, correlation of, in white troops.....	42
and pelvic diameter, correlation of, in white troops.....	42
mean, by groups and component sections, first million draft recruits.....	436
measurement of.....	186
Chest measurement (expiration), by special diseases.....	396
Chimpanzee:	
shoulder breadth of.....	204
transverse diameter of pelvis of.....	215
Chinese:	
absolute and relative height of pubic arch in, demobilization, 1919.....	202
absolute and relative height of sternal notch in, demobilization, 1919.....	199
absolute and relative leg length in, at demobilization, 1919.....	224
absolute and relative shoulder breadth in, demobilization, 1919.....	207
absolute and relative sitting heights in, demobilization, 1919.....	193
absolute and relative transverse diameter of pelvis in, at demobilization, 1919.....	219
chest circumference of.....	139
comparative frequency distribution of measurements in, at demobilization.....	236, 240
general comparison of, with other color races.....	241
index of build for.....	174
mean absolute and relative spans, demobilization, 1919.....	196
northern, ratio of pelvic diameter to stature of.....	215
mean stature of, at demobilization, 1919.....	117
mean weight in, with standard deviation for white and Negro troops, demobilization, 1919.....	136
relative shoulder breadth of.....	204
sitting height of.....	190
South, ratio of pelvic diameter to stature of.....	215
Civil War:	
and World War—	
comparative measurements at demobilization.....	242
recruits, comparison of frequency distribution of statures of.....	72
recruits, comparison of stature of.....	79
troops from different States, comparative view of mean height and mean weight of.....	124
draft recruits, native American white, chest circumference (expiration) of.....	142
recruits, American born, distribution of stature and weight of.....	74
recruits, correlation between chest circumference and stature of.....	256
veterans, index of build for, for each inch of stature.....	164
veterans (white troops) and World War veterans (white and Negro troops), index of build for.....	165
volunteers and World War troops, age distribution of.....	65
Clothing:	
of army, diversity of race size in relation to.....	47
patterns, measurement card for, demobilization, 1919.....	61
Cochin Chinese, approximate average stature of.....	47
Color races:	
chest circumferences of men of the.....	156
comparison of—	
height of pubic arch in.....	202
height of sternal notch in.....	199
leg length in.....	224
neck circumference in.....	203
shoulder breadth of.....	207
transverse pelvic diameter of.....	219

Color races—Continued.	Page.
general comparison of measurements of	241
mean index of build of	174
mean span of	196
Colorado:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919	282
with clear blue eyes, at demobilization, 1919	281
with dark brown eyes, demobilization, 1919	283
with dark brown hair, at demobilization, 1919	291
with flaxen hair, at demobilization, 1919	288
with light brown eyes, demobilization, 1919	283
with light brown hair, demobilization, 1919	290
with medium brown hair, demobilization, 1919	290
with red hair, demobilization, 1919	289
average weight of draft recruits, at mobilization, 1917–1918, and demobilization, 1919	122
comparative view of mean height and mean weight of men from	124
difference of weight of draft recruits from, at mobilization, 1917–1918, and demobiliza- tion, 1919	123
increase in stature of soldiers at demobilization over recruits, 1917–1919	76
index of build for recruits from, at mobilization, 1917–1918	166, 167
mean chest circumference (expiration), of draft recruits from	142
mean stature of draft recruits from	75
mean stature of soldiers from, at demobilization, 1919	76
relative chest circumference of recruits from	144
Colored and white troops, demobilization:	
comparative frequency distribution—	
of chest circumference (rest), by Quartermaster Corps distribution zones	571
of height, by States of nativity	566, 567, 570, 571
of waist circumference, by Quartermaster Corps distribution zones	572
of weight, by States of nativity	570
correlation between stature and waist circumference	509
correlation between stature and weight in	508
Colored troops:	
comparative frequency distribution—	
of blouse groups, by States of nativity	563
of breeches group, by States of nativity	565
demobilization, association between blouse groups—	
and antero-posterior diameter of chest	544
and neck circumference	546
and shoulder width	542
and suprapatella circumference	559
and total arm length	547
and transverse chest diameter	543
and transverse pelvic diameter	545
and weight	541
demobilization, association between breeches groups—	
and calf circumference	561
and knee height	557
and thigh circumference	558
and transverse diameter of pelvis	556
and circumference at patella	560
demobilization, correlation between—	
arm length and forearm	531
waist circumference and leg length	555
chest circumference (rest) and neck circumference	527
chest circumference (rest) and sitting height	540
chest circumference (rest) and transverse pelvis	528
chest, transverse and antero-posterior	529
height and knee height	525
leg length and knee height	526
stature and height of pubic arch	524
stature and height of sternal notch	523
stature and sitting height	521
stature and span	522
waist circumference and transverse diameter of pelvis	530
Commuter groups:	
height distribution of	109, 110
mean chest circumference of draft recruits of	150
mean height of	108

Commuter groups—Continued.	Page.
mean weight of draft recruits of	132
relative chest circumference of draft recruits of	150
weight distribution of draft recruits of	133, 134
Commuters, index of build for, by groups of sections	173
Connecticut:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919	282
with clear blue eyes, demobilization 1919	281
with dark brown eyes, demobilization, 1919	283
with dark brown hair, demobilization, 1919	291
with flaxen hair, demobilization, 1919	288
with light brown eyes, demobilization, 1919	283
with light brown hair, demobilization, 1919	290
with medium brown hair, demobilization, 1919	290
with red hair, demobilization, 1919	289
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919 ..	122
colored eyes, proportional numbers for, 1865	285
comparative view of mean height and mean weight of men from	124
comparison of index of build of men from, at demobilization in 1865 and 1919	170
increase in stature of soldiers at demobilization over stature of recruits, 1917-1919	76
index of build for recruits from, at mobilization, 1917-1918	166, 167
mean chest circumference (expiration), of draft recruits from	142
mean stature of draft recruits from	75
mean stature of soldiers from, at demobilization, 1919	76
relative chest circumference of recruits from	144
constitution, criterion of	186
Correlation:	
between arm length and forearm, white troops	262
between chest circumference and neck circumference, white troops	260
between chest circumference and transverse diameter of pelvis between cristæ, white	
troops	258
between chest circumference and transverse pelvic diameter, white troops	261
between chest circumference and weight, white troops	260
between leg length and knee height, white troops	263
between leg length and waist circumference, white troops	264
Correlations	42
Correlations, tables of. <i>See</i> , Table of Contents.	
Corsicans, average stature of	68
Cossacks, ratio of pubic height to total stature of	199
Criterion of constitution	186
Cryptorchidism, hypospadiæ, anorchism, and monorchism:	
chest circumference in drafted men with	382
correlation between height and chest circumference (expiration), in recruits with	381
correlation between height and weight in drafted men with	380
Pignet's index of build for	382
robustness in drafted men with	382
stature of drafted men with	379
weight of drafted men with	382
Danes:	
approximate average stature of	47
average stature of	68
Defects and diseases:	
height, weight and chest circumference of recruits in relation to various	296
specific, dimensions of drafted men with	299
Defects and physique, relation between	43
Delaware:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919	282
with clear blue eyes, demobilization, 1919	281
with dark brown eyes, demobilization, 1919	283
with dark brown hair, demobilization, 1919	291
with flaxen hair, demobilization, 1919	288
with light brown eyes, demobilization, 1919	283
with light brown hair, demobilization, 1919	290
with medium brown hair demobilization, 1919	290
with red hair, demobilization, 1919	289
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919 ..	122

Delaware—Continued.	Page.
comparative view of mean height and mean weight of men from	124
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build for, at demobilization, 1919.....	167
index of build for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144
Demobilization, increase in weight at, over mobilization	122, 123
Dental caries, correlation between height and chest circumference (expiration) in recruits with deficient and defective teeth and.....	362
correlation of height and weight in recruits with	361
Desert groups:	
mean chest circumference of draft recruits of.....	150
mean height of.....	108
mean weight of draft recruits of.....	132
of sections, index of build for draft recruits of.....	173
of sections, relative chest circumference of draft recruits of.....	150
weight distribution of draft recruits of.....	133, 134
Development, defective physical:	
chest circumference in drafted men with.....	375
correlation between height and chest circumference in recruits with.....	374
correlation between height and weight in drafted men.....	373
robustness in drafted men with.....	375
stature of drafted men with.....	372
weight of drafted men with.....	372
Dimensions:	
absolute, in eight European races, demobilization, 1919.....	243
associated with blouse groups, demobilization.....	273
associated with breeches groups, demobilization.....	274, 275
average, eight European races, demobilization, 1919.....	246, 247, 248, 249
average, in color races, demobilization, 1919.....	235
bodily, in relation to diseases, summary of.....	387
of drafted men with specific defects and diseases.....	299
of manikins.....	276
of white and negro troops, comparison of.....	232, 233
physical, and hemorrhoids.....	43
physical, in relation to disease.....	43
physical, their variability in relation to population.....	44
relative, in color races, demobilization, 1919.....	235
relative, in eight European races, demobilization, 1919.....	245, 250, 251, 252
summary of, of approximately 6,000 colored troops, demobilization.....	234
summary of, of approximately 100,000 white troops, demobilization.....	234
Disease, physical dimensions in relation to.....	43
Diseases and defects:	
height, weight, and chest circumference of recruits in relation to various.....	296
specific, dimensions of drafted men with.....	299
Distribution zones, Quartermaster Corps, sizes and proportions of men in.....	276
District of Columbia:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919.....	122
comparative view of mean height and mean weight of men from.....	124
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build for, at demobilization, 1919.....	167
index of build for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, demobilization, 1919.....	76
relative chest circumference of recruits from.....	144

Draft recruits (<i>see also</i> , Drafted men):	Page.
anthropometric work in connection with.....	49
chest circumference (expiration) by height.....	161
distribution of chest circumference of, by groups of sections.....	151, 152
distribution of stature and weight of.....	74
measurements of.....	51
Pignet's index, men of various heights, percentage distribution of each height.....	189
relative chest circumference of, by groups of sections.....	150
Draft recruits, first and second million:	
chest circumference (expiration) distribution by special diseases or defects.....	402, 403
height distribution by special diseases or defects.....	398, 399
relative weight and relative chest circumference of, with special diseases and defects..	410
variability of stature, associated with various defects and diseases.....	409
variability of weight, associated with various diseases and defects among.....	409
weight distribution by special diseases or defects.....	400, 401
Draft recruits, first million:	
average weight of, by States, at mobilization, 1917-1918, and demobilization, 1919....	122
correlation between height and chest circumference (expiration).....	420, 421
group 1, agricultural, North, native white.....	443
group 2, agricultural, mixed foreign and native white, North.....	446
group 3, agricultural, native white, South.....	449
group 4, agricultural, Negro, 45 per cent plus.....	452
group 5, eastern manufacturing.....	455
group 6, commuter.....	458
group 7, mining.....	461
group 8, sparsely settled.....	464
group 9, desert.....	467
group 10, maritime.....	470
group 11, mountain.....	473
group 12, mountain, whites.....	476
group 13, Indian.....	479
group 14, Mexican.....	482
group 15, native white of Scotch origin.....	485
group 16, Russian.....	488
group 17, Scandinavian.....	491
group 18, Finn.....	494
group 19, French-Canadian.....	497
group 20, German and Scandinavian.....	500
group 21, German and Austrian, 20 per cent plus.....	503
group 22, German and Austrian, 15 per cent plus.....	506
correlation between height and weight.....	417, 418, 419
group 1, agricultural, North, native white.....	442
group 2, agricultural, mixed foreign and native white, North.....	445
group 3, agricultural, native white, South.....	448
group 4, agricultural, Negro, 45 per cent plus.....	451
group 5, eastern manufacturing.....	454
group 6, commuter.....	457
group 7, mining.....	460
group 8, sparsely settled.....	463
group 9, desert.....	466
group 10, maritime.....	469
group 11, mountain.....	472
group 12, mountain whites.....	475
group 13, Indian.....	478
group 14, Mexican.....	481
group 15, native white of Scotch origin.....	484
group 16, Russian.....	487
group 17, Scandinavian.....	490
group 18, Finn.....	493
group 19, French-Canadian.....	496
group 20, German and Scandinavian.....	499
group 21, German and Austrian, 20 per cent plus.....	502, 506
group 22, German and Austrian, 15 per cent plus.....	505
correlation between weight and chest circumference (expiration).....	422, 423
group 1, agricultural, North, native white.....	444
group 2, agricultural, mixed foreign and native white, North.....	447
group 3, agricultural, native white, South.....	450
group 4, agricultural, Negro, 45 per cent plus.....	453
group 5, eastern manufacturing.....	456
group 6, commuter.....	459

Draft recruits—Continued.

correlation between weight and chest circumference—Continued.	Page.
group 7, mining.....	462
group 8, sparsely settled.....	465
group 9, desert.....	468
group 10, maritime.....	471
group 11, mountain.....	474
group 12, mountain whites.....	477
group 13, Indian.....	480
group 14, Mexican.....	483
group 15, native whites of Scotch origin.....	486
group 16, Russian.....	489
group 17, Scandinavian.....	492
group 18, Finn.....	495
group 19, French-Canadian.....	498
group 20, German and Scandinavian.....	501
group 21, German and Austrian, 20 per cent plus.....	504
group 22, German and Austrian, 15 per cent plus.....	507
distribution of, by groups of sections.....	130, 131
height and chest circumference (expiration) classes.....	158
height and weight classes.....	158
height distribution shown by groups of sections.....	109, 110
height, weight, and chest circumference (expiration) measurements showing proportionate measurements of two of them to the total of the third.....	177
index of build, calculated by three methods.....	164
mean chest circumference (expiration) of, by groups of sections.....	150
mean chest circumference (expiration) of, by sections.....	145
mean chest circumference (expiration) of, by States.....	142
mean chest girth of, by groups and component sections.....	436-441
mean height of.....	101
mean height of, by groups and component sections.....	424-429
mean height of, by groups of sections.....	108
mean stature of, by States.....	75
mean weight of.....	125
mean weight of, by groups and component sections.....	430-435
mean weight of, by groups of sections.....	132
relative chest circumference of, by States.....	144
standard deviation of chest circumference (expiration) of, by sections.....	148
total and proportionate measurements—	
by each section.....	182, 183
by groups of sections.....	181
by States.....	178, 179, 180
various heights, weights, and chest circumferences (expiration), shown for United States.....	157
weight and chest circumference (expiration) classes.....	162
weight distribution by height.....	159
weight distribution shown by groups of sections.....	133, 134
Drafted men (<i>see also</i> Draft recruits):	
dimensions of, with specific defects and diseases.....	299
standards of measurements of.....	296
with asthma—	
chest circumference of.....	357
robustness in.....	360
stature of.....	356
weight of.....	356
with astigmatism—	
chest circumference of.....	322
robustness in.....	322
stature of.....	319
weight of.....	320
with cardiac hypertrophy—	
chest circumference of.....	334
robustness in.....	334
stature of.....	331
weight of.....	331
with cryptorchidism, anorchism, monorchism, and hypospadias—	
chest circumference of.....	382
robustness in.....	382
stature of.....	379
weight of.....	382

Drafted men—Continued.

with defective and deficient teeth—	Page.
chest circumference of	363
robustness in	363
stature of	360
weight of	360
with defective physical development—	
chest circumference of	375
robustness in	375
stature of	372
weight of	372
with enlarged inguinal rings—	
chest circumference of	369
robustness in	369
stature of	369
weight of	369
with exophthalmic goiter	308
chest circumference in	309
robustness in	309
stature of	308
weight of	308
with flat-foot—	
chest circumference of	370
robustness in	370
stature of	370
weight of	370
with hemorrhoids—	
chest circumference of	356
robustness in	356
stature of	353
weight of	353
with hernia—	
chest circumference of	364
robustness in	369
stature of	363
weight of	364
with hyperopia—	
chest circumference of	318
robustness in	319
stature of	315
weight of	318
with hypertrophic tonsillitis—	
chest circumference of	326
robustness in	327
stature of	323
weight of	323
with mitral insufficiency—	
chest circumference of	338
robustness in	338
stature of	335
weight of	335
with mitral stenosis—	
chest circumference of	339
robustness in	339
stature of	338
weight of	339
with myopia—	
chest circumference of	315
robustness in	315
stature of	314
weight of	314
with overweight and obesity—	
chest circumference of	379
robustness in	379
stature of	379
weight of	379
with pulmonary tuberculosis—	
chest circumference of	301
robustness in	301
stature of	299
weight of	300

Drafted men—Continued.

with simple goiter—	Page.
chest circumference of.....	305
robustness in.....	305
stature of.....	304
with simple tachycardia—	
chest circumference of.....	330
robustness in.....	331
stature of.....	327
weight in.....	330
with underweight—	
chest circumference of.....	376
robustness in.....	376
stature of.....	375
weight of.....	376
with valvular diseases of the heart (unclassified)—	
chest circumference of.....	342
robustness in.....	345
stature of.....	342
weight of.....	342
with varicocele—	
chest circumference of.....	347
robustness in.....	348
stature of.....	345
weight of.....	346
with varicose veins—	
chest circumference of.....	347
robustness in.....	348
stature of.....	345
weight of.....	346
Dravida, calf circumference of.....	230
Dutch:	
from Holland, approximate average stature of.....	47
in general, average stature of.....	68
of the Province of Zeeland (conscripts), average stature of.....	68
English:	
absolute and relative calf circumference in, at demobilization, 1919.....	231
absolute and relative chest circumference (rest) of, demobilization, 1919.....	154
absolute and relative height of pubic arch in, demobilization, 1919.....	200
absolute and relative height of sternal notch in, demobilization, 1919.....	197
absolute and relative knee height of, at demobilization, 1919.....	225
absolute and relative leg length of, at demobilization, 1919.....	222
absolute and relative shoulder breadth of, demobilization, 1919.....	206
absolute and relative sitting heights of, and standard deviations, demobilization, 1919..	19;
absolute and relative span, with standard deviation, demobilization, 1919.....	194
absolute and relative thigh circumference of, at demobilization, 1919.....	228
absolute and relative transverse chest diameter of, demobilization, 1919.....	210
absolute and relative transverse pelvic diameter in, demobilization, 1919.....	217
absolute and relative waist circumference of, demobilization, 1919.....	199
approximate number measured, demobilization, 1919.....	243
chest circumference of.....	139
comparative frequency distribution—	
of calf circumference, at demobilization, 1919.....	231
of antero-posterior diameter in, demobilization, 1919.....	212
of chest circumference (rest), demobilization, 1919.....	155
of span, demobilization, 1919.....	195
of eye color in, demobilization, 1919.....	284
of hair color in, demobilization, 1919.....	292
of height of pubic arch in, demobilization, 1919.....	201
of height of, demobilization, 1919.....	116
of height of sternal notch in, demobilization, 1919.....	198
of knee height in, at demobilization, 1919.....	226
of length of, at demobilization, 1919.....	223
of shoulder breadth of, demobilization, 1919.....	206
of sitting height of, demobilization, 1919.....	192
of thigh circumference of, at demobilization, 1919.....	229
of transverse diameter of chest in, demobilization, 1919.....	211
of transverse pelvic diameter, demobilization, 1919.....	218
of waist circumference in, demobilization, 1919.....	214
of weight of, demobilization, 1919.....	135

English—(Continued.)	Page.
index of build of.....	173
demobilization, 1919.....	244
mean stature and standard deviation of.....	113
mean stature of.....	243
mean weight and standard deviation of.....	135
middle class, approximate average stature of.....	47
ratio of pubic height to total stature of.....	199
shoulder breadth of.....	205
sitting height of.....	190
transverse chest diameter of.....	208
Egyptians, calf circumference of.....	230
Eskimos, shoulder breadth in relation to total stature of.....	204
Estonians, average stature of.....	68
European groups, calf circumference in.....	230
Eye color:	
and hair color, distribution of, demobilization, 1919.....	295
an index of race.....	42
comparative frequency distribution of—	
by Quartermaster Corps distribution zones, based on nativity of demobilized troops.....	287
in eight European races, demobilization.....	284
in various States of nativity of demobilized men.....	573
directions for recording.....	59
distribution of.....	42, 280
in eight European races.....	284
Eye, refractive errors of:	
and chest circumference.....	43
and stature.....	43
and weight.....	43
Eyes:	
blue with brown spots—	
absolute and relative numbers of veterans with, by States of nativity, demobilization, 1919.....	282
comparative frequency distribution of, by Quartermaster Corps distribution zones.....	287
proportions of, by States.....	282
brown, proportions of, by States.....	282, 283
clear blue—	
absolute and relative numbers of veterans with, by States of nativity, demobilization, 1919.....	281
comparative frequency distribution of, by Quartermaster Corps distribution zones.....	287
in eight European races at demobilization, 1919.....	243
proportions of, by States.....	280, 281
color of, comparison of Civil War and World War data concerning.....	285
colored, proportional numbers for different States, 1865.....	285
dark brown—	
absolute and relative numbers of veterans with, by States of nativity, demobilization, 1919.....	283
comparative frequency distribution of, by Quartermaster distribution zones.....	287
light brown—	
absolute and relative numbers of veterans with, by States of nativity, demobilization, 1919.....	283
comparative frequency distribution of, by Quartermaster Corps distribution zones.....	287
Features, facial, proportions of, essential to gas-mask manufacture.....	34
Feet, flat. <i>See</i> Flat-foot.	
Finger prints:	
identification by, belongs to army anthropology.....	34
importance of, in raising and maintaining an army.....	48
Finn groups:	
height distribution of.....	109, 110
index of build for draft recruits of.....	173
mean chest circumference of draft recruits of.....	150
mean height of.....	108
mean weight of draft recruits of.....	132
relative chest circumference of draft recruits of.....	150
weight distribution of draft recruits of.....	133, 134
Finns:	
approximate average stature of.....	47
average stature of.....	68
Fiot tribe (African), pelvic diameter of.....	215

	Page.
Flat-foot:	
and weight.....	43
chest circumference in drafted men with.....	370
correlation between height and weight in recruits with.....	371
robustness in drafted men with.....	370
stature of drafted men with.....	370
weight of drafted men with.....	370
Florida:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark-brown eyes, demobilization, 1919.....	283
with dark-brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light-brown eyes, demobilization, 1919.....	283
with light-brown hair, demobilization, 1919.....	290
with medium-brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919 ..	122
comparative view of mean height and mean weight of men from.....	124
increase in stature of soldiers at demobilization over stature of recruits, 1917-1919.....	76
index of build for, at demobilization, 1919.....	167
index of build for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144
Forearm:	
average, in color races, demobilization, 1919.....	235
comparative length of, of white and Negro troops.....	40
comparison of, in white and Negro troops.....	233
length of.....	220
relative, of Bavarians.....	220
relative, of Germans.....	220
relative, of Jews.....	220
relative, of Parisians.....	220
summary of, of approximately 6,000 colored troops, demobilization.....	234
summary of, of approximately 100,000 white troops, demobilization.....	234
Forearm and arm length:	
correlation between, colored troops.....	267
correlation between, colored troops, demobilization.....	531
correlation between, white troops.....	42, 262
correlation between, white troops, demobilization.....	516
Food ration, size in relation to.....	34
Foreign and native white groups, height distribution of.....	109, 110
Formula, Pignet's.....	186
French:	
absolute and relative calf circumference of, at demobilization, 1919.....	231
absolute and relative chest circumference of, demobilization, 1919.....	154
absolute and relative height of pubic arch of, demobilization, 1919.....	200
absolute and relative height of sternal notch of, demobilization, 1919.....	197
absolute and relative knee height of, at demobilization, 1919.....	225
absolute and relative length of, at demobilization, 1919.....	222
absolute and relative shoulder breadth of, demobilization, 1919.....	206
absolute and relative sitting height of, and standard deviations, demobilization, 1919.....	191
absolute and relative span, with standard deviation, demobilization, 1919.....	194
absolute and relative thigh circumference of, at demobilization, 1919.....	228
absolute and relative transverse chest diameter of, demobilization, 1919.....	210
absolute and relative transverse pelvic diameter of, demobilization, 1919.....	217
absolute and relative waist circumference of, demobilization, 1919.....	214
approximate average stature of.....	47
approximate number measured, demobilization, 1919.....	243
average weight of adult male.....	120
chest circumference of.....	139
comparative frequency distribution—	
of antero-posterior chest diameter in, demobilization, 1919.....	212
of calf circumference in, at demobilization, 1919.....	231
of chest circumference (rest), demobilization.....	155
of eye color in.....	284

French—Continued.

comparative frequency distribution—Continued.	Page.
of hair color in, demobilization, 1919	292
of height of	116
of height of pubic arch in, demobilization, 1919	201
of knee height in, at demobilization, 1919	226
of leg length of, at demobilization, 1919	223
of shoulder breadth of, demobilization, 1919	206
of sitting height of, demobilization, 1919	192
of sternal notch in, demobilization, 1919	198
of span in, demobilization, 1919	195
of thigh circumference of, at demobilization, 1919	229
of transverse diameter of chest in, demobilization, 1919	211
of transverse pelvic diameter of	218
of waist circumference in, demobilization, 1919	214
of weight of	135
index of build of	173, 244
in general, ratio of pelvic diameter to stature of	215
mean stature and standard deviation of	113
mean stature of	243
mean weight and standard deviation of	135
minimum shoulder breadth found in	205
ratio of pubic height to total stature of	199
relative arm length in	219
relative chest circumference of	139
relative shoulder breadth of	204
sitting height of	190
thoracic index of	209
transverse chest diameter of	207, 208
French army, minimum stature in	45
French conscripts, average stature of	68
French-Canadian groups:	
height distribution of	109, 110
relative chest circumference of draft recruits of	150
mean chest circumference of draft recruits of	150
mean height of	108
mean weight of draft recruits of	132
weight distribution of draft recruits of	133, 134
French-Canadians, index of build by groups of sections	173
Frequency curves, Laplace-Charlier, stature of army conscripts and recruits, in inches, as determined by	70
Frequency distribution	71
Friesians, East, average weight of adult male	120
Gas mask, proportions of facial features essential to manufacture of	34
Georgia:	
absolute and relative number of veterans from—	
with red hair, demobilization, 1919	289
with flaxen hair, demobilization, 1919	288
with clear blue eyes, demobilization, 1919	281
with dark brown hair, demobilization, 1919	291
with medium brown hair, demobilization, 1919	290
with light brown hair, demobilization, 1919	290
with dark brown eyes, demobilization, 1919	283
with light brown eyes, demobilization, 1919	283
with blue eyes with brown spots, demobilization, 1919	282
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919	122
comparative view of mean height and mean weight of men from	124
increase in stature of soldiers at demobilization over stature of recruits, 1917-1919	76
index of build for recruits from—	
at demobilization, 1919	167
at mobilization, 1917-18	166
mean chest circumference (expiration). of draft recruits from	142
mean stature of draft recruits from	75
mean stature of soldiers from, at demobilization, 1919	76
relative chest circumference of recruits from	144
German and Austrian groups:	
height distribution of	109, 110
index of build of recruits of	173
mean chest circumference of draft recruits of	150
mean height of	108
mean weight of draft recruits of	132

German and Austrian groups—(continued)	Page.
relative chest circumference of draft recruits of.....	150
weight distribution of draft recruits of.....	133, 134
German and Scandinavian groups:	
height distribution of.....	109, 110
index of build of recruits of.....	173
mean chest circumference of draft recruits of.....	150
mean height of.....	108
mean weight of draft recruits of.....	132
relative chest circumference of draft recruits of.....	150
weight distribution of draft recruits of.....	133, 134
Germans:	
absolute and relative calf circumference of, demobilization, 1919.....	231
absolute and relative chest circumference of, demobilization, 1919.....	154
absolute and relative height of pubic arch of, demobilization, 1919.....	200
absolute and relative height of sternal notch of, demobilization, 1919.....	197
absolute and relative knee height of, at demobilization, 1919.....	225
absolute and relative length of, at demobilization, 1919.....	222
absolute and relative shoulder breadth of, demobilization, 1919.....	206
absolute and relative sitting heights of, and standard deviations, demobilization, 1919.....	191
absolute and relative span of, with standard deviation, demobilization, 1919.....	194
absolute and relative thigh circumference of, at demobilization, 1919.....	228
absolute and relative transverse chest diameter of, demobilization, 1919.....	210
absolute and relative transverse pelvic diameter of, demobilization, 1919.....	217
absolute and relative waist circumference of, demobilization, 1919.....	199
approximate number measured, demobilization, 1919.....	243
comparative frequency distribution—	
at demobilization, 1919.....	231
of antero-posterior diameter of chest of, demobilization, 1919.....	212
of chest circumference (rest), demobilization, 1919.....	155
of eye color in, demobilization, 1919.....	284
of hair color in, demobilization, 1919.....	292
of height of pubic arch in, demobilization, 1919.....	201
of height of, demobilization, 1919.....	116
of height of sternal notch in, demobilization, 1919.....	198
of knee height in, at demobilization, 1919.....	226
of leg length of, at demobilization, 1919.....	223
of shoulder breadth of, demobilization, 1919.....	206
of sitting height of, demobilization, 1919.....	192
of span of, demobilization, 1919.....	195
of thigh circumference of, at demobilization, 1919.....	229
of transverse diameter of chest in, demobilization, 1919.....	211
of transverse pelvic diameter of, demobilization, 1919.....	218
of waist circumference of, demobilization, 1919.....	214
of weight of, demobilization, 1919.....	135
index of build of.....	173, 244
mean stature and standard deviation of.....	113
mean stature of.....	243
mean weight and standard deviation of.....	135
of Bavaria, shoulder breadth of.....	204
ratio of pelvic diameter to stature of.....	215
relative forearm length of.....	220
shoulder breadth of.....	205
transverse chest diameter of.....	208
Goiter:	
and stature.....	43
and weight.....	43
exophthalmic—	
chest circumference of drafted men, with.....	309
correlation between height and chest circumference in recruits, with.....	311
correlation between height and weight in recruits, with.....	310
in drafted men, with.....	308
Pignet's index of build of drafted men, with.....	309
robustness of drafted men, with.....	309
weight of drafted men, with.....	308
simple—	
chest circumference of drafted men, with.....	305
correlation between height and chest circumference in recruits, with.....	307
correlation between height and weight in recruits, with.....	306
in drafted men with.....	304

Goiter—Continued.	
simple—Continued.	Page
Pignet's index of build of men, with.....	305
robustness of drafted men, with.....	305
weight of drafted men, with.....	305
Gorilla, transverse diameter of pelvis, in.....	215
Great Britain, standards of stature in.....	45
Greeks, average stature of.....	68
Groups of sections:	
frequency distribution of statures in the.....	111, 114, 115
index of build by.....	173
mean chest circumference by.....	149
mean weight for the different.....	127, 130, 131, 132
Hair:	
clear red—	
comparative frequency distribution of, by Quartermaster Corps distribution zones.	294
in eight European races at demobilization, 1919.....	243
dark brown—	
absolute and relative number of veterans with, by States of nativity, demobiliza-	
tion, 1919.....	291
comparative frequency distribution of, by Quartermaster Corps distribution zones.	294
distribution of.....	289
flaxen—	
absolute and relative number of veterans with, by States of nativity, demobiliza-	
tion, 1919.....	288
comparative frequency of, by Quartermaster Corps distribution zones.....	294
distribution of, by States.....	288
in eight European races at demobilization, 1919.....	243
light brown—	
absolute and relative number of veterans with, by States of nativity, demobiliza-	
tion, 1919.....	290
comparative frequency distribution of, by Quartermaster Corps distribution zones..	294
in eight European races at demobilization, 1919.....	243
medium brown—	
absolute and relative number of veterans with, by States of nativity, demobiliza-	
tion, 1919.....	290
comparative frequency distribution of, by Quartermaster Corps distribution zones..	294
red—	
absolute and relative number of veterans with, by States of nativity, demobiliza-	
tion, 1919.....	289
distribution of.....	289
smallest ratio of, found in troops of Italian origin.....	293
red and black, comparative frequency distribution of, by Quartermaster Corps dis-	
tribution zones.....	294
Hair, color:	
and eye color, distribution of, demobilization, 1919.....	295
and locality.....	43
by States.....	43
comparative frequency distribution—	
in eight races, demobilization, 1919.....	292, 293
of hair color by Quartermaster Corps distribution zones, based on nativity of de-	
mobilized troops.....	294
in various States of nativity of demobilized men.....	574
comparison of, in World War and Civil War recruits.....	291
directions for recording.....	59
distribution of.....	43, 288
distribution of by Quartermaster Corps distribution zones.....	293
Heart.	
diseases of—	
and chest circumference.....	43
.....	43
val.....	342
.....	342
.....	344
.....	343
.....	345
.....	345
.....	342
.....	342

Hebrews (<i>see also</i> , Jews):	Page.
absolute and relative calf circumference of, at demobilization, 1919	231
absolute and relative chest circumference of, demobilization, 1919	154
absolute and relative height of pubic arch of, demobilization, 1919	200
absolute and relative height of sternal notch of, demobilization, 1919	197
absolute and relative knee height of, at demobilization, 1919	225
absolute and relative length of at demobilization, 1919	222
absolute and relative pelvic diameter of, demobilization, 1919	217
absolute and relative shoulder breadth of, demobilization, 1919	206
absolute and relative sitting heights of, and standard deviations, demobilization, 1919	191
absolute and relative span of, with standard deviation, demobilization, 1919	194
absolute and relative thigh circumference of, at demobilization, 1919	228
absolute and relative transverse chest diameter of, demobilization, 1919	210
absolute and relative waist circumference of, demobilization, 1919	199
approximate number measured, demobilization, 1919	243
comparative frequency distribution—	
of anteroposterior diameter in, demobilization, 1919	212
of calf circumference in, demobilization, 1919	231
of chest circumference (rest), demobilization, 1919	155
of eye color in, demobilization, 1919	284
of hair color in, demobilization, 1919	292
of height of, demobilization, 1919	116
of height of pubic arch of, demobilization, 1919	201
of height of sternal notch of, demobilization, 1919	198
of knee height of, at demobilization, 1919	226
of leg length of, at demobilization, 1919	223
of shoulder breadth of, demobilization, 1919	206
of sitting height of, demobilization, 1919	192
of span of, demobilization, 1919	195
of thigh circumference of, at demobilization, 1919	229
of transverse diameter of chest of, demobilization, 1919	211
of transverse pelvic diameter of, demobilization, 1919	218
of waist circumference, of demobilization, 1919	214
of weight of, demobilization, 1919	135
index of build of	173, 244
mean stature and standard deviation of	113
mean stature of	243
mean weight and standard deviation of	135
shoulder breadth of	205
transverse chest diameter of	208
Height (<i>see also</i> Stature):	
chest circumference (expiration) by, first million draft recruits	161
comparative frequency distribution of—	
by States, white and colored troops, demobilization	566, 567
in each of eight races, demobilization	116
distribution—	
by groups of sections	109, 110, 114, 115
by special diseases or defects	386, 387, 388, 398, 399
in eight European races, demobilization, 1919	243
mean—	
and weight, mean, comparative view of, of men from different States, World War and Civil War	124
by groups and component sections, first million draft recruits	424-429
by groups of sections, first million draft recruits	108
by sections, first million draft recruits	101
measurement, special diseases, first and second million draft recruits	394
of pubic arch	38
pubic, comparative, of white and Negro troops	40
sitting (<i>see also</i> Sitting height)	38
comparative, of white and Negro troops	40
comparative, of color races	193
directions for measuring	57
weight, and chest circumference (expiration)—	
measurements showing proportionate measurements of two of them to the total of the third, first million draft recruits	177
of recruits in relation to various diseases and defects	296
various, shown for United States, first million draft recruits	157
weight distribution by, first million draft recruits	159, 160, 161

Height and chest circumference (expiration):	Page.
classes of, first million draft recruits.....	158
correlation between—	
first million draft recruits.....	420, 421
group 1, agricultural, North, native white, first million draft recruits.....	443
group 2, agricultural, mixed foreign and native white, North, first million draft recruits.....	446
group 3, agricultural, native white, South, first million draft recruits.....	449
group 4, agricultural, Negro, 45 per cent plus, first million draft recruits.....	452
group 5, eastern manufacturing, first million draft recruits.....	455
group 6, commuter, first million draft recruits.....	458
group 7, mining, first million draft recruits.....	461
group 8, sparsely settled, first million draft recruits.....	464
group 9, desert, first million draft recruits.....	467
group 10, maritime, first million draft recruits.....	470
group 11, mountain, first million draft recruits.....	473
group 12, mountain whites, first million draft recruits.....	476
group 13, Indian, first million draft recruits.....	479
group 14, Mexican, first million draft recruits.....	482
group 15, native whites of Scotch origin, first million draft recruits.....	485
group 16, Russian, first million draft recruits.....	488
group 17, Scandinavian, first million draft recruits.....	491
group 18, Finn, first million draft recruits.....	494
group 19, French-Canadian, first million draft recruits.....	497
group 20, German and Scandinavian, first million draft recruits.....	500
group 21, German and Austrian, 20 per cent plus, first million draft recruits.....	503
group 22, German and Austrian, 15 per cent plus, first million draft recruits.....	506
Height and knee height, correlation between, colored troops, demobilization.....	525
Height and weight:	
classes, first million draft recruits.....	158
correlation between—	
for first million draft recruits.....	417
group 1, agricultural, North, native white, first million draft recruits.....	442
group 2, agricultural, mixed foreign and native white, North, first million draft recruits.....	445
group 3, agricultural, native white, South, first million draft recruits.....	448
group 4, agricultural, Negro, 45 per cent plus, first million draft recruits.....	451
group 5, eastern manufacturing, first million draft recruits.....	454
group 6, commuter, first million draft recruits.....	457
group 7, mining, first million draft recruits.....	460
group 8, sparsely settled, first million draft recruits.....	463
group 9, desert, first million draft recruits.....	466
group 10, maritime, first million draft recruits.....	469
group 11, mountain, first million draft recruits.....	472
group 12, mountain whites, first million draft recruits.....	475
group 13, Indian, first million draft recruits.....	478
group 14, Mexican, first million draft recruits.....	481
group 15, native whites of Scotch origin, first million draft recruits.....	484
group 16, Russian, first million draft recruits.....	487
group 17, Scandinavian, first million draft recruits.....	490
group 18, Finn, first million draft recruits.....	493
group 19, French-Canadian, first million draft recruits.....	496
group 20, German and Scandinavian, first million draft recruits.....	499
group 21, German and Austrian, 20 per cent plus, first million draft recruits.....	502
group 22, German and Austrian, 15 per cent plus, first million draft recruits.....	505
Hemorrhoids:	
and physical dimensions.....	43
correlation between—	
height and chest circumference in recruits with.....	355
height and weight in recruits with.....	354
Pignet's index of build for drafted men with.....	356
robustness in drafted men with.....	356
stature of drafted men with.....	353
weight of drafted men with.....	353
Hernia:	
chest circumference of drafted men with.....	364
correlation between—	
height and chest circumference (expiration) in recruits with.....	365
height and weight in recruits with.....	366
Pignet's index of build in drafted men with.....	369

Hernia—Continued.	Page.
robustness in drafted men with.....	369
stature of drafted men with.....	363
weight of drafted men with.....	364
Hova Indians, thoracic index for.....	207, 209
Hunan, relative arm length of Lolo in.....	219
Hungarians (conscripts) average stature of.....	68
Hylobates, shoulder breadth of.....	204
Hyperopia:	
chest circumference of drafted men with.....	318
correlation between—	
height and chest circumference in recruits with.....	317
height and weight in recruits with.....	316
in drafted men.....	315
Pignet's index of build of drafted men with.....	319
robustness of drafted men with.....	319
stature of drafted men with.....	315
weight of drafted men with.....	318
Hypospadia, cryptorchidism, anorchism, and monorchism:	
chest circumference of drafted men with.....	382
correlation between—	
height and chest circumference (expiration) of drafted men with.....	381
height and weight of drafted men with.....	380
robustness of drafted men with.....	382
stature of drafted men with.....	379
weight of drafted men with.....	382
Idaho:	
absolute and relative numbers of veterans from—	
with blue eyes, with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits from, at mobilization, 1917–1918, and demobilization, 1919.....	122
comparative view of mean height and mean weight of men from.....	124
difference of weight of draft recruits from, at mobilization, 1917–1918, and demobilization, 1919.....	123
increase in stature of soldiers at demobilization over recruits, 1917–1919.....	76
index of build for—	
at demobilization, 1919.....	167
at mobilization, 1917–1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75, 76
relative chest circumference of recruits from.....	144
Illinois:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits at mobilization, 1917–1918, and demobilization, 1919.....	122
colored eyes, proportional numbers for, 1865.....	285
comparative view of mean height and mean weight of men from.....	124
comparison of index of build of men from, at demobilization in 1865 and 1919.....	170
increase in stature of soldiers at demobilization over recruits, 1917–1919.....	76
index of build for recruits from, at mobilization, 1917–1918.....	166, 167
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144

	Page.
Immigration, influence of, on stature	35, 79
Inches, table for converting into centimeters	411
Index of build (<i>see</i> Build, index of).	
Indian groups of sections:	
relative chest circumference of draft recruits of	150
sparsely settled, index of build for draft recruits of	173
sparsely settled—	
mean chest circumference of draft recruits of	150
mean height of draft recruits of	108
mean weight of draft recruits of	132
weight distribution of draft recruits of	133, 134
Indian race:	
mean stature of, at demobilization, 1919	117
mean weight of, with standard deviation for white and Negro troops, demobilization, 1919	136
Indiana:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919	282
with clear blue eyes, demobilization, 1919	281
with dark brown eyes, demobilization, 1919	283
with dark brown hair, demobilization, 1919	291
with flaxen hair, demobilization, 1919	288
with light brown eyes, demobilization, 1919	283
with light brown hair, demobilization, 1919	290
with medium brown hair, demobilization, 1919	290
with red hair, demobilization, 1919	289
average weight of draft recruits, at mobilization, 1917–1918, and demobilization, 1919	122
colored eyes, proportional numbers for, 1865	285
comparative view of mean height and mean weight of men from	124
comparison of index of build of men from, at demobilization in 1865 and 1919	170
increase in stature of soldiers at demobilization over recruits, 1917–1919	76
index of build—	
at demobilization, 1919	167
for recruits from, at mobilization, 1917–1918	166
mean chest circumference (expiration), of draft recruits from	142
mean stature—	
of draft recruits from	75
of soldiers from, at demobilization, 1919	76
relative chest circumference of recruits from	144
Indians:	
absolute and relative height of pubic arch in, demobilization, 1919	202
absolute and relative height of sternal notch in, demobilization, 1919	199
absolute and relative leg length in, demobilization, 1919	224
absolute and relative sitting height in, demobilization, 1919	198
absolute and relative shoulder breadth in, demobilization, 1919	207
absolute and relative transverse diameter of pelvis in, demobilization, 1919	219
American—	
approximate average stature of	47
relative leg length of	221
Colorado, shoulder breadth in relation to total stature of	204
comparative frequency distribution of measurements of, at demobilization	236–240
general comparison of, with other color races	241
Hova, thoracic index for	207, 209
index of build for	174
Iroquois, maximum pelvic diameter found in	215
mean absolute and relative spans, demobilization, 1919	196
Navajo—	
thoracic index of	209
transverse diameter of chest in	207
North American, sitting height of	190
sparsely settled, height distribution of	109, 110
Infantry, stature of, in Great Britain	45
Inguinal rings, enlarged:	
chest circumference of drafted men with	369
correlation between height and chest circumference (expiration), in recruits with	368
correlation between height and weight in recruits with	367
Pignet's index of build in drafted men with	369
robustness in drafted men with	369
stature of drafted men with	369
weight of drafted men with	369

Iowa:

absolute and relative numbers of veterans from—	Page.
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919..	122
colored eyes, proportional numbers for, 1865.....	285
comparative view of mean height and mean weight of men from.....	124
difference of weight of draft recruits from, at mobilization, 1917-1918, and demobiliza- tion, 1919.....	123
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144

Irish:

absolute and relative calf circumference in, demobilization, 1919.....	231
absolute and relative chest circumference of, demobilization, 1919.....	154
absolute and relative height of pubic arch in, demobilization, 1919.....	200
absolute and relative height of sternal notch in, demobilization, 1919.....	197
absolute and relative knee height of, demobilization, 1919.....	225
absolute and relative leg length of, demobilization, 1919.....	222
absolute and relative shoulder breadth of, demobilization, 1919.....	206
absolute and relative sitting heights of and standard deviations, demobilization, 1919..	191
absolute and relative span in, with standard deviation, demobilization, 1919.....	194
absolute and relative thigh circumference of, demobilization, 1919.....	228
absolute and relative transverse chest diameter of, demobilization, 1919.....	210
absolute and relative transverse pelvic diameter in, demobilization, 1919.....	217
absolute and relative waist circumference of, demobilization, 1919.....	199
approximate number measured, demobilization, 1919.....	243
comparative frequency distribution in, demobilization, 1919.....	231
comparative frequency distribution—	
of antero-posterior diameter in, demobilization, 1919.....	212
of chest circumference (rest) in, demobilization, 1919.....	155
of span in, demobilization, 1919.....	195
of eye color in, demobilization, 1919.....	284
of hair color in, demobilization, 1919.....	292
of height of pubic arch in, demobilization, 1919.....	201
of height of, demobilization, 1919.....	116
of height of sternal notch in, demobilization, 1919.....	198
of knee height in, demobilization, 1919.....	226
of shoulder breadth of, demobilization, 1919.....	206
of sitting height of, demobilization, 1919.....	192
of thigh circumference of, demobilization, 1919.....	229
of transverse diameter of chest in, demobilization, 1919.....	211
of transverse pelvic diameter in, demobilization, 1919.....	218
of waist circumference in, demobilization, 1919.....	214
of weight of, demobilization, 1919.....	135
of leg length of, demobilization, 1919.....	223
index of build of.....	173, 244
mean stature and standard deviation of.....	113
mean stature of.....	243
mean weight and standard deviation of.....	135
shoulder breadth of.....	205
transverse chest diameter of.....	208

Iroquois Indians, maximum pelvic diameter found in.....

215

Italian army, minimum stature of.....

45

Italians:

absolute and relative calf circumference in, demobilization, 1919.....	231
absolute and relative chest circumference of, demobilization, 1919.....	154
absolute and relative height of pubic arch in, demobilization, 1919.....	200
absolute and relative height of sternal notch in, demobilization, 1919.....	197

Italians—Continued.

	Page.
absolute and relative knee height of, demobilization, 1919.....	225
absolute and relative leg length of, at demobilization, 1919.....	222
absolute and relative shoulder breadth of, demobilization, 1919.....	206
absolute and relative sitting heights of, and standard deviations, demobilization, 1919.....	191
absolute and relative span of, with standard deviation, demobilization, 1919.....	194
absolute and relative thigh circumference of, demobilization, 1919.....	228
absolute and relative transverse chest diameter of, demobilization, 1919.....	210
absolute and relative transverse pelvic diameter in, demobilization, 1919.....	217
absolute and relative waist circumference of, demobilization, 1919.....	199
approximate number measured, demobilization, 1919.....	243
comparison of frequency distribution—	
at demobilization, 1919.....	195, 231
of antero-posterior diameter in, demobilization, 1919.....	212
of chest circumference (rest) in, demobilization.....	155
of eye color in, demobilization, 1919.....	284
of hair color in, demobilization, 1919.....	292
of height of pubic arch in, demobilization, 1919.....	201
of height of, demobilization, 1919.....	116
of height of sternal notch in, demobilization, 1919.....	198
of knee height, demobilization, 1919.....	226
of leg length of, demobilization, 1919.....	223
of shoulder breadth of, demobilization, 1919.....	206
of sitting height of, demobilization, 1919.....	192
of thigh circumference of, demobilization, 1919.....	229
of transverse diameter of chest in, demobilization, 1919.....	211
of transverse pelvic diameter in, demobilization, 1919.....	218
of waist circumference in, demobilization, 1919.....	214
of weight of, demobilization, 1919.....	135
index of build of.....	173, 244
in general, average stature of.....	68
mean stature and standard deviation of.....	113
mean stature of.....	243
mean weight and standard deviation of.....	135
shoulder breadth of.....	205
South, approximate average stature of.....	47
transverse chest diameter of.....	208

Japanese:

absolute and relative height of pubic arch in, demobilization, 1919.....	202
absolute and relative height of sternal notch in, demobilization, 1919.....	199
absolute and relative leg length in, demobilization, 1919.....	224
absolute and relative shoulder breadth in, demobilization, 1919.....	207
absolute and relative sitting height in, demobilization, 1919.....	193
absolute and relative transverse diameter of pelvis in, demobilization, 1919.....	219
average weight of adult male.....	120
approximate average stature of.....	47
comparative frequency distribution of measurements in, at demobilization.....	236-240
general comparison of, with other color races.....	241
index of build for.....	174
mean absolute and relative spans of, demobilization, 1919.....	196
mean stature of, at demobilization, 1919.....	117
mean weight of, with standard deviation for white and Negro troops, demobilization, 1919.....	136
ratio of pelvic diameter to stature of.....	215
relative shoulder breadth of.....	204

Jews (*see also* Hebrews):

Austrian, of Hungary, average stature of.....	68
from Russian Poland, average stature of.....	68
Polish—	
approximate average stature of.....	47
average weight of adult male.....	120
relative shoulder breadth of.....	204
shoulder breadth of.....	204
ratio of pelvic diameter to stature of.....	215
relative forearm length of.....	220
Russian, relative chest circumference of.....	139
South Russian—	
approximate average stature of.....	47, 68
average weight of adult male.....	120
Ukrainian, sitting height of.....	190

Kansas:

absolute and relative numbers of veterans from—	Page.
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits, at mobilization, 1917-1918, and demobilization, 1919..	122
comparative view of mean height and mean weight of men from.....	124
difference of weight of draft recruits from, at mobilization, 1917-1918, and demobiliza- tion, 1919.....	123
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144

Kentucky:

absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919..	123
colored eyes, proportional numbers for, 1865.....	285
comparative view of mean height and mean weight of men from.....	124
comparison of index of build of men from, at demobilization in 1865 and 1919.....	170
difference of weight of draft recruits from, at mobilization, 1917-1918, and demobiliza- tion, 1919.....	123
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144

Knee, circumference of, level of patella, directions for measuring.....

59

Knee height:

absolute and relative, in eight European races, demobilization, 1919.....	225
and breeches groups, association between, colored troops, demobilization.....	557
and breeches groups, correlation between, white troops, demobilization.....	550
and height, correlation between, colored troops, demobilization.....	525
and leg length, correlation between, colored troops, demobilization.....	526
and leg length, correlation between, white troops, demobilization.....	510
and leg length, correlation between, in white troops.....	42
and leg length, correlation between, white troops.....	263
and leg length, Negro troops.....	268
and stature, correlation between, in white troops.....	42
and stature, Negro troops, correlation between.....	265
average, in color races, demobilization, 1919.....	235
comparative frequency distribution of, in eight European races, demobilization, 1919..	226
comparison of, in white and Negro troops.....	40, 233
general discussion of.....	224
in eight European races at demobilization, 1919.....	243
mean, for Negro troops.....	224
mean, for white troops.....	224
standard deviation of.....	225
summary of, of approximately 6,000 colored troops, demobilization.....	234
summary of, of approximately 100,000 white troops, demobilization.....	234
Knee joint, height of, directions for measuring.....	58

Knee patella:	Page.
summary of, of approximately 6,000 colored troops, demobilization.....	234
summary of, of approximately 100,000 white troops, demobilization.....	234
circumference.....	232
standard deviation of.....	232
Laplace-Charlier frequency curves, stature of Army conscripts and recruits in inches, as determined by.....	70
Laplanders:	
average stature of.....	68
from Scandinavia, average stature of.....	68
ratio of pubic height to total stature of.....	199
Leg:	
circumference of—	
just above patella, directions for measuring.....	59
just below level of tuberosity of tibia, directions for measuring.....	59
index of relation of upper to lower.....	224
inside length of, from gluteal fold to tip of internal malleolus of tibia, directions for measuring.....	59
Leg length.....	39
absolute and relative, in eight European races, demobilization, 1919.....	222
absolute and relative, in five color races, demobilization, 1919.....	224
average, in color races, demobilization, 1919.....	235
comparative frequency distribution of, in each of eight European races, demobilization, 1919.....	223
comparative, of white and negro troops.....	40
comparison of, in color races.....	224
comparison of, in white and negro troops.....	233
comparison of, in eight European races.....	221, 222, 223
general discussion.....	220
in eight European races at demobilization, 1919.....	243
in relation to marching capacity.....	46
mean, of negro troops.....	221
mean, of white troops.....	221
correlated with stature.....	46
required to make long marches.....	34
relative.....	221
of African tribes.....	221
of Armenians.....	221
of Tartars.....	221
standard deviation of, for white troops.....	221
summary of, of approximately 6,000 colored troops, demobilization.....	234
summary of, of approximately 100,000 white troops, demobilization.....	234
Leg length and knee height:	
correlation between—	
in colored troops, demobilization.....	526
in white troops.....	42, 263
in negro troops.....	268
in white troops, demobilization.....	510
Leg length and waist circumference:	
correlation between—	
in colored troops, demobilization.....	555
white troops.....	42, 264
white troops, demobilization.....	548
Letts:	
approximate average stature of.....	47
relative chest circumference of.....	139
Lithuanians:	
of Lithuania (conscripts), average stature of.....	68
of Russian Poland.....	68
Lobi (African race), relative arm length of.....	220
Lolo in Hunan, relative arm length of.....	219
Locality:	
and distribution of eye color.....	42
and hair color.....	43
variability—	
in build by.....	37
in stature by.....	35, 36
in weight by.....	36

Louisiana:

absolute and relative numbers of veterans from—	Page.
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	283
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919..	122
comparative view of mean height and mean weight of men from.....	124
difference of weight of draft recruits from, at mobilization, 1917-18, and demobiliza- tion, 1919.....	123
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144

Magyars from West Hungary (conscripts), average stature of..... 68

Maine:

absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits from, at mobilization, 1917-1918, and demobilization, 1919.	122
colored eyes, proportional numbers for, 1865.....	285
comparative view of mean height and mean weight of men, from.....	124
comparison of index of build of men from, at demobilization in 1865 and 1919.....	170
increase in stature of soldiers from, at demobilization over recruits, 1917-1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144

Malay groups, calf circumference in..... 230

Manikins, dimensions of..... 276

Manufacturing area, eastern, index of build for draft recruits from..... 173

Manufacturing groups, eastern:

height distribution of.....	109, 110
mean chest circumference of draft recruits of.....	150
mean height of draft recruits of.....	108
mean weight of draft recruits of.....	132
relative chest circumference of draft recruits of.....	150
weight distribution of draft recruits of.....	133, 134

Marches, long, length of leg required to make..... 34

Maritime groups:

height distribution of.....	109, 110
mean chest circumference of draft recruits of.....	150
mean height of draft recruits of.....	108
mean weight of draft recruits of.....	132
weight distribution of draft recruits of.....	133, 134
index of build for draft recruits of.....	173
relative chest circumference of draft recruits of.....	150

Maryland:

absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283

Maryland—Continued.

absolute and relative numbers of veterans from—Continued.	Page.
with dark brown hair, demobilization, 1919.	291
with flaxen hair, demobilization, 1919.	288
with light brown eyes, demobilization, 1919.	283
with light brown hair, demobilization, 1919.	290
with medium brown hair, demobilization, 1919.	290
with red hair, demobilization, 1919.	289
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919.	122
comparative view of mean height and mean weight of men from.	124
increase in stature of soldiers at demobilization over recruits, 1917-1919.	76
index of build—	
at demobilization, 1919.	167
recruits from, at mobilization, 1917-1918.	166
mean chest circumference (expiration), of draft recruits from.	142
mean stature of draft recruits from.	75
mean stature of soldiers from, at demobilization (1919).	76
relative chest circumference of recruits from.	144
Massai of South Africa, sitting height of.	190
Massachusetts:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.	282
with clear blue eyes demobilization, 1919.	281
with dark brown eyes, demobilization, 1919.	283
with dark brown hair, demobilization, 1919.	291
with flaxen hair, demobilization, 1919.	288
with light brown eyes, demobilization, 1919.	283
with light brown hair, demobilization, 1919.	290
with medium brown hair, demobilization, 1919.	290
with red hair, demobilization, 1919.	289
average weight of draft recruits, at mobilization, 1917-1918, and demobilization, 1919.	122
colored eyes, proportional numbers for, 1865.	285
comparative view of mean height and mean weight of men from.	124
comparison of index of build of men from, at demobilization, in 1865 and 1919.	170
increase in stature of soldiers at demobilization over recruits, 1917-1919.	76
index of build—	
at demobilization, 1919.	167
for recruits from, at mobilization, 1917-1918.	166
mean chest circumference (expiration), of draft recruits from.	142
mean stature of draft recruits from.	75
mean stature of soldiers from, at demobilization (1919).	76
relative chest circumference of recruits from.	144
Mawambu pygmies, relative arm length in.	219
Means:	
of white and Negro troops, comparison of.	232
summary of.	404, 405, 406, 407
Measurement card for clothing patterns, demobilization, 1919.	61
Measurements:	
apparatus used in taking.	57
comparative, at demobilization, Civil and World wars.	242
comparative frequency distribution of, in color races at demobilization.	236
correlations between, for white and Negro troops.	253
correlation between, for white troops, demobilization.	258-264
detailed directions for.	54
directions for taking and recording.	57
for blouses.	271
for breeches.	274
general comparison of in color races.	241
height, weight, and chest circumference (expiration), distribution by States of nativity.	184
height, weight, and chest circumference (expiration), showing proportionate measurements of two of them to the total of the third, first million draft recruits.	177
instructions issued by Surgeon General relative to.	55
list of anthropologists who supervised taking, in camps.	56
Negro troops, correlation between.	264
of drafted men, standards of.	296
of draft recruits.	51
omission of.	60
physical.	45
results of standard Army physical.	64-189
special anthropological.	190
special, in Army, efforts of National Academy of Science to secure.	53

Measurements—Continued.	Page.
special uniform, orders issued relative to.....	53
specifications for arrangements required at camp for taking.....	59
statistical treatment of data obtained from.....	60
supervising personnel and camps where taken.....	56
systems used in statistical treatment of data obtained from.....	60
total and proportionate—	
by each section, first million draft recruits.....	182, 183
by groups of sections, first million draft recruits.....	181
by States, first million draft recruits.....	178, 179, 180
special diseases.....	397
Measuring rod, Seaver, used in taking measurements.....	57
Mediterranean races:	
relative transverse chest diameter of.....	208
waist circumference of.....	213
Men, demobilized:	
comparative frequency—	
of eye color in, in various States of nativity.....	573
of hair color in, in various States of nativity.....	574
Men, drafted (see Drafted men; Recruits).	
Men measured, approximate number of, in eight European races, demobilization, 1919.....	243
Mexican groups, sparsely settled:	
height, distribution of.....	109, 110
index of build for draft recruits of.....	173
mean chest circumference of draft recruits of.....	150
mean height of.....	108
mean weight of draft recruits of.....	132
relative chest circumference of draft recruits of.....	150
weight distribution of draft recruits of.....	133, 134
Michigan:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919..	122
colored eyes, proportional numbers for, 1865.....	285
comparative view of mean height and mean weight of men from.....	124
comparison of index of build of men from, at demobilization in 1865 and 1919.....	170
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration) of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	73
relative chest circumference of recruits from.....	144
Mining groups:	
height distribution of.....	109, 110
index of build for draft recruits of.....	173
mean chest circumference of draft recruits of.....	150
mean height of.....	108
mean weight of draft recruits of.....	132
relative chest circumference of draft recruits of.....	150
weight distribution of draft recruits of.....	133, 134
Minnesota:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289

Minnesota—Continued.	Page.
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919..	122
comparative view of mean height and mean weight of men from	124
difference of weight of draft recruits from, at mobilization, 1917-1918, and demobiliza- tion, 1919.....	123
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144
Mississippi:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919..	122
comparative view of mean height and mean weight of men from	124
difference of weight of draft recruits from, at mobilization, 1917-1918, and demobiliza- tion, 1919.....	123
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144
Missouri:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919..	122
colored eyes, proportional numbers for, 1865.....	285
comparative view of mean height and mean weight of men from.....	124
difference of weight of draft recruits from, at mobilization, 1917-1918, and demobiliza- tion, 1919.....	123
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144
Mitral insufficiency:	
chest circumference of drafted men with.....	338
correlation between—	
height and chest circumference in recruits with.....	337
height and weight in recruits with	336
in drafted men	335
Pignet's index of build in drafted men with.....	338
robustness in drafted men with.....	338
stature of drafted men with.....	335
weight of drafted men with.....	335

	Page.
Mitral stenosis:	
chest circumference in drafted men with.....	339
correlation between—	
height and chest circumference (expiration) in recruits with.....	341
height and weight in recruits with.....	340
in drafted men.....	338
Pignet's index of build for recruits with.....	339
robustness of drafted men with.....	339
stature of drafted men with.....	338
weight of drafted men with.....	339
Mobility, chest.....	297, 298
Mobilization, increase in weight at demobilization over.....	122, 123
Mongolians, relative pubic height of.....	200
Mongoloid groups, calf circumference in.....	230
Monkeys, lower, transverse diameter of pelvis in.....	215
Monorchism, cryptorchidism, anorchism, and hypospadia:	
chest circumference in drafted men with.....	382
weight of drafted men with.....	382
correlation between—	
height and chest circumference (expiration), in recruits with.....	381
height and weight in drafted men with.....	380
robustness in drafted men with.....	382
stature of drafted men with.....	379
Montana:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits, at mobilization, 1917-1918, and demobilization, 1919..	122
comparative view of mean height and mean weight of men from.....	124
difference in weight of draft recruits from, at mobilization 1917-1918, and demobiliza- tion, 1919.....	123
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build—	
demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of soldiers from, at demobilization, 1919.....	76
mean stature of draft recruits from.....	75
relative chest circumference of recruits from.....	144
Mountain groups:	
height distribution of.....	109, 110
index of build for draft recruits of.....	173
mean chest circumference of draft recruits of.....	150
mean height of.....	108
mean weight of draft recruits of.....	132
weight distribution of draft recruits of.....	133, 134
relative chest circumference of draft recruits of.....	150
white, height distribution of.....	109, 110
mean chest circumference of draft recruits of.....	150
mean height of.....	108
mean weight of draft recruits of.....	132
relative chest circumference of draft recruits of.....	150
weight distribution of draft recruits of.....	133, 134
Mulattoes:	
absolute and relative height of pubic arch in Negroes and, demobilization, 1919.....	202
absolute and relative leg length in Negroes and, demobilization, 1919.....	224
absolute and relative neck circumference of Negroes and, demobilization, 1919.....	203
absolute and relative shoulder breadth in Negroes and, demobilization, 1919.....	207
absolute and relative transverse diameter of pelvis in Negroes and, demobilization, 1919.....	219
American, average weight of adult male.....	120

Myopia:	Page.
chest circumference of drafted men with.....	315
correlation between—	
height and weight in recruits with.....	312
height and chest circumference in recruits with.....	313
in drafted men.....	314
Pignet's index of build of drafted men with.....	315
robustness of drafted men with.....	315
stature of drafted men with.....	314
weight of drafted men with.....	314
National Academy of Science, efforts of, to secure authorization for special measurements in Army.....	53
Nationality, rules for determining.....	60
Native white groups:	
of Scotch origin—	
mean height of.....	108
mean weight of draft recruits of.....	132
Navajo Indians:	
thoracic index of.....	209
transverse diameter of chest in.....	207
Nebraska:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919..	122
comparative view of mean height and mean weight of men from.....	124
difference of weight of draft recruits from, at mobilization, 1917-1918, and demobilization, 1919.....	123
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build—	
at demobilization.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144
Neck circumference.....	38
absolute and relative, of white and Negro troops, demobilization, 1919.....	203
and blouse groups, association between, colored troops, demobilization.....	546
and blouse groups, association between, white troops, demobilization.....	538
and chest circumference, correlation between, white troops.....	260
and chest circumference, Negro troops, correlation between.....	266
and chest circumference (rest), correlation between—	
colored troops, demobilization.....	527
white troops, demobilization.....	512
average, in color races, demobilization, 1919.....	235
comparison of in color races.....	203
comparison of, in eight European races.....	203
general discussion of.....	202
mean.....	202
level of laryngeal prominence, directions for measuring.....	58
standard deviation of.....	203
summary of, of approximately 6,000 colored troops, demobilization.....	234
summary of, of approximately 100,000 white troops, demobilization.....	234
Neck girth and chest girth, correlation of, in white troops.....	42
Negro and mulatto troops, absolute and relative neck circumference of, demobilization, 1919.....	203
Negro and white troops:	
comparison of dimensions of.....	232, 233
correlations between measurements for.....	253
mean and relative chest circumference (rest) of, demobilization, 1919.....	156

	Page.
Negro race:	
mean stature of, at demobilization, 1919.....	117
mean weight in, with standard deviation for, demobilization, 1919.....	136
Negro troops:	
calf circumference in.....	230
dimensions associated with blouse groups of, demobilization.....	273
dimensions associated with breeches groups of, demobilization.....	275
mean knee height of.....	224
mean leg length of.....	221
measurements of, correlation between.....	264
mean transverse diameter of pelvis of.....	216
relative arm length of.....	219
standard deviation—	
of arm length of.....	220
of leg length of.....	221
of pubic height of.....	202
of transverse pelvic diameter of.....	216
suprapatella circumference of.....	232
thigh circumference of.....	227
Negro-mulattoes, index of build for.....	174
Negroes:	
absolute and relative height of sternal notch in, demobilization, 1919.....	199
absolute and relative sitting heights in, demobilization, 1919.....	193
African—	
relative arm length in.....	219
thoracic index for.....	207, 209
Bugu, thoracic index for.....	207, 209
general comparison of, with other color races.....	241
mean absolute and relative spans, demobilization, 1919.....	196
pelvic diameter of.....	215
of various origins, approximate average stature of.....	47
Negroes and mulattoes:	
absolute and relative height of pubic arch in, demobilization, 1919.....	202
absolute and relative leg length of, at demobilization, 1919.....	224
absolute and relative shoulder breadth in, demobilization, 1919.....	207
absolute and relative transverse diameter of pelvis in, at demobilization, 1919.....	219
Negroes and whites, comparison of correlation between.....	268
Nevada:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
average weight of draft recruits, at mobilization, 1917-1918, and demobilization, 1919.....	122
comparative view of mean height and mean weight of men from.....	124
difference of weight of draft recruits from, at mobilization, 1917-1918, and demobilization, 1919.....	123
increase in stature of soldiers at demobilization, over recruits, 1917-1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144
New Hampshire:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289

New Hampshire—Continued.

	Page.
average weight of draft recruits, at mobilization, 1917-1918, and demobilization, 1919...	122
colored eyes, proportional numbers for, 1865.....	285
comparative view of mean height and mean weight of men from	124
comparison of index of build of men from, at demobilization in 1865 and 1919.....	170
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration) of draft recruits from	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, demobilization, 1919.....	76
relative chest circumference of recruits from.....	144

New Jersey:

absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919	291
with flaxen hair, demobilization, 1919	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919...	122
comparative view of mean height and mean weight of men from	124
comparison of index of build of men from, at demobilization in 1865 and 1919.....	170
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration) of draft recruits from	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144

New Mexico:

absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
average weight of draft recruits, at mobilization, 1917-1918, and demobilization, 1919	122
comparative view of mean height and mean weight of men from	124
difference of weight of draft recruits from, at mobilization, 1917-1918, and demobiliza- tion, 1919.....	123
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144

New York:

absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919	291
with flaxen hair, demobilization, 1919	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919.....	122

New York—Continued.	Page.
colored eyes, proportional numbers for, 1865.....	285
comparative view of mean height and mean weight of men from.....	124
comparison of index of build of men from, at demobilization in 1865 and 1919.....	170
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, demobilization, 1919.....	76
relative chest circumference of recruits from.....	144
Nordic races:	
chest circumference of.....	154
chest index of.....	210
relative chest girth of.....	154
relative sitting height of.....	191
relative transverse chest diameter of.....	208
relatively long legs of.....	200
shoulder breadth of.....	205
thigh circumference of.....	228
waist circumference of.....	213
North Carolina:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919.....	122
comparative view of mean height and mean weight of men, from.....	124
difference of weight of draft recruits from, at mobilization, 1917-1918, and demobiliza- tion, 1919.....	123
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, demobilization, 1919.....	76
relative chest circumference of recruits from.....	144
North Dakota:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits, at mobilization, 1917-1918, and demobilization, 1919.....	122
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build—	
demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
comparative view of mean height and mean weight of men from.....	124
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, demobilization, 1919.....	76
relative chest circumference of recruits from.....	144
Norwegians:	
approximate average stature of.....	47, 68
average weight of adult male.....	120

Obesity and overweight:	Page.
chest circumference in drafted men with.....	379
robustness in drafted men with.....	379
stature of drafted men with.....	379
weight of drafted men with.....	379
Ohio:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits at mobilization, 1917–1918, and demobilization, 1919..	122
colored eyes, proportional numbers for, 1865.....	285
comparative view of mean height and mean weight of men from.....	124
comparison of index of build of men from, at demobilization in 1865 and 1919.....	170
increase in stature of soldiers at demobilization over recruits, 1917–1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917–1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144
Oklahoma:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits at mobilization, 1917–1918, and demobilization, 1919...	122
comparative view of mean height and mean weight of men from.....	124
difference of weight of draft recruits from, at mobilization, 1917–1918, and demobiliza- tion, 1919.....	123
increase in stature of soldiers at demobilization over recruits, 1917–1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917–1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization (1919).....	76
relative chest circumference of recruits from.....	144
Orang outang, shoulder breadth of.....	204
Oregon:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits, at mobilization, 1917–1918 and demobilization, 1919...	122
increase in stature of soldiers at demobilization over recruits, 1917–1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917–1918.....	166
mean stature of soldiers from, at demobilization.....	76
mean stature of draft recruits from.....	75

Oregon—Continued.	Page.
comparative view of mean height and mean weight of men from.....	124
mean chest circumference (expiration), of draft recruits from.....	142
relative chest circumference of recruits from.....	144
Overweight and obesity:	
chest circumference in drafted men with.....	379
robustness of drafted men with.....	379
stature of drafted men with.....	379
weight of drafted men with.....	379
Parisians:	
ratio of pelvic diameter to stature of.....	215
relative forearm length of.....	220
shoulder breadth of.....	204
Patella, average dimensions of in color races, demobilization, 1919.....	235
Patella, circumference, and breeches groups:	
association between—	
colored troops, demobilization.....	560
white troops, demobilization.....	553
Patterns for uniforms.....	271
Pelvis, diameter of:	
absolute and relative transverse, in eight European races, demobilization, 1919.....	217
in five color races, demobilization, 1919.....	219
absolute transverse, by sections, demobilization, 1919.....	216
and chest girth, correlation of, in white troops.....	42
and waist girth, correlation of, in white troops.....	42
average, in color races, demobilization, 1919.....	235
comparative frequency distribution of transverse, in each of eight European races, demobilization, 1919.....	218
in eight European races at demobilization, 1919.....	243
mean.....	216
maximum of, found in Iroquois Indians.....	215
of men from different parts of the country, at demobilization, 1919.....	216
ratio of, to stature.....	215
transverse.....	39, 215
and blouse groups, association between, colored troops, demobilization.....	545
and blouse groups, association between, white troops, demobilization.....	537
and breeches groups, association between, colored troops, demobilization.....	556
and breeches groups, association between, white troops, demobilization.....	549
and chest circumference (rest), correlation between.....	261, 513
and chest circumference, Negro troops, correlation between.....	267
and chest circumference (rest), correlation between, colored troops, demobilization.....	528
and waist circumference, correlation between, colored troops, demobilization.....	530
and waist circumference, Negro troops, correlation between.....	267
and waist circumference, correlation between, white troops, demobilization.....	262, 515
at level of crests of ilium, directions for measuring.....	58
between crista, and chest circumference, correlation between, white troops.....	258
comparison of, in eight European races.....	216, 217
comparison of, in five color races.....	219
general discussion of.....	215
in anthropoids.....	215
in Bavarians.....	215
in chimpanzee.....	215
in gorilla.....	215
standard deviation of.....	216
summary of, of approximately 6,000 colored troops, demobilization.....	234
summary of, of approximately 100,000 white troops, demobilization.....	234
Pennsylvania:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits, at mobilization, 1917-1918, and demobilization, 1919..	122
colored eyes, proportional numbers for, 1865.....	285

Pennsylvania—Continued.

	Page.
comparative view of mean height and mean weight of men from.....	124
comparison of index of build of men from, at demobilization in 1865 and 1919.....	170
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, demobilization, 1919.....	76
relative chest circumference of recruits from.....	144
Perimeter, chest.....	186
Personnel for anthropological work in Army.....	56
Physical-examination standards.....	297
Physique and defects, relation between.....	43
Pignet's formula.....	186
Pignet's index of build:	
comparison of, for men of various heights with average chest and weight for certain per cent of men of each height.....	188, 189
for draft recruits—	
with asthma.....	360
with astigmatism.....	322
with cardiac hypertrophy.....	334
with cryptorchidism, hypospadias, anorchism, and monorchism.....	382
with defective and deficient teeth.....	363
with enlarged inguinal rings.....	369
with exophthalmic goiter.....	309
with hemorrhoids.....	356
with hernia.....	369
with hyperopia.....	319
with hypertrophic tonsillitis.....	327
with mitral insufficiency.....	338
with mitral stenosis.....	339
with myopia.....	315
with simple goiter.....	305
with simple tachycardia.....	331
with valvular diseases of the heart (unclassified).....	345
with varicocele.....	348
with varicose veins.....	348
with underweight.....	376
Pignet's index of robustness.....	186
and index of build of recruits found with specified diseases and defects.....	408
for men with pulmonary tuberculosis.....	301
Plumb line and sinker used to measure sternal notch.....	57
Poles:	
absolute and relative calf circumference in, demobilization, 1919.....	231
absolute and relative chest circumference of, demobilization, 1919.....	154
absolute and relative height of pubic arch in, demobilization, 1919.....	200
absolute and relative height of sternal notch in, demobilization, 1919.....	197
absolute and relative knee height of, demobilization, 1919.....	225
absolute and relative leg length of, demobilization, 1919.....	222
absolute and relative shoulder breadth of, demobilization, 1919.....	206
absolute and relative sitting heights of, and standard deviations, demobilization, 1919.....	191
absolute and relative span of, with standard deviation, demobilization, 1919.....	194
absolute and relative thigh circumference of, at demobilization, 1919.....	228
absolute and relative transverse chest diameter of, demobilization, 1919.....	210
absolute and relative transverse pelvic diameter in, demobilization, 1919.....	217
absolute and relative waist circumference of, demobilization, 1919.....	199
approximate number measured, demobilization, 1919.....	243
comparative frequency distribution—	
of antero-posterior diameter of chest, demobilization, 1919.....	212
of calf circumference in, demobilization, 1919.....	231
of chest circumference (rest), demobilization.....	155
of span in, demobilization, 1919.....	195
of eye color in, demobilization, 1919.....	284
of hair color in, demobilization, 1919.....	292
of height of, demobilization, 1919.....	116
of height of pubic arch in, demobilization, 1919.....	201
of height of sternal notch in, demobilization, 1919.....	198
of knee height in, at demobilization, 1919.....	226
of leg length of, at demobilization, 1919.....	223

Poles—Continued.

comparative frequency distribution—Continued.	Page.
of shoulder breadth of, demobilization, 1919.....	206
of sitting height of, demobilization, 1919.....	192
of thigh circumference of, demobilization, 1919.....	229
of transverse diameter of chest in, demobilization, 1919.....	211
of transverse pelvic diameter in, demobilization, 1919.....	218
of waist circumference in, demobilization, 1919.....	214
of weight of, demobilization, 1919.....	135
from Galicia, approximate average stature of.....	47
in general, average stature of.....	68
index of build—	
demobilization, 1919.....	173, 244
maximum shoulder breadth found in.....	205
mean stature and standard deviation of.....	113
mean stature of.....	243
mean weight and standard deviation of.....	135
ratio of pelvic diameter to stature of.....	215
ratio of pubic height to total stature of.....	199
transverse chest diameter.....	208
Population:	
characteristics of, by sections.....	80-87
consolidation of similar sections of; the series and their constituent groups.....	98
of the various sections of the United States, characteristics and composition of.....	80-87
variability of physical dimensions in relation to.....	44
Portuguese, average stature of.....	68
Primates, relative leg length of.....	200
Proportions and sizes of men in distribution zones, Quartermaster Corps.....	276
Proportions, somatic, comparison of, in eight European races.....	242
Prussia, standards of stature in.....	45
Pubic arch:	
absolute and relative height of—	
in eight European races, demobilization, 1919.....	200
in five color races, demobilization, 1919.....	202
and stature, correlation of, in white troops.....	42
comparative frequency distribution of height of, in eight European races, demobiliza- tion, 1919.....	201
comparison of height of, in color races.....	202
height of.....	38
and stature, correlation between, colored troops, demobilization.....	524
and stature, correlation between, white troops, demobilization.....	520
and stature, correlation of, white troops.....	259
and stature, Negro troops, correlation between.....	265
general discussion.....	199
summary of, of approximately 6,000 colored troops, demobilization.....	234
summary of, of approximately 100,000 white troops, demobilization.....	234
standard deviation of height of.....	200
Pubic height:	
average, in color races, demobilization, 1919.....	235
comparison of, in white and Negro troops.....	40, 233
in eight European races at demobilization, 1919.....	243
mean.....	200
ratio of, to total stature.....	199
standard deviation of, for Negro troops.....	202
Pubis, height of, directions for measuring.....	58
Pygmies, Mawambu, relative arm length in.....	219
Race:	
eye color an index of.....	42
in relation to regiment formation.....	47
variability of chest circumference according to.....	37
variability of stature in.....	36
variation in size of body according to.....	46, 47
Race size, diversity of, in relation to clothing of army.....	47
Races, color:	
average dimensions in, demobilization, 1919.....	235
chest circumference of men of.....	156
comparative frequency distribution of measurements in, at demobilization.....	236
comparative sitting heights of.....	193
comparison of height of pubic arch in.....	202
comparison of height of sternal notch in.....	199

Races, color—Continued.	Page.
comparison of leg length in.....	224
comparison of neck circumference in.....	203
comparison of shoulder breadth of.....	207
comparison of transverse pelvic diameter of.....	219
comparison of weight of.....	136
general comparison of measurements of.....	241
mean index of build of.....	174
mean span of.....	196
mean stature in, demobilization, 1919.....	117
relative dimensions in, demobilization, 1919.....	235
Races, eight European:	
absolute and relative calf circumference in, demobilization, 1919.....	231
absolute and relative height of sternal notch in, demobilization, 1919.....	197
absolute and relative knee height in, demobilization, 1919.....	225
absolute and relative leg length in, demobilization, 1919.....	222
absolute and relative shoulder height in, demobilization, 1919.....	206
absolute and relative span with standard deviation in, demobilization, 1919.....	194
absolute and relative thigh circumference in, demobilization, 1919.....	228
absolute and relative transverse pelvic diameter in, demobilization, 1919.....	217
absolute and relative waist circumference in, demobilization, 1919.....	214
absolute dimensions in, demobilization, 1919.....	243
approximate number of men measured in, demobilization, 1919.....	243
average dimensions of, demobilization, 1919.....	246, 247, 248, 249, 250, 251, 252
comparative frequency distribution—	
of antero-posterior diameter of chest in each of, demobilization, 1919.....	212
of calf circumference in, demobilization, 1919.....	231
of chest circumference (rest) in, demobilization, 1919.....	155
of hair color in, demobilization, 1919.....	292
of height of pubic arch in, demobilization, 1919.....	201
of height of sternal notch in each of, demobilization, 1919.....	198
of knee height in, demobilization, 1919.....	226
of leg length in each of, demobilization, 1919.....	223
of shoulder breadth in, demobilization, 1919.....	206
of span in each of, demobilization, 1919.....	195
of thigh circumference in, demobilization, 1919.....	229
of transverse diameter of chest in each of, demobilization, 1919.....	211
of transverse pelvic diameter in each of, demobilization, 1919.....	218
of waist circumference in, demobilization, 1919.....	214
comparison of antero-posterior chest diameter in.....	209
comparison of calf circumference in.....	230, 231
comparison of chest circumference in.....	213, 214
comparison of leg length of.....	221, 222, 223
comparison of neck circumference in.....	203
comparison of shoulder breadth in.....	205, 206
comparison of sitting height in.....	191, 192
comparison of somatic proportions of.....	242
comparison of thigh circumference of.....	227, 228, 229
comparison of transverse chest diameter in.....	208, 210
comparison of transverse pelvic diameter in.....	216, 217, 218
comparison of weight in, demobilization.....	135
comparison of span in.....	193, 194
eye color in.....	284
hair color in.....	293
index of build in, demobilization, 1919.....	244
mean chest circumference in, demobilization, 1919.....	152, 154, 155
mean stature and standard deviation of each of.....	113
relative dimensions, demobilization, 1919.....	245, 250, 251, 252
Races, Mediterranean:	
relative transverse chest diameter of.....	208
thigh circumference of.....	228
waist circumference of.....	213
Races, Nordic:	
relative transverse chest diameter of.....	208
shoulder breadth of.....	205
thigh circumference of.....	228
waist circumference of.....	213
Races, principle, approximate average stature of.....	47
Races, white and color, comparison of stature of.....	117
Ration, standard, variation with size of body.....	46

Recruits (<i>see also</i> , Draft recruits; Drafted men):	Page.
age of.....	64
and veterans, comparison of stature of, by States.....	77
average stature of, from different sections.....	97, 98, 99, 100, 101, 102, 103
British, weight of.....	119
build of.....	37, 38
Civil War, weight of.....	119
comparison of hair color in World War and Civil War.....	291
comparison of stature of, from various States, 1863-1864, and 1917-1918.....	78
distribution of frequencies of different classes of chest circumference for.....	140
draft, anthropometric work in connection with.....	49
draft, measurements of.....	51
found with specified diseases and defects, index of build and Pignet's index of robustness of.....	408
frequency distribution of, by States.....	140
frequency distribution of weight of.....	120
height, weight, and chest circumference of, in relation to various diseases and defects.....	296
index of build of, by sections, 1917-18.....	172
mean chest circumference of.....	37
mean stature of.....	34
mean stature of, at each age, 18 to 25 years, 1906-1915.....	73
mean stature of, from different States.....	74
mean weight of.....	36
mean weight of, from the different sections.....	124, 125, 126, 127
of 1917-1918 and veterans of 1919 and 1864-1865; comparison of index of build in.....	168
standard deviation of weight of.....	121
stature of, index of ability to carry weight.....	46
with asthma—	
correlation between height and chest circumference (expiration) in.....	359
height and weight in.....	358
with astigmatism—	
correlation between height and chest circumference in.....	322
correlation between height and weight in.....	321
with cardiac hypertrophy—	
correlation between height and chest circumference in.....	333
correlation between height and weight in.....	332
with cryptorchidism, hypospadias, anorchism and monorchism—	
correlation between height and chest circumference (expiration) in.....	381
correlation between height and weight in.....	380
with defective and deficient teeth and dental caries—	
correlation between height and chest circumference (expiration) in.....	362
correlation between height and weight in.....	361
with defective physical development—	
correlation between height and chest circumference (expiration) in.....	374
correlation between height and weight in.....	373
with enlarged inguinal rings—	
correlation between height and chest circumference (expiration) in.....	368
correlation between height and weight in.....	367
with exophthalmic goiter—	
correlation between height and chest circumference in.....	311
correlation between height and weight in.....	310
with flat-foot—	
correlation between height and weight in.....	371
with hemorrhoids—	
correlation between height and chest circumference (expiration) in.....	355
correlation between height and weight in.....	354
with hernia—	
correlation between height and chest circumference (expiration) in.....	365
correlation between height and weight in.....	366
with hyperopia—	
correlation between height and chest circumference in.....	317
correlation between height and weight in.....	316
with hypertrophic tonsillitis—	
correlation between height and chest circumference in.....	325
correlation between height and weight of.....	324
with mitral insufficiency—	
correlation between height and chest circumference in.....	337
correlation between height and weight in.....	336
with mitral stenosis—	
correlation between height and chest circumference (expiration) in.....	341
correlation between height and weight in.....	340

Recruits—Continued.

with myopia—	Page.
correlation between height and chest circumference in.....	313
correlation between height and weight in.....	312
with pulmonary tuberculosis—	
correlation between height and chest circumference in.....	303
correlation between height and weight in.....	302
with simple goiter—	
correlation between height and chest circumference in.....	307
correlation between height and weight in.....	306
with simple tachycardia—	
correlation between height and chest circumference in.....	329
correlation between height and weight in.....	328
with specified diseases and defects, mean stature and weight in.....	408
with underweight—	
correlation between height and chest circumference (expiration) in.....	378
correlation between height and weight in.....	377
with valvular diseases of the heart (unclassified)—	
correlation between height and chest circumference (expiration) in.....	344
correlation between height and weight in.....	343
with varicocele—	
correlation between height and chest circumference (expiration) in.....	352
correlation between height and weight in.....	351
with varicose veins—	
correlation between height and chest circumference (expiration) in.....	350
correlation between height and weight in.....	349
World War—	
and Civil War, comparison of stature of.....	79
average index of build for.....	164
correlation between stature and chest circumference of.....	256
Refraction, errors of (<i>see</i> Eye, refractive errors of).	
Regiment formation, relation of race to.....	47
Regular Army, minimum stature in, in Civil War.....	45
Rhode Island:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919...	122
comparative view of mean height and mean weight of men from.....	124
comparison of index of build of men from, at demobilization in 1865 and 1919.....	170
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144
Robustness in drafted men (<i>see also</i> , Build):	
with underweight.....	376
with asthma.....	360
with cardiac hypertrophy.....	334
with cryptorchidism, hypospadias, anorchism, and monorchism.....	382
with defective and deficient teeth.....	363
with defective physical development.....	375
with enlarged inguinal rings.....	369
with flat-foot.....	370
with hemorrhoids.....	356
with hernia.....	369
with hyperopia.....	319
with mitral insufficiency.....	338
with overweight and obesity.....	379
with astigmatism.....	322

Robustness in drafted men—Continued.	Page.
with exophthalmic goiter.....	309
with hypertrophic tonsillitis.....	327
with mitral stenosis.....	339
with myopia.....	315
with pulmonary tuberculosis.....	301
with simple goiter.....	305
with simple tachycardia.....	331
with valvular diseases of the heart (unclassified).....	345
with varicocele.....	348
with varicose veins.....	348
Pignet's index of.....	186
for men with pulmonary tuberculosis.....	301
Rumanians:	
approximate average stature of.....	47
average weight of adult male.....	120
conscripts, average stature of.....	68
from Hungary—	
approximate average stature of.....	47
average stature of.....	68
ratio of pelvic diameter to stature of.....	215
Russian groups:	
height distribution of.....	109, 110
index of build of recruits of.....	173
mean chest circumference of draft recruits of.....	150
mean height of.....	108
mean weight of draft recruits of.....	132
relative chest circumference of draft recruits of.....	150
weight distribution of draft recruits of.....	133, 134
Russians:	
chest circumference of.....	139
Great, approximate average stature of.....	47
in general, ratio of pelvic diameter to stature of.....	215
Little, approximate average stature of.....	47
white, approximate average stature of.....	47, 68
Ruthenians, approximate average stature of.....	147
Samoyedes Peninsula of Siberia, relative pubic height of inhabitants of.....	200
Scales, graduated paper metric, used in taking measurements.....	57
Scandinavian and German groups:	
height distribution of.....	109, 110
mean chest circumference of draft recruits of.....	150
mean height of.....	108
mean weight of draft recruits of.....	132
relative chest circumference of draft recruits of.....	150
weight distribution of draft recruits of.....	133, 134
Scandinavian groups:	
height distribution of.....	109, 110
index of build of recruits of.....	173
mean chest circumference of draft recruits of.....	150
mean weight of draft recruits of.....	132
relative chest circumference of draft recruits of.....	150
weight distribution of draft recruits of.....	133, 134
Scandinavians, sitting height of.....	190
Scotch:	
absolute and relative calf circumference in, demobilization, 1919.....	231
absolute and relative chest circumference in, demobilization, 1919.....	154
absolute and relative height of pubic arch in, demobilization, 1919.....	200
absolute and relative height of sternal notch in, demobilization, 1919.....	197
absolute and relative knee height of, demobilization, 1919.....	225
absolute and relative leg length of, demobilization, 1919.....	222
absolute and relative shoulder breadth of, demobilization, 1919.....	206
absolute and relative sitting height of, and standard deviations, demobilization, 1919.....	191
absolute and relative span of, with standard deviation, demobilization, 1919.....	194
absolute and relative thigh circumference of, demobilization, 1919.....	228
absolute and relative transverse chest diameter of, demobilization, 1919.....	210
absolute and relative transverse pelvic diameter in, demobilization, 1919.....	217
absolute and relative waist circumference of, demobilization, 1919.....	199
agriculturists of Galway, average stature of.....	68
approximate average stature of.....	47
approximate number measured, demobilization, 1919.....	243

Scotch—Continued.

	Page.
comparative frequency distribution—	
of antero-posterior diameter of chest in, demobilization, 1919.....	212
of calf circumference of, demobilization, 1919.....	231
of chest circumference (rest), demobilization, 1919.....	155
of eye color in, demobilization, 1919.....	284
of hair color in, demobilization, 1919.....	292
of height of, demobilization, 1919.....	116
of height of pubic arch in, demobilization, 1919.....	201
of height of sternal notch in, demobilization, 1919.....	198
of knee height in, at demobilization, 1919.....	226
of leg length of, at demobilization, 1919.....	223
of shoulder breadth of, demobilization, 1919.....	206
of sitting height of, demobilization, 1919.....	192
of span of, demobilization, 1919.....	195
of thigh circumference of, demobilization, 1919.....	229
of transverse diameter of chest in, demobilization, 1919.....	211
of transverse pelvic diameter in, demobilization, 1919.....	218
of waist circumference in, demobilization, 1919.....	214
of weight of, demobilization, 1919.....	135
in general, average stature of.....	68
index of build of.....	173, 244
mean stature and standard deviation of.....	113
mean stature of.....	243
mean weight and standard deviation of.....	135
of the north, Ayrshire, etc., average stature of.....	68
shoulder breadth of.....	205
transverse chest diameter of.....	208
Seaver measuring rod, used in taking measurements.....	57
Sections:	
average stature of recruits from different.....	100
composition of.....	88-96
frequency distribution of statures in groups of.....	111, 114, 115
Serbs:	
approximate average stature of.....	47
chest circumference of.....	139
conscripts, average stature of.....	68
Shape, main differences between white and Negro troops.....	40, 41
Shoulder breadth.....	38
absolute and relative, in eight European races, demobilization, 1919.....	206
average, in color races, demobilization, 1919.....	235
comparative frequency distribution of, in eight European races, demobilization, 1919.....	206
comparison of, in color races.....	207
comparison of, in eight European races.....	205, 206, 243
general discussion of.....	203
in relation to total stature.....	204
maximum, found in Poles.....	205
mean.....	204
minimum found in French.....	205
of anthropoid apes.....	204
of chimpanzee.....	204
of English.....	205
of Germans.....	205
of Germans of Bavaria.....	204
of Hebrews.....	205
of hylobates.....	204
of inhabitants of Admiral Islands.....	204
of Irish.....	205
of Italians.....	205
of Nordic races.....	205
of orang outang.....	204
of Parisians.....	204
of Polish Jews.....	204
of Scotch.....	205
relative—	
of Bavarians.....	204
of Belgians.....	204
of Chinese.....	204
of French.....	204
of Japanese.....	204
of Polish Jews.....	204

Shoulder breadth—Continued.	Page.
standard deviation of.....	204
summary of, of approximately 6,000 colored troops, demobilization.....	234
summary of, of approximately 100,000 white troops, demobilization.....	234
Shoulder width and blouse groups:	
association between—	
colored troops, demobilization.....	542
white troops, demobilization.....	534
Shoulders, transverse diameter of, at level of humeri, directions for measuring.....	58
Siberia, Samoyedes Peninsula of, relative pubic height of inhabitants of.....	200
Sitting height.....	38
absolute and relative, and standard deviations with coefficient of variations in eight European races, demobilization, 1919.....	191
average, in color races, demobilization, 1919.....	235
and chest circumference—	
correlation between, in Negro troops.....	266, 540
correlation between, white troops.....	266, 532, 533, 537
and stature—	
correlation between, in Negro troops.....	264
correlation between, in white troops.....	259, 517
comparative frequency distribution of, in each of eight European races, demobilization, 1919.....	192, 243
comparative, of white and Negro troops.....	40
comparison of, in white and Negro troops.....	233
general discussion.....	190
mean.....	148
standard deviation of.....	191
summary of, of approximately 6,000 colored troops, demobilization.....	234
summary of, of approximately 100,000 white troops, demobilization.....	234
Size of body:	
in relation—	
to standard ration.....	34, 46
to standards for uniforms.....	46
variation according to race.....	46, 47
Sizes and proportions of men in distribution zones, Quartermaster Corps.....	276
Somatic proportions, comparison of, in eight European races.....	242
South Carolina:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919.....	122
comparative view of mean height and mean weight of men from.....	124
difference of weight of draft recruits from, at mobilization, 1917-1918, and demobilization, 1919.....	123
increase in stature of soldiers at demobilization over stature of recruits, 1917-1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration) of draft recruits from.....	142
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144
South Dakota:	
absolute and relative numbers of veterans from—	
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
with blue eyes, with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919.....	122
comparative view of mean height and mean weight of men from.....	124

South Dakota—Continued.	Page.
difference of weight of draft recruits from, at mobilization, 1917-1918, and demobilization, 1919.....	123
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build—	
demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration) of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, demobilization, 1919.....	76
relative chest circumference of recruits from.....	144
Span.....	38
absolute.....	193, 194
absolute and relative, with standard deviation in eight European races, demobilization, 1919.....	194
average in color races, demobilization, 1919.....	235
comparative frequency distribution in each of eight European races, demobilization, 1919.....	195
comparative, of white and Negro troops.....	39, 232
comparison of, in eight European races.....	193, 194, 243
directions for measuring.....	57
general discussion.....	193
mean.....	193
of color races.....	196
relative.....	193, 194
summary of, of approximately 6,000 colored troops, demobilization.....	234
summary of, of approximately 100,000 white troops, demobilization.....	234
Span and stature:	
correlation between—	
colored troops, demobilization.....	522
white troops.....	42, 260, 518, 519, 523
Negro troops.....	265
Spaniards, average stature of.....	68
Sparsely settled groups:	
height distribution of.....	109, 110
index of build of recruits of.....	173
mean chest circumference of draft recruits of.....	150
mean weight of draft recruits of.....	132
relative chest circumference of draft recruits of.....	150
weight distribution of draft recruits of.....	133, 134
Standards:	
of height, weight, chest circumference, and mobility of chest, adopted for draft recruits.	
United States Army, 1917.....	297
of measurements of drafted men.....	296
physical examination.....	297
States, measurements by (<i>see</i> under names of States).	
Statistical perforated cards.....	63
Stature (<i>see also</i> , Height).....	34
age of volunteers and.....	35
approximate of principal races in United States.....	47
average—	
in color races, demobilization, 1919.....	235
of adult males of various nativities in United States in Civil War.....	69
of Civil War and World War compared.....	34
of recruits from different sections.....	97, 98, 99, 100, 101, 102, 103
calculated frequency distribution of, of men of United States Civil War period, France, Belgium, and Italy.....	71
chest circumference in relation to.....	48
comparative frequency distribution of, by Quartermaster Corps distribution zones, demobilization.....	568, 569
comparison of frequency distribution of, of Civil War and World War recruits.....	72
comparison of—	
in eight European races at demobilization.....	113, 243, 244
in white and color races.....	117
of native and foreign born white and colored draft recruits, 1917-1918, and white recruits, Civil War.....	79
of recruits and veterans, by States.....	77
correlated with length of leg.....	46
directions for measuring.....	57, 67

Stature—Continued.	Page.
frequency distribution of	71
by classes at mobilization and demobilization	70
gain in	34
general discussion of	66
high and low standard deviations in, in different sections	105
in drafted men—	
with flat-foot	370
with underweight	375
increase in, of soldiers at demobilization over stature of recruits, 1917-1918	76
index of build of Civil and World War veterans for each inch of	164
importance of weight in relation to	48
influence of immigration on	79
its mean, standard deviation, and coefficient of variation, for certain especially studied groups	69
limitations of, in U. S. Army in World War	46
mean	67. 74. 76
and standard deviation of each of eight European races	113
at demobilization	75
at each age, 18 to 25 years, United States Army recruits	73
by groups of similar sections	105
by States, first million draft recruits	75
by States, of soldiers at demobilization, 1919	76
comparison of, in various countries	68
comparison of, with Civil War records	67
from different States	74. 75
in five color races, demobilization, 1919	117
of first million recruits	34
of recruits	74
of troops at demobilization	34
mean weight in relation to	118
medico-military importance of	66
minimum in British Army	45
in Civil War	45
in French Army	45
in Italian Army	45
of Army conscripts and recruits, in inches, as determined by Laplace-Charlier frequency curves	70
of drafted men	296
with asthma	356
with astigmatism	319
with cardiac hypertrophy	331
with cryptorchidism, hypospadias, anorchism, and monorchism	379
with defective and deficient teeth	360
with defective physical development	372
with enlarged inguinal rings	369
with exophthalmic goiter	308
with hemorrhoids	353
with hernia	363
with hyperopia	315
with hypertrophic tonsillitis	323
with mitral insufficiency	335
with mitral stenosis	338
with myopia	314
with overweight and obesity	379
with pulmonary tuberculosis	299
with simple goiter	304
with simple tachycardia	327
with valvular diseases of the heart (unclassified)	342
with varicocele	345
with varicose veins	345
of recruits—	
comparison of, from various States, 1863-1864 and 1917-1918	78
index of ability to carry weight	46
of volunteer troops in Civil War	45
ratio of pelvic diameter to	215
standard deviation of	73
standards of—	
in Austria	45
in Belgium	45
in Great Britain	45

Stature—Continued.	
standards of—Continued.	Page.
in Prussia.....	45
in Switzerland.....	45
in the United States.....	45
summary of, of approximately 6,000 colored troops, demobilization.....	234
summary of, of approximately 100,000 white troops, demobilization.....	234
total—	
ratio of pubic height to.....	199
shoulder breadth in relation to.....	204
associated with various defects and diseases, first and second million draft recruits.....	409
by States.....	35, 36
in races.....	36
and cardiac disorders.....	43
Stature and chest circumference:	
correlation between—	
Civil War recruits.....	256
recruits, 1917-1918.....	256
Stature and defective teeth.....	43
Stature and goiter.....	43
Stature and height of pubic arch:	
correlation between—	
colored troops, demobilization.....	524
white troops.....	259, 520
Stature and height of sternal notch:	
correlation between—	
colored troops, demobilization.....	523
Negro troops.....	265
white troops.....	259, 518, 519
Stature and knee height:	
correlation of—	
Negro troops.....	265
white troops.....	42
Stature and other measurements, comparative, white and colored soldiers.....	41
Stature and pubic arch:	
correlation of—	
Negro troops.....	265
white troops.....	42
Stature and pulmonary tuberculosis.....	43
Stature and refractive errors.....	43
Stature and sitting height:	
correlation between—	
colored troops, demobilization.....	521
Negro troops.....	264
white troops.....	42, 259, 517
Stature and span:	
correlation between—	
colored troops, demobilization.....	522
Negro troops.....	265
white troops.....	42, 260, 518, 519
Stature and sternal notch, correlation of, in white troops.....	42
Stature and varicocele.....	43
Stature and varicose veins.....	43
Stature and waist circumference:	
correlation between.....	258
white and colored troops, demobilization.....	509
Stature and weight:	
correlations between.....	254
in white and colored troops, demobilization.....	508
distribution of—	
American-born Civil War recruits.....	74
draft recruits, 1917-1918.....	74
mean, index of build for.....	164
mean, of recruits with specified diseases and defects.....	408
standards of.....	297
Statures:	
frequency distribution of, in the groups of sections.....	111, 114, 115
weights associated with, in various classes of American males.....	118

	Page.
Sternal notch.....	38
absolute and relative height of—	
in eight European races, demobilization, 1919.....	197
in five color races, demobilization, 1919.....	199
average in color races, demobilization, 1919.....	235
comparative frequency distribution of height of, in each of eight European races, demobilization, 1919.....	198
comparative, of white and Negro troops.....	40
comparison of height of—	
in color races.....	199
in eight European races.....	197
in white and Negro troops.....	233
general discussion of.....	196
height of.....	196
and stature, correlation between, colored troops, demobilization.....	523
and stature, correlation between, white troops.....	42, 259, 518, 519
directions for measuring.....	58
Negro troops.....	265
summary of, of approximately 6,000 colored troops, demobilization.....	234
summary of, of approximately 100,000 white troops, demobilization.....	234
in eight European races at demobilization, 1919.....	243
mean height of.....	197
method of measuring height of.....	196
plumb line and sinker used to measure.....	57
Suprapatella:	
and breeches groups, association between, white troops, demobilization.....	552
average, in color races, demobilization, 1919.....	235
circumferences—	
and blouse groups, association between, colored troops, demobilization.....	550
general discussion.....	232
mean.....	232
standard deviation of.....	232
summary of, of approximately 6,000 colored troops, demobilization.....	234
summary of, of approximately 100,000 white troops, demobilization.....	234
Swedes:	
approximate average stature of.....	47
in general (soldiers), average stature of.....	68
of Kalmar (conscripts), average stature of.....	68
Switzerland:	
standards of stature in.....	45
French, conscripts of, average stature of.....	68
Tachycardia, simple:	
chest circumference in drafted men with.....	330
correlation between—	
height and chest circumference in recruits with.....	329
height and weight in recruits with.....	328
in drafted men.....	327
Pignet's index of build in drafted men with.....	331
robustness in drafted men with.....	331
stature of drafted men with.....	327
weight in drafted men with.....	330
Tape, cloth, metrically graduated, used in taking measurements.....	57
Tartars, relative leg length of.....	221
Teeth, defective and deficient:	
and dental caries—	
correlation between height and chest circumference (expiration) in recruits with.....	362
correlation of height and weight in recruits with.....	361
chest circumference in drafted men with.....	363
Pignet's index of build for drafted men with.....	363
robustness in drafted men with.....	363
stature of drafted men with.....	360
weight of drafted men with.....	360
Teeth, defective, and stature.....	43
Tennessee:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283

Tennessee—Continued.

absolute and relative numbers of veterans from—Continued.	Page.
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits, at mobilization, 1917-1918, and demobilization, 1919..	122
comparative view of mean height and mean weight of men from.....	124
comparison of index of build of men from, at demobilization in 1865 and 1919.....	170
difference of weight of draft recruits from, at mobilization, 1917-1918, and demobiliza- tion, 1919.....	123
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization.....	76
relative chest circumference of recruits from.....	144

Texas:

absolute and relative numbers of veterans from—	
with blue eyes, with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919..	122
comparative view of mean height and mean weight of men from.....	124
difference of weight of draft recruits from, at mobilization, 1917-1918, and demobiliza- tion, 1919.....	123
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build for, at demobilization, 1919.....	167
index of build for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144

Thigh circumference.

absolute and relative, in eight European races, demobilization, 1919.....	228
and breeches group—	
association between, colored troops, demobilization.....	558
association between, white troops, demobilization.....	551
average in color races, demobilization, 1919.....	235
comparative frequency distribution of, in eight European races, demobilization.....	229
general discussion.....	227
in eight European races.....	227, 228, 229, 243
maximum, directions for measuring.....	59
mean.....	227
standard deviation of.....	227
summary of, of approximately 6,000 colored troops, demobilization.....	234
summary of, of approximately 100,000 white troops, demobilization.....	234
Thoracic deviation, standard.....	266

Thoracic index (*see also*, Chest Diameter, Transverse):

antero-posterior diameter of chest, and.....	208
for African Negroes.....	207, 209
for Bugu Negroes.....	207, 209
for French.....	209
for Hova Indians.....	207, 209
for Navajo Indians.....	209
for various races.....	209

Thuringians of Saxony (conscripts).....	68
Thyroid disease, Army anthropology an aid in detecting.....	34

Tonsillitis, hypertrophic:	Page.
and weight.....	43
chest circumference of drafted men with.....	326
correlation between—	
height and chest circumference of recruits with.....	325
height and weight of recruits with.....	324
in drafted men.....	323
Pignet's index of build for drafted men with.....	327
robustness of drafted men with.....	327
stature of drafted men with.....	323
weight of drafted men with.....	323
Troops, white and Negro, the general comparative picture of.....	39, 41
Tuberculosis, pulmonary:	
and chest circumference.....	43
and stature.....	43
and weight.....	43
Army anthropology an aid in detecting.....	34
chest circumference of drafted men with.....	301
correlation between height and chest circumference in recruits with.....	303
correlation between height and weight in recruits with.....	302
in drafted men.....	299
Pignet's index of robustness for men with.....	301
robustness of drafted men with.....	301
stature of drafted men with.....	299
Turks from Balkans, average stature of.....	68
Ukrainians, average stature of.....	68
Underweight:	
chest circumference of drafted men with.....	379
correlation between—	
height and chest circumference in recruits with.....	376
height and weight in recruits with.....	377
Pignet's index of build for drafted men with.....	376
robustness in drafted men with.....	376
stature in drafted men with.....	375
weight of drafted men with.....	376
Uniforms:	
patterns for.....	271
size and proportions of body essential to proper cutting of.....	34
United States:	
difference of weight of draft recruits of, at mobilization, 1917-1918, and demobilization, 1919.....	123
increase in stature of soldiers at demobilization over stature of recruits, 1917-1919.....	76
mean stature, by States, of soldiers at demobilization, 1919.....	76
standards of stature in.....	45
Utah:	
absolute and relative proportion of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits, at mobilization, 1917-1918, and demobilization, 1919.....	122
comparative view of mean height and mean weight of men, from.....	124
difference of weight of draft recruits from, at mobilization, 1917-1918, and demobilization, 1919.....	123
increase in stature of soldiers, at demobilization over recruits, 1917-1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration) of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144

	Page.
Variation in two sides of body, correlation of.....	42
Varicocele:	
chest circumference of drafted men with.....	347
correlation between—	
height and chest circumference (expiration) of recruits with.....	352
height and weight of recruits with.....	351
Pignet's index of build for drafted men with.....	348
robustness of drafted men with.....	348
stature and.....	43
stature of drafted men with.....	345
weight of drafted men with.....	346
Varicose veins:	
chest circumference of drafted men with.....	347
correlation between—	
height and chest circumference (expiration) in recruits with.....	350
height and weight in recruits with.....	349
in drafted men.....	345
Pignet's index of build for drafted men with.....	348
robustness of drafted men with.....	348
stature of drafted men with.....	345
weight of drafted men with.....	346
Veins, varicose (<i>see</i> Varicose veins).	
Venetians, average stature of.....	68
Vermont:	
absolute and relative proportion of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919... 122	
colored eyes, proportional numbers for, 1865.....	285
comparative view of mean height and mean weight of men from.....	124
comparison of index of build of men from, at demobilization in 1865 and 1919.....	170
increase in stature of soldiers, at demobilization over recruits, 1917-1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration) of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144
Vertebra-ulna measurement, directions for taking.....	58
Veterans:	
and recruits, comparison of stature of, by States.....	77
build of.....	37
Civil and World Wars, index of build of.....	164, 168
Civil War, build of recruits compared with.....	38
distribution frequencies of different classes of chest circumference for.....	140
increase in mean chest circumference.....	37
Virginia:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919... 122	
comparative view of mean height and mean weight of men from.....	124
difference of weight of draft recruits from, at mobilization, 1917-1918, and demobiliza- tion, 1919.....	123

Virginia—Continued.	Page.
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build—	
demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration) of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144
Volunteers, age of, and stature.....	35
Volunteer troops, stature of, in Civil War.....	45
Waist circumference.....	39
absolute and relative, in eight European races, demobilization, 1919.....	214
and leg length, correlation between, colored troops, demobilization, 1919.....	555
and leg length—	
correlation between, white troops, demobilization.....	548
correlation between, white troops.....	42
and pelvic diameter, correlation of, in white troops.....	42
and stature—	
correlation between.....	258
correlation between, white and colored troops, demobilization.....	509
and transverse diameter of pelvis—	
correlation between, colored troops, demobilization.....	530
correlation between, Negro troops.....	267
correlation between, white troops.....	262, 515
average, in color races, demobilization, 1919.....	235
comparative frequency distribution of—	
by Quartermaster Corps distribution zones, white and colored troops, demobilization.....	572
in eight European races, demobilization, 1919.....	214
comparison of, in eight European races.....	213, 214
general discussion of.....	213
in eight European races, demobilization, 1919.....	243
mean.....	213
level of umbilicus, directions for measuring.....	58
standard deviation.....	213
summary of, of approximately 6,000 colored troops, demobilization.....	234
summary of, of approximately 100,000 white troops, demobilization.....	234
Washington:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits, at mobilization, 1917-1918, and demobilization, 1919..	122
comparative view of mean height and mean weight of men from.....	124
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
increase in stature of soldiers at demobilization over stature of recruits, 1917-1919.....	76
mean chest circumference (expiration) of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization.....	76
relative chest circumference of recruits from.....	144
Weddas, calf circumference of.....	230
Weight.....	36
at mobilization, 1917-1918, and demobilization, 1919, States arranged in order of difference of.....	123
average—	
by States, at mobilization, 1917-1918, and demobilization, 1919, first million draft recruits.....	122
in color races, demobilization, 1919.....	235
comparative frequency distribution of—	
by States of nativity, white and colored troops, demobilization.....	570
in eight European races, demobilization.....	135

Weight—Continued.	Page.
comparative, of men of different statures among white soldiers of 1865 at demobilization and white and colored soldiers at demobilization, 1919.....	254
comparison of, in the color races.....	136
comparison of, in eight European races.....	135, 244
distribution of—	
by height.....	159, 160, 161
by special diseases or defects.....	390, 391, 400, 401
by groups of sections, first million draft recruits.....	130, 131, 133, 134
frequency distribution of.....	120
of various classes of weight at mobilization, 1917–1918, and demobilization, 1919..	121
general discussion of.....	117
height, and chest circumference (expiration) measurements showing proportionate measurements of two of them to the total of the third, first million draft recruits.....	177
height and chest circumference of recruits in relation to various diseases and defects....	296
importance of—	
in diagnosis.....	48
in relation to stature.....	48
in drafted men with exophthalmic goiter.....	308
in drafted men—with flat-foot.....	370
with simple tachycardia.....	330
in eight European races, demobilization, 1919.....	243
in relation to blouse groups.....	272
increase in, at demobilization over mobilization.....	122, 123
maximum, allowed by Army regulations.....	297
mean.....	119
and height, mean, comparative view of, of men from different States, World War and Civil War.....	124
and standard deviation in each of eight European races.....	135
by groups and component sections, first million draft recruits.....	430–435
by groups of sections, first million draft recruits.....	132
by groups of sections.....	132
by sections, first million draft recruits.....	125
for different groups.....	127
for the different States.....	121
in five color races, with standard deviation for white and Negro troops, demobilization, 1919.....	136
relation to stature.....	118
of first million recruits.....	36
of recruits from the different sections.....	124, 125, 126, 127
measurement by special diseases.....	395
medico-military importance of.....	118
method for measuring.....	118
minimum, allowed by The Adjutant General.....	297
of drafted men.....	296
with asthma.....	356
with astigmatism.....	320
with cardiac hypertrophy.....	331
with cryptorchidism, anorchism, monorchism, and hypospadia.....	382
with defective and deficient teeth.....	360
with defective physical development.....	372
with enlarged inguinal rings.....	369
with hemorrhoids.....	353
with hernia.....	364
with hyperopia.....	318
with hypertrophic tonsillitis.....	323
with mitral insufficiency.....	335
with mitral stenosis.....	339
with myopia.....	314
with overweight and obesity.....	379
with pulmonary tuberculosis.....	300
with simple goiter.....	305
with underweight.....	376
with valvular diseases of the heart (unclassified).....	342
with varicocele.....	346
with varicose veins.....	346
of men at demobilization.....	119, 120
reduced variability at demobilization.....	36
standard deviation of recruits.....	121
summary of, of approximately 6,000 colored troops, demobilization.....	234

Weight—Continued.	Page.
summary of, of approximately 100,000 white troops, demobilization.....	234
variability in—	
and conditions of life.....	37
associated with various diseases and defects among first and second million draft recruits.....	409
by races.....	37
by States.....	36
Weight and blouse groups:	
association between—	
colored troops, demobilization.....	541
white troops, demobilization.....	533
Weight and chest circumference:	
correlation between.....	258
in white troops.....	42, 260
Weight and chest circumference (expiration):	
classes, first million draft recruits.....	162
correlation between—	
first million draft recruits.....	422, 423
group 1, agricultural, North, native white, first million draft recruits.....	444
group 2, agricultural, mixed foreign and white, North, first million draft recruits..	447
group 3, agricultural, native white, South, first million draft recruits.....	450
group 4, agricultural, Negro, 45 per cent plus, first million draft recruits.....	453
group 5, eastern manufacturing, first million draft recruits.....	456
group 6, commuter, first million draft recruits.....	459
group 7, mining, first million draft recruits.....	462
group 8, sparsely settled, first million draft recruits.....	465
group 9, desert, first million draft recruits.....	468
group 10, maritime, first million draft recruits.....	471
group 11, mountain, first million draft recruits.....	474
group 12, mountain whites, first million draft recruits.....	477
group 13, Indian, first million draft recruits.....	480
group 14, Mexican, first million draft recruits.....	483
group 15, native whites of Scotch origin, first million draft recruits.....	486
group 16, Russian, first million draft recruits.....	489
group 17, Scandinavian, first million draft recruits.....	492
group 18, Finn, first million draft recruits.....	495
group 19, French-Canadian, first million draft recruits.....	498
group 20, German and Scandinavian, first million draft recruits.....	501
group 21, German and Austrian, first million draft recruits.....	504
group 22, German and Austrian, first million draft recruits.....	507
Negro troops.....	265
relative, for men found with special diseases or defects in first and second million draft recruits.....	410
Weight and chest circumference (rest), correlation between, white troops, demobilization..	511
Weight and diseases of the heart.....	43
Weight and flat-foot.....	43
Weight and goiter.....	43
Weight and height:	
classes, first million draft recruits.....	158
correlation between—	
for first million draft recruits.....	417, 418, 419
group 1, agricultural, North, native white, first million draft recruits.....	442
group 2, agricultural, mixed foreign and native white, North, first million draft recruits.....	445
group 3, agricultural, native white, South, first million draft recruits.....	448
group 4, agricultural, Negro, 45 per cent plus, first million draft recruits.....	451
group 5, eastern manufacturing, first million draft recruits.....	454
group 6, commuter, first million draft recruits.....	457
group 7, mining, first million draft recruits.....	460
group 8, sparsely settled, first million draft recruits.....	463
group 9, desert, first million draft recruits.....	466
group 10, maritime, first million draft recruits.....	469
group 11, mountain, first million draft recruits.....	472
group 12, mountain whites, first million draft recruits.....	475
group 13, Indian, first million draft recruits.....	478
group 14, Mexican, first million draft recruits.....	481
group 15, native whites of Scotch origin, first million draft recruits.....	484
group 16, Russian, first million draft recruits.....	487
group 17, Scandinavian, first million draft recruits.....	490

Weight and height—Continued.	
correlation between—Continued.	Page.
group 18, Finn, first million draft recruits.	493
group 19, French-Canadian, first million draft recruits.	496
group 20, German and Scandinavian, first million draft recruits.	499
group 21, German and Austrian, 20 per cent plus, first million draft recruits.	502
group 22, German and Austrian, 15 per cent plus, first million draft recruits.	505
Weight and hypertrophic tonsillitis.	43
Weight and refractive errors.	43
Weight and stature:	
correlations between.	254
in white and colored troops, demobilization.	508
distribution of—	
American-born Civil War recruits.	74
draft recruits, 1917-1918.	74
mean—	
index of build for.	164
of recruits with specified diseases and defects.	408
standards of.	297
Weight and tuberculosis.	43
Weight and varicose veins.	43
Weights:	
associated with statures, in various classes of American males.	118
associated with statures with standard deviations, and coefficient of variation for each, in various classes of American males.	118
average, of adult males of various nationalities.	120
heights, and chest circumferences (expiration), various, shown for United States, first million draft recruits.	157
Welsh, average stature of.	68
West Virginia:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.	282
with clear blue eyes, demobilization, 1919.	281
with dark brown eyes, demobilization, 1919.	283
with dark brown hair, demobilization, 1919.	291
with flaxen hair, demobilization, 1919.	288
with light brown eyes, demobilization, 1919.	283
with light brown hair, demobilization, 1919.	290
with medium brown hair, demobilization, 1919.	290
with red hair, demobilization, 1919.	289
average weight of draft recruits, at mobilization, 1917-18, and demobilization, 1919.	122
colored eyes, proportional numbers for, 1865.	285
comparative view of mean height and mean weight of men from.	124
difference of weight of draft recruits from, at mobilization, 1917-18 and demobilization, 1919.	123
increase in stature of soldiers at demobilization over recruits, 1917-1919.	76
index of build—	
demobilization, 1919.	167
for recruits from, at mobilization, 1917-1918.	166
mean chest circumference (expiration), of draft recruits from.	142
mean stature—	
of draft recruits from.	75
of soldiers from, demobilization, 1919.	76
relative chest circumference of recruits from.	144
White and colored troops, demobilization:	
comparative frequency distribution—	
of chest circumference (rest), by Quartermaster Corps distribution zones.	571
of height, by States of nativity.	566, 567
of waist circumference, by Quartermaster Corps distribution zones.	572
of weight of, by States of nativity.	570
stature and waist circumference.	509
stature and weight in.	508
White and Negro troops:	
comparison of dimensions of.	232, 233
correlations between measurements for.	253
mean and relative chest circumference (rest), demobilization, 1919.	156
White race:	
mean stature of, at demobilization, 1919.	117
mean weight in, with standard deviation for, demobilization, 1919.	136

White troops:	Page.
absolute and relative neck circumference of, demobilization, 1919.....	203
association between blouse groups—	
and antero-posterior chest diameter, demobilization.....	536
and neck circumference, demobilization.....	538
and shoulder width, demobilization.....	534
and total arm length, demobilization.....	539
and transverse chest diameter, demobilization.....	535
and transverse pelvic diameter, demobilization.....	537
and weight, demobilization.....	533
association between breeches groups—	
and calf circumference, demobilization.....	554
and knee height, demobilization.....	550
and patella circumference, demobilization.....	553
and suprapatella, demobilization.....	552
and thigh circumference, demobilization.....	551
and transverse pelvic diameter, demobilization.....	549
calf circumference in.....	230
comparative frequency distribution—	
of blouse groups, by States of nativity.....	562
of breeches groups, by States of nativity.....	564
correlation between chest circumference—	
and sitting height, demobilization.....	532, 534
correlation between chest circumference (rest)—	
and neck circumference, demobilization.....	512
and transverse pelvic diameter, demobilization.....	513
and weight, demobilization.....	511
leg length and knee height, demobilization.....	510
length of arm and forearm in, demobilization.....	516
measurements for, demobilization.....	258-264
stature and height of pubic arch, demobilization.....	520
stature and height of sternal notch, demobilization.....	518, 519
stature and sitting height, demobilization.....	517
stature and span, demobilization.....	518, 519
transverse and antero-posterior chest, demobilization.....	514
waist circumference and leg length, demobilization.....	548
waist circumference and transverse pelvic diameter, demobilization.....	515
dimensions of, associated with breeches groups, demobilization.....	274, 275
mean knee height of.....	224
mean leg length of.....	221
mean transverse diameter of pelvis of.....	216
relative arm length of.....	219
standard deviation—	
of arm length of.....	220
of leg length of.....	221
of transverse pelvic diameter of.....	216
suprapatella, circumference of.....	232
thigh circumference of.....	227
Whites:	
absolute and relative height of pubic arch in, demobilization, 1919.....	202
absolute and relative height of sternal notch in, demobilization, 1919.....	199
absolute and relative leg length of, demobilization, 1919.....	224
absolute and relative shoulder breadth in, demobilization, 1919.....	207
absolute and relative sitting heights in, demobilization, 1919.....	193
absolute and relative transverse diameter of pelvis in, demobilization, 1919.....	219
and Negroes, comparison of correlation between.....	268
index of build for.....	174
mean absolute and relative spans of, demobilization, 1919.....	196
mountain, index of build for.....	173
native, of Scotch origin.....	109, 110, 133, 134, 150, 173
Wisconsin:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289

Wisconsin—Continued.	Page.
average weight of draft recruits at mobilization, 1917-1918, and demobilization, 1919..	122
colored eyes, proportional numbers for, 1865.....	285
comparative view of mean height and mean weight of men from.....	124
comparison of index of build of men from, demobilization in 1865 and 1919.....	170
increase in stature of soldiers at demobilization over recruits, 1917-1919.....	76
index of build—	
demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, demobilization, 1919.....	76
relative chest circumference of recruits from.....	144
World War and Civil War:	
comparative measurements at demobilization.....	242
recruits, comparison of frequency distribution of stature of.....	72, 79
comparative view of mean height and mean weight of troops from different States.....	124
volunteers, age distribution of.....	65
World War veterans, index of build for, for each inch of stature,.....	164, 165
Wyoming:	
absolute and relative numbers of veterans from—	
with blue eyes with brown spots, demobilization, 1919.....	282
with clear blue eyes, demobilization, 1919.....	281
with dark brown eyes, demobilization, 1919.....	283
with dark brown hair, demobilization, 1919.....	291
with flaxen hair, demobilization, 1919.....	288
with light brown eyes, demobilization, 1919.....	283
with light brown hair, demobilization, 1919.....	290
with medium brown hair, demobilization, 1919.....	290
with red hair, demobilization, 1919.....	289
average weight of draft recruits, at mobilization, 1917-1918, and demobilization, 1919..	122
comparative view of mean height and mean weight of men from.....	124
increase in stature of soldiers at demobilization over stature of recruits, 1917-1919.....	76
index of build—	
at demobilization, 1919.....	167
for recruits from, at mobilization, 1917-1918.....	166
mean chest circumference (expiration), of draft recruits from.....	142
mean stature of draft recruits from.....	75
mean stature of soldiers from, at demobilization, 1919.....	76
relative chest circumference of recruits from.....	144
Zones, distribution, Quartermaster Corps, sizes and proportions of men in.....	276

ADDITIONAL COPIES
OF THIS PUBLICATION MAY BE PROCURED FROM
THE SUPERINTENDENT OF DOCUMENTS
GOVERNMENT PRINTING OFFICE
WASHINGTON, D. C.
AT
\$1.75 PER COPY

LANE MEDICAL LIBRARY
STANFORD UNIV MED CTR

SEP 17 1991

STANFORD, CA 94305

UH

224

U58

1921

LANE

STORAGE

